Published in final edited form as:

Arch Sex Behav. 2013 February ; 42(2): 225–236. doi:10.1007/s10508-012-9902-1.

Sexual Orientation Disparities in Sexually Transmitted Infections: Examining the Intersection Between Sexual Identity and Sexual Behavior

Bethany G. Everett

Abstract

The terms MSM (men who have sex with men) and WSW (women who have sex with women) have been used with increasing frequency in the public health literature to examine sexual orientation disparities in sexual health. These categories, however, do not allow researchers to examine potential differences in sexually transmitted infection (STI) risk by sexual orientation identity. Using data from the National Longitudinal Survey of Adolescent Health, this study investigated the relationship between self-reported STIs and both sexual orientation identity and sexual behaviors. Additionally, this study examined the mediating role of victimization and STI risk behaviors on the relationship between sexual orientation and self-reported STIs. STI risk was found to be elevated among heterosexual-WSW and bisexual women, whether they report same-sex partners or not, whereas gay-identified WSW were less likely to report an STI compared to heterosexual women with opposite sex relationships only. Among males, heterosexual-identified MSM did not have a greater likelihood of reporting an STI diagnosis; rather, STI risk was concentrated among gay and bisexual identified men who reported both male and female sexual partners. STI risk behaviors mediated the STI disparities among both males and females, and victimization partially mediated STI disparities among female participants. These results suggest that relying solely on behavior-based categories, such as MSM and WSW, may mischaracterize STI disparities by sexual orientation.

Keywords

Sexual orientation identity; same-sex behavior; STIs; victimization; gender

INTRODUCTION

Several studies have documented elevated rates of both sexually transmitted infections (STIs) and STI risk behaviors and among MSM/MSMW (men who have sex with men/men who have sex with men and women) and WSW/WSMW (women who have sex with women/women who have sex with men and women) populations compared to men and women who engage exclusively in opposite-sex sexual relationships (Bailey, Farquhar, Owen, & Mangtani, 2004; Bell, Ompad, & Sherman, 2006; Ciesielski, 2003; Fethers, Marks, Mindel, & Estcourt, 2000; Stall, Hays, Waldo, Ekstrand, & McFarland, 2000). The terms MSM and WSW, however, may eclipse differences in sexual health behaviors and outcomes that also vary by sexual orientation identity (Muñoz-Laboy, 2004; Young & Meyer, 2005). Recently, several studies have called for researchers to examine sexual identity in conjunction with sexual behavior in order to more fully understand how STI

disparities are distributed across the population by sexual orientation (Malebranche 2008; Meyer & Wilson 2009; Moradi, Mohr, Worthington, & Fassinger, 2009; Young & Meyer 2005). To be sure, sexual behaviors are critical for understanding the spread of STIs, but behaviors do not occur in a social vacuum. Rather, they occur between persons with self-determined sexual orientation identities that may be differentially related to STI risk factors, such as victimization, in addition to STI risk factors related to sexual behaviors. Using nationally representative data, this study systematically investigated how STI risk varied at the intersection of both identity and behavior.

A large body of evidence has established that sexual orientation is multidimensional and cannot be characterized by simple dichotomies (Bauer and Jaram 2008; Diamond, 2008; Laumann, Gagnon, Michael, & Michaels, 1994; Moradi, et al., 2009; Yon-Leau & Muñoz-Laboy, 2010). As a result of this increasingly complex understanding of sexual orientation, the term “sexual minority” has been employed with greater frequency to capture a diverse population of individuals whose sexual identity, behavior, and attraction do not reflect a completely heterosexual orientation. The use of “sexual minority” as a single label for a heterogeneous population, however, has been criticized for being too broad and possibly leading to generalizations about a diverse population that are inaccurate (Nakamura, Semple, Strathdee, & Patterson, 2010; Moradi et al., 2009). To be sure, conflating sexual identity with behavior is problematic: Much research has established that identity and behavior often do not perfectly align (Bauer & Jaram, 2008; Laumann et al., 1994; Nakamura et al., 2011; Reback & Larkins, 2010; Wells, McGee, & Beautrais, 2011; Yon-Leau & Muñoz-Laboy, 2010). For example, using the National Health and Nutritional Survey 2001–2006, Xu, Maya, and Markowitz (2010a) showed that just over 7% of women reported a same-sex sexual relationship, of which 50% reported a heterosexual identity, 32% identified as bisexual, and 18% identify as gay or lesbian. Using the same data, Xu, Sternberg, and Markowitz (2010b) showed roughly 5% of males reported a same-sex relationship, of which 40% identified as heterosexual, 22% identified as bisexual, and 38% identified as gay. Few studies, however, have examined how the confluence of both of these dimensions of sexual orientation may shape STI disparities by sexual orientation.

Among males, elevated STI risk among MSM compared to MSW has been well documented (Ciesielski, 2003; Stall et al., 2000; Stall & Purcell, 2000). Studies have also found differences within the MSM population based upon whether participants report exclusively same-sex partners or both sex partners (Dodge, Jeffries, & Sandfort 2007; Jeffries & Dodge 2007; Nakamura et al., 2011) with results suggesting that MSMW have an elevated risk of reporting an STI compared to MSM (Prahbhu, Owen, Folger, & McFarland, 2004; Zule, Bobashev, Wechsberg, Costenbader, & Coomes, 2009). Few studies, however, have examined how STI risk varies at the intersection of identity and behavior. Some research has suggested that gay-MSM report more sexual partners and a higher prevalence of HIV than heterosexual-MSM (Wolitski, Jones, Wasserman, & Smith, 2006; Xu et al., 2010b). Zellner et al. (2009) examined differences in self-reported STIs and STI risk by behavior and identity among Latinos and found that, compared to heterosexual-MSW, heterosexual-MSM were more likely to report unprotected sex and had been diagnosed with an STI, but there were no significant differences in bisexual-MSMW and gay-MSM’s likelihood of reporting an STI compared to heterosexual-MSW.

Compared to males, the literature on female same-sex behavior is considerably smaller. This is, in part, due to the fact that heteronormative ideas about what “counts” as sex are often limited to penis-to-vaginal or penis-to-anus penetration (Moradi et al., 2009). The existing literature, however, suggests WSW/WSMW have higher rates of STI diagnosis and higher rates of engagement in several STI risk indicators, including intravenous drug use, exchanging sex for money, more lifetime sexual partners, and unprotected opposite-sex

relationships, compared to women who have exclusively male partners (Bailey et al., 2004; Bauer, Jairam, & Baidoonbonso, 2010; Fethers et al., 2000; Rosario, Meyer-Bahlburg, Hunter, & Gwadz, 1999; Reisner et al., 2010; Scheer et al., 2002).

Research on the intersection between sexual identity and behavior is limited and inconclusive for the female population. Some research has suggested that STI risk among WSW/WSMS does not vary by sexual orientation identity (Bell et al., 2006), while other work has found that among WSW/WSMW populations, lesbian identified women may have a lower STI risk than heterosexual-identified women (Koh, Gómez, Shade, & Rowley, 2005; Kral, Lorvick, Bluthenthal, & Watters, 1997; Xu et al., 2010a).

STI risk may vary by both sexual identity and behavior, not only because of differences in sexual behaviors between groups, but also, in part, due because of differences in exposure to victimization. Sexual behavior often takes place in private, but sexual identities are often made public. A sexual minority identity may “flag” gay, lesbian, or bisexual identified persons as targets for victimization. Indeed, several studies have shown that gay, lesbian, and bisexual persons experience higher rates of physical and sexual victimization in both adolescence and in adulthood (Austin, Roberts, Corliss, & Molnar, 2008; Doll et al. 1992; Herek, Gillis, & Cogan, 1999; Jun et al., 2010; Roberts, Austin, Corliss, Vandermorris, & Koenen, 2010; Saewyc et al., 2006), which, in turn, are associated with increased STI risk in young adulthood (Haydon, Hussey, & Halpern, 2011; Raj, Silverman, & Amaro, 2000; Thompson, Potter, Sanderson, & Malbach 1997; Wilson & Widom, 2009). Conversely, it may be more proximate STI risk behaviors, such as the type of sexual activities engaged in, condom use, or the number or sexual partners that drive STI disparities, irrespective of sexual orientation identity. The pathways through which STI risk is shaped warrant further investigation.

In summary, how STIs are distributed across the population at the intersection of both sexual identity and sexual behavior remains unclear, despite evidence that STI risk varies by both indicators. Moreover, the targeted sampling of high-risk WSW and MSW populations may provide a biased picture of STI risk among persons who engage in same-sex sexual behaviors (Meyer & Wilson, 2009; Moradi et al., 2009). This study used data from the National Longitudinal Adolescent Health and investigated (1) disparities in self-reported STIs by both sexual orientation identity and behavior-based markers of sexual orientation and (2) the mediating role of victimization and sexual risk behaviors on STI disparities.

METHOD

Participants

This study used data from Waves III and IV of the National Longitudinal Study of Adolescent Health (Add Health). The initial Add Health sample was drawn from 80 high schools and 52 middle schools, with unequal probability of selection, throughout the United States (Harris et al., 2009). Wave IV of the Add Health survey, collected between 2007 and 2008, located 92.5% of the original sample and interviewed 80.3% of the eligible participants whose ages ranged from 24 to 34 years. The sample used in this study was restricted to participants who reported at least one incident of oral, anal, or vaginal intercourse ever, and at least one male or female sexual partner. The sample excluded individuals who reported that they were “not sexually attracted to either males or females,” participants who responded “don’t know” to the sexual identity question, and participants that did not have full information for all the covariates included in the analysis. These restrictions resulted in a total sample of 7,392 female participants (94.0% of the Wave IV female sample) and 6,323 male participants (91.2% of the Wave IV male sample).

Measures

Sexual Identity-Behavior Subgroups—The measure of sexual identity-behavior subgroups incorporated two survey items. Participants were asked to choose “the description that best fits how you think about yourself: 100% heterosexual (straight), mostly heterosexual (straight), but somewhat attracted to your own sex; bisexual, that is, attracted to men and women equally; mostly homosexual (mostly gay), but somewhat attracted to the opposite sex; or 100% homosexual (gay).” Participants were also asked, “Considering all types of sexual activity, with how many male partners have you ever had sex?” and “Considering all types of sexual activity, with how many female partners have you ever had sex?” From these two questions, five mutually exclusive variables for women¹ and six mutually exclusive variables for men were created: 100% heterosexual and have only engaged in opposite-sex sexual relationship(s) (Heterosexual-WSM/MSW); 100% heterosexual and have ever engaged in same/both-sex sexual relationship(s) (Heterosexual-WSMW/MSMW); mostly heterosexual or bisexual and have only engaged in opposite-sex relationship(s) (Bisexual-WSM/MSW);² and mostly heterosexual or bisexual and have ever engaged in same/both-sex sexual relationship(s) (Bisexual-WSMW/MSMW).³ For women, one variable was created that identified women who reported a gay or mostly gay identity and have engaged in same-sex sexual relationship(s) (Gay-WSMW, N = 133). Among males, 56% of the 100% gay or mostly gay identified participants reported only having had same-sex sexual relationship(s) and 44% reported both same- and opposite-sex sexual relationship(s). Thus, two categories were created: gay-MSM (Gay-MSM, N = 94) and gay-MSMW (Gay-MSMW, N = 75).

Correlation analyses between sexual orientation identities showed that, among women, bisexual (0.43, $p < .001$) and gay (0.32, $p < .001$) identities were positively associated with same-sex behavior and heterosexual identity (-0.53, $p < .001$) was negatively associated with same-sex sex. Male correlation coefficients were similar: bisexual (0.31, $p < .001$) and gay (0.61, $p < .001$) identities were positively associated with same-sex sex and heterosexual (-0.64, $p < .001$) identity was negatively associated with same-sex sex. While these coefficients were significant, they are far from perfect correlations, suggesting that same-sex sex does not necessarily predict a gay or bisexual identity.

Self-reported STIs—STI diagnosis was derived from a question that asked participants if a doctor, nurse, or other health professional had informed them that they have had a sexually transmitted infection.

Mediating Factors—Victimization was measured using a seven item scale that summed the answers to the following questions: (1) By the time you started 6th grade, how often had your parents or other adult caregivers slapped, hit or kicked you; (2) by the time you started 6th grade, how often had one of your parents or other adult care-givers touched you in a sexual way, forced you to touch him or her in a sexual way, or forced you to have sexual relations; (3) excluding parents or caregivers, have you ever been forced in a physical or non-physical way to have any type of sexual activity against your will; (4) in the past 12

¹Only one female participant reported never having had an opposite-sex sexual relationship, therefore, all women who reported a gay or mostly gay identity were categorized as one group. Sensitivity analyses excluding this participant did not significantly alter the results; thus, this participant was included in this category.

²Sample size limitations required that some identity categories be collapsed. Preliminary analyses showed that STI risk among mostly heterosexual participants was statistically different from 100% heterosexual identified participants, but did not statistically differ from bisexual identified participants. Similarly, mostly gay and 100% gay-identified participants did not report statistically difference STI risk odds ratios and were also collapsed to provide more stable coefficient estimates. There were only 11 cases of 100% gay or mostly gay participants reporting opposite-sex only sexual relationships; therefore, these cases were excluded from the analysis.

³For clarity, mostly heterosexual and bisexual participants are referred to as “bisexual.” While in the case of this study these groups were not statistically different, it is important to note that these sexual identity labels are not perfect substitutes in all cases.

months, has someone pulled a knife or gun on you; (5) in the past 12 months, has someone shot or stabbed you; (6) in the past 12 months, has someone slapped, hit, choked, or kicked you; (7) in the past 12 months, were you beaten up?” Because of a high number of missing values for childhood physical and sexual abuse, the summed scale was divided by the number of questions the participant answered; the scale ranged from 0–1 and had an alpha of .71. The scale was recoded into a series of dummy variables that measured whether participant’s score was less than .25 (referent); .25 and .75; or .75.

Total number of partners was derived from the survey items in Wave IV that asked participants: “Considering all types of sexual activity, with how many male partners have you ever had sex?” and “Considering all types of sexual activity, with how many female partners have you ever had sex?” Anal sex was coded as a dichotomous measure derived from a survey question that asked participants “Have you ever had anal intercourse? (By anal intercourse, we mean when a man inserts his penis into his partner’s anus or butt hole).” Participants who reported having had anal intercourse were coded as yes (1) and those who did not were the reference category (0).

Several other variables were tested and excluded from the analysis. Condom use was not included as a risk factor due to the phrasing of the condom-use item on the Add Health Survey. Only participants who reported an opposite-sex sexual relationship in the last 12 months were asked “In the past 12 months, did you or your partner use any of these methods for birth control or disease prevention: condoms (rubbers).” The question did not ask about consistency, and may exclude women who engaged exclusively in same-sex sex or participants who did not have sex in the previous 12 months. Supplementary analysis that included this condom use question revealed that it was not significantly associated with self-reported STIs nor did it mediate the relationship between sexual orientation and STIs⁴. Other variables tested and excluded because they did not mediate the relationship between sexual orientation and STI risk included commercial sex, drug use in the previous 12 months, age at first sex, and number of one-time sexual partners.

Controls—This study included controls for race/ethnicity, age, education, and marital status. Age was coded as a continuous variable and ranges from 24 to 32. Race/ethnicity was coded non-Hispanic white (referent), non-Hispanic black, Hispanic, Asian, or other race. Education was coded as less than high school degree, high school degree, some college or college graduate (referent). Marital status measured whether a participant reported ever being married (yes = 1, no = referent). Because there are known differences in rates of health care utilization and STI testing by sexual orientation (Aaron, et al. 2001; Cochran et al. 2001; Wolitski, et al. 2006), this study also controlled for both of these factors. STD testing measured whether at Wave III of the survey⁵ participants had been tested to at least one sexually transmitted disease in the past 12 months (yes = 1, no = referent). Health care utilization measures whether participants have had a routine checkup in less than one year (referent), one year to less than two years, or 2 years or more.

Procedure

All analyses were conducted separately for male and female participants and used the survey commands in Stata 11.0 to correct for Add Health’s complex sampling frame. Descriptive statistics were examined for self-reported STIs and all covariates used in the analyses by sex, for the total population, and by sexual identity/behavior subpopulations. Adjusted chi square tests were conducted that compared descriptive statistics for each sexual identity-

⁴All supplementary analyses are available upon request from the author.

⁵This question was not asked at Wave IV.

behavior subgroup to the reference group (heterosexual-WSM/MSW). Logistic regression was used to examine the relationship between sexual identity-behavior subgroups and the likelihood of a self-reported STI diagnosis. All results from logistic regression are presented in terms of odds ratios, which are the exponentiated form of the coefficients.

RESULTS

Descriptive Statistics

Table 1 shows the descriptive statistics for self-reported STIs and all other covariates used in the analysis by identity-behavior subgroups for the female population. It can be seen that 75.9% of females reported a heterosexual identity and opposite-sex only sexual relationships (heterosexual-WSM) and 4.2% reported a heterosexual identity and at least one same-sex sexual relationship (heterosexual-WSMW). A total of 18.2% of females reported either a mostly heterosexual or bisexual identity, of which 50.5% reported at least one same-sex sexual relationship. Just 1.8% of females identified as gay or mostly gay.

Self-reported STI rates were significantly higher among heterosexual-WSMW (58.1%), bisexual-WSM (51.1%), and bisexual-WSMW (64.1%) compared to heterosexual-WSM (46.6%). Gay-WSMW had the lowest rates of self-reported STI (32.0%); however, this rate did not significantly differ from heterosexual-WSM. STI risk factors were also unevenly distributed by sexual orientation among females. Seventy-five percent of heterosexual-WSM women reported little or no victimization compared to 60.3% of heterosexual-WSMW, 70.3% bisexual-WSM, 65.4% gay-WSMW, and 55.8% of bisexual-WSMW. Heterosexual-WSM also reported a significantly smaller mean number of sexual partners than all other identity groups and the second lowest prevalence of anal sex after gay-WSMW.

Table 2 shows the descriptive statistics for male participants. Among males, 90.8% of the sample reported a heterosexual identity and opposite-sex only sexual relationships (heterosexual-MSW) and 2.4% reported a heterosexual identity and at least one same-sex sexual relationship (heterosexual-MSMW). A total of 4.1% of male participants identified as mostly heterosexual or bisexual, of which 45% reported at least one same-sex sexual relationship. Three percent of male participants reported a gay or mostly gay identity, of which 55.6% reported same-sex sexual relationship(s) only.

Heterosexual-MSMW (41.4%), Bisexual-MSMW (43.7%), gay-MSM (39.4%), and gay-MSMW (48.9%) reported significantly higher rates of self-reported STIs compared to heterosexual-MSW (32.0%). Fewer differences in STI risk factors were detected by sexual orientation among males compared to females. Heterosexual-MSW reported lower rates of victimization only compared to heterosexual-MSMW. However, several identity-behavior groups reported higher mean levels of sexual partners including heterosexual-MSMW (20.2), bisexual-MSMW (26.9), gay-MSM (37.5.6), and gay-MSMW (29.5) compared to heterosexual-MSW (17.3). All bisexual and gay identified males, including bisexual males who did not report same-sex sex, had a higher prevalence of reported anal sex compared to heterosexual-WSM.

STI Disparities by Sexual Orientation Among Females

The results from logistic regressions that regressed identity-behavior subgroups on self-reported STI diagnosis for females are shown in Table 3. Model 1 controlled for sociodemographic characteristics only and revealed heterosexual-WSMW ($OR = 1.75, p < .01$), bisexual-WSM ($OR = 1.47, p < .001$), and bisexual-WSMW ($OR = 2.20, p < .001$) were all more likely to report an STI diagnosis than heterosexual-WSM. Gay women were significantly less likely to report an STI than heterosexual-WSM ($OR = 0.47, p < .01$). Model 2 included the victimization scale; victimization was associated with elevated STI

risk and mediated the relationship between sexual orientation STI risk by roughly 6% for heterosexual-WSMW, 10% for bisexual-WSMW, and 4% for gay-WSMW.

Model 3 added controls for total number of sexual partners and whether the participant had had anal intercourse to the sociodemographic controls included in Model 1. These measures fully mediated the relationship between heterosexual-WSMW, bisexual-WSM, and reduced both the magnitude and significance for bisexual-WSMW STI risk ($OR=1.99, p < .05$) compared to heterosexual-WSM. Further, in Model 3, the disparity between gay-WMSW and heterosexual-WSM became larger, such that gay-WMSW ($OR = 0.34, p < .01$) were significantly less likely to report an STI compared to heterosexual-WSM. Consistent with Model 3, controlling for both victimization and sexual behaviors in Model 4 fully mediated the relationship between heterosexual-WSMW, bisexual-WSM, and bisexual-WSMW, and showed that gay-WSMW ($OR = 0.34, p < .01$) were significantly less likely to report an STI compared to heterosexual-WSM.

STI Disparities by Sexual Orientation Among Males

Estimates for STI risk among males are shown in Table 4. Model 1, which only controlled for sociodemographic characteristics, showed there was no significant difference in heterosexual-MSMW's, bisexual-MSW's, and gay-MSM's STI risk compared to heterosexual-MSW participants. Bisexual-MSMW ($OR = 1.81, p < .05$) and gay-MSMW ($OR = 2.20, p < .05$) had elevated risk of reporting an STI diagnosis compared to heterosexual-MSW. Model 2 controlled for victimization, which was independently associated with an elevated STI risk, but had almost no mediating effect on the relationship between bisexual-MSMW and self-reported STIs. Moreover, in Model 2, the odds ratio for gay-MSMW actually increased, such that gay-MSMW were almost 2.3 times as likely to have reported an STI compared to heterosexual-MSW.

Model 3 added controls for sexual behaviors. Total number of partners and anal intercourse were both positively associated with STI diagnosis and fully mediated the relationship between both bisexual-MSMW and gay-MSMW's elevated self-reported STI risk compared to heterosexual-MSMW. Model 4 controlled for all factors and showed that only gay-MSMW ($OR = 1.91, p < .05$) were more likely to report an STI than heterosexual-MSW.

DISCUSSION

Researchers have repeatedly demonstrated that sexual orientation identity and sexual behaviors do not always directly coincide with one another (Bauer & Jaram, 2009; Diamond, 2008; Laumann, et al., 1994; Moradi, Mohr et al., 2009; Yon-Leau & Muñoz-Laboy, 2010). While sexual health scholars frequently acknowledge this discrepancy, few studies have explicitly examined how STI varies at the intersection of these two indicators. This research provided new insights into the social patterning of STIs and the risk factors that mediate STI disparities in several important ways.

First, the results presented in this study showed that behavior-based measures of sexual orientation that exclude sexual identity may lead to misinformed conclusions about STI disparities. For example, compared to heterosexual identified women with opposite sex only relationships, STI risks were elevated among heterosexual and bisexual women who reported same-sex sex, but lower among gay identified women with histories of same-sex sex. Behavior-based categories also mischaracterized risk among males: No significant difference was detected in the likelihood of self-reporting an STI between heterosexual-MSM and heterosexual-MSMW, but elevated risk was detected among gay-MSMW and bisexual-MSMW. Second, this study differentiated bisexual and mostly heterosexual-identified persons who reported same-sex sex from those that reported only opposite-sex

sexual relationships. Almost 10% of the female population and 2% of the male population reported a bisexual or mostly heterosexual identity but reported never having had a same-sex sexual relationship. Previous work has shown that both-sex oriented persons are more likely to report an STI than heterosexual-identified persons; the results presented here showed that this risk varied by sexual histories. Finally, the results presented in this study demonstrated that both victimization and sexual behaviors mediated STI disparities among females, but only sexual behaviors mediated STI disparities among males.

Females

Among women, the results showed that gay-WSMW were less likely to report an STI than heterosexual-WSM. Thus, behavior-based categories of sexual orientation, which label WSWs as an “at risk” population, without considering sexual identity, mischaracterize STI risk among females. This finding was supported by Xu et al.’s (2010a) as well as Kral et al.’s (1997) research that found a lesbian identity was protective against STI risk, and contradicted Bell, et al.’s (2006) results, which suggested that STI risks did not differentiate by identity. The Bell et al. sample, however, was collected from a group of women who reported using heroin, crack, or cocaine in the previous five years, limiting the generalizability of the results to the gay female population in the United States. Some research work has suggested that gay or lesbian-identified women are less likely to be tested for STIs and less likely to get regular pap smears (Aaron et al., 2001; Cochran et al., 2001; Marrazzo, Koutsky, Kiviat, Kuypers, & Stine, 2001), which may, in part, explain the lower risk among this group. Supplementary analyses examining STI disparities with biomarker data on chlamydia and gonorrhea in Wave III of the survey, however, showed similar results to those presented in this study: gay women were not more likely to report an STI than heterosexual-WSM women.

Elevated risk of reporting an STI among WSW populations was instead concentrated among heterosexual-WSMW and bisexual-WSMW. These results were in line with other research that has shown that heterosexual-WSMW have an elevated STI risk compared to heterosexual-WSM (Bauer et al., 2010). Moreover, the findings presented in this study also showed that bisexual-WSM were also more likely to report an STI than heterosexual-WSM; behavior-based markers of sexual orientation would miss the elevated risk associated with a bisexual or mostly straight identity among women who have not engaged in same-sex sex. Elevated STI risk among both-sex oriented women found in this study corroborated other research that suggests that both-sex oriented women are at an increased STI risk compared to heterosexual-identified women (Austin et al., 2008; Fethers et al., 2000; Rosario et al., 1999). Both-sex oriented women have elevated levels of victimization, fewer political or interpersonal resources, as well as decreased access to social support systems and accurate sexual health information (Allen, Glick, Beach, & Naylor, 1998; Balsam & Mohr, 2007; Blake et al., 2001; Corliss, Austin, Roberts, & Molnar, 2009; Corliss, Shankle, & Moyer, 2007; Hutchins, 1996), which may, in part, explain the elevated risk of STI among both-sex oriented women (Mazzaferro et al., 2006). Victimization partially mediated the relationship between bisexual identified women and STI risk in this study. While supplementary analysis not shown in this study did not show that drug use mediated the relationship between sexual orientation and STI risk, other work has suggested that drug use among both-sex oriented women may increase their risk of contracting an STI (Bauer et al., 2010; Koh et al., 2005).

Differences in the number of sexual partners and prevalence of anal intercourse also mediated disparities in self-reported STIs by sexual orientation among women: controlling for these factors in Model 3 fully explained elevated STI risk among heterosexual-WSM and bisexual-WSM women, and reduced STI risk among bisexual-WSMW. Heterosexual-WSMW’s STI risk may not, in fact, be linked to behaviors engaged with women, but may be due to more frequent opposite-sex sexual relationships (Bauer et al., 2010). Supplementary

analyses not shown revealed that heterosexual-WSMW reported an average of 14.2 male partners, bisexual-WSM reported an average of 11.6 male partners, and bisexual-WSMW reported an average of 24.2 male partners compared to heterosexual-WSM, who reported an average of 7.7 male partners. Future research should continue to explore increased STI risk associated with opposite-sex sexual relationships among both-sex oriented women.

Males

Among males, much research has identified elevated risk of STI among MSM populations (Groß, Parsons, & Bimbi, 2007; Halkitis, Parsons, & Storratt, 2001; Stall & Purcell, 2000; Wolitski et al., 2006). Heterosexual-identified MSM or men on the “down-low” have often been characterized as sexually deviant and vectors of disease in heterosexual relationships (Ford, Whetten, Hall, Kaufman, & Thrasher, 2007; Saleh & Operario, 2009). The results reported in this study, however, showed that heterosexual-MSMW were not more likely to report an STI than heterosexual-MSW. Other research has also found that while the prevalence of specific risk factors may vary by identity, MSM who do not identify as gay or bisexual may not have an elevated STI risk compared to bisexual or gay identified-MSM (Bond et al., 2009; Millet, Malebranche, Mason, & Spikes, 2005; Pathela et al., 2006;). The lack of elevated STI risk among heterosexual-MSM in this study may, in part, be due to the fact that this research used a nationally representative sample of U.S. young adults as opposed to a sample drawn from high-risk populations. Alternatively, differences in STI testing may also contribute to the lack of differences in self-reported STI diagnoses by a medical professional among heterosexual-MSM as some research has suggested that MSM who have not disclosed a gay or bisexual identity are less likely to be tested for STIs (Wolitski et al., 2006).

The results showed that STI risk was concentrated among males who identified as mostly heterosexual, bisexual, or gay *and* had a history of *both* sex partners, but that the relationship between bisexual-MSMW was mediated by sexual risk behaviors. Indeed, in Model 4, the only sexual identity-behavior subgroup that was associated with elevated STI risk was gay-MSMW participants. There was no statistically significant difference in heterosexual-MSW and gay-MSM’s odds of reporting an STI. The absence of opposite-sex sexual partners among gay males may indicate early commitment to a gay identity and involvement with the gay community, which may decrease STI risk behaviors (Ramirez-Valles, 2002). While victimization and drug use (not shown) did not mediate the relationship between sexual orientation and STI disparities among males, other work has suggested that fear of violence and/or the use of drugs and alcohol contribute to an elevated risk of contracting an STI among behaviorally-bisexual males (Malebranche et al., 2010). Differences in sexual networks and locations where same-sex sex partners are sought out may also explain differences in STI risk between gay men with both-sex partners and gay men with exclusively same-sex partners (Nakamura et al., 2010). Future research should further investigate how STI risk varies among gay males by sexual histories.

Sex Differences

While both bisexual identified men and women have elevated STI risk in Model 1, several gender differences emerged in the patterns of STI disparities by sexual orientation. First, among women, a gay identity was associated with decreased risk while among men, a gay identity among behaviorally bisexual males was associated with increased STI risk. This difference may be explained by differences in STI risk associated with specific sexual behaviors. While it is indeed possible to transmit STIs through female-to-female sexual relationships (Marrazzo et al., 2001), STIs are transmitted more easily through penile-vaginal intercourse or penile-anal intercourse. Second, differences in STI risk may be associated with STI prevalence in sexual networks by sex. If STI rates are lower among

lesbian-WSMW, the chances of exposure to an STI from a sexual partner from this population is also lower. Third, these differences may also be due to differences in STI testing between gay men and women. Public health campaigns have heavily targeted gay men and MSM for STI testing and sexual health information, while gay women and WSW's have received considerably less attention regarding the spread of STI risk. Moreover, for many women, STI screenings are completed during gynecological visits required for accessing birth control. If women are not engaging in opposite-sex sex, utilizing health services for acquiring birth control may be unnecessary, decreasing the likelihood that they will interact with women's health professionals.

Conclusion and Future Directions

This study had several limitations that may be addressed in future research. First, there are several behavioral risk factors that may help explain the elevated STI risk among some sexual identity-behavior risk groups that are not included in the Add Health survey, such as the extent to which condoms are used in every sexual encounter or the sex role (i.e., receptive versus insertive anal intercourse) of the participant engaging in specific sexual behaviors. Second, this study was limited in the types of identities presented for participants to choose from. Indeed, some participants may find sexual orientation labels off-putting or irrelevant to their sexuality (Savin-Williams, 2006; Yon-Leau & Muñoz-Laboy 2010). Participants who do not identify with one of the offered identity categories were either excluded from this analysis or potentially included in an identity category that was not truly representative of their chosen identity. Third, it is difficult to assess the temporal ordering of risk factors and STI diagnosis, thus not allowing for a full meditational analysis. Supplementary analyses not shown restricted STI diagnosis to the last 12 months, however, revealed a similar pattern to the results presented. Finally, this research employed self-reported doctor diagnosis of an STI. Thus, the results may be biased towards showing greater risk among populations that were more likely to get tested. Some research, however, has suggested that self-reported doctor diagnoses are preferable to self-assessed STI risk and may not introduce a substantial amount of bias (Niccolai et al., 2005). While this study included controls for STI testing reported at Wave III of the survey, this survey item was not included in Wave IV of the Add Health survey and therefore may remain a source of bias.

This research provides evidence that it is critical to consider sexual behaviors in conjunction with sexual orientation identities when studying STI risk across the U.S. population. Indeed, these findings add to the growing body of research that suggests that behavior and identity are not interchangeable categories and that focusing on one indicator of sexual orientation alone may lead to misinformed conclusion about how STI risk is distributed across the population. More research is needed to understand sexual health needs of bisexual identified and behaviorally bisexual persons. Indeed, STI prevention efforts aimed at reducing victimization among women and addressing safe-sex practices among both men and women may benefit from a more tailored focus of the needs of these risk groups rather than grouping all identity groups who engage in same-sex sex together.

Acknowledgments

This research has benefited from the NICHD-funded University of Colorado Population Center (grant R21 HD51146) through administrative and computing support. I gratefully acknowledge support from the Eunice Shriver National Institute of Child Health and Human Development (NICHD) grant R03 HD062597 and the American Psychological Foundation (APF) Wayne F. Placek award. The content is solely the responsibility of the author and does not necessarily represent the official views of NIH, NICHD, or the APF. And I thank Dr. S. Bryn Austin, Dr. Stefanie Mollborn, Dr. Stephen Russell, and three anonymous reviewers for their insightful comments and suggestions on an earlier draft of this article.

REFERENCES

- Aaron D, Markovic N, Danielson M, Honnold J, Janosky J, Schmidt N. Behavioral risk factors for disease and preventive health practices among lesbians. *American Journal of Public Health*. 2001; 91:972–975. [PubMed: 11392943]
- Allen LB, Glick AD, Beach RK, Naylor KE. Adolescent health care experience of gay, lesbian, and bisexual young adults. *Journal of Adolescent Health*. 1998; 23:212–220. [PubMed: 9763157]
- Austin SB, Roberts AL, Corliss HL, Molnar BE. Sexual violence victimization history and sexual risk indicators in a community-based urban cohort of "mostly heterosexual" and heterosexual young women. *American Journal Public Health*. 2008; 98:1015–1020.
- Bailey JV, Farquhar C, Owen C, Mangtani P. Sexually transmitted infections in women who have sex with women. *Sexually Transmitted Infections*. 2004; 80:244–246. [PubMed: 15170014]
- Balsam KF, Mohr JJ. Adaptation to sexual orientation stigma: A comparison of bisexual and lesbian/gay adults. *Journal of Counseling Psychology*. 2007; 54:306–319.
- Bauer GR, Jairam JA. Are lesbians really women who have sex with women (WSW)? Methodological concerns in measuring sexual orientation in health research. *Women and Health*. 2008; 48:383–408.
- Bauer GR, Jairam JA, Baidooobonso SM. Sexual health, risk behaviors, and substance use in heterosexual-identified women with female sex partners: 2002 US National Survey of Family Growth. *Sexually Transmitted Diseases*. 2010; 37:531–537. [PubMed: 20502395]
- Bearman PS, Jones J, Udry JR. The National Longitudinal Study of Adolescent Health: Research design. (n.d.). Retrieved from <http://www.cpc.unc.edu/projects/addhealth/design.html>.
- Bell AV, Ompad D, Sherman SG. Sexual and drug risk behaviors among women who have sex with women. *American Journal of Public Health*. 2006; 96:1066–1072. [PubMed: 16670234]
- Blake S, Ledsky R, Lehman T, Goodenow C, Sawyer R, Hack T. Preventing sexual risk behaviors among gay, lesbian, and bisexual adolescents: the benefits of gay-sensitive HIV instruction in schools. *American Journal of Public Health*. 2001; 91:940–946. [PubMed: 11392938]
- Bond L, Wheeler DP, Millett GA, LaPollo AB, Carson LF, Liao A. Black men who have sex with men and the association of down-low identity with HIV risk behavior. *American Journal of Public Health*. 2009; 99:S92–S95. [PubMed: 19218177]
- Center for Disease Control and Prevention. HIV/STD risks in young men who have sex with men who do not disclose their sexual orientation--six US cities: 1994–2000. *Morbidity and Mortality Weekly Report*. 2003; 52:145–152.
- Ciesielski CA. Sexually transmitted diseases in men who have sex with men: An epidemiologic review. *Current Infectious Disease Reports*. 2003; 5:145–152. [PubMed: 12642001]
- Cochran S, Mays V, Bowen D, Gage S, Bybee D, Roberts S, et al. Cancer-related risk indicators and preventive screening behaviors among lesbians and bisexual women. *American Journal of Public Health*. 2001; 91:591–597. [PubMed: 11291371]
- Corliss HL, Austin SB, Roberts AL, Molnar BE. Sexual risk in "mostly heterosexual" young women: Influence of social support and caregiver mental health. *Journal of Women's Health*. 2009; 18:2005–2010.
- Corliss HL, Shankle MD, Moyer MB. Research, curricula, and resources related to lesbian gay, bisexual, and transgender health in US schools of public health. *American Journal of Public Health*. 2007; 97:1023–1027. [PubMed: 17463373]
- Diamond L. Female bisexuality from adolescence to adulthood: Results from a 10-year longitudinal study. *Development Psychology*. 2008; 44:5–14.
- Dodge B, Jeffries WL, Sandfort TGM. Beyond the down low: Sexual risk, protection, and disclosure among at-risk black men who have sex with both men and women (WSMW). *Archives of Sexual Behavior*. 2008; 37:683–696. [PubMed: 18512140]
- Doll LS, Joy D, Bartholow BN, Harrison JS, Bolan G, Douglas JM, et al. Self-reported childhood and adolescent sexual abuse among adult homosexual and bisexual men. *Child Abuse & Neglect*. 1992; 16:855–864. [PubMed: 1486514]
- Fethers K, Marks C, Mindel A, Estcourt CS. Sexually transmitted infections and risk behaviours in women who have sex with women. *Sexually Transmitted Infections*. 2000; 76:345–349. [PubMed: 11141849]

- Ford CL, Whetten KD, Hall SA, Kaufman JS, Thrasher AD. Black sexuality, social construction, and research targeting "The Down Low" ("The DL"). *Annals of Epidemiology*. 2007; 17:209–216. [PubMed: 17320787]
- Grov C, Parsons JT, Bimbi DS. Sexual risk behavior and venues for meeting sex partners: An intercept survey of gay and bisexual men in LA and NYC. *AIDS and Behavior*. 2007; 11:915–926. [PubMed: 17206536]
- Halkitis PN, Parsons JT, Stirratt MJ. A double epidemic: Crystal methamphetamine drug use in relation to HIV transmission among gay men. *Journal of Homosexuality*. 2001; 41(2):17–35. [PubMed: 11482426]
- Harris, KM.; Halpern, CT.; Whitsel, E.; Hussey, J.; Tabor, P.; Entzel, P.; Udry, JR. The National Longitudinal Study of Adolescent Health: Research design. 2009. URL: <http://www.cpc.unc.edu/projects/addhealth/design>.
- Haydon AA, Hussey JM, Halpern CT. Childhood abuse and neglect and the risk of STDs in early adulthood. *Perspectives on Sexual and Reproductive Health*. 2011; 43:16–22. [PubMed: 21388501]
- Herek GM, Gillis JR, Cogan JC. Psychological sequelae of hate-crime victimization among lesbian gay, bisexual adults. *Journal of Consulting and Clinical Psychology*. 1999; 67:945–951. [PubMed: 10596515]
- Hutchins, L. Bisexuality: Politics and community. In: Firestein, BA., editor. *Bisexuality: The psychology and politics of an invisible minority*. Thousand Oaks, CA: Sage; 1996. p. 240-259.
- Jeffries WL, Dodge B. Male bisexuality and condom use at last sexual encounter: Results from a national survey. *Journal of Sex Research*. 2007; 44:278–289. [PubMed: 17879171]
- Jun H, Austin SB, Wylie SA, Corliss HL, Jackson B, Spiegelman D, et al. The mediating effect of childhood abuse in sexual orientation disparities in tobacco and alcohol use during adolescence: Results from the Nurses' Health Study II. *Cancer Causes & Control*. 2010; 21:1817–1828. [PubMed: 20640883]
- Koh AS, Gómez CA, Shade S, Rowley E. Sexual risk factors among self-identified lesbians, bisexual women, and heterosexual women accessing primary care settings. *Sexually Transmitted Diseases*. 2005; 32:563–569. [PubMed: 16118605]
- Kral AH, Lorvick J, Bluthenthal RN, Watters JK. HIV risk profile of drug-using women who have sex with women in 19 United States cities. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology*. 1997; 16:211–217. [PubMed: 9390574]
- Laumann, EO.; Gagnon, JH.; Michael, RT.; Michaels, S. The social organization of sexuality: sexual practices in the United States. Chicago, IL: University of Chicago Press; 2000.
- Malebranche DJ. Bisexually active black men in the United States and HIV: Acknowledging more than the "Down Low". *Archives of Sexual Behavior*. 2008; 37:810–816. [PubMed: 18506612]
- Malebranche DJ, Arriola KJ, Jenkins TR, Dauria E, Patel SN. Exploring the "bisexual bridge": A qualitative study of risk behavior and disclosure of same-sex behavior among black bisexual men. *American Journal of Public Health*. 2010; 100:159–161. [PubMed: 19910348]
- Marrazzo J, Koutsky L, Kiviat N, Kuypers J, Stine K. Papanicolaou test screening and prevalence of genital human papillomavirus among women who have sex with women. *American Journal of Public Health*. 2001; 91:947–952. [PubMed: 11392939]
- Mazzaferro KE, Murray PJ, Ness RB, Bass DC, Tyus N, Cook RL. Depression, stress, and social support as predictors of high-risk sexual behaviors and STIs in young women. *Journal of Adolescent Health*. 2006; 39:601–603. [PubMed: 16982400]
- Meyer IH, Wilson PA. Sampling lesbian gay, bisexual populations. *Journal of Counseling Psychology*. 2009; 56:23–31.
- Millet G, Malebranche D, Mason B, Spikes P. Focusing "down low": Bisexual black men, HIV risk and heterosexual transmission. *Journal of the National Medical Association*. 2005; 97:708–719.
- Moradi B, Mohr JJ, Worthington RL, Fassinger RE. Counseling psychology research on sexual (orientation) minority issues: Conceptual and methodological challenges and opportunities. *Journal of Counseling Psychology*. 2009; 56:5–22.
- Muñoz-Laboy MA. Beyond "MSM": Sexual desire among bisexually-active Latino men in New York City. *Sexualities*. 2004; 7:55–80.

- Nakamura N, Semple SJ, Strathdee SA, Patterson TL. HIV risk profiles among HIV-positive, methamphetamine-using men who have sex with both men and women. *Archives of Sexual Behavior*. 2011; 40:793–801. [PubMed: 21203813]
- Niccolai LM, Kershaw TS, Lewis JB, Cicchetti DV, Ethier KA, Ickovics JR. Data collection for sexually transmitted disease diagnoses: A comparison of self-report, medical record reviews, and state health department reports. *Annals of Epidemiology*. 2005; 15:236–242. [PubMed: 15723771]
- Pathela P, Hajat A, Schillinger J, Blank S, Sell R, Mostashari F. Discordance between sexual behavior and self-reported sexual identity: A population-based survey of New York City men. *Annals of Internal Medicine*. 2006; 145:416–425. [PubMed: 16983129]
- Prabhu R, Owen CL, Folger K, McFarland W. The bisexual bridge revisited: Sexual risk behavior among men who have sex with men and women, San Francisco: 1998–2003. *Aids*. 2004; 18:1604–1606. [PubMed: 15238783]
- Raj A, Silverman JG, Amaro H. The relationship between sexual abuse and sexual risk among high school students: Findings from the 1997 Massachusetts Youth Risk Behavior Survey. *Maternal and Child Health Journal*. 2000; 4:125–134. [PubMed: 10994581]
- Ramirez-Valles J. The protective effects of community involvement for HIV risk behavior: a conceptual framework. *Health Education Research*. 2002; 17:389–403. [PubMed: 12197585]
- Reback CJ, Larkins S. Maintaining a heterosexual identity: Sexual meanings among a sample of heterosexually identified men who have sex with men. *Archives of Sexual Behavior*. 2008; 39:766–773. [PubMed: 19030977]
- Reisner S, Mimiaga M, Case P, Grasso C, O'Brien CT, Harigopal P, et al. Sexually transmitted disease (STD) diagnoses and mental health disparities among women who have sex with women screen at an urban community health center, Boston, Massachusetts, 2007. *Sexually Transmitted Diseases*. 2010; 37:5. [PubMed: 20118673]
- Roberts AL, Austin SB, Corliss HL, Vander Morris AK, Koenen KC. Pervasive trauma exposure among U.S. sexual orientation minority adults and risk of posttraumatic stress disorder. *American Journal Public Health*. 2010; 100:2433–2441.
- Rosario M, Meyer-Bahlburg HF, Hunter J, Gwadz M. Sexual risk behaviors of gay, lesbian, and bisexual youths in New York City: Prevalence and correlates. *AIDS Education and Prevention*. 1999; 11:476–496. [PubMed: 10693645]
- Saewyc EM, Skay CL, Pettingell SL, Reis EA, Bearinger L, Resnick M, et al. Hazards of stigma: The sexual and physical abuse of gay, lesbian, and bisexual adolescents in the United States and Canada. *Child Welfare*. 2006; 85:195–213. [PubMed: 16846112]
- Saleh LD, Operario D. Moving beyond "the Down Low": A critical analysis of terminology guiding HIV prevention efforts for African American men who have secretive sex with men. *Social Science & Medicine*. 2009; 68:390–395. [PubMed: 19006674]
- Savin-Williams. *The new gay teenager*. Cambridge, MA: Harvard University Press; 2006.
- Scheer S, Peterson I, Page-Shafer K, Delgado V, Gleghorn A, Ruiz J, Molitor F, McFarland W, Klausner J. the Yong Women's Survey Team. Sexual and drug use behavior among women who have sex with both women and men: Results of a population-based survey. *American Journal of Public Health*. 2002; 92:1110–1112. [PubMed: 12084692]
- Stall RD, Hays RB, Waldo CR, Ekstrand M, McFarland W. The Gay '90s: A review of research in the 1990s on sexual behavior and HIV risk among men who have sex with men. *AIDS*. 2000; 14:S101–S114. [PubMed: 11086853]
- Stall R, Purcell DW. Intertwining epidemics: A review of research on substance use among men who have sex with men and its connection to the AIDS epidemic. *AIDS and Behavior*. 2000; 4:181–192.
- Thompson NJ, Potter JS, Sanderson CA, Malbach EW. The relationship of sexual abuse and HIV risk behaviors among heterosexual adult female STD patients. *Child Abuse and Neglect*. 1997; 21:149–156. [PubMed: 9056094]
- Wells JE, McGee MA, Beautrais AL. Multiple aspects of sexual orientation: Prevalence and sociodemographic correlates in a New Zealand National Survey. *Archives of Sexual Behavior*. 2011; 40:155–168. [PubMed: 20567892]

- Wilson HW, Widom CS. Sexually transmitted diseases among adults who had been abused or neglected as children: A 30-year prospective study. *American Journal of Public Health*. 2009; 99:S197–S203. [PubMed: 19218173]
- Wolitski RJ, Jones KT, Wasserman JL, Smith JC. Self-Identification as “Down Low” among men who have sex with men (MSM) from 12, US Cities. *AIDS and Behavior*. 2006; 10:519–529. [PubMed: 16691462]
- Yon-Leau C, Muñoz-Laboy M. "I don't like to say that I'm anything": Sexuality politics and cultural critique among sexual-minority Latino youth. *Sexuality Research and Social Policy*. 2010; 7:105–117.
- Xu F, Maya SR, Markowitz LE. Women who have sex with women in the United States: Prevalence, sexual behavior and prevalence of herpes simplex virus type 2 infection-Results from National Health and Nutrition Examination Survey 2001–2006. *Sexually Transmitted Diseases*. 2010a; 37:407–413. [PubMed: 20531032]
- Xu F, Sternberg MR, Markowitz LE. Men who have sex with men in the United States: Demographic and behavioral characteristics and prevalence of HIV and HSV-2 infection. *Sexually Transmitted Diseases*. 2010b; 37:399–405. [PubMed: 20473245]
- Yon-Leau C, Muñoz-Laboy. "I don't like to say that I'm anything": Sexual politics and cultural critique among sexual-minority Latino youth. *Sexuality Research and Social Policy*. 2010; 7:105–117.
- Young RM, Meyer IH. The trouble with "MSM" and "WSW": Erasure of the sexual-minority person in public health discourse. *American Journal of Public Health*. 2005; 95:1144–1149. [PubMed: 15961753]
- Zellner JA, Martínez-Donate AP, Sañudo F, Fernández-Cerdeño A, Sipan CL, Hovell ML, Carrillo H. The interaction of sexual identity with sexual behavior and its influence on HIV risk among Latino men: Results of a community survey in Northern San Diego County, California. *American Journal of Public Health*. 2009; 99:125–132. [PubMed: 19008512]
- Zule WA, Bobashev GV, Wechsberg WM, Costenbader EC, Coomes CM. Behaviorally bisexual men and their risk behaviors with men and women. *Journal of Urban Health*. 2009; 86:48–62. [PubMed: 19513854]

Table 1

Descriptive Statistics of covariates by sex and sexual orientation, U.S. females aged 24–32

	Total Sample	100% Heterosexual		Mostly Straight/Bisexual		Mostly Gay/ 100% Gay	
		Op. Sex Only	Both Sex Sex	Op. Sex Only	Both Sex Sex	Op. Sex Only	Both Sex Sex
Self-reported STI	N=7,392	N=5,607	N=307	N=716	N=629	N=133	
	46.63	43.62	58.09 ***	51.11 **	64.19 ***	32.00	
Demographic							
Race/Ethnicity (%)							
Non-Hispanic white	68.35	67.17	65.61	74.92 *	72.54	67.47	
Non-Hispanic black	15.32	16.72	13.92	8.49 ***	10.99	18.89	
Hispanic	11.46	11.55	13.27	10.63	10.81	11.24	
Asian	3.10	3.15	4.93	4.10	1.18 **	0.77 *	
Other race/ethnicity	1.77	1.41	2.27	1.86	4.48 *	1.63	
Age(μ)	28.71	28.79	28.65	28.45	28.35	28.52	
Education (%)							
College graduate	33.97	35.82	25.54 **	38.23	20.05 ***	24.53 ***	
Some college	35.46	34.70	35.69	33.45	43.04 **	39.17	
Vocational training	9.25	9.03	7.63	10.42	10.86	7.62	
High school	13.89	13.72	19.59	11.44	13.83	20.84	
Less than high school	7.43	6.73	11.55	6.46	12.22 **	7.84	
Married, ever (%)	56.77	59.39	57.11	54.41 *	45.30 ***	15.42 ***	
Tested for STI, WII (%)	28.52	26.83	35.10 *	33.23 **	34.60 **	27.12	
Last regular medical visit (%)							
< 1 year	70.85	72.80	70.31	64.83 **	64.55 **	53.93 **	
1 year & < 2 years	13.07	12.36	15.28	18.71 **	10.22	22.16	
2 years	16.08	14.84	14.41	16.46	25.23 ***	23.91	
Victimization (%)							

	Total Sample	100% Heterosexual		Mostly Straight/Bisexual		Mostly Gay/ 100% Gay	
		Op. Sex Only	Both Sex Sex	Op. Sex Only	Both Sex Sex	Op. Sex Only	Both Sex Sex
<.25	72.33	75.43	60.34 ***	70.34 **	55.78 ***	65.39 *	
.25 and < .75	15.60	13.01	23.03 **	18.01 **	30.25 ***	19.33	
>.75	12.07	11.56	16.63 *	11.65	13.97	15.28 *	
Sexual Behaviors							
Total No. of sexual partners	10.41	7.70	15.84 ***	11.63 ***	27.77 ***	16.50 ***	
Anal sex	44.29	37.53	62.82 ***	63.64 ***	73.00 ***	33.20	

Source: National Longitudinal Survey of Adolescent Health

* *p* .05;
** *p* .01;
*** *p* .001

Descriptive Statistics of covariates by sex and sexual orientation, U.S. males aged 24–32

	Total Sample	100% Heterosexual		Mostly Straight/Bisexual		Mostly Gay/100% Gay	Mostly Gay/100% % Gay
		Op. Sex Only	Both Sex Sex	Op. Sex Only	Both Sex Sex	Same Sex Only	Both Sex Sex
Self-reported STI	N=6,323	N=5,744	N=151	N=142	N=117	N=94	N=75
	32.67	31.96	41.36 **	32.04	43.66 **	39.41 **	48.85 **
Demographic							
Race/Ethnicity (%)							
Non-Hispanic white	68.69	68.92	54.28 ***	79.56 *	73.48 *	55.16 *	67.05
Non-Hispanic black	14.47	14.59	21.17 ***	4.98 ***	12.89 **	14.52	12.49
Hispanic	11.80	11.44	19.65	10.06	11.75	22.69 *	14.39
Asian	3.08	3.12	1.66	3.11	1.88	4.33	3.86
Other race/ethnicity	1.96	1.93	3.24	2.29	0.00	3.30	2.21
Age (μ)	28.93	28.93	29.18	28.85	28.79	29.11	28.75
Education (%)							
College graduate	27.24	26.63	10.65 ***	42.26 *	37.93 *	40.91 **	49.26 **
Some college	32.46	32.45	31.38	28.50	33.04	40.14	33.60
Vocational training	9.68	9.89	10.15	7.96	7.71	6.20	1.42
High school	20.77	21.07	34.53	13.46	11.96 *	9.57 *	10.12 *
Less than high school	9.85	9.96	13.29 ***	7.82	9.36 *	3.18 **	5.6
Married, ever (%)	46.44	48.42	35.81 **	37.44	27.61 ***	2.57 ***	5.44 ***
Tested for STI, WII (%)	13.13	12.61	8.73	15.43	20.03	28.78 *	32.33
Last regular medical visit (%)							
< 1 year	47.96	47.81	48.79	46.89	41.45	61.51 ***	57.56
1 year & < 2 years	16.00	16.04	16.56	13.16	12.83	15.59	24.15
2 years	36.04	36.15	34.65	39.95	45.72	22.90 *	18.29 **
Victimization (%)							

	Total Sample	100% Heterosexual		Mostly Straight/Bisexual		Mostly Gay/100% Gay	
		Op. Sex Only	Both Sex	Op. Sex Only	Both Sex	Same Sex Only	Both Sex
< .25	77.26	77.82	52.95 ***	78.72	77.76	72.88	84.77
.25 and < .75	12.86	12.37	32.33 ***	17.19	12.23	11.7	3.70 ***
> .75	9.88	9.81	14.72	4.09 **	10.01	15.42 *	11.53
Sexual Behaviors							
Total No. of sexual partners	17.64	17.10	20.21 ***	12.09	26.92 *	37.45 ***	29.47 ***
Anal sex	45.42	43.53	37.39	51.97	79.90 ***	93.82 ***	92.34 ***

Source: National Longitudinal Survey of Adolescent Health

* p .05;

** p .01;

*** p .001

Table 3

Odds ratios for self-reported STI diagnosis among U.S. females aged 24–32.

	Model 1	Model 2	Model 3	Model 4
Sexual Identity-Behavior Subgroups (Heterosexual-WSM)				
Heterosexual-WSMW	1.75 (0.30) **	1.64 (0.29) **	1.29 (0.23)	1.24 (0.22)
Bisexual-WSM	1.47 (0.16) ***	1.42 (0.16) **	1.23 (0.14)	1.21 (0.14)
Bisexual-WSMW	2.20 (0.27) ***	1.99 (0.24) ***	1.31 (0.17) *	1.25 (0.17)
Gay-WSMW	0.47 (0.13) **	0.45 (0.12) **	0.34 (0.12) **	0.33 (0.11) **
Race/ethnicity (Non-Hispanic white)				
Non-Hispanic black	2.44 (0.22) ***	2.41 (0.21) ***	2.63 (0.23) ***	2.60 (0.22) ***
Hispanic	1.18 (0.14)	1.16 (0.14)	1.28 (0.15) *	1.26 (0.15) *
Asian	0.78 (0.18)	0.77 (0.19)	0.84 (0.19)	0.84 (0.20)
Other	1.49 (0.31)	1.43 (0.31)	1.53 (0.32) *	1.48 (0.32)
Age	1.07 (0.02) **	1.07 (0.02) **	1.06 (0.02) *	1.06 (0.02) *
Education (College graduate)				
< High school degree	1.97 (0.32) ***	1.81 (0.31) ***	1.92 (0.33) ***	1.80 (0.32) ***
High school degree	1.40 (0.16) *	1.32 (0.16) *	1.34 (0.16) *	1.29 (0.16) *
Vocational training	1.38 (0.15)	1.31 (0.14) *	1.28 (0.14) *	1.24 (0.14) *
Some college	1.39 (0.11) **	1.32 (0.11) ***	1.32 (0.11) ***	1.27 (0.10) **
Married, ever	0.62 (0.05) ***	0.62 (0.05) ***	0.68 (0.05) ***	0.68 (0.05) ***
Tested for STI, WII (not tested)	1.62 (0.10) ***	1.63 (0.10) ***	1.42 (0.09) ***	1.44 (0.09) ***
Last medical visit (< 1 year)				
1 year & > 2 years	0.84 (0.07)	0.84 (0.08)	0.85 (0.07)	0.84 (0.08)
2 years	0.86 (0.08)	0.85 (0.08)	0.83 (0.09)	0.82 (0.08)
Victimization Scale (.75)				
< .25		2.07 (0.19) ***		1.79 (0.17) ***
.25 and .75		1.27 (0.12) *		1.20 (0.12) *

	Model 1	Model 2	Model 3	Model 4
	OR (SE)	OR (SE)	OR (SE)	OR (SE)
Total number of partners			1.04 (0.00)***	1.04 (0.00)***
Anal sex, ever			1.34 (0.10)***	1.31 (0.10)***

Source: National Longitudinal Survey of Adolescent Health; OR=Odds Ratio; SE=Standard Error

* p .05;
** p .01;
*** p .001

Table 4

Odds ratios for self-reported STI diagnosis among U.S. males aged 24–32.

	Model 1	Model 2	Model 3	Model 4
Sexual Identity-Behavior Subgroups (Heterosexual-MSW)				
Heterosexual-MSMW	1.30 (0.31)	1.16 (0.29)	1.31 (0.31)	1.18 (0.28)
Bisexual-MSW	1.15 (0.31)	1.14 (0.30)	1.18 (0.32)	1.16 (0.31)
Bisexual-MSMW	1.81 (0.45) *	1.82 (0.45) *	1.59 (0.38)	1.60 (0.39)
Gay-MSM	1.38 (0.44)	1.37 (0.42)	1.09 (0.37)	1.09 (0.35)
Gay-MSMW	2.20 (0.72) *	2.30 (0.76) *	1.82 (0.60)	1.91 (0.63) *
Race/ethnicity (Non-Hispanic white)				
Non-Hispanic black	2.28 (0.27) ***	2.20 (0.26) ***	2.28 (0.27) ***	2.21 (0.27) ***
Hispanic	1.18 (0.14)	1.17 * (0.14)	1.18 (0.14)	1.17 (0.14)
Asian	0.60 (0.13) *	0.60 (0.13) *	0.64 (0.14) *	0.64 (0.14) *
Other	1.53 (0.38) *	1.48 (0.38) *	1.55 (0.38)	1.50 (0.38)
Age	1.08 (0.03) **	1.08 (0.03) **	1.07 (0.03) **	1.07 (0.03) **
Education (College graduate)				
< High school degree	1.51 (0.21) ***	1.39 (0.19) *	1.45 (0.20) **	1.35 (0.19) *
High school degree	1.25 (0.16)	1.22 (0.16)	1.26 (0.17)	1.24 (0.17)
Vocational training	1.22 (0.18)	1.18 (0.18)	1.18 (0.18)	1.15 (0.17)
Some college	1.27 (0.13) *	1.22 (0.13)	1.20 (0.13)	1.16 (0.13)
Married, ever	0.94 (0.07)	0.95 (0.07)	0.98 (0.07)	0.99 (0.07)
Tested for STI, WII (not tested)	0.84 (0.11)	0.81 (0.11)	0.75 (0.10) *	0.74 (0.10) *
Last medical visit (< 1 year)				
1 year & < 2 years	1.06 (0.12)	1.05 (0.12)	1.07 (0.12)	1.06 (0.12)
2 years	0.82 (0.08) *	0.82 (0.08) *	0.81 (0.08) *	0.81 (0.08) *
Risk Factors				
Victimization Scale (.75)		1.66 (0.17) ***		1.56 (0.16) ***
< .25				

	Model 1	Model 2	Model 3	Model 4
	OR (SE)	OR (SE)	OR (SE)	OR (SE)
.25 and < .75		1.41 (0.16) **		1.37 (0.16) **
Total number of partners			1.01 (0.00) ***	1.01 (0.02) ***
Anal sex, ever			1.33 (0.10) ***	1.32 (0.09) ***

Source: National Longitudinal Survey of Adolescent Health; OR=Odds Ratio; SE=Standard Error

*
p .05;
**
p .01;

p .001