Published in final edited form as:

ACS Comb Sci. 2012 November 12; 14(11): 621–630. doi:10.1021/co300098v.

Application of a Catalytic Asymmetric Povarov Reaction using Chiral Ureas to the Synthesis of a Tetrahydroquinoline Library

Baudouin Gerard^{†,§}, Morgan Welzel O'Shea^{†,§}, Etienne Donckele[†], Sarathy Kesavan^{†,§}, Lakshmi B. Akella^{†,§}, Hao Xu[‡], Eric N. Jacobsen[‡], and Lisa A. Marcaurelle^{†,§,*}

[†]Chemical Biology Platform, The Broad Institute of Harvard and MIT, 7 Cambridge Center, Cambridge, Massachusetts 02142

[‡]Department of Chemistry and Chemical Biology, Harvard University, 12 Oxford Street, Cambridge, Massachusetts 02138

Abstract

A 2328-membered library of 2,3,4-trisubstituted tetrahydroquinolines was produced using a combination of solution- and solid-phase synthesis techniques. A tetrahydroquinoline (THQ) scaffold was prepared via an asymmetric Povarov reaction using cooperative catalysis to generate three contiguous stereogenic centers. A matrix of 4 stereoisomers of the THQ scaffold was prepared to enable the development of stereo/structure-activity relationships (SSAR) upon biological testing. A sparse matrix design strategy was employed to select library members to be synthesized with the goal of generating a diverse collection of tetrahydroquinolines with physicochemical properties suitable for downstream discovery.

Introduction

The acid-catalyzed [4+2] cycloaddition of *N*-aryl imines and electron-rich olefins, known as the Povarov reaction,¹ is a powerful method for the synthesis of tetrahydroquinolines (THQ), a commonly occurring motif in a variety of natural products and biologically active compounds (Figure 1).² Up to three contiguous stereocenters can be generated simultaneously in the Povarov reaction, and enantioselective catalytic variants of this reaction have been identified recently.³ As part of our ongoing efforts to develop new methods for producing collections of stereochemically and skeletally diverse small molecules,^{4,5} we were interested in applying the chiral urea/ Brønsted acid co-catalyzed asymmetric Povarov reaction^{3a} (Scheme 1) to the synthesis of a THQ-based library.

*To whom correspondence should be addressed: lisa_marcaurelle@h3biomedicine.com.

§Current address: H3 Biomedicine Inc., 300 Technology Square, Cambridge, Massachusetts. 02139

At the outset of the project we faced three synthetic challenges in the production of a diverse THQ-based library: (1) adaptation of the asymmetric Povarov reaction conditions to multigram scale, (2) development of a practical strategy for accessing a matrix of stereoisomers and (3) optimization of solid-phase diversification reactions. In this paper, we describe successful solutions to these challenges in the context of the large-scale synthesis of a collection of stereoisomeric THQ scaffolds as well as the solid-phase production of a 2328-membered THQ library.

Results and Discussion

Scaffold Design

In designing a THQ scaffold for library synthesis, we required two key features: 1) a primary alcohol for loading onto solid support and 2) chemical handles for introducing appendage diversity. With these features in mind, we began by selecting the imino Diels-Alder partners for the asymmetric Povarov reaction. Use of an imine glyoxolate **5**, obtained from condensation of aniline **6** and ethyl glyoxolate **7**, as the 4- π component,⁶ was attractive for a number of reasons (Scheme 2). The use of glyoxolate ester derivatives would provide a low molecular weight scaffold bearing a functional handle for loading the products onto solid-phase supports. In addition, the primary products of the Povarov reaction (**8**) contain an epimerizable stereogenic center that could allow further diversification of the scaffold. Finally, excellent enantio- and diastereoselectivities have been demonstrated in the urea-catalyzed asymmetric Povarov reaction of glyoxylate imines.^{3a}

Various substituted anilines (4-methoxycarbonyl **6a**, 4-((*tert*-butyldimethylsilyl)oxy) **6b**, 4-bromo **6c**) were evaluated as imine precursors in the Povarov reaction with dihydrofuran as a model dienophile. The imine generated from **6a** was unstable and resulted in a low yield on large scale in the Povarov reaction. Meanwhile, reactions of **6b** were found to be poorly reproducible and became problematic with future protecting group manipulations. In contrast, **6c** offered good reactivity and reproducibility in the imine formation on large scale. Furthermore, the aryl bromide could be used as a diversity site on solid phase for cross coupling reactions.⁷ It was found that the Povarov adduct could be recrystallized and thus confirming the absolute configuration of the cycloaddition product with *endo* stereochemistry.⁸

With the diene in hand, we moved on to study the dienophile partner **9**. We focused on dienophiles featuring a handle for solid-phase diversification and those which showed good reactivity while retaining high levels of enantio- and diastereoselectivity in the Povarov reaction. As noted in previous reports, 2,3-dihydropyrrole derivative **10** is a versatile dienophile for the Povarov reaction.^{3a,9,10} We conducted a study (Table 1) to determine the most suitable protecting group for the pyrrole nitrogen under Povarov reaction conditions. As expected, a noticeable difference in reactivity was observed based on the nature of the protecting group. For example, the nosylated pyrrole **10a** proved unreactive during the Povarov reaction mostly due to poor solubility in toluene at low temperatures (-60°C) (entry 1). Meanwhile, use of Boc-protected pyrrole **10b** led to an overall decrease in yield (38%) and diastereoselectivity (from 9:1 to 7:3 *dr*) (entry 2). However, it was found that Fmoc and Cbz carbamates, **10c** and **10d** gave more suitable results with up to 80% isolated yield of the corresponding cycloadduct and with good levels of diastereoselectivity (9:1 *dr*) and enantioselectivity (up to 93:7 *er*). Thus, dienophile **10c** and **10d** were selected for the scale up of the THQ scaffolds for library production.

Large-scale reaction optimization

Having selected **5c** as diene and **10c** and **10d** as dienophiles, we next evaluated the optimization for the asymmetric Povarov reaction on multigram scale to achieve high yield and optimal levels of enantio- and diastereoselectivity suitable for library production. In preliminary studies, we found that the manner in which the imine was generated was a critical parameter for obtaining consistent results on large scale.¹¹ The imine formed *in situ* was found to be extremely sensitive to nucleophilic attack with an excess of aniline leading to side product formation.¹² A practical procedure was developed involving formation of the imine *via* slow addition of aniline **6c** into glyoxolate **7** in toluene at 0 °C, followed by isolation of the crude imine *via* solvent removal *in vacuo*.⁷

Initially, when employing previously described conditions for the Povarov reaction^{3a} (4% mol of urea catalyst **1**, 2% mol of *ortho*-nitrobenzenesulfonic acid (NBSA) as Brønsted acid in toluene at –60 °C in presence of 5 Å MS, 1 mmol scale), we observed formation of the desired product **11** with good enantio- (9:1 *er*) and diastereoselectivity (9:1 *dr*) albeit in modest yield (40%) (Table 2, entry 1). We hypothesized that residual amount of water from the Brønsted acid may hydrolyze the water-sensitive imine **5c**^{10,13} and therefore affect the overall isolated yield of **11** on large scale. After studying the effect of other Brønsted acids in the Povarov reaction, we found that the use of anhydrous *p*-toluenesulfonic acid (PTSA)¹⁴ gave comparable results (yield and enantio- and diastereoselectivity) to NBSA (Table 2, entries 2 and 3). Anhydrous PTSA was found to be easy to use and could be stored for several weeks at room temperature in a desiccator. Thus, for practical reasons we decided to use PTSA for the remainder of our optimization experiments. This modification enable us to conduct the asymmetric Povarov reaction on large scale with up to 66 mmol of **5** (17 g) in presence of **10c** as the dienophile to give moderate yield and good enantiomeric ratio (65%, 88:12 *er*, entry 5). Similar yield and enantioselectivity of **11d** was obtained when **10d** was used as dienophile in the Povarov reaction. In addition, **11d** could be easily purified *via* crystallization to give almost enantiomeric pure material. Gratifyingly, we observed no erosion of reactivity when using lower catalyst and Brønsted acid loading (2 mol% of urea catalyst and 1 mol% of PTSA (entries 3 and 4)). Similar results were obtained in the Povarov reaction when the enantiomer of the urea catalyst **1** was employed.¹⁵

Epimerization studies

In order to obtain stereochemical diversity, a practical method to epimerize the stereogenic center adjacent to the ester was sought to convert the *endo* diastereomer **11c** into the thermodynamically favored *exo* diastereoisomer **12c**. Although there was precedent for a similar epimerization strategy via the corresponding aldehyde in the synthesis of the natural product martinelline,¹⁶ we were hoping to develop a synthetic sequence to access **12c** directly and on large scale. After extensive screening of epimerization conditions,¹⁷ it was found that treatment with 0.1 M solution of sodium methoxide in methanol at 60 °C gave the desired epimerization product **14c** in 70% yield. The *endo* methyl ester **13c** that was recovered after purification could be resubjected to the same conditions to give **14c** in 90% overall yield over two cycles.

Library Scaffold Preparation

For solid-phase library production, a protecting group manipulation was required to replace the incompatible Cbz group with Fmoc. To streamline the preparation of the final cores, we decided to utilize Fmoc-pyrrole **10d** as the dienophile to generate *endo* diastereoisomers and Cbz-pyrrole **10c** to generate the remaining *exo* diastereoisomers. After Povarov cycloaddition using Fmoc-pyrrole **10d** with either **1** or its enantiomer, cycloadduct **11d** was reduced using LiBH₄ to afford the corresponding primary alcohol **15d**.¹⁸ Scaffold **15d** could

be easily recrystallized in a 9:1 mixture of benzene and dichloromethane to give enantioenriched compounds with excellent enantioselectivity (99:1 *er*). After epimerization of **11c** to **14c**, the *exo* Fmoc protected diastereoisomer was obtained *via* a stepwise processes involving: 1) ester reduction using LiBH₄ and 2) protecting group exchange from the incompatible Cbz to Fmoc in order to afford **16d**. Recrystallization of **16d** afforded enrichment of enantiopure compound up to 97:3 *er*.

During initial feasibility studies, it was found that the THQ aniline had limited reactivity upon treatment with various electrophiles, such as acyl chlorides and sulfonyl chlorides thus restricting its use as diversity site for solid-phase production. However, it was found that the aniline could be alkylated by reductive amination with a limited number of aldehydes. Formaldehyde was found to be the most reactive among the aldehydes tested. Scaffolds **15d** and **16d** were therefore methylated *via* reductive amination with excellent yields (90%) to give **17d** and **18d**. This strategy allowed us to generate eight different scaffolds, including both enantiomers of the NH and NMe *endo* and *exo* cores (Figure 2).

Solid-Phase Feasibility Studies

With the NH and NMe THQ-scaffolds in hand, we turned our attention to the development of solid-phase methods for the introduction of building blocks at the two diversity sites (secondary amine and aryl bromide). Amine capping at the first diversity site for both the NH and NMe sub-libraries involved screening with a variety of electrophiles, including isocyanates, sulfonyl chlorides, aldehydes and acids. Bis-capping was observed with the use of isocyanates in the presence of the free aniline and therefore removed from the design of the NH sub-library. While isocyanates performed well for the NMe sublibrary, the use of aldehydes was problematic resulting in significant decomposition during reductive alkylation conditions. Therefore aldehydes were removed from the NMe sublibrary design. Acids and sulfonyl chlorides performed well for both sub-libraries.

Next, the feasibility of Pd-mediated cross coupling reactions at the aryl bromide was investigated. Both Sonogashira and Suzuki cross couplings were explored. Initial screenings of the Sonogashira cross coupling led to poor conversion and produced a significant amount of side products. Therefore, we decided to focus solely on the Suzuki cross coupling for production purposes. Traditional Suzuki conditions on solid phase were attempted using 3-methoxybenzene boronic acid **19** and sulfonamides **20** and **21** as a model (Table 3, entry 1). Moderate conversion to the desired product was observed in addition to the formation of a significant amount of unknown byproducts¹⁹, possibly due to a rapid protodeboronation step²⁰ versus a slow oxidative insertion step. A variety of other conditions were screened including ligand (P(*t*-Bu)₃, PEPPSI **26**,²¹ S-Phos **24**, Pd catalyst (PdCl₂(PPh₃)₂, Pd(dba)₂), bases (TEA, K₃PO₄, Cs₂CO₃) and solvents (EtOH, PhMe, 1,4-dioxane, THF) in an attempt to increase the rate of oxidative insertion and potentially limit formation of side products without success (Table 3, entries 1–4). A recent publication from the Buchwald lab highlights the use of X-phos ligand in a preformed catalyst.²² The use of such precatalyst was reported to increase the rate of boronic acid coupling versus its decomposition due to the use of milder reaction condition. Upon treatment with the Buchwald precatalyst **27** at room temperature, we were delighted to find formation of the desired cross coupling product in high conversion without the formation of undesired side products for both NH and NMe sub-libraries (entries 5 and 6).

With this information in hand, a sparse matrix design strategy for each sub-library was implemented to select library members to be synthesized. First, a virtual library was constructed for both sub-libraries using all possible building block combinations at R₁ (amine) and R₂ (aryl bromide) using a master list of reagents (R₁ = sulfonyl chlorides, isocyanates, acids and aldehydes; R₂ = boronic acids).²³ This resulted in ~2400 compounds

per stereoisomer. Physicochemical property filters were then applied to eliminate building block combinations that led to products with undesirable properties. Property filters included the following: MW 500, ALogP -1 to 5, H-bond acceptors + donors 10, rotatable bonds 10 and TPSA 140. A total of 274 compounds per scaffold for the NH sub-library and 310 compounds per scaffold for the NMe sub-library were selected from the remaining set using chemical similarity principles, maximizing diversity but retaining near neighbors for built-in SAR.²² The reagents selected for library production are shown below (Chart 1 and Chart 2). The same set of reagents was used for each stereoisomer thereby maintaining the ability to generate SSAR for each building block combination.

Solid-Phase Library Production

With the solid-phase feasibility studies complete, we turned our attention to library production. Immobilization of the scaffold onto solid support was achieved *via* activation of the SynPhase Lanterns (L-series) with triflic acid followed by treatment with scaffolds **15–18** in the presence of excess 2,6-lutidine (Scheme 4).²⁴ The average loading levels obtained were 17 $\mu\text{mol/Lantern}$ for the NH sub-library and 13.3 $\mu\text{mol/Lantern}$ for the NMe sub-library. Removal of the Fmoc protecting group under standard conditions followed by capping with the appropriate electrophilic building blocks for NH and NMe sub-libraries yielded compounds **12**{1–42} and **13**{8–54}. Subsequent Pd-mediated cross coupling utilizing the Suzuki reaction afforded compounds **14**{1–42, 1–14} and **15**{8–54, 1–14}. Once the reaction sequence was complete, the Lanterns were subjected an aqueous sodium cyanide wash (1:1 1.0 M NaCN in water:THF solution) in order to remove any residual Pd. Release of the final compounds (**16**{1–42, 1–14} and **17**{8–54, 1–14}) from solid support was achieved via treatment with HF/pyridine. All library products were analyzed by ultra-performance liquid chromatography, and compound purity was assessed by UV detection at 210 nm. The average purity of the THQ NH library (1088 compounds) was 78%, with 71% of the library being >75% pure, while the average purity of the THQ NMe library (1240 compounds) was 75%, with 68% of the library being >75% pure. A full purity analysis is provided in Figure 3. Of note, certain nitrogen containing building blocks proved problematic during the amine capping for the NH sub-library, including acids **12**, **24**, **29** and **30**, while the *trans*-1-propenylboronic acid (**1**) generally did not perform well for either sub-library.

Library Analysis

Analysis of the THQ library (Table 4 and Figure 4) reveals that the calculated physicochemical property profile for the compound set is within the intended range for the library design (*vide supra*). Meanwhile, the structural diversity of the THQ library was analyzed in comparison to the NIH Molecular Library Repository (MLSMR) as we intended to submit a subset of these compounds to the collection at the time of the analysis. We employed multifusion similarity (MFS) maps for the comparison of each collection using extended connectivity fingerprints (ECFP₄) for molecular representation and Tanimoto coefficient as the similarity measure.²⁵ In this method, each molecule in the test set (THQ library, n = 2328) is compared to every molecule in the reference set (2011 MLSMR, n = 335834) and the largest similarity score and the mean similarity score to the reference set is obtained. The resulting mean similarity (x-axis) and maximum similarity (y-axis) values are plotted in two dimensions as a scatter plot facilitating the visual characterization and comparison. Figure 5 shows the MFS map comparing the THQ library to the MLSMR. Each data point in the map depicts a compound from the test set and its location was influenced by the reference set. (The reference compounds themselves do not appear on the plot). The maximum mean similarity of the THQ library is 0.13 (Tc) indicative of the overall structural diversity with the respect to the MLSMR reference set. Furthermore, there are no compounds with maximal similarity equal or greater than 0.57 (Tc) in the MLSMR,

illustrating that these compounds represent regions of unoccupied chemical space within this collection.

Conclusions

In summary a 2328-membered library of tetrahydroquinolines was successfully prepared using the asymmetric Brønsted acid/urea-catalyzed Povarov reaction^{3a} as a key step. Adaptation of the Povarov reaction for large scale synthesis led to the use of anhydrous PTSA as the Brønsted acid in place of NBSA and optimization of the imine formation step. These modifications enabled the large scale (>5 g) preparation of 4 stereoisomers of two THQ scaffolds with two functional handles for solid-phase diversification. *In silico* library design followed by production on SynPhase Lanterns afforded a diverse library of functionalized tetrahydroquinolines with properties suitable for downstream discovery.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was funded in part by the NIGMS-sponsored Center of Excellence in Chemical Methodology and Library Development (Broad Institute CMLD; P50 GM069721), as well as the NIH Genomics Based Drug Discovery U54 grants Discovery Pipeline RL1CA133834 (administratively linked to NIH grants RL1HG004671, RL1GM084437, and UL1DE019585). The authors would like to thank Stephen Johnston and Anita Vrcic for compound management and analytical support.

References

1. Kouznetsov VV. Recent synthetic developments in a powerful imino Diels-Alder reaction (Povarov reaction): Application to the synthesis of N-polyheterocycles and related alkaloids. *Tetrahedron*. 2009; 65:2721–2750.
2. (a) Leeson PD, Carling RW, Moore KW, Moseley AM, Smith JD, Stevenson G, Chan T, Baker R, Foster AC, Grimwood S, Kemp JA, Marshall GR, Hoogsteen K. 4-Amido-2-carboxytetrahydroquinolines. Structure-activity relationships for antagonism at the glycine site of the NMDA receptor. *J. Med. Chem.* 1992; 35:1954–1968. [PubMed: 1534584] (b) Batey RA, Simoncic PD, Lin D, Smyj RP, Lough AJ. A three-component coupling protocol for the synthesis of substituted hexahydropyrrolo[3,2-*c*]quinolines. *Chem. Comm.* 1999; 7:651–652. (c) Takamura M, Funabashi K, Kanai M, Shibasaki M. Catalytic enantioselective reissert-type reaction: Development and application to the synthesis of a potent NMDA receptor antagonist (–)-L-689,560 using a solid-supported catalyst. *J. Am. Chem. Soc.* 2001; 123:6801–6808. [PubMed: 11448184] (d) Witherup KM, Ransom RW, Graham AC, Bernad AM, Salvatore MJ, Lumma WC, Anderson PS, Pitzenberger SM, Vargatt SL. Martinelline and Martinelline Acid, Novel g-protein linked receptor antagonists from the tropical plant *Martinella iquitosensis* (Bignoniaceae). *J. Am. Chem. Soc.* 1995; 117:6682–6685. (e) Kumar A, Srivastava S, Gupta G, Chaturvedi V, Sinha S, Srivastava R. Natural product inspired diversity oriented synthesis of tetrahydroquinoline scaffolds as antitubercular agent. *ACS Comb. Sci.* 2011; 13:65–71. [PubMed: 21247127] (f) Schiemann K, Finsinger D, Zenke F, Amendt C, Knochel T, Brüge D, Buchstaller H-P, Emde U, Stahle W, Anzali S. The discovery and optimization of hexahydro-2H-pyrano[3,2-*c*]quinolines (HHPQs) as potent and selective inhibitors of the mitotic kinesin-5. *Bioorg. Med. Chem. Lett.* 2010; 20:1491–1495. [PubMed: 20149654] (g) Sridharan V, Suryavanshi PA, Menendez JC. Advances in the chemistry of tetrahydroquinolines. *Chem. Rev.* 2011; 111:7157–7259. [PubMed: 21830756] (h) Demont, EH.; Garton, NS.; Gosmini, RLM.; Hayhow, TGC.; Seal, J.; Wilson, DM.; Woodrow, MD. Tetrahydroquinolines derivatives and their pharmaceutical use. *Europe Patent PCT Int. Appl.* WO2011054841A1. 2011.
3. (a) Xu H, Zuend SJ, Woll MG, Tao Y, Jacobsen EN. Asymmetric cooperative catalysis of strong Brønsted acid-promoted reactions using chiral ureas. *Science*. 2010; 327:986–990. [PubMed:

20167783] (b) Ishitani S, Kobayashi S. Catalytic asymmetric aza Diels-Alder reactions using a chiral Lanthanide Lewis acid. Enantioselective synthesis of tetrahydroquinoline derivatives using a catalytic amount of a chiral source. *Tetrahedron Lett.* 1996; 37:7357–7360. (c) Akiyama T, Morita H, Fuchibe K. Chiral Brønsted acid-catalyzed inverse electron-demand aza Diels–Alder reaction. *J. Am. Chem. Soc.* 2006; 128:13070–13071. [PubMed: 17017784] (d) Liu H, Dagousset, Masson G, Retailleau P, Zhu JP. Chiral Brønsted acid-catalyzed enantioselective three-component Povarov reaction. *J. Am. Chem. Soc.* 2009; 131:4598–4599. [PubMed: 19334771] (e) Xie M, Liu X, Zhu Y, Zhao X, Xia Y, Lin L, Feng X. Asymmetric synthesis of tetrahydroquinolines with quaternary stereocenters through the Povarov reaction. *Chem. Eur. J.* 2011; 17:13800–13805. [PubMed: 22083970] (f) Ren L, Lei T, Ye J-X, Gong L-Z. Step-economical synthesis of tetrahydroquinolines by asymmetric relay catalytic Friedländer condensation/transfer hydrogenation. *Angew. Chem. Int. Ed.* 2012; 51:771–774. (g) Tan HR, Ng FH, Chang J, Wang J. Highly enantioselective assembly of functionalized tetrahydroquinolines with creation of an all-carbon quaternary center. *Chem. Eur. J.* 2012; 18:3865–3870. [PubMed: 22378143] (h) He L, Bekkaye M, Retailleau P, Masson G. Chiral phosphoric acid catalyzed inverse-electron-demand aza-Diels-Alder reaction of isoeugenol derivatives. *Org. Lett.* 2012; 14:3158–3161. [PubMed: 22671430]

4. (a) Schreiber SL. Target-oriented and diversity-oriented organic synthesis in drug discovery. *Science.* 2000; 287:1964–1969. [PubMed: 10720315] (b) Burke MD, Schreiber SL. A planning strategy for diversity-oriented synthesis. *Angew. Chem. Int. Ed.* 2004; 43:46–58. (c) Nielsen TE, Schreiber SL. Toward the optimal screening collection: A synthesis strategy. *Angew. Chem. Int. Ed.* 2007; 74:48–56.
5. (a) Marcaurelle LA, Comer E, Dandapani S, Duvall JR, Gerard B, Kesavan S, Lee MD IV, Liu H, Lowe JT, Marie J-C, Mulrooney CA, Pandya BA, Rowley A, Ryba TD, Suh B-C, Wei J, Young DW, Akella LB, Ross NT, Zhang Y-L, Fass DM, Reis SA, Zhao W-Z, Haggarty SJ, Palmer M, Foley MA. An aldol-based build/couple/pair strategy for the synthesis of medium- and large-sized rings: Discovery of macrocyclic histone deacetylase inhibitors. *J. Am. Chem. Soc.* 2010; 132:16962–16976. [PubMed: 21067169] (b) Gerard B, Duvall JR, Lowe JR, Murillo T, Wei J, Akella LB, Marcaurelle LA. Synthesis of a stereochemically diverse library of medium-sized lactams and sultams via S_NAr cycloetherification. *ACS. Comb. Sci.* 2011; 13:365–374. [PubMed: 21526820]
6. For references on the use of imine glyoxylate in Povarov reaction see: Alves JM, Azoia NG, Fortes GA. Regio- and stereo-selective aza-Diels–Alder reaction of ethyl glyoxylate 4-methoxyphenylimine with 1,3-dienes in the presence of $BF_3 \cdot Et_2O$. Evidence for a non-concerted mechanism. *Tetrahedron.* 2007; 63:727–734. Hermitage S, Howard JAK, Jay D, Pritchard RG, Probert MR, Whiting A. Mechanistic studies on the formal aza-Diels–Alder reactions of *N*-aryl imines: Evidence for the non-concertedness under Lewis-acid catalysed conditions. *Org. Biomol. Chem.* 2004; 2:2451–2460. [PubMed: 15326525] Borriore E, Prato M, Scorrano G, Stivanello M, Lucchini V. Synthesis and cycloaddition reactions of ethyl glyoxylate imines. Synthesis of substituted furo-[3,2-*c*]quinolines and 7*H*-indeno[2,1-*c*]quinolines. *J. Heterocycl. Chem.* 1988; 25:1831–1835.
7. Tester SA, Mata EG. Prospect of metal-catalyzed C–C forming cross-coupling reactions in modern solid-phase organic synthesis. *J. Comb. Chem.* 2008; 10:487–497. [PubMed: 18512993]
8. See Supporting Information for further details.
9. For references on the use of dihydropyrrole as dienophile in the Povarov reaction see: Hadden M, Nieuwenhuyen M, Potts D, Stevenson PJ, Thompson N. Synthesis and reactivity of hexahydropyrroloquinolines. *Tetrahedron.* 2001; 57:5615–5624.
10. For reference on synthesis of dihydropyrrole protected carbamate see: Marais W, Holzapfel CW. On a facile synthesis of melatonin and other related indoles. *Syn. Commun.* 1998; 28:3681–3691. Kraus GA, Neuenschwander K. Facile synthesis of *N*-acyl-2- pyrrolines. *J. Org. Chem.* 1981; 46:4791–4792.
11. Borriore E, Prato M, Scorrano G, Stivanello M. Synthesis and cycloaddition reactions of ethyl glyoxylate imines. Synthesis of substituted furo-[3,2-*c*]quinolines and 7*H*-indeno[2,1-*c*]quinolines. *J. Heterocycl. Chem.* 1988; 25:1831–1835.
12. This side product was identified as the aminor adduct.
13. NBSA is commercially available only as the hydrate (containing x molecules of water). To remove residual amounts of water NBSA was dissolved in a minimum amount of dry THF and azeotroped

with toluene. This sequence was repeated three times until a dark brown viscous oil was obtained. The light sensitive oil was dried under high vacuum over CaCl_2 and stored in the dark.

14. *p*-Toluenesulfonic acid was dehydrated by azeotropic distillation with toluene using a Dean-Stark apparatus. The residue was then crystallized from benzene.
15. The SSS enantiomer urea catalyst was obtained from Astatech Inc.
16. Xia C, Heng L, Ma D. Total synthesis of (+/-)-martinelline. *Tetrahedron Lett.* 2002; 43:9405–9409.
17. Keenan TP, Yaeger D, Holt DA. *Tetrahedron: Asymmetry.* 1999; 10:4331–4341.
18. (a) Khadem S, Joseph R, Rastegar M, Leek DM, Oudatchin KA, Arya P. Solution- and solid-phase approach to tetrahydroquinoline-derived polycyclics having a 10-membered ring. *J. Comb. Chem.* 2004; 6:724–734. [PubMed: 15360207] (b) Guo F, Chang BH, Rizzo CJ. An N1-hydrogen bonding model for flavin coenzyme. *Bioorg. Med. Chem. Lett.* 2002; 12:151–154. [PubMed: 11755342]
19. The molecular weight of the observed byproduct corresponded to phosphine (PR_3) incorporation as judged by LCMS.
20. (a) Kuivila HG, Reuwer JF, Mangravite JA. Electrophilic displacement reactions. XVI. Metal ion catalysis in the protodeboronation of areneboronic Acids. *J. Am. Chem. Soc.* 1964; 86:2666–2670. (b) Clapham KM, Batsanov AS, Bryce MR, Tarbit B. Trifluoromethyl-substituted pyridyl- and pyrazolylboronic acids and esters: Synthesis and Suzuki–Miyaura cross-coupling reactions. *Org. Biomol. Chem.* 2009; 7:2155–2161. [PubMed: 19421454] (c) Clapham KM, Batsanov AS, Greenwood RDR, Bryce MR, Smith AE, Tarbit B. Functionalized heteroarylpyridazines and pyridazin-3(2H)-one derivatives via palladium-catalyzed cross-coupling methodology. *J. Org. Chem.* 2008; 73:2176–2181. [PubMed: 18294000]
21. O'Brien CJ, Kantchev EAB, Valente C, Hadei N, Chass GA, Lough A, Hopkinson AC, Organ MD. Easily prepared air- and moisture-stable Pd–NHC (NHC=N-heterocyclic carbene) complexes: A reliable, user-friendly, highly active palladium precatalyst for the Suzuki–Miyaura reaction. *Chem. Eur. J.* 2006; 12:4743–4748. [PubMed: 16568494]
22. Kinzel T, Zhang Y, Buchwald SL. *J. Am. Chem. Soc.* 2010; 132, 40:14073–14075. [PubMed: 20858009]
23. For further details on the selection of reagents for master list see: Akella LB, Marcaurelle LA. Application of a sparse matrix design strategy to the synthesis of DOS libraries. *ACS. Comb. Sci.* 2011; 13:357–364. [PubMed: 21526822]
24. Ryba TD, Depew KM, Marcaurelle LA. Large scale preparation of silicon-functionalized SynPhase polystyrene Lanterns for solid-phase synthesis. *J. Comb. Chem.* 2009; 11:110–116. [PubMed: 19049425]
25. (a) Medina-Franco JL, Maggiora GM, Giulianotti MA, Pinilla C, Houghten RA. A similarity-based data-fusion approach to the visual characterization and comparison of compound databases. *Chem. Biol. Drug Des.* 2007; 70:393–412. [PubMed: 17927720] (b) Medina-Franco JL, Martinez-Mayorga K, Giulianotti MA, Houghten RA, Pinilla C. Visualization of the chemical space in drug discovery. *Curr. Comput. Aided Drug Des.* 2008; 4:322–333.

Figure 1.
Examples of natural products and biologically active compounds featuring a tetrahydroquinoline core.

Figure 2.
Matrix of 4 stereoisomers for the NH and NMe THQ scaffolds.

Figure 3.

Purity analysis for THQ library (UPLC analysis with UV detection at 210 nm). Library members are displayed as blocks of 4 stereoisomers (see legend) for both the NH and NMe sub-libraries. Reagents used for solid-phase diversification are shown on the *x*- and *y*-axes. (See Charts 1 and 2 for detailed list of reagents).

Figure 4.
Molecular weight (MW) and ALogP distribution for THQ-library members.

Figure 5. Multi-fusion similarity map comparing the THQ library (2328) to the 2011 MLSMR (335834). The reference set (MLSMR) is not shown on the map (see text for details).

Scheme 1.

Asymmetric co-catalysis of the Povarov reaction using a Brønsted acid and a chiral urea.

Scheme 2.
Retrosynthesis of THQ scaffold **8**.

Scheme 3.
Solution-phase synthesis of library scaffolds

Scheme 4.

Solid-phase synthesis of THQ library on SynPhase Lanterns

Aldehydes {1-7}: **Carboxylic Acids {8-31}:** **Sulfonyl Chlorides {32- 42} :** **Isocyanates {43-54} :**

Chart 1.
Building Blocks for Amine Capping

Boronic acids {1-13}: **Skip {14}:****Br**

Chart 2.
Building Block for Cross Coupling

Table 1

Screening of dienophiles for the asymmetric Povarov reaction

entry	R	product	<i>dr</i> ^a	<i>er</i> ^b	yield of 11 (%) ^c
1	Ns, 10a	11a/12a	—	—	N.R.
2	Boc, 10b	11b/12b	7:3	95:5	32
3	Cbz, 10c	11c/12c	9:1	93:7	58
4	Fmoc, 10d	11d/12d	9:1	84:16	80

^a Diastereoisomeric ratio (*dr*) was measured by UPLC with UV detection at 210 nm on crude reaction mixture

^b Enantiomeric ratio (*er*) was measured by SFC with UV detection at 210 nm after chromatography separation of the major diastereoisomer

^c Isolated yield of **11** after silica gel chromatography

Table 2

Large scale reaction optimization

entry	R ₁	acid	acid (mol %)	cat. (mol%)	scale (mmol)	product	<i>d^a</i>	<i>er^b</i>	yield of 11 (%) ^c
1	Cbz, 10c	NBSA	2	4	1	11c/12c	90:10	90:10	40
2	Cbz, 10c	NBSA	2	4	25	11c/12c	90:10	93:7	58
3	Cbz, 10c	PTSA	2	4	39	11c/12c	90:10	92:8	61
4	Cbz, 10c	PTSA	1	2	39	11c/12c	89:11	89:11	58
5	Cbz, 10c	PTSA	1	2	67	11c/12c	91:9	88:12	65
6	Fmoc, 10d	PTSA	1	2	35	11d/12d	90:10	84:16	89

^aDiastereoisomeric ratio (*dr*) was measured by UPLC with UV detection at 210 nm on crude reaction mixture

^bEnantiomeric ratio (*er*) was measured by SFC with UV detection at 210 nm after chromatography separation of the major diastereoisomer

^cIsolated yield of **11** after silica gel chromatography

Table 3

Solid-phase Suzuki feasibility studies

entry	SM	catalyst	base	T(°C)	solvent	product	conv (%) ^a	SM (%) ^a	byproduct (%) ^a
1	20	PdCl ₂ (PPh ₂) ₃	TEA	60	EtOH	22	60	12	18
2	20	Pd[(dba) ₂ , P(t-Bu) ₃]	K ₃ PO ₄	60	PhMe	22	70	5	25
3	20	PEPPSI	CsCO ₃	60	1,4 dioxane	22	78	9	13
4	20	Pd[(dba) ₂ , S-Phos	K ₃ PO ₄	60	PhMe	22	73	0	27
5	20	Buchwald precatalyst	K ₃ PO ₄	rt	THF	22	100	0	0
6	21	Buchwald precatalyst	K ₃ PO ₄	rt	THF	23	100	0	0

^a Conversion was measured by UPLC with UV detection at 210 nm.

Table 4

Property Analysis for the THQ Library

Property ^a	NH Scaffold ^b (<i>n</i> = 1)	NMe Scaffold ^b (<i>n</i> = 1)	NH Library (<i>n</i> = 1100)	NMe Library (<i>n</i> = 1252)
MW	283	297	420	433
ALogP	1.4	1.9	2.9	3.4
TPSA	44	36	73	65
Rotatable Bonds	1	1	3.9	3.7
HBA	3	3	4.4	4.2
HBD	3	2	2.0	1.3

^aPhysicochemical properties listed as mean values.^bProperty analysis of bare scaffolds, where R₁ and R₂ = H.