“Internal Mammary Vein Cannulation during Port-o-Cath Insertion”

Robert S. Dieter, MD, RVT^{1,2} George B. Kuzycz, MD³ Raymond A. Dieter, Jr., MD, MS^{3,4}

¹Loyola University, Maywood, Illinois

²Hines VA, Hines, Illinois

³DuPage Medical Group, Glen Ellyn, Illinois

⁴Northern Illinois Center for Surgery, Illinois

Address for correspondence and reprint requests Robert S. Dieter, MD, RVT, 2160 S. First Avenue, Maywood, IL 60153 (e-mail: rdieter@lumc.edu).

Int J Angiol 2012;21:245–246.

We read with great interest, the case report by Dr. Harish and Dr. Madhu regarding, “Inadvertent port: catheter placement in the azygous vein.”¹ The authors aptly point out that this is an uncommon occurrence during central venous access. Its diagnosis is difficult with the fluoroscopy in the anteroposterior (AP) position. Indeed, even chest X-rays in the AP projection can be of little value. We have included a case of internal mammary vein cannulation during Port-o-Cath insertion in our textbook, *Venous and Lymphatic Diseases*.² We have found that prompt diagnosis can be achieved with a venogram to confirm position. Furthermore, the lateral chest

X-ray is often beneficial in delineating the course of the catheter. Once again, we appreciate the authors for sharing their case.

References

- 1 Harish K, Madhu YC. Inadvertent port: catheter placement in the azygous vein. Int J Angiol 2012;21:103–106
- 2 Dieter RS, Kuzycz GB, Dieter RA. Venous and Lymphatic Diseases. New York: McGraw-Hill; 2011; Chap 16:205