Reply to comment on Bakker et al.: Change of Böhler's angle during conservatively-treated displaced intra-articular calcaneal fractures

Jens A. Halm · Berdine Bakker ·
Esther M. M. Van Lieshout · Tim Schepers

Received: 7 December 2012 / Accepted: 9 December 2012 / Published online: 29 December 2012
© Springer-Verlag Berlin Heidelberg 2012

Dear Editor,

We thank Jiang et al. for their interest in our manuscript. Herein, we would like to reply to their comments on Bakker et al.: Change of Böhler's angle during conservatively-treated displaced intra-articular calcaneal fractures [1].

1. With respect to the patient characteristics, we looked at age and gender. Data on smoking habits and diabetes were not available to us for analysis. In the literature, the time patients remain non-weight bearing ranges between six and 12 weeks. It is our protocol to keep these patients non-weight bearing for 12 weeks, which we expect is enough for fracture healing even in patients who smoke or have diabetes.
2. In the study a HET (High Energy Trauma) was defined as per the Advanced Trauma Life Support Criteria [2]:
 - a. a fall from a height of more than three metres
 - b. car accident with speed above 50 km/h (restrained victim)
 - c. ejection from a car
 - d. vehicle rollover on collision

- e. deformation of vehicle by more than 50 cm
- f. pedestrian victim versus vehicle at speed above 20 km/h
- g. bicycle/motorcycle versus vehicle at speed above 30 km/h

All other trauma was considered to be of a low energy type.

3. As stated correctly in the letter by Jiang et al. the analysis of differences in Böhler's angle between male ($n=30$) and female patients ($n=8$), revealed no statistical differences. Neither the univariate nor the multivariable analysis showed a statistically significant difference. This might, of course, be due to the low number of female participants, but other factors such as osteoporosis cannot be ruled out.
4. All patients were treated with early range of motion exercises and 12 weeks non-weight bearing.
5. We cannot correct for patient noncompliance, but one might speculate that patients with bilateral calcaneal fractures may be more compliant with the non-weight bearing regime being wheelchair-bound. On the other hand one might argue that, for the same reason, patients with bilateral calcaneal fracture tend to bear weight earlier to avoid a wheelchair. As per protocol the bilateral fractures were also kept non-weight bearing for 12 weeks.
6. The average follow-up was indeed about a half year. This is because after this period people were released from outpatient follow-up. Quite a large

J. A. Halm · B. Bakker · E. M. M. Van Lieshout · T. Schepers (✉)
Department of Surgery–Traumatology, Erasmus MC, University
Medical Center Rotterdam, Room H822-k, P.O. Box 2040, 3000
CA Rotterdam, The Netherlands
e-mail: t.schepers@erasmusmc.nl

proportion of patients, especially those in whom conservative treatment is preferred, are subsequently lost to follow-up. In surgically treated patients with longer follow-up a decrease in Böhler's angle is seen even after one year [3, 4]. We would like to encourage others to look at conservatively treated patients with a longer follow-up and to evaluate patients with a minimally displaced fracture with an initial angle of more than 15°, as these were not available for our current study. Secondly, it would be interesting to see whether or not this secondary decrease has an effect on outcome.

References

1. Bakker B, Halm JA, Van Lieshout EMM, Schepers T (2012) The fate of Bohler's angle in conservatively-treated displaced intra-articular calcaneal fractures. *Int Orthop* 36:2495–2499
2. American College of Surgeons (2004) Advanced trauma life support student manual. 7th edn. American College of Surgeons Committee on Trauma, Chicago, 14
3. Hyer CF, Atway S, Berlet GC, Lee TH (2010) Early weight bearing of calcaneal fractures fixated with locked plates: a radiographic review. *Foot Ankle Spec* 3:320–323
4. Rak V, Ira D, Masek M (2009) Operative treatment of intra-articular calcaneal fractures with calcaneal plates and its complications. *Indian J Orthop* 43:271–280