Practical Diagnostic Testing for Human Immunodeficiency Virus

J. BROOKS JACKSON^{1,2*} AND HENRY H. BALFOUR, JR.^{1,3}

Departments of Laboratory Medicine and Pathology¹ and Pediatrics,³ University of Minnesota Medical School, Minneapolis, Minnesota 55455,¹ and American Red Cross—St. Paul Region, St. Paul, Minnesota 55107²

INTRODUCTION	124
ANTIBODY TESTING	124
EIA	125
Western Blot	126
IFA	127
RIPA	127
Future HIV Antibody Tests	128
Clinical Significance and Use of HIV Antibody Testing	128
HIV ANTIGEN DETECTION	129
HIV CULTURE	131
HYBRIDIZATION TECHNIQUES	132
BIOHAZARDS OF HIV TESTING	133
CONCLUSION	134
ACKNOWLEDGMENTS	135
LITERATURE CITED	135

INTRODUCTION

The first case of acquired immunodeficiency syndrome (AIDS) was reported in June 1981 (22). As of August 1987, more than 40,000 cases of AIDS have been reported in the United States according to the Centers for Disease Control (CDC) (30). More than half of these patients have died, and the remainder will probably die within a couple of years unless effective treatment is developed. The cause of this deadly disease is a human retrovirus now known as human immunodeficiency virus type 1 (HIV). This virus was discovered by Montagnier's group in 1983, using electron microscopy, reverse transcriptase assay of culture supernatant fluid, immunofluorescence, and radioimmunoprecipitation (7). These results were later confirmed, and the virus was further characterized by Gallo's group in 1984 (47, 98, 107, 109). HIV is now known to infect preferentially CD4 lymphocytes (T-helper cells) which results in severe immunologic impairment, leading to opportunistic infections, cancer, and death (62, 68, 126, 128). HIV can also infect macrophages and possibly other cell types in the central nervous system, which may lead to severe dementia (49, 69, 70). On the other hand, having integrated into the deoxyribonucleic acid (DNA) of host cells, the virus can remain latent for years without causing any signs or symptoms. Nevertheless, 76% of individuals who have been infected at least 7 years have developed AIDS or AIDS-related complex (ARC) (N. A. Hessel, G. W. Rutherford, P. M. O'Malley, L. S. Doll, W. W. Darrow, and H. W. Jaffe, Third International Conference on Aids, p. 1, 1987). It is possible that all HIV-infected people who do not die of other causes first will eventually succumb to AIDS.

Exposure to body fluids such as semen, blood, and cervical secretions appears necessary for transmission (25). Consequently, most AIDS patients initially were sexually active homosexual men, blood transfusion recipients, hemo-

philia patients, intravenous drug abusers who shared needles, and infants of infected mothers (25). However, as the prevalence of HIV infection increases, the general population will become at greater risk primarily through heterosexual transmission, as apparently has been the case in Africa (99).

Fear that HIV infection could be transmitted by blood transfusion prompted the development of a diagnostic routine screening test for HIV antibodies in the serum of blood donors who do not recognize or admit their own risk of infection (107). In addition to protecting the blood supply, the test has been useful for other prevention efforts in high-risk groups and in making the diagnosis of AIDS, because the disease has many different clinical presentations.

The CDC now estimates that 1.5 million Americans are infected and predicts that there will be 324,000 AIDS cases in the United States by the end of 1991 (119). Moreover, another retrovirus, named HIV-2, which also causes AIDS, has recently been reported in Africa (33). The existence of an additional pathogenic human retrovirus could increase the CDC estimates. The need for clinical laboratories to develop expertise in the diagnostic testing for HIV will only become greater. In this review, we discuss the principles, procedures, and merits of the different HIV detection assays and address the biohazards of HIV testing to the laboratory worker.

ANTIBODY TESTING

Several types of assays have been developed in the past 4 years for the detection of antibody to HIV. The enzyme-linked immunosorbent assay (EIA) is the most widely used serologic test for HIV antibody detection in the world today. At least seven commercial EIAs have been licensed in the United States since the first one in March 1985. The EIA is used primarily as a screening test of blood donors and individuals at risk for AIDS. Individuals who are repeatedly

* Corresponding author.

reactive (at least two of three tests positive) by the EIA for HIV antibodies in their sera or plasma should then be retested for HIV antibody with a supplemental assay, such as the Western blot (29). Less commonly used as a supplemental test is the indirect immunofluorescence assay (IFA) or radioimmunoprecipitation assay (RIPA). Positive reactivity by one of these assays confirms the presence of HIV antibody, which is indicative of past exposure to HIV proteins and presumably current HIV infection. However, it should be noted that, while these tests are used to confirm positive EIA results, they are not necessarily more sensitive or specific than the EIA, because they are technically difficult and subjective in interpretation and the procedures and reagents vary considerably in different laboratories (17). In fact, of these supplemental assays, only one commercial Western blot kit (E. I. du Pont de Nemours & Co., Inc., Wilmington, Del.) has been licensed as of July 1987.

With this in mind, we now describe the methods and clinical utility of these assays for HIV antibody detection.

EIA

The EIA test for HIV antibody was developed in the research laboratory soon after HIV was discovered to be the causative agent of AIDS. Nearly all commercial kits use the same method, in which partially purified viral antigens derived from whole disrupted virus in cell culture are bound to polystyrene beads or to plastic microtiter wells. The serum or plasma sample is incubated with the fixed antigens for 30 min to 2 h and then washed. If any immunoglobulin G (IgG) antibody attaches to the bead or well, it will be detected when an anti-IgG antibody is added. Typically, the anti-IgG antibody is labeled with an enzyme (i.e., alkaline phosphatase, horseradish peroxidase) so that, after any unbound antibody is washed away, an enzymatic reaction will produce a color proportional to the amount of human IgG present when a substrate is added. Some EIA kits use a mixture of antibodies to the heavy chains and light chains of human immunoglobulin so that IgM antibodies to HIV may also be detected. Positive and negative controls are run simultaneously to determine the absorbance level or cutoff value above which a positive reaction is defined. Most EIA kits are automated and take <4 h to perform.

Initially, a single sample from a serum or plasma specimen is tested. In patients who have recently been transfused with blood or fluid expanders, a pretransfusion specimen should be used for testing to avoid dilution of HIV antibody to undetectable levels (28). If the test result is positive, two samples from the initial specimen are typically tested. If at least two of three samples show positive reactivity, the specimen is considered to be repeatedly reactive and a supplemental assay such as a Western blot assay should be performed (81). No repeatedly reactive units of blood should be used for transfusion regardless of the supplemental test result (23), even though these donors are unlikely to be infected (19, 39). Patients or blood units that are nonreactive in at least two of three EIAs are considered to be noninfectious, and supplemental testing is not necessary unless the individual is in a high-risk group for HIV infection.

The sensitivity and specificity of the commercial EIAs are quite good if one uses the Western blot as the "gold" standard. In a comparison study of six different commercial EIA kits, the sensitivity of the assays ranged from 98.1 to 100.0% based on 163 Western blot-positive individuals (101). In testing 4,999 Western blot-negative blood donors, 0.06 to 1.3% were initially reactive and 0.0 to 0.42% were repeatedly

reactive, giving a specificity of 99.6 to 100.0%. However, the predictive value of a positive test was highly dependent on the prevalence of HIV infection in the population tested. In AIDS and ARC patients, the predictive value of a positive test was 100% for all six EIA kits, but ranged from 5 to 100% in the blood donor group. In another study, the predictive value of a negative EIA test ranged from 99.99% in the general population to as low as 76.9% in members of high-risk groups depending on the cutoff used for positive reactivity and the incidence of HIV infection (112). While the EIA kits are very sensitive compared with the Western blot, it should be remembered that these assays detect antibody to HIV, not antigen. Therefore, HIV-infected individuals may be missed by antibody tests prior to seroconversion. Seroconversion usually occurs within 2 to 3 months (35, 39, 45, 60, 72, 80) but can take up to 5 months (118) or even longer if the individual is immunocompromised (3). One recent study reports that latent HIV infection may occur for up to 34 months or longer without overt seroconversion as detected by EIA screening tests (100). Moreover, some EIA and Western blot methods detect only IgG antibody to HIV. Therefore, some current screening and supplemental Western blot assays will not detect HIV-infected individuals prior to the switch from IgM to IgG antibody (10, 93). The time to switch from IgM to IgG HIV antibody has been reported to take as long as 41 weeks (9, 10). Therefore, follow-up testing of individuals at high risk for HIV infection is important. In addition, it should be noted that current EIAs detect antibody to the predominant HIV strain. These tests were positive in only 5 (24%) of 21 patients infected with HIV-2 (34). Fortunately, no HIV-2 infections in blood donors or patients have been reported in the United States.

On the other hand, false-positive EIAs can occur. In general, false-positive results are inversely related to the degree of HIV EIA reactivity. In a blood bank screening program, highly reactive specimens by EIA were strongly associated with a positive Western blot or culture (86.7%), in contrast to moderately and weakly reactive specimens (1.9%) (124). These false-positive results can be due to intrinsic variability in test performance, heat inactivation of serum, HLA antibodies with specificity for HLA DQ3 and DR4 antigens (found in the H-9 cell line used to grow stock virus), and probably other contaminating antigens from cell lines (12, 71). False-positive EIA results have been reported to be as high as 19% in hemophilia patients (113), 13% in alcoholic patients with hepatitis (83), 4% in hemodialysis patients (96), and 24% in patients who have a positive rapid plasma reagin test (41). Therefore, confirmatory testing of all EIA repeatedly reactive specimens is essential.

In addition to the commercial EIA kits which detect a number of different HIV antibodies, nonlicensed competitive EIAs (Abbott Laboratories, North Chicago, Ill.) which are specific for the detection of antibody to the core antigens or the envelope antigens have been developed (53). These assays have beads coated with recombinant core antigen or with recombinant envelope antigen. Serum samples are incubated with the beads and enzyme-conjugated human antibody to HIV core or envelope antigen for 16 to 22 hs. Beads are washed and substrate is added. The intensity of the color formed is inversely related to the amount of antibody to the HIV core or envelope antigen in the sample. Antibodies to envelope antigens are most consistently found in patients with HIV infection, whereas p24 antibody titers appear to decline as HIV-infected individuals clinically deteriorate (6, 18, 53, 75, 76). Declining p24 antibody titers therefore may have prognostic significance (125).

FIG. 1. Schematic of HIV virion and genome and an example of a strongly positive Western blot showing virus-specific bands. LTR, Long terminal repeat. Courtesy of Dupont Medical Products, Wilmington, Del.

Western Blot

The Western blot assay for HIV antibody is a technique in which partially purified whole virus is electrophoresed in a polyacrylamide gel, transferred onto nitrocellulose, and incubated with patient sera (117). HIV antibody is then detected by using an anti-human IgG antibody conjugated with an enzyme which will produce a colored band upon addition of substrate. Molecular weight standards and positive and negative control sera are run simultaneously, to allow identification of viral bands, which are usually read by the naked eye. This assay utilizes essentially the same principle as the EIA except that it has the advantage of identifying antibodies specific for different HIV viral antigens (Fig. 1). It appears that antibodies to the HIV core (*gag*) protein (p24)

and its precursor (p55) occur earliest in infection and tend to decrease or become undetectable with clinical deterioration (39, 53, 74, 76, 125). On the other hand, antibodies to the primary envelope (*env*) gene product (gp160) and its processed fragments (gp120) and the transmembrane protein (gp41) can be detected in nearly all HIV-infected patients regardless of clinical stage (6, 37, 67). Antibodies to the polymerase gene products (p31, p51, and p66) are also commonly detected. Other antibodies to the TAT_{III} protein (p14), the 3'-orf protein (p27), the *sor* protein (p23), and other *gag* proteins (p15, p9, p7) are detected less consistently (42).

Hence, the Western blot will yield a profile of bands characteristic of HIV when reacted with antibody-positive serum (Fig. 1). In addition, the Western blot in our laboratory is more sensitive than the EIA by a 2-log dilution of

serum. However, the disadvantages are that it is technically difficult, relatively expensive, and subjective in interpretation and requires approximately 24 h to perform. Partially purified HIV antigen is commercially available for those laboratories that want to prepare nitrocellulose strips for use in testing (E. I. du Pont de Nemours; Genetic Systems, Seattle, Wash.; Litton Bionetics, Charleston, S.C.). Premade nitrocellulose strips with separated HIV antigens are also commercially available (Bio-Rad Laboratories, Richmond, Calif.; E. I. de Pont de Nemours; Protatek International, St. Paul, Minn.). Recently, a commercially available Western blot kit with premade strips and all necessary reagents and controls has been licensed for use (E. I. du Pont de Nemours).

The definition of a "positive" Western blot is somewhat arbitrary and is by no means uniform throughout the United States. In our laboratory we use the following definition: negative, no bands characteristic for HIV are present; positive, an envelope antibody (gp41, gp120, or gp160) and at least one other HIV characteristic band. Bands characteristic for HIV are p15, p17, p24, p31, gp41, p51, p55, p66, gp120, and gp160. The CDC uses a similar definition except that the presence of the gp41 band alone can define the specimen as Western blot positive.

All other band patterns that do not fit into the above categories are considered indeterminate, and we repeat the tests in duplicate. If on retesting the results are still indeterminate, we recommend that the patient be retested 2 to 3 months later since reactivity to one or two core antigens (p17, p24, p55) exclusively may represent early infection with HIV (39, 53), cross-reactivity with another retrovirus (HTLV-I or HIV-2) (34), or a poorly understood cross-reaction commonly found in healthy people (11, 17, 36) and patients with HLA antibodies, bilirubinemia, connective tissue disease, and polyclonal gammopathies (product insert from Biotech/Du Pont HIV Western Blot Kit, Wilmington, Del.). The recently licensed Western blot kit requires that no bands be present for a blot to be interpreted as negative and that a band must be present at p24, p31, and either gp41 or gp160 to be interpreted as positive. Any pattern that does not meet these criteria is considered indeterminate. This strict criterion for a positive blot greatly increases the specificity of the assay, which is mainly intended for use in screening blood donors. However, this strict interpretation decreases the sensitivity of the assay from nearly 100% in AIDS patients to 56% since only 60% of AIDS patients have antibody to p24 and 83% have antibody to p31 (product insert from Biotech/Du Pont HIV Western Blot Kit). While this interpretation increases the number of supplemental tests, it greatly reduces the number of false positives in low-risk groups. Prior to this licensed test, false positives in low-risk groups have been estimated to be about 1 per 20,000 when used in conjunction with the EIA as a screening test (84).

The true rate of not detecting an HIV-infected individual (false-negative rate) with the Western blot is presently not known, but certainly is not zero. There have been reports of culture-positive Western blot-negative individuals (55, 105), and in one CDC study the Western blot was negative in 2 (8%) of 25 culture-positive EIA-positive blood donors (124). Presumably these people either had not developed antibody or had very low titers of antibody. Negative Western blots or blots showing reactivity only at p24 or p31 have also been reported in patients infected with HIV-2 (34). It may become necessary to use both HIV and HIV-2 antigens in the EIA and Western blot in areas where HIV-2 infection is signifi-

cant. It has been estimated that the false-negative rate in the blood donor population is approximately 1 in 250,000 donors when the Western blot is used in conjunction with the EIA (15). In high-risk groups, on the other hand, the false-negative rate is probably significantly higher. A reduction in false-negative and false-positive Western blots may be possible with the recent development of recombinant HIV antigens for use in immunoblotting due to the purity of these antigens.

IFA

The indirect IFA for HIV antibody is a fast and reliable supplemental test (13, 46, 64, 77, 92, 98, 106). Typically, uninfected and HIV-infected malignant T-cell lines (H9 or HUT 78) in log growth phase are applied to the wells of a fluorescence slide, air dried, fixed in acetone for 10 min, and then stored at -20°C or colder until use. To perform the assay, slides are warmed to room temperature, 10 μl of an appropriate serum dilution (1:10 or 1:20) is applied to each well, and slides are incubated in a humid chamber for 30 min, washed three times for 5 min in buffered saline, and air dried. Then 10 μl of anti-human fluorescein isothiocyanate-conjugated IgG is applied to each well. The slide is incubated for 30 min, washed three times for 5 min, counterstained with 0.001% Evans Blue, air dried, mounted, and read under a high-power ($\times 400$) fluorescence microscope. Known positive and negative control sera should be included in each assay. Specimens are interpreted as positive by the degree of fluorescence intensity, percentage of fluorescent cells, and the type of fluorescence pattern observed. There appears to be excellent agreement between IFA and Western blot. Sandstrom et al. in a comparison study found that all 68 Western blot-negative sera were also negative by IFA and only 1 of 36 sera positive in Western blotting was negative by IFA (106). Three additional samples from this patient were all negative by both tests. Gallo et al. found complete agreement between EIA and IFA on sera from 88 AIDS patients (46). However, only 18 (36%) of 50 blood donor specimens reactive by EIA were positive by IFA and Western blot. Specimens negative by IFA were negative by Western blot as well. It should be pointed out that 20 (5%) of 407 specimens in this study required absorption with uninfected H9 cells due to nonspecific staining and that only 11 (0.27%) of 4,000 gave nonspecific reactions that could not be removed by cell absorption. All 11 specimens were Western blot negative and 6 possessed antinuclear antibody.

The above results suggest that, compared with the Western blot, the IFA is equally sensitive and specific as a supplemental test. The IFA also has the advantage over Western blot of being performed in <2 hs; it also requires less technical expertise and can be run on either serum or plasma specimens. Moreover, since nearly all HIV-infected individuals have antibodies to the cell-associated proteins gp160 and gp120, the IFA will more readily detect these antibodies. gp160 and gp120 are expressed in infected cells, but are generally not present in purified preparations of virus used in EIA kits and in some Western blot procedures (37). However, interpretation of the IFA is more subjective, and uninfected cells may be required for serum absorption.

RIPA

The RIPA is primarily a research technique in which HIV-infected cells in their peak log phase of growth are

exposed to radioactive isotopes (e.g., [³⁵S]methionine) for a few hours, lysed with RIPA buffer, and then ultracentrifuged to obtain a soluble cell lysate containing radiolabeled viral proteins (2, 6–8, 38, 65, 67, 109). Cell lysates are cleared first with negative control serum bound to protein A-Sepharose beads and then reacted with patient serum preabsorbed with protein A-Sepharose beads. The immunoprecipitates are eluted in sample buffer by boiling at 100°C and then separated by electrophoresis on sodium dodecyl sulfate-polyacrylamide gels. The banding patterns are similar to those of Western blot except that the gp120 and gp160 proteins are more often apparent by RIPA, whereas gp41 and core proteins are detected more effectively by Western blot (32, 45). In addition, RIPA appears to result in fewer false positives in low-risk blood donors who have isolated bands to core proteins by Western blot (32, 45). Therefore, RIPA is probably slightly more sensitive and specific than Western blot and should be considered as a possible supplemental test for serum samples that are indeterminate by Western blot or IFA (32, 37). However, RIPA is mainly confined to the research laboratory because it is cumbersome and expensive and requires the maintenance of infected cell lines and handling of radioisotopes.

Future HIV Antibody Tests

Second generation assays for HIV antibody that use recombinant antigens or chemically synthesized peptides have already been developed and are awaiting Food and Drug Administration approval (16, 20, 51, 94, 108). In one study, an EIA that uses a single molecularly cloned expressed envelope polypeptide had a sensitivity of 99.9% and a specificity of 99.1% when compared with Western blot (16). The sensitivity and specificity of another EIA that uses a 12-amino acid gp41 peptide were reported to be 99.2 and 100%, respectively (51). A dot enzyme immunoassay, using a recombinant envelope protein (sensitivity, 98.2%) and taking about 30 min to perform, has been developed (20). A sensitive assay for the detection of HIV antibodies in saliva has also recently been reported (94). Of extreme practical importance is the recent development of a latex agglutination test, using recombinant envelope antigen. By this method less than 5 min is required to test serum for HIV antibody. Its reported sensitivity and specificity compared with Western blot are 100% each (104).

These diagnostic assays have the potential for greater specificity, are less costly and less hazardous to manufacture than whole virus preparations, and may take less time to perform. On the other hand, an assay using a single peptide, even with a highly conserved amino acid sequence, may fail to detect antibodies directed against other HIV antigens in the early phase of seroconversion or fail to detect antibodies directed against related retroviruses such as HIV-2 (50). Therefore, assays that use several synthetic or recombinant antigens from different HIV strains may be more appropriate. On the other hand, the use of individual synthetic peptides offers possibilities for the determination of type-specific antibodies to different HIV strains (50, 89).

Clinical Significance and Use of HIV Antibody Testing

HIV antibody testing is generally used for screening blood donors and for the diagnosis of HIV infection. Because the EIA is sensitive, specific, fast, and inexpensive, it is the screening test of choice. All repeatedly reactive specimens

should be retested with a supplemental test such as Western blot or IFA. A positive supplemental test is not diagnostic of AIDS or ARC, but only indicates past exposure to HIV. However, the individual is presumably infected and considered to be capable of transmitting HIV. It appears that roughly one-third of individuals with positive supplemental tests will develop AIDS within 7 years and another third will experience some symptoms (Hessol et al., Third International Conference on AIDS, 1987). Antibody-positive individuals should be counselled for the purpose of preventing further transmission (i.e., altering sexual behavior, refraining from blood or semen donations, warning others of the need to take appropriate precautions if there is potential contact with patient's body fluids, reconsidering pregnancy, and avoiding needle sharing).

Blood donors who are repeatedly reactive by EIA, but are negative by supplemental testing, are most likely not infected (19). However, for safety reasons blood from these donors should not be used for transfusion unless the donors are subsequently found to be negative by both EIA and a licensed Western blot on two separate occasions at least 6 months apart. In addition, each serum specimen should be negative by at least two EIA kits that differ in the cell line used as a source of viral antigen. Due to the very low probability that EIA-positive/Western blot-negative individuals in low-risk groups are infected, these individuals are generally not informed of a "false positive" result that could cause tremendous anxiety. On the other hand, some blood banks do inform regular donors because they find it less ethical to continue drawing units of blood which will be discarded. However, if a person that is EIA positive/Western blot-negative has been engaging in high-risk behavior or is symptomatic, retesting in 2 to 3 months for antibody or even a blood culture for HIV antigen or reverse transcriptase may be warranted.

Individuals who are repeatedly reactive by EIA and indeterminate on supplemental testing should also be retested in 2 to 3 months or tested by another supplemental assay or both. Generally, these people are informed of their indeterminate result as it may signify early infection, especially in people at high risk.

The question of who should be tested is very controversial. Mandatory testing for HIV antibody is currently required for blood donors, federal prisoners, military recruits, active-duty military personnel, foreign-service officers, and aliens seeking to immigrate to the United States. In addition, the Public Health Service recommendations concerning who should be counseled and tested for HIV antibody is reprinted here from *Morbidity and Mortality Weekly Reports* (29).

1. Persons who may have sexually transmitted disease. All persons seeking treatment for a sexually transmitted disease, in all health-care settings including the offices of private physicians, should be routinely* counseled and tested for HIV antibody. *"Routine counseling and testing" is defined as a policy to provide these services to all clients after informing them that testing will be done. Except where testing is required by law, individuals have the right to decline to be tested without being denied health care or other services.

2. IV-drug abusers. All persons seeking treatment for IV-drug abuse or having a history of IV-drug abuse should be routinely counseled and tested for HIV antibody. Medical professionals in all health-care settings, including prison clinics, should seek a history of IV-drug abuse from patients and should be aware of its

implications for HIV infection. In addition, state and local health policy makers should address the following issues:

- Treatment programs for IV-drug abusers should be sufficiently available to allow persons seeking assistance to enter promptly and be encouraged to alter the behavior that places them and others at risk for HIV infection.

- Outreach programs for IV-drug abusers should be undertaken to increase their knowledge of AIDS and of ways to prevent HIV infection, to encourage them to obtain counseling and testing for HIV antibody, and to persuade them to be treated for substance abuse.

3. Persons who consider themselves at risk. All persons who consider themselves at risk for HIV infection should be counseled and offered testing for HIV antibody.

4. Women of childbearing age. All women of childbearing age with identifiable risks for HIV infection should be routinely counseled and tested for HIV antibody, regardless of the health-care setting. Each encounter between a health-care provider and a woman at risk and/or her sexual partners is an opportunity to reach them with information and education about AIDS and prevention of HIV infection. Women are at risk for HIV infection if they:

- Have used IV drugs.
- Have engaged in prostitution.
- Have had sexual partners who are infected or are at risk for infection because they are bisexual or are IV-drug abusers or hemophiliacs.

- Are living in communities or were born in countries where there is a known or suspected high prevalence of infection among women.

- Received a transfusion before blood was being screened for HIV antibody but after HIV infection occurred in the United States (e.g., between 1978–1985 [26]). Educating and testing these women before they become pregnant allows them to avoid pregnancy and subsequent intrauterine perinatal infection of their infants (30–50% of the infants born to HIV-infected women will also be infected).

All pregnant women at risk for HIV infection should be routinely counseled and tested for HIV antibody. Identifying pregnant women with HIV infection as early in pregnancy as possible is important for ensuring appropriate medical care for these women; for planning medical care for their infants; and for providing counseling on family planning, future pregnancies, and the risk of sexual transmission of HIV to others.

All women who seek family planning services and who are at risk for HIV infection should be routinely counseled about AIDS and HIV infection and tested for HIV antibody. Decisions about the need for counseling and testing programs in a community should be based on the best available estimates of the prevalence of HIV infection and the demographic variables of infection.

5. Persons planning marriage. All persons considering marriage should be given information about AIDS, HIV infection, and the availability of counseling and testing for HIV antibody. Decisions about instituting routine or mandatory premarital testing for HIV antibody should take into account the prevalence of HIV infection in the area and/or population group as well as other factors and should be based upon the likely cost-effectiveness of such testing in preventing further spread of infection. Premarital testing in an area with a

prevalence of HIV infection as low as 0.1% may be justified if reaching an infected person through testing can prevent subsequent transmission to the spouse or prevent pregnancy in a woman who is infected.

6. Persons undergoing medical evaluation or treatment. Testing for HIV antibody is a useful diagnostic tool for evaluating patients with selected clinical signs and symptoms such as generalized lymphadenopathy; unexplained dementia; chronic, unexplained fever or diarrhea; unexplained weight loss; or diseases such as tuberculosis as well as sexually transmitted diseases, generalized herpes, and chronic candidiasis.

Since persons infected with both HIV and the tubercle bacillus are at high risk for severe clinical tuberculosis, all patients with tuberculosis should be routinely counseled and tested for HIV antibody. Guidelines for managing patients with both HIV and tuberculous infection have been published.

The risk of HIV infection from transfusions of blood or blood components from 1978–1985 was greatest for persons receiving large numbers of units of blood collected from areas with high incidences of AIDS. Persons who have this increased risk should be counseled about the potential risk of HIV infection and should be offered antibody testing.

7. Persons admitted to hospitals. Hospitals, in conjunction with state and local health departments, should periodically determine the prevalence of HIV infections in the age groups at highest risk for infection. Consideration should be given to routine testing in those age groups deemed to have a high prevalence of HIV infection.

8. Persons in correctional systems. Correctional systems should study the best means of implementing programs for counseling inmates about HIV infection and for testing them for such infection at admission and discharge from the system. In particular, they should examine the usefulness of these programs in preventing further transmission of HIV infection and the impact of the testing programs on both the inmates and the correctional system. Federal prisons have been instructed to test all prisoners when they enter and leave the prison system.

9. Prostitutes. Male and female prostitutes should be counseled and tested and made aware of the risks of HIV infection to themselves and others. Particularly prostitutes who are HIV-antibody positive should be instructed to discontinue the practice of prostitution. Local or state jurisdictions should adopt procedures to assure that these instructions are followed.

In addition, hemophiliac patients who have received factor concentrates since 1977 and males who have had intercourse with another male since 1977 should be routinely considered for HIV antibody testing and for counseling.

We should emphasize that reporting of HIV test results should comply with state laws and HIV test results should be kept strictly confidential. In addition, individuals with confirmed HIV-positive test results should be provided counseling and medical referral.

HIV ANTIGEN DETECTION

In 1986, commercial tests (Abbott Laboratories; E. I. du Pont de Nemours) became available for the detection of HIV antigen in serum, plasma, and cerebrospinal fluid (CSF), using EIA technology (52). The principle of HIV antigen

detection in these kits involves incubating the patient's specimen overnight with a polystyrene bead or in a plastic microwell coated with polyclonal anti-HIV antibody (Fig. 2). The bead or microwell is then rinsed and incubated with a rabbit or goat antibody to HIV antigen to form a sandwich. After the bead or microwell is washed, an enzyme-conjugated anti-goat or anti-rabbit antibody can then be added to increase the test sensitivity and specificity. After a final wash, the appropriate substrate can be added. If the antigen-antibody-enzyme complex is present, a color change will be detected by an increase in absorbance at a specific wavelength. Dilutions of known amounts of HIV antigen can be used to generate a standard curve for quantitating the amount of free HIV antigen in the patient's specimen. The detectable limit for this assay is between 50 and 100 pg/ml (52, 53).

If a specimen is found to be repeatedly reactive, a specific antibody neutralization assay should be performed to confirm the presence of HIV antigen. This assay is also commercially available (Abbott Laboratories) and involves preincubating the specimen or control with human antibody to HIV before running the HIV antigen EIA. If HIV antigen is present, the neutralizing antibody will form an immune complex which prevents binding of the HIV antigen to the HIV antibody-coated bead or microwell. This results in a reduction of absorbance when compared with the specimen in which human serum nonreactive for antibody to HIV is used in place of neutralizing antibody. A specimen is considered to be a confirmed positive for HIV antigen if the reduction in absorbance of the neutralized specimen when compared with a matched non-neutralized control is 50% or greater and if the non-neutralized control absorbance is greater than or equal to the calculated cutoff.

Serum, plasma, CSF, and cell culture supernatant have been shown to be suitable specimens for HIV antigen detection by EIA (52). However, testing of lipemic or hemolyzed specimens should be avoided. If specimens are to be stored, they may be refrigerated at 2 to 8°C for as long as 1 week or also frozen. Multiple freeze-thaw cycles of the specimen should be avoided.

The assay requires an overnight incubation step and generally takes 24 to 30 h for the entire procedure. If a test is repeated and then the specimen is neutralized for confirmation, it may take 3 to 4 days to generate a positive test result. The cost of reagents for research laboratories is approximately \$4.00 to \$8.00 per test, so obtaining a confirmed positive test result may cost at least \$20.00 for reagents alone.

While the primary antibody used to capture the HIV

FIG. 2. Diagram of HIV antigen EIA design. Courtesy of Abbott Laboratories, North Chicago, Ill. HPRO, Horseradish peroxidase; OPD, o-phenylenediamine.

antigen is polyclonal, the specific antibody for the p24/55 core proteins appears to have 10,000 times more affinity than the antibodies for gp120/41 envelope proteins (53). Consequently, the current antigen EIA kits primarily detect free p24 antigen. It is important to emphasize that these tests appear to detect free p24 antigen. In patients who have a high titer of p24 antibody, p24 antigen appears to be complexed with antibody and is difficult to detect. This concept is supported by a study in which 35 (70%) of 50 Western blot-positive patients with absent p24 antibody levels had detectable HIV antigen versus 3 (8%) of 38 patients with detectable levels of p24 antibody (75). Moreover, patients who are initially antigen positive after exposure often become antigen negative when they seroconvert and develop p24 antibody (1). Likewise, it is common for patients to become antigen positive as their p24 antibody falls during progression from the asymptomatic state to AIDS (66). It is not clear whether this increased positivity is due to greater antigen production as the disease progresses or solely due to falling antibody titers or both. A recent study suggests that the production of detectable HIV antigen reflects enhanced expression of viral genes and that loss of p24 antibody is secondary (75).

The ability to detect directly the presence of HIV in serum or CSF is a significant improvement over antibody testing in several respects. First, the direct detection of HIV antigen indicates active HIV infection as opposed to past exposure to HIV proteins. Second, serum antigen detection allows for the diagnosis of HIV infection prior to antibody seroconversion, which may take longer than 1 year from time of exposure to occur (3). Third, the detection of HIV antigen in serum has prognostic significance. Among HIV antigen-positive, asymptomatic, antibody-positive homosexual males, 23.9% progressed to AIDS over a 21-month period compared with 1.3% of HIV antigen-negative, asymptomatic, antibody-positive homosexual males (F. DeWolf, J. Goudsmit, D. A. Paul, J. M. A. Lange, C. Hooykaas, and R. A. Coutinho, Third International Conference on AIDS, p. 19, 1987).

However, the sensitivity of the HIV serum antigen assay is poor if one assumes that most if not all HIV antibody-positive individuals are infected. The sensitivity of the test improves with clinical progression of disease, showing 4% positivity in asymptomatic patients, 56% in ARC patients, and 70% in AIDS patients (66). Likewise, the CSF of five of nine adults with AIDS-related encephalopathy and five of five children with AIDS had detectable HIV antigen (52). However, none was detected in the CSF of 13 symptomless HIV antibody-positive controls. The specificity is much better than the sensitivity. In testing serum of blood donors, 9 of 2,096 (0.43%) were repeatedly reactive but none of their specimens could be neutralized, indicating false-positive reactions (115).

While the HIV antigen EIA is not an appropriate screening test due to its poor sensitivity, there are several clinical situations in which it may be useful.

(i) *Individuals in high-risk groups prior to antibody seroconversion.* It is not known what percentage of individuals will express HIV antigen in serum during the interval between exposure and seroconversion. However, at least eight individuals have been reported to be antigen positive 4 to 16 weeks before HIV antibody appeared (1, 52). HIV antigenemia has also been temporally linked to acute HIV infection before the appearance of HIV antibody (123). This test may be especially helpful if positive in an HIV antibody-negative pregnant woman in a high-risk group for HIV infection.

(ii) *HIV antibody-positive asymptomatic individuals for prognostic purposes.* As noted above, asymptomatic individuals who were serum antigen positive were 18 times more likely to progress to AIDS over a 21-month period (DeWolf et al., Third International Conference on AIDS, 1987). Such patients could be selected for clinical trials, using drugs active against HIV such as zidovudine (formerly known as azidothymidine).

(iii) *HIV-infected patients receiving antiviral drug therapy.* Mean HIV antigen levels have been shown to decrease significantly over time in patients treated with zidovudine compared with placebo-treated controls (31). Monitoring of HIV antigen levels may be useful in measuring the antiviral effect of drug therapy.

(iv) *HIV antibody-positive infants.* HIV antibody may be passively transferred in utero from HIV-infected mothers without HIV transmission to the infant. A positive HIV antigen test on the infant's serum is useful in confirming HIV infection (86).

(v) *Differentiating AIDS from primary immunodeficiency syndromes in HIV antibody-negative children.* Since antibody response may be blunted or absent in children with congenital immunodeficiency syndromes, the diagnosis of AIDS may be missed if only HIV antibody testing is performed (14). HIV antibody-negative children with an immunodeficiency syndrome should be checked for HIV antigen, especially if they received a transfusion prior to antibody testing or the mother is HIV antibody positive.

(vi) *Individuals with indeterminate supplemental antibody tests.* Most healthy individuals in low-risk groups with indeterminate Western blots are probably not infected (120), but a negative antigen test may provide some reassurance, especially if the Western blot or RIPA indicates the absence of antibodies to p24/55.

HIV CULTURE

In 1983, HIV was first propagated by using phytohemagglutinin (PHA) and T-cell growth factor in cocultures of lymph node cells from a homosexual with lymphadenopathy and T lymphocytes from a healthy adult donor and from umbilical cord blood of newborn humans (7). The presence of HIV was detected by electron microscopy, showing lymphocytes with characteristic immature particles with dense (C-type) budding at the plasma membrane, and high levels of reverse transcriptase (RT) activity in the culture supernatant fluid on day 15 of coculture.

Subsequently, Gallo et al. confirmed these findings and developed a cell culture system, using clones (H4, H9) of a neoplastic T-cell line (HT) for the continuous production of HIV (47, 98). The presence of HIV was further confirmed by serological analysis (107, 109) and by IFA, using fixed cells of HT-infected clones and rabbit antiserum to HIV or patient serum (98).

Levy and Shimabukuro reported the recovery of HIV by RT assay to be about 68% (53 of 78) in AIDS patients, 41% (7 of 17) in HIV antibody-positive asymptomatic homosexual males, and 0% (0 of 85) in antibody-negative healthy heterosexuals, using cocultures of patient peripheral mononuclear cells (PMC) and PHA-stimulated human PMC from clinically healthy individuals (78). In 64% of the positive cultures, RT activity was detected within 9 to 15 days from initiation of culture and in an additional 21% after 21 days. From a diagnostic viewpoint these results were rather discouraging, because the sensitivity of culture is relatively

poor, the time to positivity is relatively long, the RT assay is tedious, and the cost of reagents and labor is several hundred dollars. Moreover, RT activity is not specific for HIV and may be produced by other retroviruses such as human T-cell lymphotropic virus type I (HTLV-I) (97). Fortunately, refinements in HIV cell culture techniques and the development of sensitive HIV antigen detection EIAs in the last year have dramatically improved the recovery rate of HIV and decreased the time and cost of HIV culture. In our laboratory, we use the following coculture technique to recover HIV from the blood of infected patients.

(i) Patient PMC are separated from 30 ml of heparinized blood on a Ficoll-Paque (Pharmacia, Piscataway, N.J.) gradient within 8 h of collection.

(ii) A total of 10^7 PMC are placed in a 25-cm² tissue culture flask containing a total of 15 ml of RPMI 1640 medium, 20% fetal calf serum, 5% interleukin-2, 5 µg of Polybrene per ml, 200 U of penicillin per ml, 200 µg of streptomycin per ml, and 5×10^6 PHA-stimulated peripheral mononuclear cells from an HIV antibody-negative blood donor. Donor PMC are PHA stimulated (4 µg/ml) for 2 to 4 days in the above medium without interleukin-2 before addition to patient PMC.

(iii) Cocultures are incubated at 37°C in a 5% CO₂ atmosphere for up to 28 days.

(iv) Approximately 8 ml of culture medium above the settled PMC is removed twice a week for testing and replaced with an equal volume of fresh medium. An additional 3×10^6 PHA-stimulated donor PMC are added once a week.

Using this cell culture technique and one of the commercial HIV antigen EIAs (Abbott Laboratories) described above, we have been able to detect HIV antigen in the culture supernatants of cocultured PMC in 33 of 33 (100%) AIDS patients, 32 of 33 (97%) HIV antibody-positive asymptomatic homosexual males, 14 of 14 (100%) HIV antibody-positive symptomatic hemophiliacs, and in 0 of 18 HIV antibody-negative hemophiliacs. In terms of time to positivity, HIV antigen was detected in 46% of positive culture supernatant fluids by day 3 of culture, in 94% by day 10, and in 100% by day 21. We are now testing these patients' sera for HIV antigen.

In our laboratory, it appears that HIV can replicate more quickly and consistently in mitogen-stimulated donor PMC than in cell lines. However, it is necessary to refeed the culture every week or two with freshly stimulated PMC to sustain HIV replication.

In addition, the HIV antigen EIA has the following advantages over the RT assay for detection of HIV in culture. (i) Radioactive isotopes and a scintillation counter are not required. (ii) Pelletting of supernatant with an ultracentrifuge or polyethylene glycol is not required. (iii) Less sample volume (200 µl versus 1 to 3 ml) is required. (iv) Less technical hands-on time is required (4 versus 5 h). (v) Five to ten times the number of samples can be tested per run. (vi) The cost of HIV antigen EIA supplies per patient assay is approximately one-fifth that of the RT assay. (vii) The HIV antigen EIA is as sensitive and probably more sensitive than RT in detecting HIV (95) (viii) The RT assay is much more difficult to standardize. Consequently, many laboratories have discontinued it.

However, the RT assay has the theoretical advantage of detecting other HIV strains (HIV-2) or other retroviruses that the current HIV antigen EIAs may miss (33, 97). There are a number of published RT assay procedures for those laboratories interested in detecting RT activity in concen-

trated culture supernatant fluid (5, 63, 78, 97, 103). The principle of most of these RT assays is as follows.

RT is an enzyme unique to a group of viruses known as retroviruses, of which the HIV is a member. RT is a ribonucleic acid (RNA)-dependent DNA polymerase encoded for by the *pol* gene and requires Mg^{2+} . In the RT assay, an RNA template primer, polyriboadenylate: oligodeoxythymidylate₁₂₋₁₈ (a long adenine strand as the template and a short thymidine strand as the primer), is mixed with radioactive tritium, [3H]thymidine, a buffer containing Mg^{2+} , dithiothreitol as a reductant, Triton X-100 as a detergent to disrupt whole virus particles, and a sample of concentrated culture supernatant. This mixture is incubated at 37°C for 1 h. If RT is present in the sample, the radioactive thymidine will be incorporated onto the template. Cold acid is added to stop the reaction and precipitate the strands. After an incubation on ice, the reaction mixture is poured over a filter to trap the strands and washed to remove unbound [3H]thymidine. The filters are put into scintillation vials, dried, and counted in scintillation cocktail. Activity (in counts per minute) over a certain limit is indicative of HIV presence in the sample. To detect contaminating cellular DNA polymerases which could result in a false positive, the RT assay is run simultaneously, using a DNA template primer, polydeoxyadenylate: oligodeoxythymidylate₁₂₋₁₈, instead of the RNA template primer.

The clinical utility of isolating HIV from the peripheral mononuclear cells of different patient groups is similar to that of the serum HIV antigen assay (see above). However, culture is more sensitive. In addition, one study reported a strong association between clinical status and the rate of replication of HIV isolates *in vitro*, suggesting that culture may be useful for determining strain virulence (5). Moreover, time to positivity in culture may correlate with the patient's viral load since it has been shown to increase from 7 to 22 days with a 10^{-4} dilution of stock virus (102). Whether viral load is related to clinical progression of disease is not yet known.

HIV culture may also be useful in monitoring drug therapy. The authors of one study reported a significant drop in their ability to isolate HIV from mononuclear cells in patients receiving an antiviral drug called ampligen (21).

The drawbacks to HIV culture are that it is expensive, it may take weeks, and it creates the potential for exposure of personnel to high concentrations of virus.

HIV can also be cultured from other body fluids, including CSF (59), serum (85), cervical secretions (121), saliva (56, 58), semen (61, 127), plasma (40, 44), tears (43), breast milk (116), and brain tissue (79). However, the percentage of HIV recovery is less from these fluids: 73% in CSF from patients with AIDS-related neurologic syndromes (59), 4% in saliva of AIDS patients (58), and 70% in the plasma of AIDS patients (40). The clinical utility in culturing these fluids is not clear, although one study suggests that recovery of HIV from CSF in patients with neurologic disorders indicates HIV as the probable cause since HIV was not recovered from CSF of AIDS patients who did not have neurologic problems (59). Another study suggests that virus isolation from plasma is more reliable than antigen detection for the demonstration of HIV in primary infection prior to antibody seroconversion (44).

HYBRIDIZATION TECHNIQUES

An alternative diagnostic detection system for HIV is that of molecular hybridization. When an HIV virion attaches to

the cell surface, two messenger RNA transcripts along with the RNA-dependent DNA polymerase RT enter the cell (48). The RT synthesizes unintegrated linear DNA duplexes, using the HIV RNA as a template. Some of these linear DNA molecules circularize, migrate to the nucleus, and integrate randomly into the host genome, becoming the 9.7-kilobase proviral form of HIV (111). The proviral DNA may then be transcribed into RNA for the genetic material of new virions or for translation into HIV proteins.

The HIV genome apparently does not contain sequences homologous to normal human DNA (111) or to HTLV-I or HTLV-II proviral DNA to any appreciable extent (122). Therefore, one can probe infected cells by using labeled HIV DNA or RNA sequences to detect HIV nucleic acids by one or more hybridization techniques.

However, the sensitivity of these techniques is poor if one attempts to detect HIV directly in patient cells or tissue. Using Southern hybridization, Shaw et al. detected HIV DNA sequences in the PMC of 1 of 15 AIDS patients and 0 of 7 ARC patients (111). In lymph node cells, 4 of 16 AIDS patients and 3 of 18 ARC patients were positive. None of five Kaposi sarcoma tissue specimens were positive. These data indicate that there is less than one HIV DNA copy per 10 cells (the detection limit of their technique). The data from Harper et al., using *in situ* hybridization directly on PMC and lymph node cells, support this conclusion. They were able to detect viral DNA in 6 of 7 (86%) lymph node and 7 of 14 (50%) peripheral blood samples from AIDS/ARC patients, using a ^{35}S -labeled HIV-specific RNA probe after a 2-day exposure to autoradiographic emulsion (57). The number of labeled cells in these preparations was very small, constituting <0.01% (<1 in 10,000), and in some cases <0.001% (<1 in 100,000). Cells treated with ribonuclease before hybridization exhibited essentially no label, indicating that the grains had been specific for viral RNA and not integrated HIV DNA. Infected lymphocytes appeared to contain 20 to 300 copies of RNA per cell. Koenig et al. showed that *in situ* hybridization could also be used to detect HIV RNA sequences in mononucleated and multinucleated macrophages in brain sections of demented patients, although the diagnosis was more easily made by the positive HIV serology (69).

To increase the sensitivity of HIV detection, Monroe et al. amplified the virus first in cell cocultures. They then used a cytoplasmic dot-blot technique to detect HIV RNA sequences, using a ^{32}P -labeled 9-kilobase proviral DNA fragment (87). In half of the cocultivations that resulted in virus isolation, the RNA dot-blot was positive several days before any RT activity in the supernatant was detected. However, the RNA dot-blot was positive in only 20% of positive samples by day 8 of culture, 50% by day 12, 73% by day 15, and 100% by day 19. This time to positivity was poor compared with that of the HIV antigen EIA detection system described earlier despite the fact that the RNA dot-blot could detect one infected cell per 10^3 when blotting 5×10^6 cells.

Recently, an *in vitro* DNA amplification procedure plus an oligonucleotide-based detection procedure have been developed that might obviate the need for culture (73). This nucleic acid amplification procedure, or polymerase chain reaction, uses two primers of known sequence, positioned 10 to 300 base pairs apart, that are complementary to the plus and minus strands of highly conserved regions of target HIV DNA. After denaturation of the target DNA and annealing of the primers, template-directed incorporation of deoxynucleoside triphosphates occurs when a heat-stable DNA polymerase is added. The newly synthesized DNA strands may

then serve as templates themselves. Repeated cycles of denaturation, primer annealing, and extension result in a several hundred thousandfold increase of the copies of the region flanked by the primers. These amplified sequences are then detected by an oligomer restriction technique in which an end-labeled oligonucleotide probe hybridizes in solution to a region of the amplified sequence. This newly formed duplex forms a specific endonuclease cleavage site which yields an oligonucleotide of defined size when the specific restriction enzyme for this site is added. Using this technique, Kwok et al. were able to generate a diagnostic oligomer in 17 (89%) of 19 DNA samples from RT-positive cell cultures of seropositive AIDS or ARC patients (73). In comparison, only 11 (61%) of 18 samples were positive by Southern blot analysis, using a nick-translated probe. Whether this technique will be able to amplify and detect HIV with similar sensitivity in patient samples without culture remains to be seen. It is possible that the number of HIV DNA genomes in some individuals is too low to amplify or the genome may have sequence alterations in the region complementary to the primers and probes which prevent hybridization. On the other hand, this polymerase chain reaction-oligomer restriction procedure, which can be performed in a day, has the potential to detect directly latent HIV infection in individuals who lack serological markers.

In summary, molecular hybridization methods can be used to detect HIV if viral RNA is being actively synthesized or if the HIV proviral DNA can be greatly amplified through the use of a polymerase chain reaction procedure. However, these methods are technically difficult and their sensitivity and specificity in a large number of patient tests are not presently known.

BIOHAZARDS OF HIV TESTING

HIV has been isolated from a number of body fluids and tissues as mentioned previously. HIV has also been recovered from a dried preparation of H9-infected cells after 24 h and from an air-dried preparation of plasma containing $7 \log_{10}$ 50% tissue culture infectious doses of HIV after 3 days (102). However, data defining the risk of HIV infection to health care workers from exposure to HIV-infected body fluids or cultures, while preliminary, are reassuring.

Transmission of HIV infection by accidental inoculation with a needle contaminated with blood from an HIV antibody-positive patient has been documented by seroconversion in at least four health care workers with no other risk factors for HIV infection (4, 88, 90, 114). More recently, three instances of health care workers who appeared to acquire HIV infections from skin or mucous membrane exposure to blood from HIV-infected patients have been reported (27). Still, the risk of HIV transmission from needlesticks or mucous membrane exposure to HIV-infected blood appears to be low. In a CDC cooperative surveillance project, none of 82 health care workers who had open wounds or mucous membranes exposed to HIV-infected blood have seroconverted, and only 1 of 298 (0.3%) health care workers seroconverted after needlestick injuries or cuts with sharp objects contaminated with HIV-infected blood (82). In contrast, the risk of acquiring hepatitis B virus infection following a needlestick from a hepatitis B virus carrier ranges from 6 to 30% (54, 110).

In 1985, two different reagent production laboratories reported that several laboratory workers may have been inadvertently exposed to an aerosol of concentrated HIV; one worker was cut by a piece of glass from a broken carboy

that contained HIV-infected cells and culture fluid. None of the potentially exposed persons had shown evidence of seroconversion after 6 months in one incident and 12 months in the other as a result of these occupational exposures (24). A technologist in our laboratory who has handled over 500 positive cultures in the last 2 years has not seroconverted nor have the other four technologists and four nurses in our AIDS Treatment Evaluation Unit who voluntarily agree to be tested every 3 months for HIV antibody. However, officials at the National Institutes of Health recently reported that a person who worked in an HIV propagation laboratory became HIV antibody positive. This individual apparently had none of the known risk factors for contracting HIV, and accidental exposure was not reported (91).

To minimize the risk of HIV transmission in the laboratory the following recommended precautions printed here from *Morbidity and Mortality Weekly Report* should be adhered to (24).

1. Biosafety Level (BSL) 2 standards and special practices, containment equipment, and facilities as described in the CDC-NIH biosafety manual are recommended for activities involving clinical specimens, body fluids, or tissues from humans or laboratory animals that may contain HIV. These are the same practices recommended for all clinical specimens. Emphasis is placed on the following practices, which are included in the manual:

- a. Use of syringes, needles and other sharp instruments should be avoided if possible. Used needles and cutting instruments should be discarded into a puncture-resistant container with a lid. Needles should not be resheathed, purposely bent, broken, removed from disposable syringes, or otherwise manipulated by hand.

- b. Gloves should be worn by all personnel engaged in activities that may involve skin contact with potentially infectious fluids, tissues, or cultures and by laboratory workers with dermatitis or other lesions on the hands who may have direct or indirect contact with potentially infectious materials. Handwashing with soap and water should be a routine practice immediately after direct contact with potentially infectious materials and on completion of work, even when gloves are worn.

- c. Generation of aerosols, splashes, and spills of potentially infectious materials should be avoided in procedures involving body fluids or tissues, during necropsy of cadavers, and in similar procedures on animals experimentally infected with HIV. Laboratory workers should use a biological safety cabinet when propagating the virus to further reduce the risk of exposure. Although the major precautions are listed here, the CDC-NIH biosafety manual contains additional related precautions. In all instances, the laboratory director is responsible for assessing the biosafety level to be used.

- d. Human serum from any source that is used as a control or reagent in a test procedure should be handled at BSL 2. Appended to this Agent Summary Statement is a statement (Addendum 1) issued by CDC on the use of all human control or reagent sera shipped to other laboratories. The Food and Drug Administration requires that manufacturers of human serum reagents use a similarly worded statement.

- e. Animal BSL 2 practices, containment equipment, and facilities are recommended for activities involving nonhuman primates experimentally infected with HIV.

Laboratory coats, gowns, or uniforms should be worn by laboratory workers, as is customary for other BSL 2 or 3 practices, depending on the nature of the work, concentration of the virus, and volume of material being handled. Because many animals bite, and some throw feces, urine, or expectorate at humans, animal-care personnel must wear coats, protective gloves, coveralls or uniforms, and face shields as appropriate to protect the skin and mucous membranes of the eyes, nose, and mouth from potential exposure to these substances when working with animals likely to manifest such behavior.

2. Activities such as growing research-laboratory-scale amounts of HIV or related viruses or virus-producing cell lines, working with concentrated virus preparations, or conducting procedures that may produce droplets or aerosols should be performed in a BSL 2 facility with the additional practices and containment equipment recommended for BSL 3.

3. Activities involving industrial-scale, large-volume, or high-concentration production and manipulation of HIV are to be conducted with BSL 3 requirements.

4. All laboratory glassware, equipment, disposable materials, and wastes suspected or known to contain HIV must be decontaminated, preferably in an autoclave, before washing, discarding, etc. Incineration of solid wastes may be used as an alternative method of disposal.

5. There is no evidence that laboratory clothing soiled with materials known or suspected to contain HIV poses a transmission hazard, and the handling of such clothing is covered under BSL 2 practices. However, to be consistent with BSL 3 recommendations, when laboratory clothing becomes contaminated with HIV preparations, it should be decontaminated before being laundered or discarded.

6. Work surfaces should be decontaminated at the end of each day on completion of procedures or when overtly contaminated. Many commonly used chemical disinfectants with such active ingredients as sodium hypochlorite, formaldehyde, glutaraldehyde, or phenols, can be used to decontaminate laboratory work surfaces; they can also be used to decontaminate some laboratory instruments, specific areas of contaminated laboratory clothing, and spills of infectious materials. Prompt decontamination of spills and other overt contamination should be standard practice.

7. The prudent and recommended approach to handling human serum known or suspected to contain HIV is to use the same precautions that should be used routinely to prevent transmission of blood borne infections, including hepatitis B. Available data on the effectiveness of heat to destroy HIV suspected or known to be present in human serum are at variance because of variations in volume of serum, concentration of the virus, temperature, and duration of exposure to heat. Similarly, results of chemical analyses or antibody assays may vary when sera are heated before testing according to the analysis or assay being performed. However, there is agreement that testing heated serum for HIV antibody by enzyme immunoassays often yields false-positive results.

8. No HIV vaccine has been developed, and no drugs have been shown to be completely safe and effective for therapy. As part of an ongoing medical surveillance program for employees, all laboratory workers before

being assigned to activities with a high potential for exposure should have a serum sample obtained and stored at -40°C (-40°F) for possible future testing. Subsequent serum samples should be obtained and stored in accordance with laboratory policy or following an inadvertent laboratory exposure involving materials described above. When indicated, these serum specimens should be tested by a qualified laboratory using currently recommended procedures for HIV antibody. Furthermore, the physician requesting serologic testing of these serum specimens must first obtain informed consent from the laboratory worker and describe the confidentiality safeguards available to protect test results. The laboratory workers whose serum specimens are to be tested should understand how the test results are to be used, the implications of a positive or negative test result, and the limits, if any, of the confidentiality safeguards. An employee whose serum HIV antibody test is reactive and whose subsequent test and evaluation confirm the presence of HIV infection should be counseled to follow the Public Health Service recommendations for preventing transmission.

9. In addition to HIV, other primary, as well as opportunistic, pathogenic agents may be present in the body fluids and tissues of persons who are antibody positive or have AIDS-related complex or AIDS. Laboratory workers should follow accepted bio-safety practices to ensure maximum protection against inadvertent laboratory infection with agents other than HIV that may also be present in clinical specimens.

Addendum

CDC cautionary notice for all human serum samples used as controls or reagents:

Warning:

Because no test method can offer complete assurance that laboratory specimens do not contain HIV, hepatitis B virus, or other infectious agents, this specimen(s) should be handled at the BSL 2 as recommended for any potentially infectious human serum or blood specimen in the CDC-NIH manual, Biosafety in Microbiological and Biomedical Laboratories, 1984, pages 11-13.

One or more of the following statements should be included with the above warning statement:

This specimen is negative for antibody to HIV.

This specimen is positive for hepatitis B surface antigen (HBsAg).

This specimen is positive for antibody for HIV.

This specimen has NOT been tested for hepatitis B surface antigen (HBsAg).

This specimen has NOT been tested for antibody to HIV.

This specimen has been heated at 56°C (133°F) for 30 minutes (which will not inactivate HBsAg but will inactivate HTLV-III/LAV).

CONCLUSION

The demand for HIV testing will continue to grow. As the AIDS epidemic spreads, the availability of fast sensitive/specific diagnostic and prognostic tests will be required. EIAs for screening and Western blot assays and IFA for confirmation are sensitive and specific and are now commercially

available for most laboratories. Other confirmation assays, using RIPA or culture techniques, will be in greater demand as the number of "indeterminate" results increases. While these sophisticated assays may be suitable only for research or reference laboratories, it is important that virologists and other laboratorians involved in HIV testing understand the principles and limitations of these assays as well as the associated biohazards and appropriate safety measures.

ACKNOWLEDGMENTS

This work was supported by Public Health Service contract NO1-AI-72626 from the National Institutes of Health and by a grant from the Minnesota Medical Foundation.

LITERATURE CITED

- Allain, J., Y. Laurian, D. A. Paul, and D. Senn. 1986. Serological markers in early stages of human immunodeficiency virus infection in hemophiliacs. *Lancet* ii:1233-1236.
- Allan, J. S., J. E. Coligan, F. Barin, J. G. Sodroski, and T. H. Lee. 1985. Major glycoprotein antigens that induce antibodies in AIDS patients are encoded by HTLV-III. *Science* 228:1091-1094.
- Anderson, K. C., B. C. Gorgone, R. G. Marlink, R. Ferriani, M. E. Essex, P. M. Benz, and J. E. Groopman. 1986. Transfusion-acquired human immunodeficiency virus infection among immunocompromised persons. *Ann. Intern. Med.* 105:519-527.
- Anonymous. 1984. Needlestick transmission of HTLV-III from a patient infected in Africa. *Lancet* ii:1276-1277.
- Asjo, B., L. Morfeldt-Manson, J. Albert, G. Biberfeld, A. Karlsson, K. Lidman, and E. M. Fenyo. 1986. Replicative capacity of human immunodeficiency virus from patients with varying severity of HIV infection. *Lancet* ii:660-662.
- Barin, F., M. F. McLane, J. S. Allan, and T. H. Lee. 1985. Virus envelope protein of HTLV-III represents major target antigen for antibodies in AIDS patients. *Science* 228:1094-1096.
- Barré-Sinoussi, F., J. C. Chermann, F. Rey, M. T. Nugeyre, S. Chamaret, J. Gruest, C. Dautet, C. Axler-Blin, F. Vezinet-Brun, C. Rouzioux, W. Rosenbaum, and L. Montagnier. 1983. Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 220:868-871.
- Barré-Sinoussi, F., U. Mathur-Wagh, F. Rey, F. Brun-Vezinet, S. R. Yancovitz, C. Rouzioux, L. Montagnier, D. Mildvan, and J. C. Chermann. 1985. Isolation of lymphadenopathy-associated virus (LAV) and detection of LAV antibodies from US patients with AIDS. *J. Am. Med. Assoc.* 253:1737-1739.
- Bedarida, G., G. Cambie, F. D'Agostino, M. G. Ronsivale, E. Berto, M. E. Grisi, P. R. Crocchiolo, and E. Magni. 1986. Anti-IgM screening for HIV. *Lancet* ii:1456.
- Bedarida, G., G. Cambie, F. D'Agostino, J. B. Ronsivale, E. Berto, M. E. Grisi, and E. Magni. 1986. HIV IgM antibodies in risk groups who are seronegative on ELISA testing. *Lancet* ii:570-571.
- Biberfeld, G., U. Bredberg-Raden, B. Bottiger, P. Putkonen, J. Blomberg, P. Juto, and G. Wadell. 1986. Blood donor sera with false-positive Western blot reactions to human immunodeficiency virus. *Lancet* ii:289-290.
- Blanton, M., K. Balakrishnan, U. Dumaswala, K. Zelenski, and T. J. Greenwalt. 1987. HLA antibodies in blood donors with reactive screening tests for antibody to the immunodeficiency virus. *Transfusion* 27:118-119.
- Blumberg, R. S., E. G. Sandstrom, T. J. Paradis, D. N. Nuemeyer, M. G. Sarngadharan, K. L. Hartshorn, R. E. Byington, M. S. Hirsch, and R. T. Schooley. 1986. Detection of human T-cell lymphotropic virus type III-related antigens and anti-human T-cell lymphotropic virus type III antibodies by anticomplementary immunofluorescence. *J. Clin. Microbiol.* 23:1072-1077.
- Borkowsky, W., K. Krasinski, D. Paul, T. Moore, D. Bebenroth, and S. Chandwani. 1987. Human-immunodeficiency-virus infections in infants negative for anti-HIV by enzyme-linked immunoassay. *Lancet* i:1168-1171.
- Bove, J. R. 1987. Transfusion-associated hepatitis and AIDS. *N. Engl. J. Med.* 317:242-245.
- Burke, D. S., B. L. Brandt, R. R. Redfield, T. Lee, R. M. Thorn, G. A. Beltz, and C. Hung. 1987. Diagnosis of human immunodeficiency virus infection by immunoassay using a molecularly cloned and expressed virus envelope polypeptide. *Ann. Intern. Med.* 106:671-676.
- Burke, D. S., and R. R. Redfield. 1986. False-positive Western blot tests for antibodies to HTLV-III. *J. Am. Med. Assoc.* 256:347.
- Burke, D. S., R. R. Redfield, P. Putman, and S. S. Alexander. 1987. Variations in Western blot banding patterns of human T-cell lymphotropic virus type III/lymphadenopathy-associated virus. *J. Clin. Microbiol.* 25:81-84.
- Cable, R. G., R. M. Kakaiya, S. C. Roberts, and C. R. Martin. 1986. Follow-up testing of blood donors found to be enzyme immunoassay positive/Western blot negative for HTLV-III antibody. *J. Am. Med. Assoc.* 256:40-41.
- Carlson, J. R., S. C. Mertens, J. L. Yee, M. B. Gardner, E. J. Watson-Williams, J. Ghayeb, M. B. Jennings, and R. J. Biggar. 1987. Rapid, easy, and economical screening test for antibodies to human immunodeficiency virus. *Lancet* i:361-362.
- Carter, W. A., D. R. Strayer, I. Brodsky, M. Lewin, M. G. Pellegrino, L. Einck, H. F. Henriques, G. L. Simon, D. M. Parenti, R. G. Scheib, R. S. Schulof, D. C. Montefiori, W. E. Robinson, W. M. Mitchell, D. J. Volsky, D. Paul, H. Paxton, W. A. Meyer, K. Kariko, N. Reichenbach, R. J. Suhadolnik, and D. H. Gillespie. 1987. Clinical, immunological, and virological effects of amplitgen, a mismatched double-stranded RNA, in patients with AIDS or AIDS-related complex. *Lancet* i:1286-1292.
- Centers for Disease Control. 1981. Pneumocystis pneumonia—Los Angeles. *Morbidity and Mortality Weekly Report* 30:250-252.
- Centers for Disease Control. 1985. Provisional public health service inter-agency recommendations for screening donated blood and plasma for antibody to the virus causing acquired immunodeficiency syndrome. *Morbidity and Mortality Weekly Report* 34:1-5.
- Centers for Disease Control. 1986. Human T-lymphotropic virus type III/lymphadenopathy associated virus: agent summary statement. *Morbidity and Mortality Weekly Report* 35:540-549.
- Centers for Disease Control. 1986. Update: acquired immunodeficiency syndrome—United States. *Morbidity and Mortality Weekly Report* 35:757-766.
- Centers for Disease Control. 1987. Human immunodeficiency virus infection in transfusion recipients and their family members. *Morbidity and Mortality Weekly Report* 36:137-140.
- Centers for Disease Control. 1987. Update: Human immunodeficiency virus infections in health-care workers exposed to blood of infected patients. *Morbidity and Mortality Weekly Report* 36:286-289.
- Centers for Disease Control. 1987. Human immunodeficiency virus infection transmitted from an organ donor screened for HIV antibody—North Carolina. *Morbidity and Mortality Weekly Report* 36:306-308.
- Centers for Disease Control. 1987. Public health service guidelines for counseling and antibody testing to prevent HIV infection and AIDS. *Morbidity and Mortality Weekly Report* 36:509-515.
- Centers for Disease Control. 1987. Update: acquired immunodeficiency syndrome—United States. *Morbidity and Mortality Weekly Report* 36:522-526.
- Chaisson, R. E., J. Allain, and P. A. Volberding. 1986. Significant changes in HIV antigen level in the serum of patients treated with azidothymidine. *N. Engl. J. Med.* 315:1610-1611.
- Chiodi, F., U. Bredberg-Raden, G. Biberfeld, B. Bottiger, J. Albert, B. Asjo, E. M. Fenyo, and E. Norrby. 1987. Radioimmuno-precipitation and Western blotting with sera of human immunodeficiency virus infected patients: a comparative study. *AIDS Res. Human Retroviruses* 3:165-176.

33. Clavel, F., D. Guetard, F. Brun-Vezinet, S. Chamaret, M. Rey, M. O. Santos-Ferreira, A. G. Laurent, C. Dauguet, C. Katlama, C. Rouzioux, D. Klatzmann, J. L. Champalimaud, and L. Montagnier. 1986. Isolation of a new human retrovirus from West African patients with AIDS. *Science* 233:343-346.
34. Clavel, F., K. Mansinho, S. Chamaret, D. Guetard, V. Favier, J. Nina, M. Santos-Ferreira, J. Champalimaud, and L. Montagnier. 1987. Human immunodeficiency virus type 2 infection associated with AIDS in West Africa. *N. Engl. J. Med.* 316:1180-1185.
35. Cooper, D. A., J. Gold, P. Maclean, B. Donovan, R. Finlayson, T. G. Barnes, H. M. Michelmore, P. Brooke, and R. Penny. 1985. Acute AIDS retrovirus infection: definition of a clinical illness associated with seroconversion. *Lancet* i:537-540.
36. Courouge, A., J. Muller, and D. Richard. 1986. False-positive Western blot reactions to human immunodeficiency virus in blood donors. *Lancet* ii:921-922.
37. Essex, M., J. Allan, P. Kanki, M. F. McLane, G. Malone, L. Kitchen, and T. H. Lee. 1985. Antigens of human T-lymphotropic virus type III/lymphadenopathy-associated virus. *Ann. Intern. Med.* 103:700-703.
38. Essex, M., M. F. McLane, T. H. Lee, N. Tachibana, J. I. Mullins, J. Kreiss, C. K. Kasper, M. C. Poon, A. Landay, S. F. Stein, D. P. Francis, C. Cabradilla, D. N. Lawrence, and B. L. Evatt. 1983. Antibodies to human T-cell leukemia virus membrane antigens (HTLV-MA) in hemophiliacs. *Science* 221:1061-1064.
39. Esteban, J. I., J. W. Shih, C. C. Tai, A. J. Bodner, J. W. D. Kay, and H. J. Alter. 1985. Importance of Western blot analysis in predicting infectivity of anti-HTLV-III/LAV positive blood. *Lancet* ii:1083-1086.
40. Falk, L. A., D. Paul, A. Landay, and H. Kessler. 1987. HIV isolation from plasma of HIV-infected persons. *N. Engl. J. Med.* 316:1547-1548.
41. Fleming, D. W., S. L. Cochi, R. S. Steece, and H. F. Hull. 1987. Acquired immunodeficiency syndrome in low-incidence areas. *J. Am. Med. Assoc.* 258:785-787.
42. Franchini, G., M. Robert-Guroff, A. Aldovini, N. C. Kan, and F. Wong-Staal. 1987. Spectrum of natural antibodies against five HTLV-III antigens in infected individuals: correlation of antibody prevalence with clinical status. *Blood* 69:437-441.
43. Fujikawa, L. S., S. Z. Salahuddin, A. G. Palestine, H. Masur, R. B. Nussenblatt, and R. C. Gallo. 1985. Isolation of human T-lymphotropic virus type III from the tears of a patient with the acquired immunodeficiency syndrome. *Lancet* ii:529-530.
44. Gaines, H., J. Albert, M. Von Sydow, A. Sonnerborg, F. Chiodi, A. Ehrnst, O. Strannegard, and B. Asjo. 1987. HIV antigenemia and virus isolation from plasma during primary HIV infection. *Lancet* i:1317-1318.
45. Gaines, H., M. Von Sydow, A. Sonnerborg, J. Albert, J. Czajkowski, P. O. Pehrson, F. Chiodi, L. Moberg, E. V. Fenyo, B. Asjo, and M. Forsgren. 1987. Antibody response in primary human immunodeficiency virus infection. *Lancet* i:1249-1253.
46. Gallo, D., J. L. Diggs, G. R. Shell, P. J. Dailey, M. N. Hoffman, and J. L. Riggs. 1986. Comparison of detection of antibody to the acquired immune deficiency syndrome virus by enzyme immunoassay, immunofluorescence, and Western blot methods. *J. Clin. Microbiol.* 23:1049-1051.
47. Gallo, R. C., S. Z. Salahuddin, M. Popovic, G. M. Shearer, M. Kaplan, B. F. Haynes, T. J. Palker, R. Redfield, J. Oleske, B. Safai, G. White, P. Foster, and P. D. Markham. 1984. Frequent detection and isolation of cytopathic retroviruses (HTLV-III) from patients with AIDS and at risk for AIDS. *Science* 224:500-502.
48. Gallo, R. C., and F. Wong-Staal. 1985. A human T-lymphotropic retrovirus (HTLV-III) as the cause of the acquired immunodeficiency syndrome. *Ann. Intern. Med.* 103:679-689.
49. Gartner, S., P. Markovits, D. M. Markovitz, M. H. Kaplan, R. C. Gallo, and M. Popovic. 1986. The role of mononuclear phagocytes in HTLV-III/LAV infection. *Science* 233:215-219.
50. Gnann, J. W., J. B. McCormick, S. Mitchell, J. A. Nelson, and M. B. A. Oldstone. 1987. Synthetic peptide immunoassay distinguishes HIV type 1 and HIV type 2 infections. *Science* 237:1346-1349.
51. Gnann, J. W., P. L. Schwimmbeck, J. A. Nelson, A. B. Truax, and M. B. A. Oldstone. 1987. Diagnosis of AIDS by using a 12-amino acid peptide representing an immunodominant epitope of the human immunodeficiency virus. *J. Infect. Dis.* 156:261-267.
52. Goudsmit, J., F. DeWolf, D. A. Paul, L. G. Epstein, J. M. A. Lange, W. J. A. Krone, H. Spellman, E. C. Wolters, J. Van Der Noordaa, J. M. Oleske, H. J. Van Der Helm, and R. A. Coutinho. 1986. Expression of human immunodeficiency virus antigen (HIV-Ag) in serum and cerebrospinal fluid during acute and chronic infection. *Lancet* ii:177-180.
53. Goudsmit, J., J. M. A. Lange, D. A. Paul, and G. J. Dawson. 1987. Antigenemia and antibody titers to core and envelope antigens in AIDS, AIDS-related complex, and subclinical human immunodeficiency virus infection. *J. Infect. Dis.* 155:558-560.
54. Grady, G. F., V. A. Lee, A. M. Prince, G. L. Gitnick, K. A. Fawaz, G. N. Vyas, M. D. Levitt, J. R. Senior, J. T. Galambos, T. E. Bynum, J. W. Singleton, B. F. Clowdus, K. Akdamar, R. D. Aach, E. I. Winkelman, G. M. Schiff, and T. Hersch. 1978. Hepatitis B immune globulin for accidental exposures among medical personnel: final report of a multicenter controlled trial. *J. Infect. Dis.* 138:625-638.
55. Groopman, J. E., P. I. Hartzband, L. Shulman, S. Z. Salahuddin, M. G. Sarngadharan, M. F. McLane, M. Essex, and R. Gallo. 1985. Antibody seronegative human T-lymphotropic virus type III (HTLV-III)-infected patients with acquired immunodeficiency syndrome or related disorders. *Blood* 66:742-744.
56. Groopman, J. E., S. Z. Salahuddin, M. G. Sarngadharan, P. D. Markham, M. Gonda, A. Sliski, and R. C. Gallo. 1984. HTLV-III in saliva of people with AIDS-related complex and healthy homosexual men at risk for AIDS. *Science* 226:447-448.
57. Harper, M. E., L. M. Marselle, R. C. Gallo, and F. Wong-Staal. 1986. Detection of lymphocytes expressing human T-lymphotropic virus type III in lymph nodes and peripheral blood from infected individuals by in situ hybridization. *Proc. Natl. Acad. Sci. USA* 83:772-776.
58. Ho, D. D., R. E. Byington, R. T. Schooley, T. Flynn, T. R. Rota, and M. S. Hirsch. 1985. Isolation of HTLV-III from 83 saliva and 50 blood samples from 71 seropositive homosexual men. *N. Engl. J. Med.* 313:1606.
59. Ho, D. D., R. Rota, R. T. Schooley, J. C. Kaplan, J. D. Allan, J. E. Groopman, L. Resnick, D. Felsenstein, C. A. Andrews, and M. S. Hirsch. 1985. Isolation of HTLV-III from cerebrospinal fluid and neural tissues of patients with neurologic syndromes related to the acquired immunodeficiency syndrome. *N. Engl. J. Med.* 313:1493-1497.
60. Ho, D. D., M. G. Sarngadharan, L. Resnick, F. Dimarzo-Veronese, T. R. Rota, and M. S. Hirsch. 1985. Primary human T-lymphotropic virus type III infection. *Ann. Intern. Med.* 103:880-883.
61. Ho, D. D., R. T. Schooley, T. R. Rota, J. C. Kaplan, and T. Flynn. 1984. HTLV-III in the semen and blood of a healthy homosexual man. *Science* 226:451-453.
62. Hymes, K. D. 1986. Kaposi's sarcoma in AIDS. *Lab. Med.* 17:668-673.
63. Jackson, B., K. Sannerud, F. Rhame, R. Tsang, and H. H. Balfour, Jr. 1987. Comparison of reverse transcriptase assay with the Retro-Tek viral capture assay for detection of human immunodeficiency virus. *Diagn. Microbiol. Infect. Dis.* 7:185-192.
64. Kaminsky, L. S., T. McHugh, D. Stites, P. Volberding, G. Henle, W. Henle, and J. A. Levy. 1985. High prevalence of antibodies to acquired immune deficiency syndrome (AIDS)-associated retrovirus (ARV) in AIDS and related conditions but not in order disease states. *Proc. Natl. Acad. Sci. USA* 82:5535-5539.
65. Kanki, P. J., F. Barin, S. M'Boup, J. S. Allan, J. O. Pomet-Lemonne, R. Marlink, M. F. McLane, T. Lee, B. Arbelille, F. Denis, and M. Essex. 1986. New human T-lymphotropic retrovirus related to simian T-lymphotropic virus type III (STLV-

- III) *AGM*. *Science* 232:238-243.
66. Kenny, C., J. Parkin, G. Undershill, N. Shah, B. Burnell, E. Osborne, and D. J. Jeffries. 1987. HIV antigen testing. *Lancet* i:565-566.
67. Kitchen, L. W., F. Barin, J. L. Sullivan, M. F. McLane, D. B. Brettler, P. H. Levine, and M. Essex. 1984. Aetiology of AIDS-antibodies to human T-cell leukemia virus (type III) in hemophiliacs. *Nature (London)* 312:367-369.
68. Knowles, D. M. 1986. Malignant lymphomas occurring in association with acquired immunodeficiency syndrome. *Lab. Med.* 17:674-678.
69. Koenig, S., H. E. Gendelman, J. M. Orenstein, M. C. Dal Canto, G. H. Pezeshkpour, M. Yungbluth, F. Janotta, A. Aksamit, M. A. Martin, and A. S. Fauci. 1986. Detection of AIDS virus in macrophages in brain tissue from AIDS patients with encephalopathy. *Science* 233:1089-1093.
70. Koyanagi, Y. S., Miles, R. T., Mitsuyasu, J. E., Merrill, H. V., Vinters, and I. S. Y. Chen. 1987. Dual infection of the central nervous system by AIDS viruses with distinct cellular tropisms. *Science* 236:819-822.
71. Kuhn, P., S. Seidl, and G. Holzberger. 1985. HLA DR4 antibodies cause positive HTLV-III antibody ELISA results. *Lancet* i:1222-1223.
72. Kumar, P., J. E. Pearson, D. H. Martin, S. H. Leech, P. D. Buisseret, H. C. Bezbak, F. M. Gonzalez, J. R. Royer, H. Z. Streicher, and W. C. Saxinger. 1987. Transmission of human immunodeficiency virus by transplantation of a renal allograft, with development of the acquired immunodeficiency syndrome. *Ann. Intern. Med.* 106:244-245.
73. Kwok, S., D. H. Mack, K. B. Mullis, B. Poiesz, G. Ehrlich, D. Blair, A. Friedman-Kien, and J. J. Sninsky. 1987. Identification of human immunodeficiency virus sequences by using in vitro enzymatic amplification and oligomer cleavage detection. *J. Virol.* 61:1690-1694.
74. Lange, J., R. A. Coutinho, W. J. A. Krone, L. F. Verdonch, S. A. Danner, J. Van Der Noordaa, and J. Goudsmit. 1986. Distinct IgG recognition patterns during progression of subclinical and clinical infection with lymphadenopathy associated virus/human T lymphotropic virus. *Br. Med. J.* 292:228-230.
75. Lange, J., and J. Goudsmit. 1987. Decline of antibody reactivity to HIV core protein secondary to increased production of HIV antigen. *Lancet* i:448.
76. Lange, J., D. A. Paul, H. G. Huisman, F. DeWolf, H. Van Den Berg, R. A. Coutinho, S. A. Danner, J. Van Der Noordaa, and J. Goudsmit. 1986. Persistent HIV antigenemia and decline of HIV core antibodies associated with transition to AIDS. *Br. Med. J.* 293:1459-1462.
77. Levy, J. A., A. D. Hoffman, S. M. Kramer, J. A. Landis, J. M. Shimabukuro, and L. S. Oshiro. 1984. Isolation of lymphocytopathic retroviruses from San Francisco patients with AIDS. *Science* 225:840-842.
78. Levy, J. A., and J. Shimabukuro. 1985. Recovery of AIDS-associated retroviruses from patients with AIDS or AIDS-related conditions and from clinically healthy individuals. *J. Infect. Dis.* 152:734-738.
79. Levy, J. A., J. Shimabukuro, H. Hollander, J. Mills, and L. Kaminsky. 1985. Isolation of AIDS-associated retroviruses from cerebrospinal fluid and brain of patients with neurological symptoms. *Lancet* ii:586-588.
80. Marlink, R. G., J. S. Allan, M. F. McLane, M. Essex, K. C. Anderson, and J. E. Groopman. 1986. Low sensitivity of ELISA testing in early HIV infection. *N. Engl. J. Med.* 315:1549.
81. Martin, P. W., D. R. Burger, S. Caouette, A. S. Goldstein, and F. Peetoom. 1986. Importance of confirmatory tests after strongly positive HTLV-III screening tests. *N. Engl. J. Med.* 314:1577-1578.
82. McCray, E. 1986. Occupational risk of the acquired immunodeficiency syndrome among health care workers. *N. Engl. J. Med.* 314:1127-1132.
83. Mendenhall, C. L., G. A. Roselle, C. J. Grossman, S. D. Rouster, R. E. Weesner, and U. Dumaswala. 1986. False positive tests for HTLV-III antibodies in alcoholic patients with hepatitis. *N. Engl. J. Med.* 314:921-922.
84. Meyer, K. B., and S. G. Pauker. 1987. Screening for HIV: can we afford the false positive rate? *N. Engl. J. Med.* 317:238-241.
85. Michaelis, B. A., and J. A. Levy. 1987. Recovery of human immunodeficiency virus from serum. *J. Am. Med. Assoc.* 257:1327.
86. Mok, J. Q., C. Giaquinto, A. DeRossi, I. Grosch-Worner, A. E. Ades, and C. S. Peckham. 1987. Infants born to mothers seropositive for human immunodeficiency virus. *Lancet* i:1164-1167.
87. Monroe, J. E., C. Andrews, J. L. Sullivan, and C. Mulder. 1987. Use of cytoplasmic dot-blot hybridization to detect human immunodeficiency virus RNA sequences in cultures of peripheral blood. *J. Infect. Dis.* 155:320-322.
88. Neisson-Vernant, C., S. Arfi, D. Mathez, J. Leibowitch, and N. Monplaisir. 1986. Needlestick HIV seroconversion in a nurse. *Lancet* ii:814.
89. Norrby, E., G. Biberfeld, F. Chiodi, A. von Gegerfeldt, A. Naucier, E. Parks, and R. Lerner. 1987. Discrimination between antibodies to HIV and to related retroviruses using site-directed serology. *Nature (London)* 329:248-250.
90. Oksenhendler, E., M. Harzic, J. LeRoux, C. Rabian, and J. P. Clauvel. 1986. HIV infection with seroconversion after a superficial needlestick injury to the finger. *N. Engl. J. Med.* 315:582.
91. Palca, J. 1987. Safety issues raised by worker's infection with AIDS virus. *Nature (London)* 329:191.
92. Pan, L. Z., C. Cheng-Mayer, and J. A. Levy. 1987. Patterns of antibody response in individuals infected with the human immunodeficiency virus. *J. Infect. Dis.* 155:626-632.
93. Parry, J. V., and P. P. Mortimer. 1986. Place of IgM antibody testing in HIV serology. *Lancet* ii:979-980.
94. Parry, J. V., K. R. Perry, and P. P. Mortimer. 1987. Sensitive assays for viral antibodies in saliva: an alternative to tests on serum. *Lancet* ii:72-75.
95. Paul, D. A., L. A. Falk, H. A. Kessler, R. M. Chase, B. Blaauw, D. S. Chudwin, and A. L. Landay. 1987. Correlation of serum HIV antigen and antibody with clinical status in HIV-infected patients. *J. Med. Virol.* 22:357-363.
96. Peterman, T. A., G. R. Lang, N. J. Mikos, S. L. Solomon, C. A. Schable, P. M. Feorino, J. A. Britz, and J. R. Allen. 1986. HTLV-III/LAV infection in hemodialysis patients. *J. Am. Med. Assoc.* 255:2324-2326.
97. Poiesz, B. J., F. W. Ruscetti, A. F. Gazdar, P. A. Bunn, J. D. Minna, and R. C. Gallo. 1980. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc. Natl. Acad. Sci. USA* 77:7415-7419.
98. Popovic, M., M. G. Sarngadharan, E. Read, and R. C. Gallo. 1984. Detection, isolation, and continuous production of cytopathic retroviruses (HTLV-III) from patients with AIDS and pre-AIDS. *Science* 224:497-500.
99. Quinn, T. C., J. M. Mann, J. W. Curran, and P. Piot. 1986. AIDS in Africa: an epidemiologic paradigm. *Science* 234:955-963.
100. Ranki, A., S. Valle, M. Krohn, J. Anttonen, J. Allain, M. Leuther, G. Franchini, and K. Krohn. 1987. Long latency precedes overt seroconversion in sexually transmitted human immunodeficiency-virus infection. *Lancet* ii:589-593.
101. Reesink, H. W., P. N. Lelie, J. G. Huisman, W. Schaasberg, M. Gonsalves, C. Aaij, I. N. Winkel, J. A. Van Der Does, A. C. Hekker, J. Desmyter, and J. Goudsmit. 1986. Evaluation of six enzyme immunoassays for antibody against human immunodeficiency virus. *Lancet* ii:483-486.
102. Resnick, L., K. Veren, S. Z. Salahuddin, S. Tondreau, and P. D. Markham. 1986. Stability and inactivation of HTLV-III/LAV under clinical and laboratory environments. *J. Am. Med. Assoc.* 255:1887-1891.
103. Rey, M. A., B. Spire, D. Dormont, F. Barre-Sinoussi, L. Montagnier, and J. C. Chermann. 1984. Characterization of the RNA dependent DNA polymerase of a new human T lymphotropic retrovirus (lymphadenopathy associated virus). *Bio-*

- chem. Biophys. Res. Commun. 121:126-133.
104. Riggins, C. H., G. A. Beltz, C. H. Hung, R. M. Thorn, and D. J. Marciani. 1987. Detection of antibodies to human immunodeficiency virus by latex agglutination with recombinant antigen. *J. Clin. Microbiol.* 25:1772-1773.
105. Salahuddin, S. Z., J. E. Groopman, P. D. Markham, M. G. Sarngadharan, R. R. Redfield, M. F. McLane, M. Essex, A. Sliski, and R. C. Gallo. 1984. HTLV-III in symptom-free seronegative persons. *Lancet* ii:1418-1420.
106. Sandstrom, E. G., R. T. Schooley, D. D. Ho, R. Byington, M. G. Sarngadharan, M. E. MacLane, M. Essex, R. C. Gallo, and M. S. Hirsch. 1985. Detection of human anti-HTLV-III antibodies by indirect immunofluorescence using fixed cells. *Transfusion* 25:308-312.
107. Sarngadharan, M. G., M. Popovic, L. Bruch, J. Schupbach, and R. C. Gallo. 1984. Antibodies reactive with human T-lymphotropic retroviruses (HTLV-III) in the serum of patients with AIDS. *Science* 224:506-508.
108. Schulz, T. F., J. M. Aschauer, P. Hengster, C. Larcher, H. Wachter, B. Fleckenstein, and M. P. Dierich. 1986. Envelope gene-derived recombinant peptide in the serodiagnosis of human immunodeficiency virus infection. *Lancet* ii:111-112.
109. Schupbach, J., M. Popovic, R. V. Gilden, M. A. Gonda, M. G. Sarngadharan, and R. C. Gallo. 1984. Serological analysis of a subgroup of human T-lymphotropic retroviruses (HTLV-III) associated with AIDS. *Science* 224:503-505.
110. Seeff, L. B., E. C. Wright, H. J. Zimmerman, H. J. Alter, A. A. Dietz, B. F. Felsher, J. D. Finkelstein, P. Garcia-Pont, J. L. Gerin, H. B. Greenlee, J. Hamilton, P. V. Holland, P. M. Kaplan, T. Kiernan, R. S. Koff, C. M. Leevy, V. J. McAuliffe, N. Nath, R. H. Purcell, E. R. Schiff, C. C. Schwartz, C. H. Tamburro, Z. Vlahcevic, R. Zemel, and D. S. Zimmon. 1978. Type B hepatitis after needle-stick exposure: prevention with Hepatitis B immune globulin. *Ann. Intern. Med.* 88:285-293.
111. Shaw, G. M. B. H. Hahn, S. K. Arya, J. E. Groopman, R. C. Gallo, and F. Wong-Staal. 1984. Molecular characterization of human T-cell leukemia (lymphotropic) virus type III in the acquired immune deficiency syndrome. *Science* 226:1165-1171.
112. Sivak, S. L., and G. P. Wormser. 1986. Predictive value of a screening test for antibodies to HTLV-III. *Am. J. Clin. Pathol.* 85:700-703.
113. Smith, D. M., S. Dewhurst, S. Shepherd, D. J. Volsky, and J. C. Goldsmith. 1987. False-positive enzyme-linked immunosorbent assay reactions for antibody to human immunodeficiency virus in a population of midwestern patients with congenital bleeding disorders. *Transfusion* 27:112.
114. Stricof, R. L., and D. L. Morse. 1986. HTLV-III/LAV seroconversion following a deep intramuscular needlestick injury. *N. Engl. J. Med.* 314:1115.
115. Stute, R. 1987. HIV antigen detection in routine blood donor screening. *Lancet* i:566.
116. Thiry, L., S. Sprecher-Goldberger, T. Jonckheer, J. Levy, P. Van De Perre, P. Henrivaux, J. Cogniaux-LeClerc, and N. Clumeck. 1985. Isolation of AIDS virus from cell-free breast milk of three healthy virus carriers. *Lancet* ii:891-892.
117. Tsang, V. C. W., J. M. Peralta, and A. R. Simons. 1983. Enzyme-linked immunoelectrotransfer blot techniques (EITB) for studying the specificities of antigens and antibodies separated by gel electrophoresis. *Methods Enzymol.* 92:377-391.
118. Ulstrup, J. C., K. Skaug, K. J. Figenschau, I. Orstavik, J. N. Brunn, and G. Petersen. 1986. Sensitivity of Western blotting (compared with ELISA and immunofluorescence) during seroconversion after HTLV-III infection. *Lancet* i:1151-1152.
119. U.S. Public Health Service. 1986. Coolfont Report: a PHS plan for prevention and control of AIDS and the AIDS virus. *Public Health Rep.* 101:341-348.
120. Van der Poel, C. L., H. W. Reesink, T. Tersmette, P. N. Lelie, H. Huisman, and F. Miedema. 1986. Blood donations reactive for HIV in Western blot, but non-infective in culture and recipients of blood. *Lancet* ii:752-753.
121. Vogt, M. W., D. J. Witt, D. E. Craven, R. Byington, D. F. Crawford, R. T. Schooley, and M. S. Hirsch. 1986. Isolation of HTLV-III/LAV from cervical secretions of women at risk for AIDS. *Lancet* i:525-527.
122. Wain-Hobson, S., P. Sonigo, O. Danos, S. Cole, and M. Alizon. 1985. Nucleotide sequence of the AIDS virus, LAV. *Cell* 40:9-17.
123. Wall, R. A., D. W. Denning, and A. Amos. 1987. HIV antigenemia in acute HIV infection. *Lancet* i:566.
124. Ward, J. W., A. J. Grindon, P. M. Feorino, C. Schable, M. Parvin, and J. R. Allen. 1986. Laboratory and epidemiologic evaluation of an enzyme immunoassay for antibodies to HTLV-III. *J. Am. Med. Assoc.* 256:357-361.
125. Weber, J. N., P. R. Clapham, R. A. Weiss, D. Parker, C. Roberts, J. Duncan, I. Weller, C. Carne, R. S. Tedder, A. J. Pinching, and R. Cheingsong-Popov. 1987. Human immunodeficiency virus infection in two cohorts of homosexual men: neutralizing sera and association of anti-gag antibody with prognosis. *Lancet* i:119-121.
126. Wetherbee, R. E. 1986. Infections in AIDS. *Lab. Med.* 17:679-684.
127. Zagury, D., J. Bernard, J. Leibowitch, B. Safai, J. E. Groopman, M. Feldman, M. G. Sarngadharan, and R. C. Gallo. 1984. HTLV-III in cells cultured from semen of two patients with AIDS. *Science* 226:449-451.
128. Zolla-Pazner, S. 1986. Immunologic abnormalities in infections with the human immunodeficiency virus. *Lab. Med.* 17:685-689.