Published in final edited form as:

Org Lett. 2012 December 7; 14(23): 6056–6059. doi:10.1021/ol302947r.

Palladium-Catalyzed Carbonylative Cyclization/Arylation Cascade for 2-Aroyl Indolizine Synthesis

Zhou Li, Dmitri Chernyak, and Vladimir Gevorgyan*

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607-7061

Abstract

An efficient synthesis of densely substituted 2-aroyl indolizines via the palladium-catalyzed carbonylative cyclization/arylation is reported. This transformation proceeds via the *5-endo-dig* cyclization of 2-propargylpyridine triggered by an aroyl Pd-complex. It produced diversely substituted 2-aroyl indolizines in good-to-excellent yield.

Indolizines have attracted noteworthy attention in recent years due to their profound biological effects.¹ Both naturally-occurring and synthetic indolizines have shown great potential in pharmaceutical research as cytotoxins,² anti-inflammatory agents,³ and 5-HT₃ receptor antagonists⁴. In this regard, transformations that utilize readily available substrates to provide access to diversely substituted indolizines, especially those bearing an electron withdrawing group at the C-2 position⁵, are in high demand. The classic Tschichibabin reaction provides straightforward access to C-2 substituted indolizines via the condensation of picolines and α -bromo-acetophenone derivatives. However, the limited availability of starting materials restricts the substitution pattern of the product.⁶ The [3+2] cycloaddition of pyridinium ylides with alkynes provides another viable route to the indolizine core. However, the regioselectivity issue, as well as the moderate yields caused by necessary oxidation of the formed intermediate, limits its synthetic application.⁷ The Morita-Baylis-Hillman reaction can also be utilized for the preparation of indolizin-2-yl ketone from an appropriate Michael acceptor. However, it suffers from the narrow range of starting materials that can be used and the usually low reactivity of the substrates.⁸ In addition to the traditional condensation methods, transition metal catalysis has been widely used for the construction of diversely functionalized, especially heteroatom-substituted indolizines, under mild conditions.⁹ Nonetheless, the selective introduction of functionality at the C-2 position is still a challenging task.¹⁰ Along this line, our group has recently reported a synthesis of 2-aryl indolizines via the palladium catalyzed arylative *5-endo-dig* cyclization of 2-propargylpyridine (Scheme 1, eq. 1).¹¹ Herein, we report a Pd-catalyzed cascade carbonylative cyclization/arylation approach to diversely substituted 2-aroyl indolizines that proceeds in good-to-excellent yields (eq. 2).

vlad@uic.edu.

Supporting Information Available Detailed experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Continuing with our efforts in the synthesis of diversely functionalized indolizines, we thought that the employment of benzoyl chloride instead of an aryl halide as electrophile in this cascade cyclization would provide 2-benzoyl indolizine.¹² However, the reaction of pivaloate **1a** with benzoyl chloride in the presence of a Pd catalyst failed to produce even trace amount of **2a** (eq. 3).

Thus, we thought of alternative methods to introduce the benzoyl function at the C-2 position of indolizine. Since (1) an acyl palladium species can be formed via a migratory insertion of carbon monoxide into an aryl-palladium bond; and (2) there have been reported examples on synthesis of diaryl ketones via the palladium catalyzed carbonylative cyclization/arylation cascades,¹³ we anticipated that under a CO atmosphere, the *in situ* formed acyl palladium species might trigger the expected carbonylative cyclization/arylation cascade to obtain **2**.

(3)

To test this idea, we examined the carbonylative cyclization of pivaloate **1a** and different aryl halides under continuous 10 psi CO supply. Initially we found that the carbonylative cyclization of **1a** and iodobenzene in the presence of Pd(PPh₃)Cl₂ catalyst yielded **2a** in quantitative yield (Table 1, entry 1). However, aryl halides bearing electron-withdrawing groups, such as methyl *p*-iodobenzoate, were not competent reactants under these conditions (entry 2).¹⁴ On the contrary, the combination of Pd(OAc)₂ catalyst, PCy₃ ligand and triethylamine base afforded benzoate **2f** in 90% yield (entry 3). Under the same conditions, the yield of **2a** was only 19% along with a substantial amount of palladium black produced, which implied decomposition of an excessively reactive catalyst (entry 4). The yield of **2a** was improved to 51% by switching ligand to PPh₃ (entry 5). The reaction that was performed in a sealed Schlenk tube, which simplified the reaction setup and allowed a higher pressure (20 psi and above), provided **2a** in 86% yield at 20 psi CO pressure (entry 6). Finally the screening of different solvents under a reduced temperature revealed that acetonitrile was the optimal choice over DMF and toluene (entries 7–9).

With the optimized conditions in hand, the scope of reaction was examined. It was found that a series of iodoarenes bearing different substituents at various positions smoothly underwent this carbonylative cyclization with **1a**, yielding indolizines **2a** to **2i** in moderate to excellent yields (Table 2, entries 1–9). The cyclization of 3-*n*-hexyl, *n*-butyl and cyclohexenyl substituted pivaloates also provided the expected products in good yield (entry 10 to 15). Substrates possessing a functionalized pyridine ring at C-5 produced indolizines **2p**, **2q** and pyrrolo[1,2-*a*]quinoline **2r** in moderate-to-good yields (entries 16–18). On the contrary, cyclization of 3-methyl substituted pyridine provided 8-methyl **2s** in a relative low yield (entry 19). In addition to various pivaloates, a TBDMS ether was equally effective in this reaction, yielding **2t** in 80% yield (entry 20). More interestingly, the cyclization of 1-(1-pyridin-2-yl-propargyl)morpholine resulted in the 1-morpholin-1-yl indolizine **2u** in 71% yield, which provided a convenient access to the C-2 substituted 1-amino-indolizines.¹⁵

Presumably, this palladium catalyzed carbonylative cyclization starts with the formation of the aroyl palladium species **3** (Scheme 2) via a migratory insertion of CO into the aryl-palladium bond, followed by its coordination of the triple bond of **1** to the palladium center.

The *5-endo-dig* cyclization leads to the formation of indolizinium intermediate **4**, which after a proton loss-aromatization produces heteroaryl palladium species **5**. The reductive elimination of **5** releases the product 2-benzoylindolizine **2** to regenerate the Pd catalyst.

In summary, the Pd-catalyzed carbonylative cyclization/arylation approach to 2-aryl indolizines has been developed. This new method allows for the efficient and selective synthesis of important⁵ 2-aryl indolizines from readily available propargyl pyridines, iodoarenes under a carbon monoxide atmosphere.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The support of the National Institute of Health (GM-64444 and 1P50 GM-086145) is gratefully acknowledged.

References

1. (a) Flitsch, W. Comprehensive Heterocyclic Chemistry, II. Vol. 8. Pergamon Press; Oxford: 1984. p. 237-248. (b) Fan H, Peng J, Hamann MT, Hu JF. Chem Rev. 2008; 108:264. [PubMed: 18095718]
2. (a) Ishibashi F, Tanabe S, Oda T, Iwao M. J Nat Prod. 2002; 65:500. [PubMed: 11975488] (b) Pla D, Marchal A, Olsen CA, Francesch A, Cuevas C, Albericio F, Alvarez M. J Med Chem. 2006; 49:3257. [PubMed: 16722644] (c) Chittchang M, Batsomboon P, Ruchirawat S, Ploypradith P. ChemMedChem. 2009; 4:457. [PubMed: 19152364]
3. Oslund RC, Cermak N, Gelb MH. J Med Chem. 2008; 51:4708. [PubMed: 18605714]
4. Bermudez J, Fake CS, Joiner GF, Joiner KA, King FD, Miner WD, Sanger GJ. J Med Chem. 1990; 33:1924. [PubMed: 2362270]
5. For biological activity of indolizines bearing an electron withdrawing group at C-2 position, see: Shen YM, Lv PC, Chen W, Liu PG, Zhu HL, Zhang MZ. Euro J Med Chem. 2010; 45:3184–3190. Wu XW, Wu ZP, Chen JW, Zhang W, Gu LQ, Huang ZS, An LK, Wang LX, Zhang HB. Euro J Med Chem. 2011; 46:4625. Warui DM, Baranger AM. J Med Chem. 2009; 52:5462. [PubMed: 19691339]
6. (a) Tschitschibabin AE. Ber Dtsch Chem Ges. 1927; 60:1607. (b) Kostik EL, Abiko A, Oku A. J Org Chem. 2001; 66:2618. [PubMed: 11304179] (c) Liu Y, Sun JW. J Org Chem. 2012; 77:1191. [PubMed: 22185652]
7. (a) Katritzky AR, Qiou G, Yang B, He HY. J Org Chem. 1999; 64:7618. (b) Bora U, Saikia A, Boruah RC. Org Lett. 2003; 5:435. [PubMed: 12583737] (c) Rotaru AV, Druta ID, Oeser T, Müller TJJ. Helv Chim Acta. 2005; 88:1798.
8. (a) Bode ML, Kaye PT. J Chem Soc Perkin Trans 1. 1990:2612. (b) Basavaiah D, Rao AJ. Chem Commun. 2003:604. (c) Specowius V, Bendrath F, Winterberg M, Khurshid A, Langer P. Adv Synth Catal. 2012; 354:1163.
9. (a) Kel'in AV, Sromek AW, Gevorgyan V. J Am Chem Soc. 2001; 123:2074. [PubMed: 11456838] (b) Seregin I, Gevorgyan V. J Am Chem Soc. 2006; 128:12050. [PubMed: 16967938] (c) Liu Y, Song Z, Yan B. Org Lett. 2007; 9:409. [PubMed: 17249774] (d) Chuprakov S, Gevorgyan V. Org Lett. 2007; 9:4463. [PubMed: 17892296] (e) Hardin AR, Sarpong R. Org Lett. 2007; 9:4547. [PubMed: 17845052] (f) Barluenga J, Lonzi G, Riesgo L, López LA, Tomás M. J Am Chem Soc. 2010; 132:13200. and Ref. (11), (15). [PubMed: 20825189]
10. For the regioselectivity of the direct C-H functionalization of indolizine, see: Renard M, Gubin J. Tetrahedron Lett. 1992; 33:4433. Shipman M. Science of Synthesis. 2000; 10:777. Park CH, Ryabova V, Seregin IV, Sromek AW, Gevorgyan V. Org Lett. 2004; 6:1159. [PubMed: 15040747] Seregin IV, Ryabova V, Gevorgyan V. J Am Chem Soc. 2007; 129:7742. [PubMed: 17539646] Petit A, Flygare J, Miller AT, Winkel G, Ess DH. Org Lett. 2012; 14:3680.

11. (a) Chernyak D, Skontos C, Gevorgyan V. *Org Lett.* 2010; 12:3242. [PubMed: 20545315] (b) Chernyak D, Gevorgyan V. *Org Lett.* 2010; 12:5558. [PubMed: 21058673]
12. For palladium catalyzed acylation reactions using acyl chlorides as electrophile, see: Tohda Y, Sonogashira K, Hagihara N. *Synthesis.* 1977:777. Negishi, E-i; Bagheri, V.; Chatterjee, S.; Luo, F-T.; Miller, JA.; Stoll, AT. *Tetrahedron Lett.* 1983; 24:5181. Renaldo AF, Labadie JW, Stille JK. *Org Synth.* 1989; 67:86. Kabalka GW, Malladi RR, Tejedor D, Kelley S. *Tetrahedron Lett.* 2000; 41:999.
13. For selected reports on the palladium catalyzed carbonylative cyclization/arylation cascade, see: Arcadi A, Cacchi S, Carnicelli V, Marinelli F. *Tetrahedron.* 1994; 50:437. Dai G, Larock RC. *Org Lett.* 2001; 4:193. [PubMed: 11796048] Hu Y, Zhang Y, Yang Z, Fathi R. *J Org Chem.* 2002; 67:2365. [PubMed: 11925257] Dai D, Larock RC. *Org Lett.* 2002; 4:193. [PubMed: 11796048]
14. See Supporting Information for details.
15. For synthesis of 2-nonsubstituted 1-aminoindolizine via the A^3 coupling reaction, see: Yan B, Liu Y. *Org Lett.* 2007; 9:4323. [PubMed: 17854200] Bai Y, Zeng J, Ma J, Gorityala BK, Liu XW. *J Comb Chem.* 2010; 12:696. [PubMed: 20831266]

Scheme 1.
Indolizine Synthesis via Palladium Catalyzed Cyclization

Scheme 2.
Proposed Mechanism

Table 1

Optimization of Conditions

entry	ligand	base	solvent	temp.	CO ^a	yield (%) ^b
1	Pd(PPh ₃) ₂ Cl ₂ /PPh ₃	K ₂ CO ₃	DMF	100	A	2a > 99 ^c
2	Pd(PPh ₃) ₂ Cl ₂ /PPh ₃	K ₂ CO ₃	DMF	100	A	2f decomp. ^c
3	Pd(OAc) ₂ /PCy ₃	NEt ₃	DMF	80	A	2f 90
4	Pd(OAc) ₂ /PCy ₃	NEt ₃	DMF	80	A	2a 19
5	Pd(OAc) ₂ /PPh ₃	NEt ₃	DMF	80	A	2a 51
6	Pd(OAc) ₂ /PPh ₃	NEt ₃	DMF	80	B	2a 86
7	Pd(OAc) ₂ /PPh ₃	NEt ₃	DMF	70	B	2a 65
8	Pd(OAc) ₂ /PPh ₃	NEt ₃	toluene	70	B	2a 45
9	Pd(OAc) ₂ /PPh ₃	NEt ₃	MeCN	70	B	2a 95

^a A: Reactions were conducted in flask with continuous 10 psi CO supply. B: Reactions were conducted in sealed Schlenk tube with 20 psi CO.^b Isolated yields of 0.5 mmol reactions.^c 1 equivalent of TBAI was added.

Table 2

Reaction Scope of Carbonylative Cyclization.

Reaction scheme: A substituted indole derivative (1) with substituents R¹, R², and R³ reacts under conditions ^a to form a carbonyl compound (2) with substituents R¹, R², and R³.

entry	2	yield (%) ^b
1	2a	95
2	2b	77
3	2c	78
4	2d	99
5	2e	59
6	2f	61
7	2g	69

entry	2	yield (%) ^b
8	<p>2h</p>	64
9	<p>2i</p>	64
10	<p>2j</p>	80
11	<p>2k</p>	80
12	<p>2l</p>	64
13	<p>2m</p>	81
14	<p>2n</p>	67

entry	2	yield (%) ^b
15	<p>2o</p>	72
16	<p>2p</p>	90
17	<p>2q</p>	85
18	<p>2r</p>	53
19	<p>2s</p>	35
20	<p>2t</p>	80
21	<p>2u</p>	71

^aConditions: Pd(OAc)₂ 5 mol %, PPh₃ 10 mol %, Ar-I 1.5 equiv, triethylamine 2 equiv, MeCN 0.2 M, 70 °C, CO 20 psi in sealed Schlenk tube, 12 h.

^bIsolated yields of 0.5 mmol reactions.