Published in final edited form as:

Stem Cells. 2013 March ; 31(3): 607–611. doi:10.1002/stem.1294.

Long-term functional engraftment of mesenchymal progenitor cells in a mouse model of accelerated aging

Lakshman Singh^{1,†}, Tracy A. Brennan^{1,†}, Jung-Hoon Kim¹, Kevin P. Egan¹, Emily A. McMillan¹, Qijun Chen², Kurt D. Hankenson^{3,4}, Yi Zhang⁵, Stephen G. Emerson⁶, F. Brad Johnson², and Robert J. Pignolo^{1,4}

¹Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

²Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

³Department of Animal Biology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA 19104

⁴Department of Orthopaedic Surgery, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

⁵Department of Medicine, University of Michigan School of Medicine, Ann Arbor, MI 48109

⁶Department of Medicine, Columbia University Medical Center, New York, NY, 10032

Abstract

Age-related osteoporosis is characterized by a decrease in bone-forming capacity mediated by defects in the number and function of osteoblasts. An important cellular mechanism which may in part explain osteoblast dysfunction that occurs with aging is senescence of mesenchymal progenitor cells (MPCs). In the telomere-based *Wn^{-/-}Terc^{-/-}* model of accelerated aging, the osteoporotic phenotype of these mice is also associated with a major decline in MPC differentiation into osteoblasts. To investigate the role of MPC aging as a cell-autonomous mechanism in senile bone loss, transplantation of young wild-type whole bone marrow into *Wn^{-/-}Terc^{-/-}* mutants was performed and the ability of engrafted cells to differentiate into cells

Corresponding Author: Robert J. Pignolo, M.D., Ph.D., Departments of Medicine and Orthopaedic Surgery, Perelman School of Medicine, University of Pennsylvania, 424B Stemmler Hall, 36th Street and Hamilton Walk, Philadelphia, PA 19104, Telephone: 215-746-8138, FAX: 215-573-2133, pignolo@mail.med.upenn.edu.

[†]Equal contributions.

Disclosure of Potential Conflicts of Interest

Authors declare no conflicts of interest.

Author Contributions:

Lakshman Singh: collection and/or assembly of data; data analysis and interpretation; manuscript writing; final approval of manuscript.

Tracy Brennan: collection and/or assembly of data; data analysis and interpretation; manuscript writing; final approval of manuscript.

Jung-Hoon Kim: collection and/or assembly of data; data analysis and interpretation; manuscript writing; final approval of manuscript.

Kevin P. Egan: collection and/or assembly of data; final approval of manuscript.

Emily A. McMillan: collection and/or assembly of data; final approval of manuscript.

Qijun Chen: collection and/or assembly of data; provision of study materials; final approval of manuscript.

Kurt D. Hankenson: data analysis and interpretation; manuscript writing; final approval of manuscript.

Yi Zhang: collection and/or assembly of data; data analysis and interpretation; manuscript writing; final approval of manuscript.

Stephen G. Emerson: data analysis and interpretation; manuscript writing; final approval of manuscript.

F. Brad Johnson: collection and/or assembly of data; provision of study materials; data analysis and interpretation; manuscript writing; final approval of manuscript.

Robert J. Pignolo: conception and design; financial support; collection and/or assembly of data; data analysis and interpretation; manuscript writing; final approval of manuscript.

of the osteoblast lineage was assessed. We found that whole bone marrow transplantation (BMT) in $Wrn^{-/-}Terc^{-/-}$ mice resulted in functional engraftment of MPCs up to 42 weeks, which was accompanied by a survival advantage, as well as delays in microarchitectural features of skeletal aging.

Keywords

Telomere; telomere dysfunction; aging; osteoporosis; mesenchymal stem cells; engraftment

Introduction

Skeletal aging of bone is characterized by loss of mineral content and microarchitectural changes that decrease bone strength and predispose bone to damage from repeated loading [1, 2].

Osteoporosis is common in the Werner and dyskeratosis congenita premature aging syndromes, both characterized by telomere dysfunction [3, 4]. One of the targets of WRN helicase is telomeric DNA, but need for WRN at telomeres is minimized in mice by long telomeres and abundant telomerase, making Wrn knockout mice relatively unaffected [5, 6]. However, combining Wrn mutation with shortened telomeres of telomerase ($Terc$) knockout mice results in an accelerated aging model [5, 7]. Deficiencies in $Wrn^{-/-}Terc^{-/-}$ mutant mice cause a low bone mass phenotype due to impaired osteoblast differentiation in the context of intact osteoclast differentiation [8, 9]. This impaired differentiation is associated with telomere dysfunction, as measured by the association of DNA damage proteins with telomeres in mesenchymal progenitor cells (MPCs) isolated from double mutant mice [9]. MPCs from $Wrn^{-/-}Terc^{-/-}$ mutants have a reduced *in vitro* lifespan but also display impaired osteogenic potential with dysfunctional telomeres independently of proliferative state [9].

Here we test the hypothesis that MPC aging contributes to bone loss in an accelerated aging mouse model that recapitulates many aspects of age-related bone loss.

Materials and Methods

Detailed Material and Methods are described in the supporting information.

Results

Wild-type whole bone marrow transplantation (BMT) into $Wrn^{-/-}Terc^{-/-}$ mutants confers a survival advantage

To test the role of telomere-based MPC aging on age-related osteoporosis, $Wrn^{-/-}Terc^{-/-}$ mutants were transplanted at 3 months of age with whole BM from young wild type donors. At 10.5 months after transplantation or when animals exhibited signs of significant distress and impending demise (whichever occurred first), mutants were sacrificed for analysis of functional MPC engraftment and concomitant measurements of skeletal microarchitectural features. As shown in Fig. 1, transplanted animals exhibited a survival advantage, having a mean life span about 30% longer than untransplanted controls (12.89 ± 0.21 months versus 10 ± 0.57 months). This difference is particularly remarkable given that telomerase-deficient mice are hypersensitive to ionizing radiation and otherwise would have been expected to incur substantial harm from irradiation associated with BMT [10].

Long-term functional engraftment of MPCs in $Wrn^{-/-}Terc^{-/-}$ mice

Enhanced green fluorescent protein-positive (GFP⁺) wild-type mice were used as donors in all BMT experiments. GFP⁺ MPCs in BM aspirates were identified from transplanted animals by fluorescent immunohistochemistry and represented 54.0 ± 7.1 percent of plastic adherent stromal cells after 30 hours in culture. Expanded MPC cultures from young wild-type donor animals are essentially CD45⁻ Sca-1⁺ cells and can differentiate in vitro into osteoblasts and adipocytes (Fig. S1). CD45⁻ Sca-1⁺ MPC are present after long-term BMT in $Wrn^{-/-}Terc^{-/-}$ mice (Fig. 2).

MPCs demonstrate functional engraftment as differentiated GFP⁺ osteoblasts and osteocytes in bone sections from recipient $Wrn^{-/-}Terc^{-/-}$ mice (Fig. 3, Fig. S2). MPC functional engraftment is present up to 10.5 months after BMT. In femur sections from transplanted animals 20 ± 8 percent of cortical osteocytes and 6 ± 2 percent of trabecular osteocytes were derived from engrafted precursors. Among endocortical and trabecular bone-lining osteoblasts, 15 ± 6 percent and 5 ± 1 percent were from precursors of donor origin, respectively. Differentiation of engrafted MPCs was most evident as endocortical osteocytes (Fig. S2).

Delays in microarchitectural features of skeletal aging in $Wrn^{-/-}Terc^{-/-}$ mutants after BMT

Despite being ~30% older than non-transplanted double mutants, $Wrn^{-/-}Terc^{-/-}$ BMT recipients had preserved or improved measures of bone microarchitecture. Transplanted animals showed no statistically significant changes in trabecular bone volume/total volume, trabecular number, or in cortical thickness (Fig. 4). Interestingly, there was a statistically significant increase in the ratio of cortical area to total area, suggesting that transplanted mice were able to improve endocortical bone mass. Similarly, $Wrn^{-/-}Terc^{-/-}$ BMT recipients had preserved osteoblasts/bone surface and no increase in the number of osteoclasts/bone surface [Fig. S3].

Discussion

Although the causal role(s) of telomere dysfunction in age-related osteoporosis are not completely established, there is evidence of its importance. Telomere lengthening mechanisms are not present in human BM MPCs [11]. Exogenous telomerase expression extends in vitro proliferative capacity, accelerates osteogenic differentiation, and enhances bone formation upon subcutaneous transplantation into mice [12, 13]. In addition, progeroid syndromes on which the $Wrn^{-/-}Terc^{-/-}$ mutants are based (Werner syndrome and dyskerostis congenita, respectively) display premature osteoporosis [3, 4].

Although we cannot exclude that hematopoietic precursors including hematopoietic stem cells (HSCs), transplanted along with MPCs, may support mesenchymal engraftment and/or differentiation into cells of the osteoblast lineage, it is unlikely that CD45⁻ Sca-1⁺ MPCs are derived directly from HSCs. However, there may be an early common BM progenitor for hematopoietic cells and osteoblasts delineated as Lin⁻Sca-1⁺cKit⁺CD45⁺. [14, 15].

Soluble hematopoietic factors alone may be sufficient to exert effects on bone remodeling, or development of osteoblasts may depend on the proximity of hematopoietic cells. Sca-1, a cell-surface molecule also expressed on HSCs, appears to be necessary to maintain self-renewal of MPCs and suggests that the latter is plausible. In support of this, Sca-1 knockout mice develop age-dependent osteoporosis [16].

There could be systemic effects of BMT which may influence skeletal engraftment and differentiation of MPCs. Also, microenvironmental factors (including oxidative stress) may

be at play. For example, we previously showed that osteoblast differentiation of MPCs from $Wrn^{-/-}Terc^{-/-}$ mutants is rescued by reducing oxidative stress [9]. Increased oxidative stress favoring senescence in BM MPCs has also been postulated based on proteome screening of these cells from young and old rodents [17]. Muscle-derived stem/progenitor cells from young wild-type mice transplanted into a murine progeria model extended life span and improved degenerative changes in tissues where donor cells are not detected [18]. Although we cannot be sure that extra-skeletal effects of BMT are responsible for life span extension in $Wrn^{-/-}Terc^{-/-}$ mutants, the fact that $GFP^{+}Sca-1^{+}CD45^{-}$ MPCs are present in bone tissue after long-term transplantation suggests that they play a role in maintenance of bone microarchitectural features over the period of extended survival.

To the extent that bone loss with physiologic aging involves telomere-based aging, our data indicates MPC senescence is a contributory mechanism. However, BMT as a therapeutic strategy may be limited by inadequate MPC engraftment [19]. It is debatable whether donor MPCs from human or mouse sources have sustained engraftment in host BM; however, our and other reports indicate that this is so [20–26]. Therefore, it is reasonable to suggest decreased bone regeneration with age may be partially reversed by transplantation of young donor MPCs.

$Wrn^{-/-}Terc^{-/-}$ mutants may have altered BM stroma which permits effective engraftment of donor MPCs. Thus BMT, performed for other reasons in telomere-based accelerated aging syndromes (e.g. aplastic anemia), may also confer beneficial effects in stabilizing or delaying premature osteoporosis. In fact, aplastic anemia in individuals with telomere-based progeroid syndromes may actually occur due to defective (telomerase-deficient) stroma which cannot support HSCs. Whole BMT in these patients may be successful [27] because it theoretically will correct both the stromal as well as the hematopoietic defects.

Conclusions

Replacement of aging MPCs with young cells results in delays or amelioration of aspects of skeletal aging. $Wrn^{-/-}Terc^{-/-}$ recipients of whole BMT have functional reconstitution of MPCs and stable or improved bone microarchitectural features compared to much younger, non-transplanted mice.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Grant support: National Institutes of Health/National Institute on Aging grant R01AG028873 (R.J.P.), University of Pennsylvania Institute on Aging pilot grant award (F.B.J., R.J.P.), Penn Center for Musculoskeletal Disorders P30 AR050950 pilot grant award (R.J.P.).

This work was supported by National Institutes of Health/National Institute on Aging grant R01AG028873 (R.J.P.), University of Pennsylvania Institute on Aging pilot grant awards (F.B.J., R.J.P.) and a Penn Center for Musculoskeletal Disorders pilot grant award (R.J.P.).

References

1. Marie PJ, Kassem M. Extrinsic mechanisms involved in age-related defective bone formation. *J Clin Endocrinol Metab.* 2011; 96:600–609. [PubMed: 21209032]
2. Kassem M, Marie PJ. Senescence-associated intrinsic mechanisms of osteoblast dysfunctions. *Aging Cell.* 2011; 10:191–197. [PubMed: 21210937]

3. Mason PJ, Wilson DB, Bessler M. Dyskeratosis congenita -- a disease of dysfunctional telomere maintenance. *Curr Mol Med*. 2005; 5:159–170. [PubMed: 15974869]
4. Hofer AC, Tran RT, Aziz OZ, et al. Shared phenotypes among segmental progeroid syndromes suggest underlying pathways of aging. *J Gerontol A Biol Sci Med Sci*. 2005; 60:10–20. [PubMed: 15741277]
5. Du X, Shen J, Kugan N, et al. Telomere shortening exposes functions for the mouse Werner and Bloom syndrome genes. *Mol Cell Biol*. 2004; 24:8437–8446. [PubMed: 15367665]
6. Lombard DB, Beard C, Johnson B, et al. Mutations in the WRN gene in mice accelerate mortality in a p53-null background. *Mol Cell Biol*. 2000; 20:3286–3291. [PubMed: 10757812]
7. Chang S, Multani AS, Cabrera NG, et al. Essential role of limiting telomeres in the pathogenesis of Werner syndrome. *Nat Genet*. 2004; 36:877–882. [PubMed: 15235603]
8. Pignolo RJ, Suda RK, McMillan EA, et al. Defects in telomere maintenance molecules impair osteoblast differentiation and promote osteoporosis. *Aging Cell*. 2008; 7:23–31. [PubMed: 18028256]
9. Wang H, Chen Q, Lee SH, et al. Impairment of osteoblast differentiation due to proliferation-independent telomere dysfunction in mouse models of accelerated aging. *Aging Cell*. 2012; 11:704–713. [PubMed: 22621437]
10. Wong KK, Chang S, Weiler SR, et al. Telomere dysfunction impairs DNA repair and enhances sensitivity to ionizing radiation. *Nat Genet*. 2000; 26:85–88. [PubMed: 10973255]
11. Bernardo ME, Zaffaroni N, Novara F, et al. Human bone marrow derived mesenchymal stem cells do not undergo transformation after long-term in vitro culture and do not exhibit telomere maintenance mechanisms. *Cancer Res*. 2007; 67:9142–9149. [PubMed: 17909019]
12. Simonsen JL, Rosada C, Serakinci N, et al. Telomerase expression extends the proliferative life-span and maintains the osteogenic potential of human bone marrow stromal cells. *Nat Biotechnol*. 2002; 20:592–596. [PubMed: 12042863]
13. Yudoh K, Matsuno H, Nakazawa F, et al. Reconstituting telomerase activity using the telomerase catalytic subunit prevents the telomere shorting and replicative senescence in human osteoblasts. *J Bone Miner Res*. 2001; 16:1453–1464. [PubMed: 11499868]
14. Dominici M, Pritchard C, Garlits JE, et al. Hematopoietic cells and osteoblasts are derived from a common marrow progenitor after bone marrow transplantation. *Proc Natl Acad Sci U S A*. 2004; 101:11761–11766. [PubMed: 15282377]
15. Olmsted-Davis EA, Gugala Z, Camargo F, et al. Primitive adult hematopoietic stem cells can function as osteoblast precursors. *Proc Natl Acad Sci U S A*. 2003; 100:15877–15882. [PubMed: 14673088]
16. Bonyadi M, Waldman SD, Liu D, et al. Mesenchymal progenitor self-renewal deficiency leads to age-dependent osteoporosis in Sca-1/Ly-6A null mice. *Proc Natl Acad Sci U S A*. 2003; 100:5840–5845. [PubMed: 12732718]
17. Kasper G, Mao L, Geissler S, et al. Insights into mesenchymal stem cell aging: involvement of antioxidant defense and actin cytoskeleton. *Stem Cells*. 2009; 27:1288–1297. [PubMed: 19492299]
18. Lavasani M, Robinson AR, Lu A, et al. Muscle-derived stem/progenitor cell dysfunction limits healthspan and lifespan in a murine progeria model. *Nat Commun*. 2012; 3:608. [PubMed: 22215083]
19. Lee WY, Cho SW, Oh ES, et al. The effect of bone marrow transplantation on the osteoblastic differentiation of human bone marrow stromal cells. *J Clin Endocrinol Metab*. 2002; 87:329–335. [PubMed: 11788670]
20. Horwitz EM, Prockop DJ, Fitzpatrick LA, et al. Transplantability and therapeutic effects of bone marrow-derived mesenchymal cells in children with osteogenesis imperfecta. *Nat Med*. 1999; 5:309–313. [PubMed: 10086387]
21. Jones OY, Good RA, Cahill RA. Nonmyeloablative allogeneic bone marrow transplantation for treatment of childhood overlap syndrome and small vessel vasculitis. *Bone Marrow Transplant*. 2004; 33:1061–1063. [PubMed: 15048144]
22. Meuleman N, Vanhaelen G, Tondreau T, et al. Reduced intensity conditioning haematopoietic stem cell transplantation with mesenchymal stromal cells infusion for the treatment of

- metachromatic leukodystrophy: a case report. *Haematologica*. 2008; 93:e11–13. [PubMed: 18166768]
23. Pereira RF, O'Hara MD, Laptev AV, et al. Marrow stromal cells as a source of progenitor cells for nonhematopoietic tissues in transgenic mice with a phenotype of osteogenesis imperfecta. *Proc Natl Acad Sci U S A*. 1998; 95:1142–1147. [PubMed: 9448299]
24. Pozzi S, Lisini D, Podesta M, et al. Donor multipotent mesenchymal stromal cells may engraft in pediatric patients given either cord blood or bone marrow transplantation. *Exp Hematol*. 2006; 34:934–942. [PubMed: 16797421]
25. Reyes M, Li S, Foraker J, et al. Donor origin of multipotent adult progenitor cells in radiation chimeras. *Blood*. 2005; 106:3646–3649. [PubMed: 16099880]
26. Villaron EM, Almeida J, Lopez-Holgado N, et al. Mesenchymal stem cells are present in peripheral blood and can engraft after allogeneic hematopoietic stem cell transplantation. *Haematologica*. 2004; 89:1421–1427. [PubMed: 15590390]
27. Ghavamzadeh A, Alimoghadam K, Nasser P, et al. Correction of bone marrow failure in dyskeratosis congenita by bone marrow transplantation. *Bone Marrow Transplant*. 1999; 23:299–301. [PubMed: 10084264]

Figure 1. Overall survival advantage of $Wn^{-/-}Terc^{-/-}$ mice after bone marrow transplantation (BMT). Kaplan-Meier plot of $Wn^{-/-}Terc^{-/-}$ mutants with (n=13) and without (n=8) wild-type whole BMT. $p=0.00035$ by log-rank test.

Figure 2.

Long-term engraftment of Sca-1⁺CD45⁻ MPCs after BMT. Serial bone sections were stained with the indicated antibodies at left. Representative examples of engrafted MPCs in a *Wn^{-/-}Terc^{-/-}* recipient are shown by arrows. Scale bar = 20 μ m.

Figure 3. Donor MPCs differentiate into bone-lining osteoblasts (and subsequently osteocytes) that are incorporated into bone. Arrowheads indicate bone-lining cells. Arrows indicate osteocytes located in their lacunae. Scale bar = 10 μ m.

Figure 4.

Microarchitectural features of skeletal aging are delayed in $Wrn^{-/-}Terc^{-/-}$ mutants after BMT. (Top panels) Representative micro-CT 3-dimensional reconstructions of trabecular and cortical bone in $Wrn^{-/-}Terc^{-/-}$ mutants without (n=6) and with BMT (n=6) are shown. (Bottom panels) Quantification of trabecular and cortical bone parameters in $Wrn^{-/-}Terc^{-/-}$ mutants demonstrates preservation of microarchitectural features with BMT. Note that BMT recipients were about 30% older than non-BMT controls. BV, bone volume; TV, total volume; Tb.N, trabecular number; Ct Th, cortical thickness; Ct.Ar, cortical area; Tt.Ar, total area. ****, $p < 0.002$.