CORRESPONDENCE

The Care of Preterm Infants With Birth Weight Below 1250 g: Risk-Adjusted Quality Benchmarking as Part of Validating a Caseload-Based Management System

by Dr. rer. medic. Marcus Kutschmann, Dr. med. Sven Bungard, Joachim Kötting, Andrea Trümner, Prof. Dr. med. Christoph Fusch and Dr. med. Christof Veit in volume 31–32/2012

Room for Improvement

The publication (1) shows that the mortality risk for preterm infants with a birth weight below 1250 g is considerably and significantly lower in centers treating 30 or more cases every year than in centers with fewer than 30 cases/year. If all preterm infants had been treated in such hospitals then mortality for the entire group would likely have been lower. This primary result must not be overshadowed by more detailed results.

The heterogeneity of hospitals with 30 or more cases is, however, considerable. If the quality results were known for individual hospitals, it might be assumed that—if timely data were available and quality results were consistent over time—then, as a second step, overall mortality would drop to even lower levels.

If the quality results regarding mortality and morbidity could be modeled with greater accuracy (for example by improved risk adjustment, as suggested by Vogtmann et al. [2]), then a further improvement in the results (third step) could be expected, if cases were concentrated in higher-quality hospitals.

The primary result, that centers with more than 30 cases/year on average achieve better results, has the advantage of being generally useful in terms of the management and steering of service provision. In contrast the improvement potentials, if more accurate quality results were obtained and known for individual hospitals, could be more difficult to realize, as e.g. the results of individual centers are subject to uncertain changes over time. Other generalizable characteristics should also be determined (for example, characteristics of the structural quality, such as provision of certain intervention options, qualified staff, etc). It can be difficult to show that such characteristics are reliably associated with adequately risk-adjusted quality results, and infer causality on this basis. In a scenario where such characteristics are lacking, Kutschmann et al. (1) at least showed that the number of cases is a relevant criterion.

The Federal Joint Committee (*Gemeinsamer Bundesausschuss*, G-BA) was well advised to set a minimum caseload, which ensures centralization and better quality care – and therefore the likelihood of survival of preterm infants.

DOI: 10.3238/arztebl.2013.0116a

REFERENCES

1. Kutschmann M, Bungard S, Kötting J, Trümner A, Fusch C, Veit C: The care of preterm infants with birth weight below 1250 g: risk-adjusted quality benchmarking as part of validating a caseload-based management system. *Dtsch Arztebl Int* 2012; 109(31–32): 519–26.
2. Vogtmann C, Koch R, Gmyrek D, Kaiser A, Friedrich A: Risk-adjusted intraventricular hemorrhage rates in very premature infants – towards quality assurance between neonatal units. *Dtsch Arztebl Int* 2012; 109(31–32): 527–33.

Dipl.-Med. Hans-Werner Pfeifer, Leiter Referat Qualitätssicherung GKV-Spitzenverband, Berlin, Hans-Werner.Pfeifer@gkv-spitzenverband.de

Conflict of interest statement

The author declares that no conflict of interest exists according.

One-Sided Calculation

The study reported by Kutschmann et al (1) shows that the mortality risk for very low birth weight infants is 34% higher in hospitals with less than 30 preterm infants < 1250 g birth weight per year compared to hospitals with higher patient volumes. Since the risk adjusted mortality in many small centers was below average, whereas it was above average in some larger centers, the authors conclude that the caseload is not a suitable indicator of the quality of care. According to the laws of stochastic processes, however, the wide variance of mortality in small centers may actually be the result of the small caseload per center. The lower mortality rate in some small hospitals over a time period of three years may thus be merely an effect of a lower caseload, not an argument against the regionalization of care for very premature infants, which makes sense for various other reasons too.

In spite of risk adjustment, mortality was systematically calculated to the disadvantage of perinatal centers.

Not including preterm infants admitted more than 24 hours after the birth or those referred to other units (17.5% of the cohort) seemingly leads to lower mortality rates in smaller centers (preterm infants that were transferred to larger centers mostly have severe complications) and a seeming increase in mortality in larger centers (since it is mostly healthy and stable babies that are referred to smaller centers from larger ones).

In adjusting the risk, parameters were considered that include poor basic care given to preterm infants (baseline deficit, maximum FiO₂) as a risk-increasing factor. These variables reflect, however, the competence of the obstetric and neonatology team; including these into the risk adjustment blurs differences in the treatment quality of different centers.

Unfortunately, the differences in mortality were not stratified by gestational age, although the differences between large and small centers have been demonstrated to increase with falling gestational age (2–4). Such stratification might have provided some answers

to the question of the gestational age threshold below which stronger regionalization of the care of preterm infants might be effective.

DOI: 10.3238/arztebl.2013.0116b

REFERENCES

1. Kutschmann M, Bungard S, Kötting J, Trümner A, Fusch C, Veit C: The care of preterm infants with birth weight below 1250 g: risk-adjusted quality benchmarking as part of validating a caseload-based management system. *Dtsch Arztebl Int* 2012; 109(31–32): 519–26.
2. Bartels DB, Wypij D, Wenzlaff P, Damann O, Poets CF: Hospital volume and neonatal mortality among very low birth weight infants. *Pediatrics* 2006; 117: 2206–14.
3. Teig N, Wolf HG, Bücker-Nott HJ: Mortalität bei Frühgeborenen < 32 Schwangerschaftswochen in Abhängigkeit von Versorgungsstufe und Patientenvolumen in Nordrhein-Westfalen. *Z Geburtshilfe Neonatol* 2007; 211: 118–22.
4. Trotter A, Pohlandt F: Aktuelle Ergebnisqualität der Versorgung von Frühgeborenen < 1500 g Geburtsgewicht als Grundlage für eine Regionalisierung der Risikogeburten. *Z Geburtshilfe Neonatol* 2010; 214: 55–61.

Dr. med. Norbert Teig, Abteilung für Neonatologie und pädiatrische Intensivmedizin, Universitätskinderklinik und Perinatalzentrum Bochum
norbert.teig@ruhr-uni-bochum.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Unanswered Questions

The study reported by Kutschmann and colleagues (1) of the discriminative ability of a minimum caseload of 30 cases per year shows that minimum caseload reflects differences in the quality of care given by different hospitals to an unsatisfactory degree.

However, the study has methodological weaknesses. In calculating the logistic regression model, the authors did not consider the different levels of data (individual level and hospital level), and cluster effects were not corrected for. This may result in an overestimate of the association between caseload and mortality; for this reason, a multilevel model would have been more suitable (2).

Furthermore, some questions remain unanswered after reading the article. In addition to the effect of caseload on the mortality of preterm infants, it would have been interesting to investigate the extent to which the caseload of < 30 is even of relevance in explaining the different mortality rates at different hospitals. To this end one would have to calculate the proportion of overall mortality that is due to hospital-specific factors and not to individual factors, and determine how much of the proportion of hospital-specific factors is explained by the caseload. This can also be shown in a multilevel model (3). The relevance of several other quality indicators mentioned in the article—such as staffing structures and equipment—could also be captured in such a model.

Furthermore it would have been interesting not only to investigate the discriminative ability of a minimum caseload of 30 cases/year but also the degree to which a useful threshold value can be deduced at all. The possible model for this could be a study by the Institute

for Quality and Efficiency in Health Care (*Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen, IQWiG*) of the threshold values for total knee joint endoprosthesis (4).

DOI: 10.3238/arztebl.2013.0117a

REFERENCES

1. Kutschmann M, Bungard S, Kötting J, Trümner A, Fusch C, Veit C: The care of preterm infants with birth weight below 1250 g: risk-adjusted quality benchmarking as part of validating a caseload-based management system. *Dtsch Arztebl Int* 2012; 109(31–32): 519–26.
2. Hox JJ: Why do we need special multilevel analysis techniques. *Multilevel Analysis: Techniques and Applications*. New Jersey: Lawrence Erlbaum Associates 2002.
3. Snijders TAB, Bosker RJ: Discrete dependent variables: In: *Multilevel analysis. An introduction to basic and advanced multilevel modeling*. 2nd edition. London: Sage Publications 2012; 289–322.
4. IQWiG: Entwicklung und Anwendung von Modellen zur Berechnung von Schwellenwerten bei Mindestmengen für die Knie-Totalendoprothese. Abschlussbericht B05/01a. Köln: Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG), 2005.

Dr. med. Melanie Eßer, MPH
Bayerische Arbeitsgemeinschaft für Qualitätssicherung, München

Prof. Dr. med. Rüdiger von Kries, Institut für Soziale Pädiatrie und Jugendmedizin, Ludwig-Maximilians-Universität München

Prof. Dr. rer. nat. Ulrich Mansmann
Institut für Medizinische Informationsverarbeitung, Biometrie und Epidemiologie, Ludwig-Maximilians-Universität München

Conflict of interest statement

The authors declare that no conflict of interest exists.

Quality Analysis Is Overdue

Kutschmann et al (1) analyzed neonatal data from 7400 preterm infants from Bavaria, Baden-Württemberg, Hesse, Lower Saxony, and North Rhine–Westphalia for the years 2007–2009 and found that the risk-adjusted mortality risk in smaller centers with a lower annual caseload of less than 30 preterm babies with a birth weight of less than 1500 g was significantly higher than in centers with a higher caseload (odds ratio 1.34). This result confirms earlier results from Lower Saxony. Bartels et al. (2) studied the mortality rates of all live-born infants. The study reported by Kutschmann et al. (1) analyzed only treatment data submitted by the hospitals (admissions to pediatric centers that were reported to the neonatal units). In their discussion they refer to other authors who compared a similar dataset with the official birth statistics and found that 33% of deaths in preterm babies with a birth weight below 1000 g were not included in the neonatal unit reports (3) and they state that this data gap had been closed since 2010. However, this latter assumption does not seem to be correct, because in Baden-Württemberg, for example, only 56 of the 99 liveborn preterm infants included in the official statistic in 2010 had been reported to the neonatal units. According to the official statistics, 28 of these babies died, but the neonatal units reported only 13 deaths. In Bavaria in 2010, only 34 of 69 officially registered preterm babies with a birth weight below 500 g were included in the neonatal unit reports; of the 39 officially registered deaths, only 15 made it

into the neonatal reports. A look at the Scandinavian countries shows that such data capture is feasible, improves the data quality, and can provide relevant insights.

DOI: 10.3238/arztebl.2013.0117b

REFERENCES

1. Kutschmann M, Bungard S, Kötting J, Trümner A, Fusch C, Veit C: The care of preterm infants with birth weight below 1250 g: risk-adjusted quality benchmarking as part of validating a caseload-based management system. *Dtsch Arztebl Int* 2012; 109(31–32): 519–26.
2. Bartels DB, Wypij D, Wenzlaff P, Dammann O, Poets CF: Hospital volume and neonatal mortality among very low birth weight infants. *Pediatrics* 2006; 117: 2206–14.
3. Hummler HD, Poets C: Mortalität sehr unreifer Frühgeborener – Erhebliche Diskrepanz zwischen Neonatallerhebung und amtlicher Geburten-/Sterbestatistik. *Z Geburtshilfe Neonatol* 2011; 215: 10–7.
4. The EXPRESS Group: One-year survival of extremely preterm infants after active perinatal care in Sweden. *JAMA* 2009; 301: 2225–33.

Prof. Dr. med. Helmut Hummler

Sektion Neonatologie und Pädiatrische Intensivmedizin
Klinik für Kinder- und Jugendmedizin, Universitätsklinikum Ulm
helmut.hummler@uniklinik-ulm.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

The aim of any quality-based coordination of care has to be high-quality, supraregionally homogeneous healthcare. If no quality data are available then agreeing a minimum caseload is at the very least a protective measure against the risks entailed in ad-hoc provision of care, which in critical situations provides neither the necessary experience and routines nor the necessary structures and processes. This is confirmed by the significantly better quality of care offered by larger departments. However, wherever quality data are available thanks to mandatory documentation by neonatal units, it is not clear why one should rely on caseload alone. The results show how differentiated the outcome quality above a minimum caseload—using the example of mortality—has to be assessed. Concentration alone does not seem a convincing concept. Including three years of results permits a classification of the quality of care delivered by individual centers that is robust enough to accommodate accidental fluctuations.

We agree with Pfeifer as well as Vogtmann et al. (1) that the necessary risk adjustment should not be limited to the parameters gestational age and severe malformations. Our analyses showed nine risk factors that are relevant for the adjustment model. These include, after thorough consideration, administration of FiO₂ and base excess at admission—parameters that at this early stage reflect the condition of the preterm infants rather than the care given by the center and can therefore be counted as risk factors.

Using a logistic regression model in the non-hierarchical form aimed to adjust for all factors that reflect exclusively the risk profile of the preterm infants

and are not affected by the care given in the centers. By including the centers in a multilevel model we would have corrected for the non-specific but present effect of hospital care. Such an approach would have run counter to our study's objective—namely, to conduct a risk-adjusted comparison between hospitals. Undoubtedly it would be interesting to investigate the way in which factors such as staffing structures, equipment, the organization, process and internal quality management, and the quality culture affect mortality in preterm infants and how these correlate to the caseload. But this information was not available to us. However, one objective of our study was to evaluate, for the given data, the potential use of neonatal data. Looking for a threshold value would make sense only if a minimum quality standard was defined and if, in the context of theoretical modeling to date, a monotonically decreasing or increasing effect on the result could be assumed. The direct evaluation of individual centers according to the outcome quality that was actually achieved on the basis of the available neonatal data seemed the more pragmatic approach. The extent to which the data for 2010—when the new survey of neonatal data started—are sufficiently complete is not known to us as we do not have access to these data. This will need to be clarified in the coming years by taking birth records into account.

The exclusion of preterm infants admitted to neonatal units more than 24 hours after birth and of infants referred to other units was unavoidable. Children admitted 24 hours after birth were likely to be pretreated. Therefore, a clear assignment of treatment quality to a single neonatal unit was not possible. Moreover, including children admitted more than 24 hours after birth would have been likely to give rise to the accusation of artificially increasing mortality rates in large centers with these high-risk infants. However, mortality cannot be calculated for infants referred to other units. Therefore, the risk-adjusted comparison between preterm infants exclusively treated in one center seemed to us the most clearly defined and therefore fairest approach.

DOI: 10.3238/arztebl.2013.0118

REFERENCES

1. Vogtmann C, Koch R, Gmyrek D, Kaiser A, Friedrich A: Risk-adjusted intraventricular hemorrhage rates in very premature infants – towards quality assurance between neonatal units. *Dtsch Arztebl Int* 2012; 109(31–32): 527–33.
2. Kutschmann M, Bungard S, Kötting J, Trümner A, Fusch C, Veit C: The care of preterm infants with birth weight below 1250 g: risk-adjusted quality benchmarking as part of validating a caseload-based management system. *Dtsch Arztebl Int* 2012; 109(31–32): 519–26.

Dr. rer. medic. Marcus Kutschmann, Dipl.-Stat., Dr. med. Christof Veit
BQS Institut für Qualität und Patientensicherheit,
m.kutschmann@bqs-institut.de

Conflict of interest statement

The BQS Institute for Quality and Patient Safety (BQS Institut für Qualität & Patientensicherheit) received financial support from the German Hospital Association (DKG, Deutsche Krankenhausgesellschaft e. V.) for the analysis on which the authors' reply is based.

Dr Kutschmann and Dr Veit are employees of the BQS Institute for Quality and Patient Safety.