Transcription Factor GATA-4 Regulates Cardiac Muscle-Specific Expression of the α -Myosin Heavy-Chain Gene

JEFFERY D. MOKKENTIN,^{1,2,3} DHANANJAYA V. KALVAKOLANU,^{4,5} AND BRUCE E. MARKHAM^{1,2,3*}

Departments of Physiology¹ and Microbiology,⁴ Cardiovascular Research Center,² Cancer Center,⁵ and Molecular and Cellular Biology Training Program,³ Medical College of Wisconsin, Milwaukee, Wisconsin 53226

Received 18 January 1994/Returned for modification 1 March 1994/Accepted 29 April 1994

The α -myosin heavy-chain (α -MHC) gene is the major structural protein in the adult rodent myocardium. Its expression is restricted to the heart by a complex interplay of *trans*-acting factors and their *cis*-acting sites. However, to date, the factors that have been shown to regulate expression of this gene have also been found in skeletal muscle cells. Recently, transcription factor GATA-4, which has a tissue distribution limited to the heart and endodermally derived tissues, was identified. We recently found two putative GATA-binding sites within the proximal enhancer of the α -MHC gene, suggesting that GATA-4 might regulate its expression. In this study, we establish that GATA-4 interacts with the α -MHC GATA sites to stimulate cardiac muscle-specific expression. Mutation of the GATA-4-binding sites either individually or together decreased activity by 50 and 88% in the adult myocardium, respectively. GATA-4-dependent enhancement of activity from a heterologous promoter was mediated through the α -MHC GATA sites. Coinjection of an α -MHC promoter construct with a GATA-4 expression vector permitted ectopic expression in skeletal muscle but not in fibroblasts. Thus, the lack of α -MHC expression in skeletal muscle correlates with a lack of GATA-4. GATA-4 DNA binding activity was significantly up-regulated in triiodothyronine- or retinoic acid-treated cardiomyocytes. Putative GATA-4-binding sites are also found in the regulatory regions of other cardiac muscle-expressed structural genes. This indicates a mechanism whereby triiodothyronine and retinoic acid can exert coordinate control of the cardiac phenotype through a *trans*-acting regulatory factor.

The myosin heavy chain (MHC) is the major contractile protein in the heart. While there are several MHC isoforms encoded by a multigene family, only α -MHC and β -MHC are expressed in cardiac muscle cells. In developing rodents, α -MHC mRNA is expressed between 7.5 and 8 days postcoitum in the cardiac tube but diminishes as the ventricular chambers form between 8 and 9 days postcoitum and remains at low levels until just after birth (24, 25). β -MHC is the predominant MHC isoform in the ventricles during fetal development. At birth, a time coincident with a dramatic increase in circulating thyroid hormone, α -MHC levels increase to as much as 95% of the total MHC in the heart and remain high throughout normal adult life (24).

The α -MHC gene is developmentally and hormonally regulated at the transcriptional level by a complex interplay of multiple elements (12, 24, 33, 34). Elements sufficient to direct significant expression in cardiac myocytes are located in the proximal regulatory region (from position –374 to –39) even though upstream elements contribute to the overall regulation of this gene (4, 12, 37, 48). These sequences are also sufficient to maintain tissue-restricted expression. Within the proximal enhancer-promoter, a number of elements that participate in transcriptional control have been identified. These include a site that interacts with myocyte-specific enhancer-binding factor (MEF-2) (19, 34), an E-box motif (EB1) that binds α -MHC binding factor 2 (BF-2) (33), sites that interact with an M-CAT binding factor (MCBF) and an A-rich binding factor (ARF) (36), and a second E-box and a CArG box (35). In addition, triiodothyronine (T_3) regulates α -MHC gene expression through the interaction of the thyroid regulatory element with

thyroid hormone receptor (9, 11–13, 16, 29). These sites interact with regulatory factors that appear to be present in several cell types. However, significant expression of the α -MHC gene is restricted to atrial and ventricular cardiomyocytes and a layer of the lung referred to as the pulmonary myocardium (45).

Cardiac muscle-specific expression of the myosin light chain-2 (MLC-2) gene is dependent upon a 28-bp region (HF-1) that interacts with factors which also appear to be present in several cell types (21, 38, 55). The MBF-2 region of the rat α -MHC gene interacts with factors (MEF-2 and BF-2) that are similar if not identical to those that interact with the MLC-2 HF-1 region (34). Thus, these factors may represent a regulatory mechanism of cardiac muscle-specific expression that is common between these and other genes. However, expression of the α -MHC gene in cardiomyocytes is not strictly dependent upon MEF-2, since a reconstituted α -MHC promoter did not require this site for cardiac muscle-specific expression (33). This suggests that additional elements participate in the tissue-restricted expression of the α -MHC gene.

Recently, a family of GATA DNA binding proteins that interact with DNA sequence elements that contain the conserved motif 5'-GATA-3' have been identified. Molecular cloning of the cDNAs encoding the GATA binding proteins has demonstrated that these proteins contain one or two zinc finger DNA binding domains (2, 22, 47, 52). A novel member of this family, GATA-4, was found to be expressed in the heart and in endodermally derived tissues (2). GATA-4 is a tissue-specific, retinoic acid (RA)-inducible, and developmentally regulated transcription factor. GATA-4 is present in the *Xenopus* presumptive cardiac ventral mesoderm prior to formation of the cardiac tube (18) and is present in the mouse embryo at a similar stage (49). The early expression of GATA-4 in the heart suggests a possible role for this regula-

* Corresponding author. Mailing address: Department of Physiology, Medical College of Wisconsin, 8701 Watertown Plank Rd., Milwaukee, WI 53226. Phone: (414) 266-5617. Fax: (414) 257-8570.

TABLE 1. Oligonucleotides used in this study

Oligonucleotide	Base sequence ^a
PRE-D.....	5'-TCCTGTCTCTCTATCTGCCCATCGGCCCTTTG-3'
D mut.....	5'-TCCTGTCTCTCTCTgcaTGCCCTgaGGCCCTTTG-3'
D5' mut.....	5'-TCCTGTCTCTCTCTgcaTGCCCATCGGCCCTTTG-3'
D3' mut.....	5'-TCCTGTCTCTCTCTATCTGCCCTgaGGCCCTTTG-3'
D site.....	5'-TCTATCTGCCCATCGGCC-3'
MCS.....	5'-TCGCGACTCGAGCCCGGATCGATGCGCCGCCACGTG-3'
BF-2.....	5'-AGGGCCATGTGGGTAG-3'
PRE-B.....	5'-AGGGCCATGTGGGTAGGGAGGTGGTGTGAGACGG-3'
SBS.....	5'-TCAAAGGGAAACTGAGTCATGCAC-3'
PCR anti.....	5'-GCTTGGGATGGAGAAATCTCGG-3'

^a Mutated bases are indicated by lowercase letters.

tory factor in cardiogenesis (2, 18). However, no target promoter has been identified.

We report here that the α -cardiac MHC gene is a target for transcription factor GATA-4. Two sites in the α -MHC enhancer bound to a factor in cardiomyocyte nuclear extracts that was indistinguishable from GATA-4 as indicated by DNA binding specificity and interactions with antibody against GATA-4. GATA-4 DNA binding activity was increased in cardiomyocytes treated with RA or T₃. Mutation of the two GATA-4 binding sites individually or together decreased activity by approximately 50 and 88%, respectively, when activity in the adult rat heart was assayed. The two GATA-4 sites were capable of mediating GATA-4-regulated transcription from a heterologous promoter. Coinjection of a GATA-4 expression vector with an α -MHC-reporter construct into skeletal muscle resulted in a significant induction of activity, indicating that the lack of α -MHC gene expression in skeletal muscle is due, in part, to a lack of GATA-4. These results indicate that GATA-4 plays an important role in the cardiac muscle-specific expression of the α -MHC gene. Since GATA-4 is expressed prior to α -MHC in the early stages of cardiac development, our results suggest the possibility that GATA-4 is responsible, in part, for the transient expression of the α -MHC gene in the embryonic heart tube. The presence of GATA-4 in the embryonic heart and the demonstration of its ability to regulate cardiac muscle-specific expression support the hypothesis that GATA-4 plays an important role in cardiac development.

MATERIALS AND METHODS

Oligonucleotides. The sequence of each oligonucleotide used in this study is shown in Table 1. The sense strand is shown unless otherwise indicated, although for binding studies, the complementary strand was also used. All oligonucleotides were purchased from Operon Technologies (Alameda, Calif.). PRE-D spans a sequence from position -283 to -249 from the rat α -MHC promoter (26). PCR anti contains sequences from position +17 to -5. The PRE-B, MCS, and SBS oligonucleotides have been reported previously (28, 33). Other oligonucleotides contain subsets of the PRE-D sequence or mutations of PRE-D.

Cell culture, transfection, and reporter gene assays. The conditions for isolation and culturing of 1- to 2-day-old neonatal rat cardiomyocytes were as described previously (33). Cardiomyocytes were cultured in serum-free Ham's F12 medium supplemented as previously described (11). After 24 h in culture, cardiomyocytes remained untreated or were treated with all-*trans* RA or T₃ to final concentrations of 100 and 10 nM, respectively. The mouse lung fibroblast cell line L929 was cultured in Dulbecco's modified Eagle's medium supple-

mented with 10% fetal bovine serum. DNA transfections were carried out by using a modified calcium phosphate method (5). Equimolar ratios of reporter and effector DNAs were transfected. To correct for variations in the transfection process, 2 μ g of plasmid pRSVCAT was cotransfected in each case. Cells (5×10^5) were trypsinized, mixed with the calcium phosphate: DNA precipitates, and plated in 100 mM culture dishes. Medium (3 ml) was added to each plate, and the plates were incubated at 37°C for 24 h in a CO₂ incubator. Subsequently, the cells were shocked with 15% dimethyl sulfoxide, prepared in Hanks' balanced salt solution, for 90 s and washed thoroughly. Fresh medium (10 ml) was added to the plates, and incubation was continued for an additional 24 h. Cells were harvested 48 h posttransfection and lysed, and luciferase and chloramphenicol acetyltransferase (CAT) assays were performed as described below.

Site-directed mutagenesis and reporter gene construction. Mutations were introduced into specific sites in the 3,300 bp of the α -MHC gene present in plasmid pSV0MCAT (12). Mutagenesis was performed by the rolling-circle PCR mutagenesis method (14). Conditions for mutagenesis were as described previously, and the subsequent analysis of the mutant constructs was performed as described previously (33). The oligonucleotide D5' mut, D3' mut, or D5'-3' mut was used as the forward PCR primer to introduce mutations into pSV0MCAT. Antisense oligonucleotide PRE-B was used as the reverse PCR primer in the rolling-circle mutagenesis procedure (Table 1).

Construct pMHC598 was created by blunt-end ligation of a 615-bp PCR fragment derived from the rat α -MHC promoter (from position -598 to +17) into the pGL basic luciferase reporter plasmid (Promega). The PCR mixture contained 1 U of Vent DNA polymerase, 1 $\times$ Vent buffer (New England Biolabs, Beverly, Mass.), 100 pmol each of phosphorylated SBS sense and PCR anti oligonucleotides (Table 1), 200 μ M deoxynucleoside triphosphates, and 10 ng of pSV0MCAT template DNA. Reaction conditions consisted of denaturation at 96°C for 30 s, annealing at 60°C for 30 s, and extension at 72°C for 60 s and for 35 cycles.

Eukaryotic expression vectors pMT₂ and pMT₂-GATA4 were a gift of David Wilson and are described elsewhere (2). pGLPREdx2 was generated by blunt-end ligation of two double-stranded (ds) PRE-D oligonucleotides 40 bp upstream from the simian virus 40 early promoter at the *Sma*I site in the pGL-promoter luciferase plasmid (Promega). The insert integrity, sequence, and orientation of each clone was confirmed by DNA sequencing.

In vivo gene transfer by direct injection. Clones to be assayed for activity were injected directly into the left apex of the adult heart as previously described (34). Briefly, male Sprague-Dawley rats (225 to 250 g) were anesthetized with

0.15 ml of a Ketaset-acepromazine mixture (Ketaset [100 mg/ml], 10 ml, and promACE [Aveco; 10 mg/ml], 2.2 ml) per 100 g of body weight and ventilated with a respirator. A left lateral thoracotomy was performed to expose the beating heart. An 80- μ g portion of the construct pSV0MCAT, pSV0MCAT-D3' mut, pSV0MCAT-D5' mut, or pSV0MCAT-D5'-3' mut was injected along with 10 μ g of pRSV β gal (for transfection normalization) in 80 μ l of saline. Injections were performed with a 28-gauge needle. DNA was also injected into the tongues of the animals receiving the cardiac injections to assay for expression in skeletal muscle. Immediately after the surgical procedure the animals were given penicillin G at 30,000 U/100 g of body weight and allowed to recover for 6 days. After 6 days, rats that had received cardiac or skeletal muscle injections were sacrificed by lethal injection of pentobarbital, and the tissues were collected.

A similar protocol was used for coinjection experiments when pMHC598 and pMT₂ or pMT₂-GATA-4 were assayed for expression in the heart and skeletal muscle. An 80- μ g portion of pMHC598 and 16 μ g of either pMT₂ or pMT₂-GATA-4 were injected along with 10 μ g of pRSVCAT in 80 μ l of saline. After 6 days, rats that had received injections were sacrificed as described above, and the tissues were collected.

Luciferase, CAT, and β -galactosidase assays. For injection experiments with the CAT reporter gene, extracts were prepared essentially as described previously (34). Heart extracts were prepared by mincing the tissue in 600 μ l of nuclear lysis buffer and then subjecting it to two 10-s bursts of sonication with a Branson Sonifier 250 with a microtip probe at a setting of 1 (Branson Ultrasonics, Danbury, Conn.). Cellular debris was pelleted by centrifugation at $10,000 \times g$ for 10 min at 4°C, and CAT and β -galactosidase assays were performed as described previously (10, 40).

For injection experiments using the luciferase reporter gene, extracts were prepared by mincing heart or skeletal muscle extract in 900 μ l of cell lysis buffer (1% Triton X-100, 100 mM Tris-Cl [pH 7.8], 2 mM EDTA, 2 mM dithiothreitol, 2 μ g of aprotinin per ml, 2 μ g of leupeptin per ml, 1 μ g of pepstatin per ml, 100 μ g of phenylmethylsulfonyl fluoride per ml) and then homogenizing it in a 4-ml-volume Dounce homogenizer (up and down 10 to 12 times) at 4°C until the tissues were well dispersed. Cellular debris was pelleted by centrifugation at $10,000 \times g$ for 10 min at 4°C. To assay luciferase activity in L929 cells, plates were washed in phosphate-buffered saline and the cells were scraped into 200 μ l of cell lysis buffer. The cellular debris was removed by centrifugation, and the supernatant was assayed. Protein concentration was determined by the method of Bradford (3). Luciferase assays were performed as previously described (7) with minor modifications. Cardiac protein extract (400 μ g), skeletal muscle protein extract (1,000 μ g), or 50 μ l of L929 cell extract was added to 300 μ l of luciferase reaction buffer (4 mM ATP [pH 7.0], 30 mM Tricine [pH 7.8], 15 mM MgSO₄, 10 mM dithiothreitol) in clear tubes (12 by 47 mm) suitable for luminometry. The samples were placed in a luminometer (EG&G Berthold model LB 953 distributed by Wallace, Gaithersburg, Md.), and 100 μ l of 1 mM D-luciferin was injected and mixed with them. Data were collected over 4-s intervals.

Preparation of nuclear extracts and in vitro transcription and translation of GATA-4. Nuclear extracts were prepared from cultured cardiomyocytes as described previously (8) with minor modifications (33). Extracts from COS cells and COS cells transfected with pMT₂-GATA-4 were a gift from David Wilson. For preparation of GATA-4 in vitro, the 1.9-kb GATA-4 insert in pMT₂-GATA-4 was removed by *Eco*RI digestion and cloned into the *Eco*RI site in pBluescript SK⁻

(Stratagene). Linearized template was used to prepare GATA-4 mRNA by using an in vitro transcription kit (Stratagene) according to the manufacturer's recommendations. RNA was translated by using an in vitro translation kit (Stratagene).

EMSAs and dimethyl sulfate (DMS) and diethyl pyrocarbonate (DEPC) interference assays. Electrophoretic mobility shift assays (EMSAs) were performed as described earlier (33). Unless otherwise indicated, 12 μ g of protein extract and 2 μ g of poly(dI-dC) · (dI-dC) were used in each assay. The probes used in each experiment are indicated in the figure legends. Reaction mixtures were allowed to incubate at room temperature for 15 min. The samples were loaded on a nondenaturing acrylamide gel (5% acrylamide, 0.083% bisacrylamide, 0.5 $\times$ Tris-borate-EDTA) and were run at 25 mA for 2 h. The gel was dried and autoradiographed. Supershift experiments were performed exactly as described for a standard EMSA except that 1 to 5 μ l of antiserum was added immediately after the protein extract was added.

DEPC footprinting was performed essentially as described previously (44). DEPC-treated probe was used in reactions described for gel shift experiments except that the reactions were scaled up 25-fold. Complex isolation and electrophoresis conditions are described in reference 33. For methylation interference experiments, radioactive sense and antisense strands of PRE-D were methylated with DMS (30), annealed with an equal amount of an unlabeled opposite strand, and used in a 25-fold scaleup of gel shift conditions described above. Unbound probe and the band corresponding to the specific interaction were excised, electroeluted from the gel, ethanol precipitated, washed, and dried. Isolated DNA was treated with piperidine and run on a 20% polyacrylamide-8.3 M urea gel. Gels were exposed to X-ray film for 12 to 18 h.

RESULTS

Identification of factors in cardiomyocyte nuclear extract that interact with PRE-D. PRE-D (Fig. 1A) was identified during experiments to catalogue regions of the α -MHC proximal promoter that interacted with a factor or factors present in nuclear extracts prepared from primary cultured neonatal rat cardiomyocytes. EMSAs suggested that this ds oligonucleotide interacted with one or more factors (Fig. 1B). Competition experiments demonstrated that a 50-fold molar excess of unlabeled PRE-D, but not an equal amount of an unrelated ds oligonucleotide (MCS), competed for binding of the factors that formed the complex (GATA-4; see below) seen in Fig. 1B. To further characterize the interaction with PRE-D, we performed DMS and DEPC interference footprint analysis. The DMS footprint demarcated an interaction on the antisense strand over an 18-bp sequence from position -253 to -270 in the α -MHC promoter (Fig. 2A). Methylation of each guanine residue in this region, with the exception of the one at position -261, interfered with factor binding. No interaction was seen on the sense strand. Since the DMS footprint on the antisense strand occurred close to the 5' end, it was not clear whether the diminished signal at the 5' end was due to interference or to loss of the small fragment during processing. To test this, a ds oligonucleotide (D site) that lacked the G at position -253 was prepared. The D site oligonucleotide did not compete for factor binding as well as the PRE-D oligonucleotide, suggesting that the G at position -253 is part of the binding site (Fig. 1B). The DEPC footprint further delineated the interaction(s), implicating all of the A residues on both strands of the promoter between positions -253 and -270 in factor binding. The footprints are shown schematically in Fig. 2B. The faster-

FIG. 1. Identification of a factor that interacts with PRE-D in a sequence-specific manner. (A) DNA sequence of the antisense strand of the α -MHC gene proximal regulatory element D (PRE-D) spanning positions -283 to -249. The boxes represent the two putative binding sites for GATA-4. D3' mut and D5' mut are identical to PRE-D except at the underlined bases. D site is an oligonucleotide containing a subset of PRE-D sequences. (B) EMSA of 32 P-labeled PRE-D ds oligonucleotide and cardiomyocyte nuclear protein extract. The competitor oligonucleotides indicated were added at a 50:1 molar excess.

migrating complex seen in Fig. 1B footprinted identically to the slower-migrating complex (data not shown) in a manner consistent with its being a cleavage product of a protein that formed the slower-migrating complex.

Visual analysis of the footprinted region suggested some symmetry of binding. On the antisense strand, beginning at position -254, the footprinted guanines were separated from the footprinted G, A, and A on the sense strand by 2 bp. This same pattern was seen to repeat beginning at position -262. Also, the more upstream portion of the footprint was positioned over the sequence 5' AGATAG 3', which is a perfect consensus site for the transcription factor GATA-4 (2). A second possible binding site for GATA-binding proteins, 5' GATG 3', was present in the downstream portion of the footprint (20, 31).

Assuming that the footprinted region contained two GATA-4 sites, we made ds oligonucleotides that were identical to PRE-D except that the individual GATA sites were mutated. These two oligonucleotides were used at a 50-fold molar excess to compete for factor binding to the intact PRE-D probe in EMSAs (Fig. 1B). While both competed somewhat for factor binding, neither of the mutant ds oligonucleotides

FIG. 2. (A) DMS and DEPC interference footprints of the protein-DNA interaction between the PRE-D oligonucleotide and a factor present in cardiomyocyte nuclear extracts. B, bound oligonucleotide fraction; F, free fraction. The asterisks show the positions of the bases that interfere with factor binding. (B) Schematic representation of the DMS and DEPC footprints. Open circles represent bases identified by DEPC interference footprinting while closed circles represent bases identified by both DEPC and DMS footprints.

was as effective as intact PRE-D in competition experiments. The one containing the intact upstream site (D3' mut) was consistently a better competitor than the one containing the downstream site (D5' mut) (Fig. 1), suggesting that the upstream site may have a higher affinity for factor binding than the downstream site. At a 200-fold molar excess, all of the competitors shown in Fig. 1B, with the exception of MCS, competed equally well for factor binding (data not shown).

GATA-4 interacts with two sites in the α -MHC proximal enhancer-promoter. To determine if the factor interacting with the GATA sites within PRE-D was GATA-4, preimmune serum and antiserum directed against mouse GATA-4 were obtained and used in supershift EMSAs. The anti-GATA-4 serum did not cross-react significantly with other members of the GATA-binding protein family (35, 49). Cardiomyocyte nuclear extract was probed with PRE-D in the presence of preimmune or anti-GATA-4 serum. The resulting complexes were resolved on an EMSA gel (Fig. 3). In the presence of immune serum, the band labeled GATA-4 was diminished, with corresponding increases in two supershifted bands. Five microliters of antiserum completely shifted the reaction, indicating that the complex migrating to the GATA-4 position was completely dependent upon the presence of GATA-4 or a GATA-4-related protein. The complex was not affected by the presence of preimmune serum.

To further compare GATA-4 activity with the GATA binding activity found in rat cardiomyocytes, supershift EMSA reactions performed with extracts from pMT₂-GATA-4-trans-

FIG. 3. Cardiomyocyte-specific DNA binding activity is dependent upon GATA-4. In EMSA reactions, cardiomyocyte nuclear extract was incubated with labeled PRE-D alone or with the indicated amounts of either anti-GATA-4 or preimmune (Pre-imm.) serum. The position of the GATA-4-dependent complex is indicated. SS indicates the positions of the supershifted bands.

fected COS cells (2) and cardiomyocytes were run on the same gel (Fig. 4, lanes 1 through 6). On the basis of electrophoretic mobility, the GATA-4-dependent complex in cardiomyocytes was indistinguishable from the mouse GATA-4 produced from the pMT₂-GATA-4 expression vector in COS cells. Untransfected COS cells did not show any GATA binding activity as previously reported (2). When the reaction mixtures containing cardiomyocyte extracts and the transfected COS cell extracts were incubated with the anti-GATA-4 serum, both reactions were supershifted to similar extents, with the corresponding supershifted bands migrating to the same positions. The faster-migrating complex was also present in both cardiomyocyte and transfected COS cell nuclear extracts. This band was partially supershifted and completely supershifted in the cardiomyocyte and transfected COS cell extracts, respectively.

To determine if GATA-4 acted alone or in combination with other factors present in cardiomyocytes and COS cells, the GATA-4-dependent complex from cardiomyocytes was compared with the complex that formed with in vitro-translated mouse GATA-4 (Fig. 4, lanes 7 through 12). When PRE-D was incubated with mouse GATA-4 prepared from reticulocyte lysate, the mobility of the resulting complex was indistinguishable from the similar activity in cardiomyocytes. No such interaction was present in sham-translated reticulocyte lysate. Anti-GATA-4 serum supershifted the GATA-4-dependent bands from both cardiomyocyte extracts and the in vitro translation reaction. The similar electrophoretic mobilities and supershift interactions of the GATA-4-dependent complexes from three different sources suggest that GATA-4 is present in cardiomyocytes and that the GATA-4-dependent complex contains only GATA-4.

The presence of tandem sites and the fact that both sites

FIG. 4. The GATA-4-dependent complexes with PRE-D from three different sources have the same mobility. EMSAs were performed by using PRE-D as a probe with extracts or in vitro-translated GATA-4 as indicated above the lanes. Lanes 1 through 3 contain cardiomyocyte nuclear extract, GATA-4-transfected COS cell extract, and COS cell nuclear extract, respectively, with the labeled PRE-D oligonucleotide in the presence of 1 μ l of preimmune (Pre-imm.) serum. Lanes 4 through 6 are the same except that they contain 1 μ l of anti-GATA-4 serum instead of preimmune serum. Lanes 7 through 9 show cardiomyocyte nuclear extract (Cardio.), in vitro-translated GATA-4 reticulocyte lysate (Ret.-GATA-4), and sham-translated reticulocyte lysate (Ret.), respectively, with the labeled PRE-D oligonucleotide in the presence of 1 μ l of preimmune serum. Lanes 10 through 12 are the same as lanes 7 through 9 except that they contain 1 μ l of anti-GATA-4 serum instead of preimmune serum. SS, positions of the supershifted bands.

were footprinted from the same EMSA band (Fig. 1) suggested that there might be a cooperative interaction between the proteins binding to these sites. If this were correct, we would expect to see a faster-migrating complex in EMSA gels, at low protein concentrations, corresponding to a single occupied site. Repeated attempts to obtain such a complex were unsuccessful. To substantiate this result, we used ds oligonucleotides identical to PRE-D except that one of the two GATA binding sites was mutated (D5' mut and D3' mut). The complexes formed with these single-site oligonucleotides were indistinguishable in mobility from the complex formed when both sites were present (Fig. 5). Since the probes were of nearly equal specific activities, these results indicate that GATA-4 has a higher affinity for the 5' GATA-binding site than for the 3' site. This is consistent with the competition data shown in Fig. 1. The complexes formed with each of the oligonucleotides were GATA-4 dependent, since they supershifted with anti-GATA-4 serum but not with preimmune serum (Fig. 5). These results suggest that the GATA-4 band in EMSAs represents two separate GATA-4 interactions with PRE-D, one with the upstream GATA site and the other with the downstream GATA site.

Mutation of the α -MHC GATA-4-binding sites diminishes expression from the α -MHC gene. To determine if GATA-4 participated in the regulation of the α -MHC gene, the two

FIG. 5. Mobility of the GATA-4-dependent complex associated with PRE-D is not altered when one of the GATA-binding sites is mutated. EMSAs were performed with PRE-D or with identical oligonucleotides except for mutations in the 3' GATA site (D3' mut) or the 5' GATA site (D5' mut). These reactions were done in the presence of 1 μ l of preimmune (lanes 1 through 3) or anti-GATA-4 (lanes 4 through 6) serum. The position of the GATA-4 band and those of the corresponding supershifted bands (ss) are indicated.

GATA-4-binding sites in PRE-D were mutated alone or together within the context of the 2.9-kb promoter in pSV0MCAT. Initially, a mutant with extensive changes throughout the area that produced the footprint was constructed. This mutant contained mutations in both GATA-binding sites (D5'-3' mut). D5'-3' mut was injected into the hearts of adult rats as described above. After 6 days, the area of injection was excised and assayed for CAT activity. Compared with that of wild-type pSV0MCAT ($20,595 \pm 3,254$ cpm per 50 μ g of protein [34]), the mutations in D5'-3' mut reduced activity by 88% (Fig. 6). The mutations in D5'-3' mut also eliminated all interactions with the PRE-D oligonucleotide (data not shown). When the upstream (D5' mut) and downstream (D3' mut) GATA sites were mutated individually, activity from the α -MHC promoter was reduced by approxi-

Construct	-250	-280	Percent of pSV0MCAT activity	N
Wildtype	5'CAAAGGGCCGATGGGCAGATAGAGGAGAGACAGGA		100% $\pm$ 15	9
D3' mut	TCA		45% $\pm$ 6	7
D5' mut		TGC	53% $\pm$ 4	7
D5'-3' mut	TCA	TGC	12% $\pm$ 3	9

FIG. 6. Transcriptional activity of the α -MHC enhancer-promoter and GATA-4 site-directed mutants in vivo. The antisense sequence shown is from the α -MHC gene in the region where the mutations were introduced, and the base substitutions for each mutant are represented below the sequences. The position relative to the transcriptional start site of the α -MHC gene is indicated (26). The activity of each construct is given as the percentage of pSV0MCAT activity $\pm$ 1 standard deviation in the adult rat heart (34).

FIG. 7. The α -MHC GATA sites mediate GATA-4-dependent transcriptional enhancement from a heterologous promoter but not from the α -MHC promoter in nonmuscle cells. Mouse L929 fibroblast cells were cotransfected with equimolar amounts of reporter plasmid pGLPREDx2 or pMHC598 and either pMT₂ or pMT₂-GATA-4, incubated for 48 h, and assayed for luciferase activity. In a control experiment, the parent construct containing the simian virus 40 early promoter but lacking the PRE-D sites (pGL-promoter) was also transfected with pMT₂-GATA-4. The activity is given as relative luciferase units divided by CAT activity $\times 10^{-2}$. The error bars in the graph represent the standard error of the mean for each construct tested ($n \geq 4$).

mately 50% (Fig. 6). With either site mutated individually, GATA-4 was still capable of interacting with PRE-D (Fig. 5). These results indicated that the loss of either GATA site reduced but did not prevent α -MHC gene expression. Loss of both sites had much more serious consequences.

To determine if the GATA sites within the α -MHC regulatory region were capable of mediating enhanced transcription in the presence of GATA-4, two copies of PRE-D were cloned upstream of the simian virus 40 early promoter in the vector pGL2-promoter to form pGLPREDX2. The ability of GATA-4 to transactivate this promoter construct was examined by cotransfection experiments. pGL2-promoter and pGLPREDX2 were assayed for activity in the mouse fibroblast cell line L929, in the presence of either pMT₂ or pMT₂-GATA-4 (Fig. 7). Only basal activity was observed from pGL-promoter and pGLPREDX2 in L929 cells when cells were cotransfected with pMT₂. In contrast, activity from pGLPREDX2 was increased approximately 16-fold in the presence of pMT₂-GATA-4. Activity from pGL-promoter was not affected by pMT₂-GATA-4. These results indicate that GATA-4 can act through the α -MHC GATA-binding sites to transactivate transcription from a heterologous promoter.

GATA-4 induces expression of the α -MHC gene in skeletal muscle. As discussed above, all of the factors previously shown to participate in the regulation of the α -MHC gene have been found in skeletal muscle, even though this gene is not expressed at significant levels in skeletal muscle cells. The tissue-restricted expression of GATA-4 and its participation in the regulation of α -MHC gene expression suggested the possibility that this transcription factor was responsible, in part, for the cardiac specificity of expression. If the deficiency of GATA-4 in skeletal muscle was sufficient to prevent α -MHC expression, then the presence of a GATA-4 expression vector in skeletal muscle cells should stimulate expression from an α -MHC promoter-reporter construct. The region of the α -MHC promoter from position -598 to +17 was cloned adjacent to a luciferase reporter gene (pMHC598). This re-

FIG. 8. GATA-4 induces high-level expression from an α -MHC-reporter construct in skeletal muscle. pMHC598 was injected alone or with either pMT₂ or pMT₂-GATA-4 into the tongues of rats and assayed for luciferase activity. A second reporter construct, pMHC598mut, was identical to pMHC598 except that both GATA-binding sites were mutated as shown in Fig. 6. pMHC598mut was coinjected with pMT₂-GATA-4. pGL basic is the parent vector of pMHC598 that contains no promoter. The activity is given as relative luciferase units divided by CAT activity $\times 10^{-2}$. The error bars represent the standard error of the mean for each construct assayed. The number of samples was 5 for pMHC598 and pMT₂, 6 for pMHC598mut and pMT₂-GATA-4, and 11 for pMHC598 and pMT₂-GATA-4.

porter construct was injected alone or coinjected with pMT₂ or pMT₂-GATA-4 into skeletal muscle of adult rats (Fig. 8). When pMHC598 was injected alone, it produced a level of activity that was only slightly above that of the background expression from the pGL basic vector. pGL basic has no promoter driving expression. When pMHC598 was coinjected with pMT₂, activity from pMHC598 actually decreased to below background levels (see below). However, when the reporter was coinjected with pMT₂-GATA-4, there was a dramatic increase in activity, which represented an eightfold increase in expression over that of pMHC598 alone (Fig. 8). The same experiment was done in the hearts of the same animals. In the heart, the presence of the pMT₂-GATA-4 did not up-regulate expression from pMHC598 (data not shown), suggesting that the levels of GATA-4 in the heart were not rate limiting for α -MHC expression.

In initial coinjection experiments, we observed that both pMT₂ and pMT₂-GATA-4 squelched α -MHC promoter activity even at equimolar ratios of expression vector and reporter. At a 1:4 pMT₂-to-pMHC598 ratio, squelching of activity from the α -MHC promoter was less severe, and this ratio of expression vector to reporter was used in the injection experiments in Fig. 8.

Next we addressed whether the observed stimulation of the α -MHC promoter in skeletal muscle was mediated through the GATA-4-binding sites. Both GATA-4 sites in pMHC598 were eliminated by site-directed mutagenesis, as was done with pSV0MCAT (see above). The resulting promoter construct (pMHC598mut) was coinjected with pMT₂-GATA-4 into skeletal muscle and subsequently assayed for activity. As seen in Fig. 8, pMT₂-GATA-4 did not stimulate expression of the mutant promoter in skeletal muscle, indicating that GATA-4 induces expression solely through the binding sites that are present in PRE-D.

Finally, it was of interest to determine if the stimulatory effect of GATA-4 on the α -MHC promoter required other

FIG. 9. GATA-4 DNA binding activity is up-regulated in cardiomyocytes treated with RA or T₃. EMSAs were done by using PRE-D (lanes 1 through 4) or PRE-B (lanes 5 through 8) (Table 1) (33) to probe cardiomyocyte nuclear extracts prepared from untreated, RA-treated, or T₃-treated cells as indicated above the lanes. The same extracts were used for both probes. The positions of the GATA-4-dependent and BF-2-dependent interactions are indicated.

muscle transcription factors. To address this issue, pMHC598 was assayed for promoter activity alone or in combination with pMT₂ or pMT₂-GATA-4 in mouse L929 cells (Fig. 7). In L929 cells, no squelching was observed. pMT₂-GATA-4 did not stimulate transcription from pMHC598 in these nonmuscle cells. This is in contrast to the stimulation of the heterologous promoter construct pGLPREDX2 in the same cell type or when pMT₂-GATA-4 was coinjected with pMHC598 into skeletal muscle. These results suggest that expression from the α -MHC enhancer-promoter requires GATA-4 but that GATA-4, by itself, is not sufficient to enhance expression of the α -MHC gene.

Thyroid hormone or RA treatment up-regulates GATA-4 DNA binding activity in cardiomyocytes. To further characterize the GATA-4 binding activity in cardiomyocytes, primary neonatal rat cardiomyocytes were isolated and cultured in serum-free Ham's F12 medium. The plates were divided into three groups. Group 1 received no additional treatment. Group 2 was supplemented with all-*trans* RA to a final concentration of 100 nM. Group 3 was supplemented with T₃ to a final concentration of 10 nM. After 48 h of treatment, nuclear extracts were prepared from each of these groups and probed for GATA-4 activity. Results of a representative experiment are shown in Fig. 9. The levels of GATA-4 DNA binding activity were clearly higher in the T₃-treated and RA-treated cells (Fig. 9). These same extracts were probed with the binding site for BF-2. The level of BF-2 binding activity in the T₃-treated cells was not different from that in untreated cells, and the level of binding activity in the RA-treated cells was somewhat decreased. When these experiments were quantitated on a PhosphorImager, it was found that GATA-4 DNA binding activity was increased 3-fold by T₃ and 1.8-fold by RA compared with that of untreated extract. The decrease in BF-2 activity with RA treatment did not occur

in most nuclear extract preparations and probably indicates some variability in the extract preparations. However, since the same RA extract was used for the experiments shown in Fig. 9, a 1.8-fold increase in GATA-4 activity stimulated by RA probably reflects a level of induction lower than that which actually occurred.

DISCUSSION

In this study, we established that GATA-4 regulates expression of the α -MHC gene through an interaction with GATA-binding sites within the proximal regulatory region. This activity is necessary to explain the tissue-restricted expression of the α -MHC gene. GATA-4 was found to act through the α -MHC GATA-4-binding sites to enhance expression from the α -MHC gene and from a heterologous promoter. Previous work has suggested that the lack of a positive factor in skeletal muscle rather than the presence of a negative regulator prevents the expression of the α -MHC gene in that cell type (12). Our study substantiates this hypothesis by demonstrating that the introduction of GATA-4 into a skeletal muscle milieu dramatically increases α -MHC promoter activity.

GATA-4 interacts with the GATA-binding sites within the α -MHC regulatory region. The factor present in nuclear extracts of neonatal rat cardiomyocytes that interacts with the two α -MHC GATA-binding sites was identified as GATA-4 (Fig. 1 through 5). Supershift EMSA (Fig. 3) and immunoprecipitation experiments (data not shown) demonstrated that the complex formed on the GATA-binding sites in the α -MHC enhancer was dependent upon GATA-4 or a closely related factor. It was possible, however, that GATA-4 interacted with an additional factor(s) to form the complex with the GATA-binding site. To examine this possibility, we compared the mobilities of the GATA-4-dependent complexes from extracts of cardiomyocytes, from COS cells transfected with the GATA-4 expression vector, and from *in vitro*-transcribed and -translated GATA-4. Our results demonstrated that the mobilities of these complexes were indistinguishable (Fig. 4). While it is formally possible that there is a factor common to cardiomyocytes, COS cells, and rabbit reticulocyte lysate that forms a complex with GATA-4, it is more likely that GATA-4 alone is responsible for the interaction.

Previously published work has not established that GATA-4 is present in myocytes of the heart (2, 18). The primary cultures used in this study are differentially plated to enrich for cardiomyocytes (11), suggesting that the GATA-4 activity is present in the muscle cells of the heart. Further, the functional studies of the adult heart indicate that GATA-4 is present in cardiomyocytes, since injection of naked DNA into the heart and into skeletal muscle results in uptake only in myocytes (19, 23, 50, 51). Thus, the data presented here suggest that GATA-4 is present in cardiomyocytes.

There appears to be no cooperative binding of GATA-4 molecules to the two GATA-binding sites in the α -MHC enhancer *in vitro*. The mobilities of the complexes formed on the wild-type and single-site mutant oligonucleotides were identical (Fig. 5), and we were unable to demonstrate the presence of a doubly occupied site at higher protein concentrations. Thus, GATA-4 may interact with either site, but not both sites simultaneously, in *in vitro* binding reactions. While only one of the two GATA-binding sites is occupied at any time, our results suggest that the presence of two GATA-4 sites enables α -MHC to recruit GATA-4 more efficiently. This interpretation is consistent with the DEPC footprint, since exposure of the gel represented by the autoradiogram in Fig.

2A for a longer period shows that each site is only partially protected.

GATA-4 regulates transcription of the α -MHC gene. When one or both of the GATA-binding sites were mutated within the context of the α -MHC promoter fusion construct pSV0MCAT, activity was decreased to 50% and 12% of the wild-type level, respectively, in adult rat heart muscle (Fig. 6). The difference in activity between the single- and double-site mutants is consistent with the interpretation of the binding data that suggests a lack of functional cooperative interaction between the two GATA-4 sites. There was also a direct correlation between loss of functional activity and loss of GATA-4 binding to oligonucleotides containing these mutant sites. This suggests that this zinc finger protein is involved in transcriptional regulation of the rat α -MHC gene. The α -MHC GATA-binding sites were capable of supporting GATA-4-enhanced expression from a heterologous promoter (Fig. 7), indicating that the GATA-binding sites are positive regulatory elements. Taken together, these results indicate that one role of GATA-4 in the heart is to enhance expression of the α -MHC gene, and they suggest that GATA-4 may be an important component of the molecular mechanism controlling transcription of several genes in the heart (see Table 2).

GATA-4 plays a role in cardiac muscle-specific expression of the α -MHC gene. The α -MHC gene is regulated by a number of factors, including BF-2 and MEF-2 (33, 34), that are common to skeletal and cardiac muscle, but significant levels of α -MHC mRNA are found only in the heart. GATA-4 was first described as a transcription factor present in the heart and in endodermally derived tissues (2). Because of the tissue-restricted distribution of this protein and its ability to regulate expression of the α -MHC gene, we felt that GATA-4 might be involved in the cardiac muscle-specific expression of this gene. We tested this hypothesis by performing experiments involving coinjection of an α -MHC-reporter construct and a GATA-4 expression vector into the tongue. The tongue was selected because of the ease of injection and its large proportion of skeletal muscle cells. Control experiments indicated that the α -MHC enhancer-promoter, by itself, had a low level of basal activity in skeletal muscle cells found in the tongue (Fig. 8). This was similar to the low level of α -MHC gene expression previously observed with several skeletal muscle and non-muscle cell types (12, 13, 17, 48). In the presence of the GATA-4 expression vector, the α -MHC promoter was dramatically up-regulated (Fig. 8). These results establish that the lack of GATA-4 in skeletal muscle cells is responsible, in part, for the lack of expression of the α -MHC gene there. A similar result was also seen in preliminary experiments with other skeletal muscle cells (35).

The GATA-4-stimulated expression from pMHC598 in skeletal muscle was in contrast to its effect in L929 cells. GATA-4 failed to stimulate expression from pMHC598 in L929 cells, even though the expression vector pMT₂-GATA-4 is active in this nonmuscle cell line (Fig. 7). Therefore, the lack of activity from the α -MHC promoter in L929 cells, in the presence of GATA-4, is most likely due to a lack of certain muscle-specific transcription factors. These results strongly suggest that GATA-4 must interact with other muscle regulatory factors to stimulate transcription of the α -MHC gene.

Cardiac muscle-specific gene expression. The first well-described pathway of cardiac muscle-specific gene expression was that of the MLC-2a gene involving the HF-1 region (21, 38, 55). HF-1 is similar to the MBF-2 region of the α -MHC gene, suggesting that MBF-2 may also participate in cardiac muscle-restricted expression of the α -MHC gene (34). The HF-1 and MBF-2 regions of their respective genes each contain a

TABLE 2. Putative GATA-binding sites in cardiac tissue-expressed genes

Gene ^a	Sequence	Reference
r α -MHC	-256 CCGATGGGCA -265	This work
r α -MHC	-264 CAGATAGAGG -273	This work
r β -MHC	-270 GGGATATTTT -261	46
r TnCc/s	-92 AGGATATTCC -83	39
r MLC-2	-229 ACCATGCTCT -238	15
m MLC-1 _A	-141 GTGATAATAC -132	6
	-130 AAGATGTCAG -121	
	-348 TTGATAACCC -357	
m MLC-1 _V	-1079 AAGATAGAGA -1070	6
	-994 AAGATAGAGA -985	
	-736 TGGATACAGC -745	
m MCK	-93 CAGATGAGCT -102	43
r ANF	-282 AAGATAACCA -273	42
r ANF	-124 CTGATAACTT -115	42
h/m c-actin	-482 AAGATGAGAA -473	32
h c-actin	-332 TGGATGCTCT -341	32
m c-actin	-127 TAGATGGTGC -118	32

^a m, r, c, s, and h in designations refer to mouse, rat, cardiac, slow, and human, respectively.

^b MCK, muscle creatine kinase.

^c ANF, atrial natriuretic factor.

putative DNA recognition sequence that interacts with MEF-2, implicating this protein in cardiac muscle-restricted expression. In addition, mutation of the MEF-2 site in the mouse α -MHC gene permitted ectopic expression of this gene in transgenic mice (1). However, only the BF-2 portion of the MBF-2 region was necessary to enhance expression from a 325-bp α -MHC promoter fragment, and this expression was cardiac muscle restricted (33). Therefore, we suspected that the cardiac specificity of α -MHC gene expression was also controlled by a factor(s) other than MEF-2. The identification of GATA-4 as a regulator of cardiac muscle-specific expression of the α -MHC gene represents a second component of cardiac muscle-restricted gene expression.

Interestingly, much of the work describing the function of the MLC-2 HF-1 region in determining cardiac specificity, including experiments using the transgenic mouse model, has been done in the context of 250 bp of the proximal MLC-2 enhancer-promoter (21, 38, 55). Within that 250-bp region is a putative GATA-binding site (15) (see Table 2). Similarly, there is a GATA-binding site in the DNA recognition site for cardiac enhancer factor 2, which has been implicated in cardiac muscle-specific expression of the troponin C gene (39). GATA-binding sites can also be found in the regulatory regions of several additional genes expressed in the heart (Table 2). The presence of potential GATA-4 binding sites in the regulatory regions of several cardiac tissue-expressed genes and the demonstration that GATA-4 regulates at least one of these genes suggest that GATA-4 is an important regulator of cardiac muscle-specific gene expression.

Developmental implications of GATA-4 regulatory activity and a role for thyroid hormone and RA in cardiac development. While the regulation of the α -MHC gene results from the interaction of multiple elements, it is dependent upon GATA-4. GATA-4 is expressed in the embryonic mouse heart tube at day 7.5 postcoitum and remains at high levels throughout development (49). Between days 7.5 and 8 postcoitum, the α -MHC gene is transiently expressed in the heart tube (25). This temporal pattern of expression is consistent with the possibility that GATA-4 is responsible, in part, for the early expression of α -MHC mRNA in the developing heart.

GATA-4 may also regulate the early expression of the β -MHC, atrial myosin light-chain 1 (MLC-1_A), and ventricular MLC-1 (MLC-1_V) genes in the heart, since these genes are expressed concomitantly with the α -MHC gene (25) and each has putative GATA-4 sites in its regulatory region (Table 2).

Previous work has demonstrated that expression of GATA-4 is regulated by RA in F9 embryonal carcinoma cells (2). We show here that the DNA binding activity of GATA-4 in cardiomyocytes is up-regulated by treatment with RA or T₃ (Fig. 9). RA and T₃ have also been shown to regulate the expression of the α -MHC gene (12, 27). These agents can affect gene expression through interactions of their receptors and coregulators with DNA regulatory elements in target genes (recently reviewed in reference 54). This was presumed to be the exclusive mechanism by which RA and T₃ regulated the expression of the α -MHC gene. However, the identification of an RA-inducible factor that regulates the α -MHC gene indicates a more complex control pathway.

It is interesting to speculate that RA and/or T₃ might induce GATA-4 in the developing heart, which, in turn, is responsible for up-regulating the expression of α -MHC and other genes. The available data indicate that mRNAs encoding RA receptors are first detectable in the heart later than those encoding GATA-4 and α -MHC (41). However, no data concerning the initial appearance of thyroid hormone receptors or retinoic acid X receptors (RXR) in the heart are available. These results suggest that RA and T₃ may exert their influences at later developmental stages but do not rule out the possibility of a role for these agents in early stages of cardiogenesis.

ACKNOWLEDGMENTS

We thank David Wilson for his generous gifts of GATA-4 anti-serum, COS cell extracts, pMT₂, and pMT₂-GATA-4; D. Wilson, G. Lyons, and G. Engelmann for helpful discussions; E. Sternberg for reviewing the manuscript; and Allen Cowley, Jr., for the use of some equipment. We also thank Jane Wolf for technical assistance.

This work was supported by Public Health Service grant HL43662 from the National Heart, Lung, and Blood Institute to B.E.M. J.D.M. is a predoctoral fellow of the American Heart Association (Wisconsin Affiliate). B.E.M. is an Established Investigator of the American Heart Association.

REFERENCES

- Adolph, E. A., A. Subramaniam, P. Cserjesi, E. N. Olson, and J. Robbins. 1993. Role of MEF-2 in transcriptional regulation of the α -cardiac myosin heavy chain gene. *J. Biol. Chem.* 268:5349-5352.
- Arceci, R. J., A. A. J. King, M. C. Simon, S. H. Orkin, and D. B. Wilson. 1993. Mouse GATA-4: a retinoic acid-inducible GATA-binding transcription factor expressed in endodermally derived tissues and heart. *Mol. Cell. Biol.* 13:2235-2246.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72:248-254.
- Buttrick, P. M., M. L. Kaplan, R. N. Kitsis, and L. A. Leinwand. 1993. Distinct behavior of cardiac myosin heavy chain gene constructs in vivo. *Circ. Res.* 72:1211-1217.
- Chen, C., and H. Okayama. 1987. High-efficiency transformation of mammalian cells by plasmid DNA. *Mol. Cell. Biol.* 7:2745-2752.
- Cohen, A., P. J. R. Barton, B. Robert, I. Garner, S. Alanson, and M. E. Buckingham. 1988. Promoter analysis of myosin alkali light chain genes expressed in mouse striated muscle. *Nucleic Acids Res.* 16:10037-10052.
- de Wet, J. R., K. V. Wood, M. DeLuca, D. R. Helinski, and S. Subramani. 1987. Firefly luciferase gene: structure and expression in mammalian cells. *Mol. Cell. Biol.* 7:725-737.
- Dignam, J. D., R. M. Lebowitz, and R. G. Roeder. 1983. Accurate transcription initiation by RNA polymerase II in a soluble extract

- from isolated mammalian nuclei. *Nucleic Acids Res.* **11**:1475–1489.
9. Glass, C. K., S. M. Lipkin, O. V. Devary, and M. G. Rosenfeld. 1989. Positive and negative regulation of gene transcription by retinoic acid-thyroid hormone receptor heterodimers. *Cell* **59**: 697–708.
 10. Gorman, C. M., L. F. Moffat, and B. H. Howard. 1982. Recombinant genomes which express chloramphenicol acetyltransferase in mammalian cells. *Mol. Cell. Biol.* **2**:1044–1051.
 11. Gustafson, T. A., J. J. Bahl, B. E. Markham, W. R. Roeske, and E. Morkin. 1987. Hormonal regulation of myosin heavy chain and α -actin expression in cultured fetal rat heart myocytes. *J. Biol. Chem.* **262**:13316–13322.
 12. Gustafson, T. A., B. E. Markham, J. J. Bahl, and E. Morkin. 1987. Thyroid hormone regulates the expression of a transfected α -myosin heavy chain fusion gene in fetal heart cells. *Proc. Natl. Acad. Sci. USA* **84**:3122–3126.
 13. Gustafson, T. A., B. E. Markham, and E. Morkin. 1986. Effects of thyroid hormone on α -actin and myosin heavy chain gene expression in cardiac and skeletal muscles of the rat: measurements of mRNA content using synthetic oligonucleotide probes. *Circ. Res.* **59**:194–201.
 14. Hemsley, A., N. Arnheim, M. D. Toney, G. Cortopassi, and D. J. Galas. 1989. A simple method for site directed mutagenesis using polymerase chain reaction. *Nucleic Acids Res.* **17**:6545–6551.
 15. Henderson, S. A., M. Spencer, A. Sen, C. Kumar, M. A. Q. Siddiqui, and K. R. Chien. 1989. Structure, organization and expression of the rat cardiac myosin light chain-2 gene. *J. Biol. Chem.* **264**:18142–18148.
 16. Izumo, S., and V. Mahdavi. 1988. Thyroid hormone receptor α isoforms generated by alternative splicing differentially activate myosin HC gene transcription. *Nature (London)* **334**:539–542.
 17. Izumo, S., B. Nadal-Ginard, and V. Mahdavi. 1986. All members of the MHC multigene family respond to thyroid hormone in a highly tissue specific manner. *Science* **231**:597–600.
 18. Kelly, C., H. Blumberg, L. I. Zon, and T. Evans. 1993. GATA-4 is a novel transcription factor expressed in the endocardium of the developing heart. *Development* **118**:817–827.
 19. Kitsis, R. N., P. M. Buttrick, E. M. McNally, M. L. Kaplan, and L. A. Leinwand. 1991. Hormonal modulation of a gene injected into rat heart *in vivo*. *Proc. Natl. Acad. Sci. USA* **88**:4138–4142.
 20. Ko, L. J., and J. D. Engel. 1993. DNA-binding specificities of the GATA transcription factor family. *Mol. Cell. Biol.* **13**:4011–4022.
 21. Lee, K. J., R. S. Ross, H. A. Rockman, A. N. Harris, T. X. O'Brien, M. van Bilsen, H. E. Shubeita, R. Kandolf, G. Brem, J. Price, S. M. Evans, H. Zhu, W.-M. Franz, and K. R. Chien. 1992. Myosin light chain-2 luciferase transgenic mice reveal distinct regulatory programs for cardiac and muscle-specific expression of a single contractile protein gene. *J. Biol. Chem.* **267**:15875–15885.
 22. Lee, M.-E., D. H. Temizer, J. A. Clifford, and T. Quertermous. 1991. Cloning of the GATA-binding protein that regulates endothelin-1 gene expression in endothelial cells. *J. Biol. Chem.* **266**:16188–16192.
 23. Leinwand, L. A., and J. M. Leiden. 1991. Gene transfer into cardiac myocytes *in vivo*. *Trends Cardiovasc. Med.* **1**:271–276.
 24. Lompre, A.-M., B. Nadal-Ginard, and V. Mahdavi. 1984. Expression of the cardiac ventricular α - and β -myosin heavy chain genes is developmentally and hormonally regulated. *J. Biol. Chem.* **259**:6437–6446.
 25. Lyons, G. E., S. Schiaffino, D. Sassoon, P. Barton, and M. Buckingham. 1990. Developmental regulation of myosin gene expression in mouse cardiac muscle. *J. Cell Biol.* **111**:2427–2436.
 26. Mahdavi, V., A. P. Chambers, and B. Nadal-Ginard. 1984. Cardiac α - and β -myosin heavy chain genes are organized in tandem. *Proc. Natl. Acad. Sci. USA* **81**:2626–2630.
 27. Markham, B. E. 1990. Synergistic regulation of the α -myosin heavy chain gene by triiodothyronine and retinoic acid in primary cardiomyocytes, abstr. 78. Council. High Blood Pressure Research, 44th Annu. Conf. Sci. Sessions. 1990.
 28. Markham, B. E., J. J. Bahl, T. A. Gustafson, and E. Morkin. 1987. Interaction of a protein factor within a thyroid hormone sensitive region of rat α -myosin heavy chain gene. *J. Biol. Chem.* **262**: 12856–12862.
 29. Markham, B. E., R. W. Tsika, J. J. Bahl, P. G. Anderson, and E. Morkin. 1990. An anti-sense *c-erbA* clone inhibits thyroid hormone-induced expression from the α -myosin heavy chain promoter. *J. Biol. Chem.* **265**:6489–6493.
 30. Maxam, A. M., and W. Gilbert. 1980. Sequencing DNA by labeling the end and breaking at bases: DNA segments, end-labels, cleavage reactions, polyacrylamide gels, and strategies. *Methods Enzymol.* **65**:499–560.
 31. Merika, M., and S. H. Orkin. 1993. DNA-binding specificity of GATA family transcription factors. *Mol. Cell. Biol.* **13**:3999–4010.
 32. Minty, A., and L. Kedes. 1986. Upstream regions of the human cardiac actin gene that modulate its transcription in muscle cells: presence of an evolutionarily conserved repeated motif. *Mol. Cell. Biol.* **6**:2125–2136.
 33. Molkentin, J. D., R. S. Brogan, S. M. Jobe, and B. E. Markham. 1993. Expression of the α -myosin heavy chain gene in the heart is regulated in part by an E-box dependent mechanism. *J. Biol. Chem.* **268**:2602–2609.
 34. Molkentin, J. D., and B. E. Markham. 1993. Myocyte-specific enhancer-binding factor (MEF-2) regulates α -cardiac myosin heavy chain gene expression *in vitro* and *in vivo*. *J. Biol. Chem.* **268**:19512–19520.
 35. Molkentin, J. D., and B. E. Markham. Unpublished data.
 36. Molkentin, J. D., and B. E. Markham. Submitted for publication.
 37. Morkin, E., J. J. Bahl, and B. E. Markham. 1989. Control of cardiac myosin heavy chain gene expression by thyroid hormone, p. 381–389. *In* L. Kedes and F. Stockdale (ed.), *Cellular and molecular biology of muscle development*. Alan R. Liss, Inc., New York.
 38. Navankasattusas, S., H. Zhu, A. V. Garcia, S. M. Evans, and K. R. Chien. 1992. A ubiquitous factor (HF-1a) and a distinct muscle factor (HF-1b/MEF-2) form an E-box-independent pathway for cardiac muscle gene expression. *Mol. Cell. Biol.* **12**:1469–1479.
 39. Parmacek, M. S., A. J. Vora, T. Shen, E. Barr, F. Jung, and J. M. Leiden. 1992. Identification and characterization of a cardiac-specific transcriptional regulatory element in the slow/cardiac troponin C gene. *Mol. Cell. Biol.* **12**:1967–1976.
 40. Rosenthal, N. 1987. Identification of regulatory elements of cloned genes with functional assays. *Methods Enzymol.* **152**:704–720.
 41. Ruberte, E., P. Dolle, P. Chambon, and G. Morriss-Kay. 1991. Retinoic acid receptors and cellular retinoid binding proteins. II. Their differential pattern of transcription during early morphogenesis in mouse embryos. *Development* **111**:45–60.
 42. Seidman, C. E., D. W. Wong, K. D. Bloch, and J. G. Seidman. 1988. Cis-acting sequences that modulate atrial natriuretic factor gene expression. *Proc. Natl. Acad. Sci. USA* **85**:4104–4108.
 43. Sternberg, E. A., G. Spizz, W. M. Perry, D. Vizard, T. Weil, and E. N. Olson. 1988. Identification of upstream and intragenic regulatory elements that confer cell-type-restricted and differentiation-specific expression on the muscle creatine kinase gene. *Mol. Cell. Biol.* **8**:2896–2909.
 44. Sturm, R., T. Baumrucker, R. Franza, Jr., and W. Herr. 1987. A 100-kD HeLa cell octamer binding protein (OBP100) interacts differently with two separate octamer related sequences within the SV40 enhancer. *Genes Dev.* **1**:1147–1160.
 45. Subramaniam, A., W. K. Jones, J. Gulick, S. Wert, J. Neumann, and J. Robbins. 1991. Tissue-specific regulation of the α -myosin heavy chain gene promoter in transgenic mice. *J. Biol. Chem.* **266**:24613–24620.
 46. Thompson, W. R., B. Nadal-Ginard, and V. Mahdavi. 1991. A myoD1-independent muscle-specific enhancer controls the expression of the α -myosin heavy chain gene in skeletal and cardiac muscle cells. *J. Biol. Chem.* **266**:22678–22688.
 47. Tsai, S.-F., D. I. K. Martin, L. I. Zon, A. D. D'Andrea, G. G. Wong, and S. H. Orkin. 1989. Cloning of cDNA for the major DNA-binding protein of the erythroid lineage through expression in mammalian cells. *Nature (London)* **339**:446–451.
 48. Tsika, R. W., J. J. Bahl, L. A. Leinwand, and E. Morkin. 1990. Thyroid hormone regulates expression of a transfected human α -myosin heavy-chain fusion gene in fetal rat heart cells. *Proc. Natl. Acad. Sci. USA* **87**:379–383.
 49. Wilson, D. B. (Washington University). 1993. Personal communication.

50. **Wolff, J. A., R. W. Malone, P. Williams, W. Chong, G. Acsadi, A. Jani, and P. L. Felgner.** 1990. Direct gene transfer into mouse muscle *in vivo*. *Science* **247**:1465–1468.
51. **Wolff, J. A., P. Williams, G. Acsadi, S. Jiao, A. Jani, and W. Chong.** 1991. Conditions affecting direct gene transfer into rodent muscle *in vivo*. *BioTechniques* **11**:474–485.
52. **Yamamoto, M., L. J. Ko, M. W. Leonard, H. Beug, S. H. Orkin, and J. D. Engel.** 1990. Activity and tissue-specific expression of the transcription factor NF-E1 multigene family. *Genes Dev.* **4**:1650–1662.
53. **Yu, Y.-T., R. E. Breithart, L. B. Smoot, Y. Lee, V. Mahdavi, and B. Nadal-Ginard.** 1992. Human myocyte-specific enhancer factor 2 comprises a group of tissue-restricted MADS box transcription factors. *Genes Dev.* **6**:1783–1798.
54. **Zhang, X.-K., and M. Pfahl.** 1993. Regulation of retinoid and thyroid hormone action through homodimeric and heterodimeric receptors. *Trends Endocrinol. Metab.* **4**:156–162.
55. **Zhu, H., A. V. Garcia, R. S. Ross, S. M. Evans, and K. R. Chien.** 1991. A conserved 28-base-pair element (HF-1) in the rat cardiac myosin light-chain-2 gene confers cardiac-specific and α -adrenergic-inducible expression in cultured neonatal rat myocardial cells. *Mol. Cell. Biol.* **11**:2273–2281.