Published in final edited form as:

J Theor Biol. 2012 October 21; 311: 19–27. doi:10.1016/j.jtbi.2012.07.001.

The contribution of age structure to cell population responses to targeted therapeutics

Pierre Gabriel^{a,*}, Shawn P. Garbett^b, Vito Quaranta^b, Darren R. Tyson^b, and Glenn F. Webb^c

^aUniversité Pierre et Marie Curie-Paris 6, UMR 7598 LJLL, BC187, 4, place de Jussieu, F-75252 Paris cedex 5, France

^bDepartment of Cancer Biology, Vanderbilt University, 2220 Pierce Ave., Nashville, TN 37232, United States

^cDepartment of Mathematics, Vanderbilt University, 1326 Stevenson Center, Nashville, TN 37240, United States

Abstract

Cells grown in culture act as a model system for analyzing the effects of anticancer compounds, which may affect cell behavior in a cell cycle position-dependent manner. Cell synchronization techniques have been generally employed to minimize the variation in cell cycle position. However, synchronization techniques are cumbersome and imprecise and the agents used to synchronize the cells potentially have other unknown effects on the cells. An alternative approach is to determine the age structure in the population and account for the cell cycle positional effects post hoc. Here we provide a formalism to use quantifiable lifespans from live cell microscopy experiments to parameterize an age-structured model of cell population response.

Keywords

Cell cycle; Intermitotic time; Renewal equation; Exponentially modified Gaussian

1. Introduction

When examined individually in time lapse microscopy experiments, cells grown in culture display variability in the length of their cell cycles (between mitotic events), and this variability is represented by intermitotic time (IMT) distributions (Dowling et al., 2005; Lee and Perelson, 2008; Liu et al., 2004). These distributions are usually obtained from asynchronously dividing populations of cells, which achieve a steady-state age structure when the population is growing exponentially. In experimental studies that examine the effects of perturbations on cellular proliferation, it is often desirable to determine whether the perturbation is affecting cells in particular stages within their cell cycle, i.e. whether the perturbation has cell cycle-specific effects. Since we do not know where a cell is in the cycle, an alternative is to define the age of a cell as the time elapsed from its last division and deal with this measurable quantity instead of the cell cycle phase. In this paper we adopt this definition and provide a formalism to convert data obtained as IMT distributions to

parameterize an age-structured population model and, thus, identifying the contribution of age structure to the response of the cell population to perturbation.

Models of age-structured populations using partial differential equations, such as those originally described by A. Lotka, A.G. McKendrick, W.O. Kermack, and F. von Foerster, are well adapted to model the dynamical features of experimental cultures of cells transiting the cell cycle with variable IMT. These models have been widely studied from a mathematical perspective (Ainseba et al., 2011; Ainseba and Benosman, 2011; Calsina and Saldaña, 1995; Clément et al., 1987; Diekmann et al., 1984; Engel and Nagel, 2000; Metz and Diekmann, 1983; Perthame, 2007; Picart et al., 2011; Webb, 1985), but the application of the model to experimental data has been hampered by an inability to determine the age-dependent model parameters. The usual approach for parameter estimation is to solve numerically an inverse problem (see Ackleh, 1999; Banks et al., 2007, to appear; Doumic et al., 2009, 2010; Engl et al., 1994; Gyllenberg et al., 2002; Picart and Ainseba, 2011; Pilant and Rundell, 1991a,b; Rundell, 1989, 1993; Perthame and Zubelli, 2007 on this question for structured population models), but this requires extensive input data and is specific to a given situation. A much more convenient approach is to assume that the distributed parameters lie in a class of functions with only a few constants (power laws or other forms Gabriel, 2011) and obtain the parameters from the fit of the functions to experimental data.

One function that often provides a better fit to IMT data than others, such as log-normal, inverse normal or gamma functions, is an exponentially modified Gaussian (EMG) (Golubev, 2010; Tyson et al., in press). Under conditions in which the IMT distribution can be explained by an EMG model, we submit that the age-dependent division rate can be identified as an error function. Starting from this observation, we present a simple method to recover the parameters of this error function from parameters fitted to the experimental IMT data. Once reliably parameterized, the age-structured model can be used to make predictions about cell age-dependent effects of perturbations, for example, whether cells arrest during their cell cycle in response to treatment with antiproliferative compounds.

Individual based models (IBM) have also been used to track individual cell behavior in proliferating cell populations (Alarcón et al., 2003; Mallet and Pillis, 2006; Gerlee and Anderson, 2008; Patel et al., 2001; Powathil et al., 2012; Quaranta et al., 2008, 2005; Ramis-Conde et al., 2008; Ribba et al., 2004). IBM have advantages in their direct connection to discrete events and to cell-cell interactions. IBM are usually readily implementable to computer simulations, but may require many repeated runs to access an average outcome. IBM are usually difficult to analyze theoretically with respect to parametric input, particularly with highly sensitive parameters. The advantages of continuum differential equations models, such as the cell age structured models we develop here, are their tractability for theoretical analysis and their reproducibility for specific parametric input. The practical difficulties of determining this parametric input are the problem we focus upon here. IBM and differential equations models should be viewed as companion approaches, which complement and compare their insights and predictions for proliferating cell population behavior, when individual cell variation is a primary consideration.

2. The age-structured model

As with human populations, we can associate an age to each individual cell in a cell population. We define the age of a cell as the time elapsed from its last mitosis. Starting from this definition, we derive an age-structured model giving the dynamics of populations of proliferating and quiescent cells. This model is adapted to investigate the effect of treatment by the drug erlotinib on *in vitro* PC-9 cancer cell lines. It has been recently shown in Tyson et al. (in press) that the main effect of erlotinib on cancer cells is to induce entry

into quiescence. In Tyson et al. (in press) a system of ordinary differential (without age structure) is used to model these experiments. We hypothesize here that erlotinib induced quiescence is linked to the age of the cells involved. Many age-structured models with quiescence can be found in the literature (see Arino et al., 1997; Bekkal Brikci et al., 2008a,b; Gyllenberg and Webb, 1987,1990, 1991 for examples). Here we present a simple age-structured model with quiescence which allows us to explain observed delays in response to erlotinib. We start from the observation that there is no effect of the treatment on total population growth during the first twenty hours (see Tyson et al., in press and Fig. 7). This time corresponds almost exactly to the minimal age of division observed for PC-9 cells. It suggests that erlotinib acts only during a specific phase of the cell cycle, which based on its biological activity would be expected to be in G1. The model we present is based on this idea and considers a fractional rate f of cells that become quiescent in an age-dependent manner, where the fraction is assumed to reflect the dose of erlotinib used for the treatment.

We start from the McKendrick-Von Foerster model which is widely used to model the cell cycle (see Metz and Diekmann, 1983; Perthame, 2007; Webb, 1985 and references). The partial differential equation in this model provides the evolution of the density $p(t,a)$ of cells with age, or “cell cycle phase”, a at time t . To take into account the quiescent cells, we introduce the quantity $Q(t)$ which represents the density of quiescent cells at time t . It evolves according to an ordinary differential equation coupled to the equation on $p(t,a)$. We obtain the system

$$\begin{cases} \frac{\partial}{\partial t} p(t,a) + \frac{\partial}{\partial a} p(t,a) + \beta(a) p(t,a) + \mu p(t,a) = 0, & t \geq 0, a > 0, \\ p(t,0) = 2(1-f) \int_0^\infty \beta(a) p(t,a) da, \\ p(0,a) = p_0(a), \\ \frac{d}{dt} Q(t) = 2f \int_0^\infty \beta(a) p(t,a) da - \mu Q(t), \\ Q(0) = Q_0. \end{cases} \quad (1)$$

In this model, the proliferating cells age one-to-one with time at speed $da/dt = 1$, and divide with rate $\beta(a)$. When a cell divides at mitosis, it produces two daughters which are either proliferating with age $a=0$, or quiescent. This is taken into account by the boundary condition at $a=0$ and the first term in the equation for $Q(t)$, where the parameter $f \in [0,1]$ represents the fraction of proliferating cells which become quiescent. The coefficient μ is a death rates. The number of proliferating cells at time t with age between a_1 and a_2 is $\int_{a_1}^{a_2} p(t,a) da$, and the total number of cells at time t is $N(t) = \int_0^\infty p(t,a) da + Q(t)$.

The division rate β has a probabilistic interpretation: the probability that a cell did not divide by age a is given by

$$\mathbb{P}(a) = e^{-\int_0^a \beta(a') da'}.$$

Since all the proliferating cells must divide at some time by definition, the division rate has to satisfy

$$\lim_{a \rightarrow +\infty} \int_0^a \beta(a') da' = +\infty. \quad (2)$$

The model is completed with initial data $p_0(a)$ and Q_0 . We choose the time $t=0$ to be the beginning of the erlotinib treatment, so at this time there are no quiescent cells ($Q_0 = 0$). The age distribution of the proliferating cells is assumed to be at equilibrium (i.e.

$p_0(a) = \text{const } \widehat{p}(a)$, see Section 3 for the mathematical definition of $\widehat{p}$). The experimental

values of the total population $N(t)$ along time are plotted after normalization by the initial value $N(0)$ on a log-scale (see Fig. 7). Because of this normalization, we consider an initial distribution such that $\int_0^\infty p_0(a) da = 1$ which leads to $p_0(a) = \widehat{p}(a)$ because of the definition of $\widehat{p}$.

We want to compare the solutions of model (1) to the experimental observations presented in Tyson et al. (in press). The first step is to estimate the different parameters of the model. To estimate the value of the fraction f for different doses of treatment, we use data available in Tyson et al. (in press). The fraction F of quiescent cells is estimated by examination of whether cells treated with erlotinib divide or not before the end of the experiment. We want to use this experimental fraction F to estimate the coefficient f of the model. For the sake of simplicity, assume that the death rate μ is 0 in (1). In this case, at the end of the labeling period t_0 , the quantity of labeled quiescent cells corresponds to $Q(t_0)$ and the quantity of proliferating cells corresponds to $\int_0^{t_0} p(t, 0) dt$. Thus, the fraction F of quiescent cells at t_0 is

$$F = \frac{Q(t_0)}{Q(t_0) + \int_0^{t_0} p(t, 0) dt}. \quad (3)$$

Now we compute this quantity from model (1), keeping in mind that we have assumed no mortality. We have

$$Q(t_0) = \int_0^{t_0} \frac{dQ}{dt}(t) dt = 2f \int_0^{t_0} \int_0^\infty \beta(a) p(t, a) da dt$$

and

$$\int_0^{t_0} p(t, 0) dt = 2(1-f) \int_0^{t_0} \int_0^\infty \beta(a) p(t, a) da dt.$$

Finally we obtain

$$\frac{Q(t_0)}{Q(t_0) + \int_0^{t_0} p(t, 0) dt} = \frac{2f}{2f + 2(1-f)} = f. \quad (4)$$

So when there is no death rate, the experimental fraction F corresponds exactly to the coefficient f . In our simulations we consider positive a death rate μ as suggested by the results in Section 4.2. But because the numerical value we recover for μ is very small ($\mu \ll 1$ in Fig. 6), we can consider that f is still well-approximated by F and we use the fractions available in Tyson et al. (in press) to parameterize f .

The identification of the coefficients β and μ is much more delicate because $\beta(a)$ is a distributed function. The two following sections are devoted to presenting a method to recover these coefficients from IMT distributions. Then these coefficients are used to compare the solutions of Eq. (1) to experimental data in Section 5.

3. Modelling the intermitotic time

Since the age of a cell is defined as the time elapsed from its last mitosis, the IMT of a cell is its age at division. This definition allows us to interpret the IMT distributions in terms of a dynamic age-structured population model.

Experimental IMT distributions can be seen as histograms which represent, for a given population, the density of cells with a certain age of division (see Fig. 1 for an example taken from Tyson et al., in press). The age of division is distributed into N_a bins of width Δa . For all i between 1 and N_a , the height H_i of the i th bar represents the density of cells with an age of division in the window $[i\Delta a, (i+1)\Delta a]$. The histogram $(H_i)_{i=1}^{N_a}$ is normalized to represent a density:

$$\Delta a \sum_{i=1}^{N_a} H_i = 1. \quad (5)$$

We briefly explain here the method used in Tyson et al. (in press) to build IMT histograms (we refer to this paper for more details). The data are obtained using extended temporally resolved automated microscopy (ETRAM) in which cell nuclei are fluorescently labeled, imaged by automated time lapse fluorescence microscopy, and tracked as individual cells from the resultant image stacks. Cells are subjected to various microenvironmental conditions (such as the addition of a drug) at a specific time during image acquisition (set to time t_0) and the effect of the perturbation from that point in time is followed across the entire population and individual cells within it. The duration of observation (T) is chosen large enough to observe that almost all cycling cells divide before this final time. The data are organized into the bins to obtain a histogram, which is then normalized to represent a density of cells.

The next step is to describe the IMT distributions in terms of an age-structured model and use this interpretation to parameterize Eq. (1). We want to estimate the coefficients β and μ , and for this we fix $f=0$ in (1). This corresponds to the situation when there is no erlotinib treatment and thus all the cells are proliferating. We recover in this case the original McKendrick–Von Foerster model:

$$\begin{cases} \frac{\partial}{\partial t} p(t, a) + \frac{\partial}{\partial a} p(t, a) + \beta(a) p(t, a) + \mu p(t, a) = 0, & t \geq 0, a > 0 \\ p(t, 0) = 2 \int_0^\infty \beta(a) p(t, a) da, \\ p(0, a) = p_0(a). \end{cases} \quad (6)$$

It is a transport equation which satisfies a maximum principle, namely if the initial distribution $p_0(a)$ is nonnegative (positive) then the distribution $p(t, a)$ remains nonnegative (positive) for all time $t > 0$. The solutions of (6) have a remarkable behavior as time evolves, in that the age structure equilibrates no matter what the initial age structure of the population may be. This effect is known as *asynchronous exponential growth*, and its interpretation is that the population disperses over time to a limiting asymptotic equilibrium age structure where the fraction of the population in any age range $[a_1, a_2]$ satisfies

$$\lim_{t \rightarrow \infty} \frac{\int_{a_1}^{a_2} p(t, a) da}{\int_0^\infty p(t, a) da}$$

equals to a constant independent of the initial age structure (Webb, 1987). Moreover, the solutions to this equation, as $t \rightarrow \infty$, are known to behave like a separated variables solution, that is, $p(t, a) = a$ function of time only $\times$ a function of age only. More precisely, consider the eigenvalue problem:

$$\begin{cases} \lambda \widehat{p}(a) + \partial_a \widehat{p}(a) + \beta(a) \widehat{p}(a) + \mu \widehat{p}(a) = 0, \\ \widehat{p}(0) = 2 \int_0^\infty \beta(a) \widehat{p}(a) da, \\ \widehat{p}(\cdot) > 0, \quad \int \widehat{p}(a) da = 1. \end{cases} \quad (7)$$

This problem has a unique solution given by

$$\widehat{p}(a) = \widehat{p}(0) e^{-\int_0^a (\beta(a') + \mu + \lambda) da'},$$

where $\lambda > 0$ is the unique value such that

$$1 = 2 \int_0^\infty \beta(a) e^{-\int_0^a (\beta(a') + \mu + \lambda) da'} da \quad (8)$$

and

$$\widehat{p}(0) = \left(\int_0^\infty e^{-\int_0^a (\beta(a') + \mu + \lambda) da'} da \right)^{-1}.$$

Then we can prove that, for large times,

$$p(t, a) \sim \text{const} \widehat{p}(a) e^{\lambda t}$$

(see Appendix A for more details and references). If the population of cells proliferates over a sufficiently long time, we can assume that this asymptotic behavior is reached and use it to investigate the IMT distributions. An experimental observation that the total population (independent of age structure) is growing exponentially is an indicator that the population has effectively reached the equilibrium age distribution, which can be checked using ETRAM for other time series data collection (see Fig. 2).

Now we give a continuous expression of the IMT distribution in terms of the age-structured model. The age distribution of the cells relative to time t_0 (time of perturbation) is given by a truncation of the equilibrium age distribution $\widehat{p}$:

$$\overline{p}_0(a) = \begin{cases} \rho e^{-\int_0^a (\beta(a') + \mu + \lambda) da'} & \text{if } 0 \leq a \leq t_0, \\ 0 & \text{if } a > t_0, \end{cases}$$

where ρ is a scaling constant. We then follow this age distribution along time and obtain for $t > 0$

$$\overline{p}(t, a) = \begin{cases} \rho e^{-\int_0^a (\beta(a') + \mu + \lambda) da'} e^{\lambda t} & \text{if } t \leq a \leq t + t_0, \\ 0 & \text{otherwise.} \end{cases}$$

According to the age-structured model, and because of the normalization (5), the IMT distribution satisfies

$$I_T(a) := C_T^{-1} \int_0^T \beta(a) \bar{p}(t, a) dt, \quad C_T := \int_0^\infty \int_0^T \beta(a) \bar{p}(t, a) dt da, \quad (9)$$

by definition of the division rate β . The fact that no cell can divide in a time less than t_0 means

$$\forall a \leq t_0, \quad \beta(a) = 0. \quad (10)$$

Under this condition, for T large, the function I_T is close to the function:

$$I_\infty(a) := C_\infty^{-1} \beta(a) e^{-\int_0^a (\beta(a') + \mu) da'} \quad (11)$$

where

$$C_\infty := \int_0^\infty \beta(a) e^{-\int_0^a (\beta(a') + \mu) da'} da. \quad (12)$$

This convergence is made mathematically precise by the following claim (the proof is given in Appendix B for a more general case in which it is not assumed that $p(t, a)$ is close to the equilibrium age distribution):

Claim 1. *Under Assumptions (2) and (10), we have the convergence*

$$\int_0^\infty |I_T(a) - I_\infty(a)| da \xrightarrow{T \rightarrow \infty} 0.$$

Remark 2. We can easily prove that, under the additional assumption

$$\lim_{a \rightarrow +\infty} \beta(a) e^{-\int_0^a (\beta(a') + \mu) da'} = 0, \quad (13)$$

we also have

$$\sup_{a \geq 0} |I_T(a) - I_\infty(a)| \xrightarrow{T \rightarrow +\infty} 0.$$

Condition (13) is satisfied from Assumption (2), if for example, β bounded or monotonic.

Because in the experimental protocol T is chosen large enough to observe no dividing cells at the end, we can approximate I_T by I_∞ which has a simple expression in terms of the division and death rates (11).

4. From the intermitotic time to the division rate

In this section we explain how we can recover the parameters β and μ from the IMT distribution I_∞ . We first present the method in the case when there is no death ($\mu = 0$). This simplification is useful to give an formula inversion which gives the rate β in terms of I_∞ . Then we extend the method to include possible death rates.

4.1. The case $\mu = 0$

In the case $\mu = 0$, the constant C_∞ is equal to 1 and we have

$$I_{\infty}(a) = \beta(a) e^{-\int_0^a \beta(a') da'}.$$

This expression can be inverted to recover the division rate from the IMT distribution (see Hinow et al., 2007; Billy et al., in press)

$$\beta(a) = \frac{I_{\infty}(a)}{\int_a^{\infty} I_{\infty}(a') da'}. \quad (14)$$

We start from the fitting of the experimental IMT distributions, which are observed to be positively skewed. From a fitting procedure for I_{∞} , we use Eq. (14) to recover the division rate of the McKendrick–Von Foerster equation.

First consider as in Hinow et al. (2007) that the IMT distribution is a shifted gamma function (see also Bernard et al., 2003 and references therein). Setting

$$I_{\infty}(a|m, \sigma) = \begin{cases} 0 & \text{if } 0 \leq a \leq m, \\ \frac{a-m}{\sigma^2} e^{-(a-m)/\sigma} & \text{if } a \geq m, \end{cases} \quad (15)$$

we can solve explicitly Eq. (14) and we find

$$\beta(a) = \begin{cases} 0 & \text{if } 0 \leq a \leq m, \\ \frac{a-m}{\sigma(\sigma+a-m)} & \text{if } a \geq m. \end{cases} \quad (16)$$

So by fitting an experimental IMT distribution with a shifted gamma function, we obtain two parameters m and σ which allows reconstruction of the division rate of the renewal equation.

To have a smoother transition at the minimum age of division m , one can consider a second shifted gamma function:

$$I_{\infty}(a|m, \sigma) = \begin{cases} 0 & \text{if } 0 \leq a \leq m, \\ \frac{(a-m)^2}{2\sigma^3} e^{-(a-m)/\sigma} & \text{if } a \geq m. \end{cases} \quad (17)$$

Then the corresponding β is

$$\beta(a) = \begin{cases} 0 & \text{if } 0 \leq a \leq m, \\ \frac{1}{\sigma} \frac{(a-m)^2}{2\sigma^2 + 2\sigma(a-m) + (a-m)^2} & \text{if } a \geq m. \end{cases} \quad (18)$$

The different functions (15)–(18) are plotted in Fig. 3

It has been observed that an exponentially modified Gaussian (EMG) is often a better model for IMT distributions than the gamma function (Golubev, 2010; Tyson et al., in press). An EMG is defined as the convolution of a Gaussian with a decreasing exponential, but after solving it can be written with three parameters as

$$I_{\infty}(a|\beta_0, m, \sigma) = \beta_0 \text{Erfc}\left(\frac{m-a}{\sigma}\right) e^{-2\beta_0((\beta_0\sigma^2/2)-m+a)}. \quad (19)$$

where the (complementary) error function is defined by

$$\operatorname{Erfc}(z) = 1 - \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt.$$

Replacing I_∞ by an EMG in Eq. (14), we cannot compute explicitly the expression for β . But by numerical comparison, we obtain a division rate β that is essentially indistinguishable from an error function (see Fig. 4).

Instead of fitting the IMT distribution by an EMG and then fitting the corresponding β by an error function, we may directly assume that β is an error function:

$$\beta(a) = \beta_0 \operatorname{Erfc}\left(\frac{m-a}{\sigma}\right). \quad (20)$$

We can then explicitly derive a new fitting formula for I_∞ due to Eq. (11)

$$I_\infty(a|\beta_0, m, \sigma) = \beta_0 \operatorname{Erfc}\left(\frac{m-a}{\sigma}\right) e^{-\int_0^a \beta_0 \operatorname{Erfc}\left(\frac{m-a'}{\sigma}\right) da'}, \quad (21)$$

where the integral in the exponential can be computed as

$$\int_0^a \operatorname{Erfc}\left(\frac{m-a'}{\sigma}\right) da' = m \operatorname{Erfc}\left(\frac{m}{\sigma}\right) - \frac{\sigma}{\sqrt{\pi}} e^{-(m/\sigma)^2} - (m-a) \operatorname{Erfc}\left(\frac{m-a}{\sigma}\right) + \frac{\sigma}{\sqrt{\pi}} e^{-((m-a)/\sigma)^2}. \quad (22)$$

Using this formula for the IMT instead of an EMG formula, the fitting parameters provide immediately the division rate β .

When we know the Malthusian growth parameter (population growth rate λ) from experimental data (see Fig. 2), it is possible to recover a division rate β such that relation (8) is satisfied. This is an important step in order to parameterize and apply an age-structured model to experimental data. We notice the relationship of β in terms of $I_\infty(a)$ is

$$2 \int_0^\infty I_\infty(a) e^{-\lambda a} da = 2 \int_0^\infty \beta(a) e^{-\int_0^a \beta(a') da'} e^{-\lambda a} da = 1.$$

To take advantage of this relationship between the IMT data, $\beta(a)$, and the Malthusian parameter λ at the equilibrium distribution, we define a new histogram $(\tilde{H}_i)_{1 \leq i \leq N}$ by

$$\forall i, \quad \tilde{H}_i := \frac{2H_i e^{-\lambda a_i}}{\Delta a \sum_{i=1}^N 2H_i e^{-\lambda a_i}}, \quad (23)$$

where $a_i := (i + \frac{1}{2}) \Delta a$ is the mean age of the i th bar. Thus defined, the new histogram incorporates information about λ and satisfies the relation:

$$\Delta a \sum_{i=1}^\infty \tilde{H}_i = 1. \quad (24)$$

We then fit this new histogram $(\tilde{H}_i)$ with the model

$$\tilde{I}_\infty(a) := 2I_\infty(a) e^{-\lambda a} \quad (25)$$

instead of fitting (H_i) with $I_{\infty}(a)$. Because of the normalization (24), we expect that fitting provides parameters such that $\int_0^{\infty} \tilde{I}_{\infty}$, and this relation can be checked numerically *a posteriori*.

Method and example: We divide the process into three steps and illustrate it by an example. Each required fitting step can be performed using the freely available Ezyfit Matlab toolbox <<http://www.fast.u-psud.fr/ezyfit/S>>.

Step 1: Determine equilibrium IMT distribution—

- a. Obtain a histogram (H_i) for the experimental IMT distribution of control (untreated) cells. *Here we use an IMT distribution obtained in Tyson et al. (in press) for PC-9 lung cancer cells using ETRAM* (see Fig. 1).
- b. Plot the time evolution of the total population on a $\log_e$ -scale, verify a linear fit and obtain the slope as the experimental value for the Malthusian parameter λ of the cell population (see Fig. 2 for an example).
- c. Construct the new histogram $(\tilde{H}_i)$ from (H_i) and λ by using the definition (23).

Step 2: Obtain parameters from model fit to IMT distribution—

- a. Choose a form for I_{∞} as a gamma function (15) or (17), or as the new EMG form (21).
- b. Fit the histogram $(\tilde{H}_i)$ with the corresponding form $\tilde{I}_{\infty}$ from definition (25). *For PC-9 cancer cells we choose the form (21), because in Tyson et al. (in press) it was observed that the IMT distribution appeared to be an EMG* (see Fig. 5 for the example).
- c. Verify that the correlation coefficient R^2 of the IMT data and the chosen form $\tilde{I}_{\infty}$ is close to 1.

Step 3: Parameterize age-structured model—If numerical integral $\int_0^{\infty} \tilde{I}_{\infty}(a)$ is close to 1, the fitting parameters provide a good approximation of the division rate β , which can then be propagated through the population, based on the choice of I_{∞} . *In the example, β is given by Eq. (20) with the numerical parameters of Fig. 5.*

4.2. Introducing a death rate

If we include a death rate $\mu > 0$ in the model, then there does not exist an inversion formula as Eq. (14) to recover β analytically. So we consider that β is a rational fraction given by Eq. (16) or (18), or an error function given by Eq. (20), and we derive the corresponding model I_{∞} from Eq. (11). The death rate μ will appear in the expression of I_{∞} as an additional fitting parameter and will be determined together with the parameters of β by the fitting of experimental IMT distributions. Note that a fixed rate μ based on measured data can also be used if available to avoid the addition of a free parameter.

Unlike the case $\mu = 0$ for which $C_{\infty} = 1$, we have for a positive death rate that $C_{\infty} < 1$. This constant is a function of the fitting parameters (see Eq. (12)), but we do not have an analytic expression of this function in general. A solution to this problem is to use the Malthusian parameter λ , which incorporates information about μ and $\beta(a)$, to define a form $\tilde{I}_{\infty}$ as in Section 4.1. Indeed we obtain, because of relation (8),

$$\tilde{I}_{\infty}(a) := 2\beta(a) e^{-\int_0^a (\beta(a') + \mu) da'} e^{-\lambda a},$$

which does not involve C_{∞} . Then, the division rate $\beta(a)$ is obtained from $\tilde{I}_{\infty}(a)$ as before using the modified histogram $(\tilde{H}_i)$ defined from (H_i) by (23).

Example. We fit the same distribution as in Section 4.1 still considering that β is an error function. With a constant death rate μ , we obtain the four parameters model:

$$\tilde{I}_{\infty}(a|\beta_0, m, \sigma, \mu) = 2\beta_0 \operatorname{Erfc}\left(\frac{m-a}{\sigma}\right) e^{-\int_0^a \beta_0 \operatorname{Erfc}\left(\frac{m-a'}{\sigma}\right) + \mu da'}, \quad (26)$$

where the integral $\int_0^a \beta_0 \operatorname{Erfc}\left(\frac{m-a'}{\sigma}\right) da'$ is given by Eq. (22). The fitting provides new parameters for β and a positive death rate μ (see Fig. 6). The correlation R^2 is slightly better than in Fig. 5 and the integral $\int_0^{\infty} \tilde{I}_{\infty}(a) da$ is significantly closer to 1. So we can assume that mortality has to be considered for this cell line.

5. Numerical simulation and discussion

Once the parameters accurately estimated, we use them to compare the behavior of the solution of Eq. (1) to the experimental data in Tyson et al. (in press). Since the model does not have analytic solutions, we need to perform numerical simulations. Many numerical methods are available for structured population equations (see for instance Ackleh and Ito, 1997; Angulo and López-Marcos, 2000, 2004; Douglas and Milner, 1987; Gabriel and Tine, 2010; Kostova, 2003). Here we use a scheme based on the method of characteristics, as in Angulo and López-Marcos (2000, 2004) and Kostova (2003), for its anti-dissipative properties.

We can see in Fig. 7 that the numerical simulations are very similar to the experimental curves. In particular, the delay of 20 h before the effect of the treatment on the growth of the population is apparent, and this 20 h period pulses two more times as the population approaches a new equilibrium distribution during the total time of the experiment. We have thus developed an age structured model that can explain the dynamic effects of erlotinib on PC-9 cells, which are intrinsically dependent on the age of proliferating cells. The model is also accurate in a quantitative point of view. The quiescent fractions f which are used in the numerical simulations are not chosen arbitrarily to obtain the adequate behavior. They are linked to the treatment dose of erlotinib and estimated from experimental data in a rigorous way so that they are realistic.

The model has been chosen as simple as possible to be able to recover all the parameters from experimental IMT distributions. This simplicity leads to qualitative differences between observed data and simulations in Fig. 7. Experimental observation suggests the population reaches equilibrium when treated with drug whereas simulated population size is still increasing at the end of the experiment. An age-structure for quiescent cells together with an age-dependent rate of death should be considered to explain this plateau effect. Experimental data also suggest that drug treatment increases population growth rate above that of untreated cells briefly, but model does not capture this. This could be obtained by considering a lower death rate for quiescent cells than for proliferating cells. But for such more complex models, additional experimental data and a new parameter estimation method would be necessary to estimate the death rates.

6. Conclusion

Linking experimental observation of cell behavior between the single-cell and population scales has recently been described using newly developed mathematical models (Tyson et al., in press). However, this approach does not take into account the possible cell age-dependent effects of a perturbation on cell behavior, such as would be expected if the effects occur in cells at a specific position in the cell cycle. Since these studies were performed with asynchronously dividing cell populations, it is evident that the mathematical models of these experiments should have age structure as a primary feature. In fact, to fit the data, the authors had to use an artificial time offset to account for the age-structured effects. Here we provide a formalized approach to accurately account for cell age-dependent effects on cellular behavior. A major difficulty in the parameterization of age-structured models is the determination of the age dependent division rate. Our study provides a method for the quantitative recovery of this rate by fitting experimental IMT distributions to special forms, such as gamma functions or exponential modified Gaussians. This model, once successfully parameterized, is very useful for simulating and analyzing age-dependent phenomena in cell population dynamics. We have presented one such application for the *in vitro* treatment of cancer cells by erlotinib. This example shows the utility of age-structured population models in explaining the connection of drug therapy to phenomena such as cell cycle phase entry into quiescence. The method we have presented can be implemented readily to many issues in cell population behavior when there is experimental data based on cell age.

Acknowledgments

The research stays of P. Gabriel at Vanderbilt University have been financially supported by a grant of the Fondation Pierre Ledoux Jeunesse Internationale.

Appendix A. Convergence to the equilibrium

The long-time behavior $p(t, a) \sim \text{const } \widehat{p}(a) e^{\lambda t}$ can be proven by using semi-groups methods (Metz and Diekmann, 1983; Webb, 1985) or General Relative Entropy techniques (Michel et al., 2005; Perthame, 2007). We detail here a result provided by General Relative Entropy.

Consider ϕ the unique solution to the adjoint eigenvalue problem:

$$\begin{cases} \lambda \phi(a) - \partial_a \phi(a) + \beta(a) \phi(a) + \mu \phi(a) = 2\phi(0) \beta(a), \\ \phi(\cdot) \geq 0, \quad \int \widehat{p}(a) \phi(a) da = 1. \end{cases}$$

Then the General Relative Entropy method allows us to prove that

$$\int_0^\infty |p(t, a) e^{-\lambda t} - \rho_0 \widehat{p}(a)| \phi(a) da \xrightarrow{t \rightarrow \infty} 0,$$

where

$$\rho_0 = \int_0^\infty \phi(a) p_0(a) da.$$

Appendix B. Convergence of the IMT distribution

Theorem 3. Suppose that Assumptions (2) and (10) are satisfied and consider an initial distribution $\bar{p}_0$ such that $\bar{p}_0(a) = 0$ for all $a > t_0$. Then we have the convergence

$$\int_0^\infty |I_T(a) - I_\infty(a)| da \xrightarrow{T \rightarrow \infty} 0,$$

where I_T and I_∞ are defined in (9) and (11).

For the sake of simplicity, the proof of Theorem 3 is given in the case $\mu = 0$. But it can be easily adapted to the case with a death rate.

Proof of Theorem 3 in the case $\mu = 0$. First we extend the initial distribution and the division rate $\beta(a)$ to negative ages by setting $\bar{p}_0(a) = \beta(a) = 0$ for $a < 0$. Since the daughters of the labeled cells are not tracked, the age distribution $\bar{p}(t, a)$ of labeled cells satisfies a transport equation without boundary condition. Thus it writes, using the characteristic method,

$$\forall t, a \geq 0, \quad \bar{p}(t, a) = \bar{p}_0(a - t) e^{-\int_0^t \beta(a-s) ds}.$$

Introducing this expression in the definition of $I_T(a)$ we obtain, with changes of variables,

$$\begin{aligned} I_T(a) &= C_T^{-1} \int_0^T \beta(a) \bar{p}_0(a - t) e^{-\int_0^t \beta(a-s) ds} dt \\ &= C_T^{-1} \int_0^T \beta(a) \bar{p}_0(a - t) e^{-\int_{a-t}^a \beta(a') da'} dt \\ &= C_T^{-1} \int_{a-T}^a \beta(a) \bar{p}_0(u) e^{-\int_0^a \beta(a') da'} du \\ &= C_T^{-1} \beta(a) e^{-\int_0^a \beta(a') da'} \int_{a-T}^a \bar{p}_0(u) e^{\int_0^u \beta(a') da'} du. \end{aligned}$$

Since the support of $\bar{p}_0$ is included in $[0, t_0]$ and $\beta(a) = 0$ for $a \in [0, t_0]$ due to Assumption (2), we have for all $u \in \mathbb{R}$

$$\bar{p}_0(u) e^{\int_0^u \beta(a') da'} = \bar{p}_0(u),$$

so $I_T(a)$ writes

$$I_T(a) = C_T^{-1} \beta(a) e^{-\int_0^a \beta(a') da'} \int_{a-T}^a \bar{p}_0(u) du.$$

Define the primitive

$$P(a) := \int_0^a \bar{p}_0(u) du.$$

This function is nondecreasing and satisfies $P(a) = 0$ for $a < 0$ and $P(a) = P(t_0) = \int_0^\infty \bar{p}_0$ for $a > t_0$ so we have

$$\int_{a-T}^a \bar{p}_0(u) du = P(a) - P(a-T) = \begin{cases} P(a) & \text{if } 0 \leq a \leq t_0, \\ P(t_0) & \text{if } t_0 \leq a \leq T, \\ P(t_0) - P(a-T) & \text{if } T \leq a \leq T+t_0, \\ 0 & \text{if } a \geq T+t_0. \end{cases}$$

Because $\beta(a) = 0$ for $a > t_0$, we obtain by integration of $I_T(a)$ on $\mathbb{R}^+$

$$\begin{aligned} 1 &= C_T^{-1} P(t_0) \int_0^T \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &\quad + C_T^{-1} \int_T^{T+t_0} \beta(a) e^{-\int_0^a \beta(a') da'} (P(t_0) - P(a-T)) da \end{aligned}$$

which gives

$$\begin{aligned} C_T - P(t_0) &= \int_0^T \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &= \int_T^{T+t_0} \beta(a) e^{-\int_0^a \beta(a') da'} (P(t_0) - P(a-T)) da \\ &\leq P(t_0) \int_T^{T+t_0} \beta(a) e^{-\int_0^a \beta(a') da'} da \xrightarrow{T \rightarrow +\infty} 0. \end{aligned}$$

Since

$$\int_0^\infty \beta(a) e^{-\int_0^a \beta(a') da'} da = 1,$$

we conclude that

$$C_T \xrightarrow{T \rightarrow +\infty} P(t_0) = \int_0^\infty \bar{p}_0(a) da.$$

Then we have

$$\begin{aligned} \int_0^\infty |I_T(a) - I_\infty(a)| da &\leq |C_T^{-1} P(t_0) - 1| \int_0^T \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &\quad + \int_T^{T+t_0} |C_T^{-1} [P(t_0) - P(a-T)] - 1| \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &\quad + \int_{T+t_0}^\infty \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &\leq |C_T^{-1} P(t_0) - 1| \int_0^T \beta(a) e^{-\int_0^a \beta(a') da'} da \\ &\quad + K \int_T^\infty \beta(a) e^{-\int_0^a \beta(a') da'} da \xrightarrow{T \rightarrow \infty} 0 \end{aligned}$$

and it ends the proof of Theorem 3.

References

Ackleh AS. Parameter identification in size-structured population models with nonlinear individual rates. *Math. Comp. Modelling.* 1999; 30(9-10):81–92.

- Ackleh AS, Ito K. An implicit finite difference scheme for the nonlinear size-structured population model. *Numer. Funct. Anal. Optim.* 1997; 18(9-10):865–884.
- Ainseba B, Benosman C. CML dynamics: optimal control of age-structured stem cell population. *Math. Comput. Simul.* 2011; 81(10):1962–1977.
- Ainseba B, Picart D, Thiéry D. An innovative multistage, physiologically structured, population model to understand the European grapevine moth dynamics. *J. Math. Anal. Appl.* 2011; 382(1):34–46.
- Alarcón T, Byrne H, Maini P. A cellular automaton model for tumour growth in inhomogeneous environment. *J. Theor. Biol.* 2003; 225(2):257–274. [PubMed: 14575659]
- Angulo O, López-Marcos JC. Numerical integration of nonlinear size-structured population equations. *Ecol. Modelling.* 2000; 133(1-2):3–14.
- Angulo O, López-Marcos J. Numerical integration of fully nonlinear size-structured population models. *Appl. Numer. Math.* 2004; 50(3-4):291–327.
- Arino O, Sánchez E, Webb GF. Necessary and sufficient conditions for asynchronous exponential growth in age structured cell populations with quiescence. *J. Math. Anal. Appl.* 1997; 215(2):499–513.
- Banks HT, Banks JE, Dick LK, Stark JD. Estimation of dynamic rate parameters in insect populations undergoing sublethal exposure to pesticides. *Bull. Math. Biol.* 2007; 69:2139–2180. [PubMed: 17453304]
- Banks HT, Sutton KL, Thompson WC, Bocharov G, Doumic M, Schenkel T, Arguilaguet J, Giest S, Peligero C, Meyerhans A. A new model for the estimation of cell proliferation dynamics using CFSE data. *J. Immunol. Methods*, to appear.
- Bekkal Brikci F, Clairambault J, Perthame B. Analysis of a molecular structured population model with possible polynomial growth for the cell division cycle. *Math. Comput. Modelling.* 2008a; 47(7-8):699–713.
- Bekkal Brikci F, Clairambault J, Ribba B, Perthame B. An age-and-cyclin-structured cell population model for healthy and tumoral tissues. *J. Math. Biol.* 2008b; 57(1):91–110. [PubMed: 18064465]
- Bernard S, Pujo-Menjouet L, Mackey MC. Analysis of cell kinetics using a cell division marker: mathematical modeling of experimental data. *Biophys. J.* 2003; 84(5):3414–3424. [PubMed: 12719268]
- Billy F, Clairambault J, Fercoq O, Gaubert S, Lepoutre T, Ouillon T, Saito S. Synchronisation and control of proliferation in cycling cell population models with age structure. *Math. Comput. Simulation*. in press.
- Calsina A, Saldaña J. A model of physiologically structured population dynamics with a nonlinear individual growth rate. *J. Math. Biol.* 1995; 33:335–364.
- Clément, P.; Heijmans, HJAM.; Angenent, S.; van Duijn, CJ.; de Pagter, B. *One-Parameter Semigroups*, CWI Monographs. Vol. 5. North-Holland Publishing Co.; Amsterdam: 1987.
- Diekmann O, Heijmans HJAM, Thieme HR. On the stability of the cell size distribution. *J. Math. Biol.* 1984; 19:227–248.
- Douglas J, Milner F. Numerical methods for a model of population dynamics. *Calcolo.* 1987; 24:247–254.
- Doumic M, Perthame B, Zubelli J. Numerical solution of an inverse problem in size-structured population dynamics. *Inverse Problems.* 2009; 25(4):045008.
- Doumic M, Maia P, Zubelli J. On the calibration of a size-structured population model from experimental data. *Acta Biotheor.* 2010; 58:405–413. [PubMed: 20676731]
- Dowling MR, Milutinovic D, Hodgkin PD. Modelling cell lifespan and proliferation: is likelihood to die or to divide independent of age. *J. R. Soc. Interface.* 2005; 2(5):517–526. [PubMed: 16849210]
- Engel, K-J.; Nagel, R. *One-parameter semigroups for linear evolution equations*. In: Brendle, S.; Campiti, M.; Hahn, T.; Metafune, G.; Nickel, G.; Pallara, D.; Perazzoli, C.; Rhandi, A.; Romanelli, S.; Schnaubelt, R., editors. *Graduate Texts in Mathematics*. Vol. 194. Springer-Verlag; New York: 2000.
- Engl H, Rundell W, Scherzer O. A regularization scheme for an inverse problem in age-structured populations. *J. Math. Anal. Appl.* 1994; 182(3):658–679.

- Gabriel P. The shape of the polymerization rate in the prion equation. *Math. Comput. Modelling.* 2011; 53(7-8):1451–1456.
- Gabriel P, Tine LM. High-order WENO scheme for polymerization-type equations. *ESAIM Proc.* 2010; 30:54–70.
- Gerlee P, Anderson A. A hybrid cellular automaton model of clonal evolution in cancer: the emergence of the glycolytic phenotype. *J. Theor. Biol.* 2008; 250(4):705–722. [PubMed: 18068192]
- Golubev A. Exponentially modified Gaussian (EMG) relevance to distributions related to cell proliferation and differentiation. *J. Theor. Biol.* 2010; 262(2):257–266. [PubMed: 19825376]
- Gyllenberg M, Webb GF. Age-size structure in populations with quiescence. *Math. Biosci.* 1987; 86(1):67–95.
- Gyllenberg M, Webb GF. A nonlinear structured population model of tumor growth with quiescence. *J. Math. Biol.* 1990; 28(6):671–694. [PubMed: 2250141]
- Gyllenberg, M.; Webb, GF. *Mathematical Population Dynamics*. Vol. 131. New Brunswick, NJ: 1991. Quiescence in structured population dynamics: applications to tumor growth; p. 45-62. 1989). *Lecture Notes in Pure and Applied Mathematics* Dekker, New York
- Gyllenberg M, Osipov A, Päivarintä L. The inverse problem of linear age-structured population dynamics. *J. Evol. Equation.* 2002; 2(2):223–239.
- Hinow P, Wang S, Arteaga C, Webb G. A mathematical model separates quantitatively the cytostatic and cytotoxic effects of a HER2 tyrosine kinase inhibitor. *Theor. Biol. Med. Modelling.* 2007; 4(1):14.
- Kostova T. An explicit third-order numerical method for size-structured population equations. *Numer. Methods Partial Differential Equation.* 2003; 19(1):1–21.
- Lee H, Perelson A. Modeling *t* cell proliferation and death in vitro based on labeling data: generalizations of the Smith–Martin cell cycle model. *Bull. Math. Biol.* 2008; 70:21–44. [PubMed: 17701260]
- Liu D, Umbach DM, Peddada SD, Li L, Crockett PW, Weinberg CR. A random-periods model for expression of cell-cycle genes. *Proc. Natl. Acad. Sci. USA.* 2004; 101(19):7240–7245. [PubMed: 15123814]
- Mallet D, Pillis LD. A cellular automata model of tumor-immune system interactions. *J. Theor. Biol.* 2006; 239(3):334–350. [PubMed: 16169016]
- Metz, JAJ.; Diekmann, O., editors. *The Dynamics of Physiologically Structured Populations*. *Lecture Notes in Biomathematics*. Vol. 68. Springer-Verlag; Berlin: 1983. 1986. papers from the Colloquium held in Amsterdam
- Michel P, Mischler S, Perthame B. General relative entropy inequality: an illustration on growth models. *J. Math. Pures Appl.* 2005; 84(9):1235–1260.
- Patel AA, Gawlinski ET, Lemieux SK, Gatenby RA. A cellular automaton model of early tumor growth and invasion: the effects of native tissue vascularity and increased anaerobic tumor metabolism. *J. Theor. Biol.* 2001; 213(3):315–331. [PubMed: 11735284]
- Perthame, B. *Transport equations in biology*, *Frontiers in Mathematics*. Birkhäuser Verlag; Basel: 2007.
- Perthame B, Zubelli J. On the inverse problem for a size-structured population model. *Inverse Problems.* 2007; 23(3):1037–1052.
- Picart D, Ainseba B. Parameter identification in multistage population dynamics model. *Nonlinear Anal.: Real World Appl.* 2011; 12(6):3315–3328.
- Picart D, Ainseba B, Milner F. Optimal control problem on insect pest populations. *Appl. Math. Lett.* 2011; 24(7):1160–1164.
- Pilant M, Rundell W. Determining a coefficient in a first-order hyperbolic equation. *SIAM J. Appl. Math.* 1991a; 51(2):494–506.
- Pilant M, Rundell W. Determining the initial age distribution for an age structured population. *Math. Popul. Stud.* 1991b; 3(1):3–20. [PubMed: 12343114]

- Powathil GG, Gordon KE, Hill LA, Chaplain MA. Modelling the effects of cell-cycle heterogeneity on the response of a solid tumour to chemotherapy: biological insights from a hybrid multiscale cellular automaton model. *J. Theor. Biol.* 2012; 308:1–19. [PubMed: 22659352]
- Quaranta V, Weaver AM, Cummings PT, Anderson AR. Mathematical modeling of cancer: the future of prognosis and treatment. *Clin. Chim. Acta.* 2005; 357(2):173–179. [PubMed: 15907826]
- Quaranta V, Rejniak KA, Gerlee P, Anderson AR. Invasion emerges from cancer cell adaptation to competitive microenvironments: quantitative predictions from multiscale mathematical models. *Semin. Cancer Biol.* 2008; 18(5):338–348. [PubMed: 18524624]
- Ramis-Conde I, Drasdo D, Anderson AR, Chaplain MA. Modeling the influence of the e-cadherin–catenin pathway in cancer cell invasion: a multiscale approach. *Biophys. J.* 2008; 95(1):155–165. [PubMed: 18339758]
- Ribba, B.; Alarcón, T.; Marron, K.; Maini, P.; Agur, Z. The use of hybrid cellular automaton models for improving cancer therapy. In: Sloot, P.; Chopard, B.; Hoekstra, A., editors. *Cellular Automata. Lecture Notes in Computer Science.* Vol. 3305. Springer; Berlin, Heidelberg: 2004. p. 444–453.
- Rundell W. Determining the birth function for an age structured population. *Math. Popul. Stud.* 1989; 1(4):377–395. [PubMed: 12281955]
- Rundell W. Determining the death rate for an age-structured population from census data. *SIAM J. Appl. Math.* 1993; 53(6):1731–1746.
- Tyson DR, Garbett SP, Frick PL, Quaranta V. A method to quantify population-level cell proliferation dynamics from single-cell data. *Nat. Methods.* in press.
- Webb, GF. *Theory of nonlinear age-dependent population dynamics*, Monographs and Textbooks in Pure and Applied Mathematics. Vol. 89. Marcel Dekker Inc.; New York: 1985.
- Webb GF. An operator-theoretic formulation of asynchronous exponential growth. *Trans. Am. Math. Soc.* 1987; 303(2):751–763.

HIGH LIGHTS

- ▶ We present a method to recover the division rate of an age-structured model from experimental data.
- ▶ This method is based on the fitting of experimental intermitotic time distributions.
- ▶ We present an application of the results concerning the treatment of cancer cells by erlotinib.

Fig. 1.
 Plot of an histogram $(H_i)_{1 \leq i \leq N_a}$ representing the IMT distribution of a population of PC-9 lung cancer cells. The data are from Tyson et al. (in press) and the bin width is $\Delta a = \frac{10}{8}$.

Fig. 2.

The evolution of the total population $N(t)$ is plotted on log-scale (blue) (the data are from Tyson et al., in press). The data are well-fitted with a line of slope $\lambda = 0.022$ (correlation coefficient: $R^2 = 0.9967$). This is evidence that the total population is growing exponentially fast with exponential constant λ . Thus, $N(t) = \exp^{\lambda t} N(0)$, and if t^* is the population doubling time, then $N(t^*) = 2N(0)$, and $t^* = \ln 2 / \lambda$. (For interpretation of the references to color in this figure caption, the reader is referred to the web version of this article.)

Fig. 3. The two different gamma functions and their corresponding division rate $\beta(a)$ are plotted for coefficients $m=17$ and $\sigma=2$.

Fig. 4.

Left: The division rate $\beta(a)$ is obtained numerically from (14) by using an EMG with coefficients $m=22$, $\sigma=2$ and $\beta_0=0.2$ to fit I_∞ (19). Right: The formula for β , as an error function with these same three parameters, is numerically indistinguishable from the numerically obtained β ($R^2 = 1$).

Fig. 5.

Fitting of experimental data in Tyson et al. (in press) with the model (21). The fitting parameters are $\beta_0 = 0.14204$, $m = 24.456$ and $\sigma = 3.3451$. The correlation coefficient is $R^2 = 0.95363$ and the integral of $\tilde{I}_\infty(a|\beta_0, m, \sigma)$ is $\int_0^\infty \tilde{I}_\infty(a) da \approx 1.0983$. The formula for the age dependent division rate is $\beta(a) = \beta_0 \text{Erfc}((m-a)/\sigma)$.

Fig. 6.

Fitting of experimental data with the model (26). The fitting parameters are $\beta_0 = 0.17879$, $m = 25.007$, $\sigma = 3.6141$ and $m = 0.00333$. The correlation coefficient is $R^2 = 0.95611$ and the integral of $\tilde{I}_\infty(a|\beta_0, m, \sigma)$ is $\int_0^\infty \tilde{I}_\infty(a) da \approx 1.0132$. The formula for the age dependent division rate is $\beta(a) = \beta_0 \text{Erfc}((m-a)/\sigma)$.

Fig. 7.

Left: Experimental data without erlotinib, with a 50 nM dose and with a 5000 nM dose. The total quantity $N(t)$ is plotted on a log-scale. Right: Numerical simulation of model (1) for $f=0$, $f=0.6$ and $f=0.84$. The curves represent the evolution of $\ln(P(t)+Q(t))$ where

$P(t) := \int_0^\infty p(t,a) da$ is the total quantity of proliferating cells at time t . They are obtained by solving Eq. (1) numerically with the parameters of Fig. 6 and with $\nu = 0.004$.