Published in final edited form as:

J Perinat Med. 2011 May ; 39(3): 227–236. doi:10.1515/JPM.2011.035.

Listeriosis in Human Pregnancy: a systematic review

Ronald F. Lamont, BSc, MB, ChB, MD, FRCOG^{1,2}, Jack Sobel, MD³, Shali Mazaki-Tovi, MD^{1,2}, Juan Pedro Kusanovic, MD^{1,2}, Edi Vaisbuch, MD^{1,2}, Sun Kwon Kim, MD, PhD¹, Niels Uldbjerg, MD DMSc¹, and Roberto Romero, MD^{1,2,4}

¹Perinatology Research Branch, NICHD/NIH/DHHS, Bethesda, Maryland and Detroit, Michigan, USA

²Wayne State University School of Medicine, Department of Obstetrics and Gynecology, Detroit, Michigan, USA

³Wayne State University School of Medicine, Department of Infectious Diseases, Detroit, Michigan, USA

⁴Center for Molecular Medicine and Genetics, Wayne State University, Detroit, Michigan, USA

Abstract

Listeria is commonly found in processed and prepared foods and listeriosis is associated with high morbidity and mortality. Preventative measures are well prescribed and monitoring and voluntary recall of contaminated products has resulted in a 44% reduction in the prevalence of perinatal listeriosis in the USA. Pregnant women are at high risk for listeriosis, but symptoms are non-specific and diagnosis is difficult. The intracellular life-cycle of *Listeria* protects the bacterium from host innate and adaptive immune responses. Antibiotic treatment requires agents able to penetrate, distribute, and remain stable within host cells. Prolonged use of high-dose ampicillin can significantly improve neonatal outcome.

Keywords

Listeriosis; *Listeria monocytogenes*; infection; foodborne; pregnancy; newborn; neonate; Epidemics

INTRODUCTION

Since the 1960s, and following the introduction and widespread use of refrigerators, processed foods, and the extended shelf-life of foods, listeriosis due to *Listeria monocytogenes* has become more widespread. Although rare in pregnancy, because of the potential for serious feto-maternal consequences, it is important that practicing obstetricians are familiar with the diagnosis, pathogenesis, treatment and prevention of *Listeria* infection. Accordingly, to review this problem, we conducted a computerized search using PubMed, Embase, Cinahl, Lilacs, ISI Web of Science, the Cochrane Central Register of Controlled Trials (all from inception or 1960 to September 2010), and Research Registries of ongoing trials using a combination of key and text words and Medical Subject Headings related to *Listeria monocytogenes*, listeriosis, Pregnancy, Newborn, Neonate, Infectious Disease Transmission, Vertical, Epidemics, and Food-borne infection.

Address correspondence to: Ronald F. Lamont, BSc, MB, ChB, MD, FRCOG and Roberto Romero, M.D., Perinatology Research Branch, NICHD, NIH, DHHS, Wayne State University/Hutzel Women's Hospital, 3990 John R, Box 4, Detroit, MI 48201, USA, Telephone (313) 993-2700, Fax: (313) 993-2694, rlamont@med.wayne.edu and prbchiefstaff@med.wayne.edu.

EPIDEMIOLOGY

In 1997, listeriosis was estimated to be responsible for 500 deaths annually in the USA [99], and in 2000 the prevalence was estimated to be 4 per million population [20]. In Europe, rates vary between 0.1 and 11.3 per million population, and approximately 20% involve neonatal infection [117]. Though no clear seasonal pattern exists, illness in humans tends to present in late summer and early autumn [81,98]. When the disease occurs in children, up to 54% have no apparent immunocompromising condition [22,151].

Listeriosis is mainly a food-borne infection, and sporadic cases as well as epidemics have been linked to contaminated food [18,43,85,87,130,144]. Human listeriosis typically occurs sporadically, but occasional nosocomial clusters have been reported [41,83,139]. In the USA, routine screening and surveillance often leads to a recall of food products due to concerns with respect to listerial contamination [72]. Since the same serotypes cause disease in humans and animals [123], it was originally believed that the disease was caused by direct transmission from animals; however, this has only rarely been documented [14]. In sporadic outbreaks, 11% of all food samples retrieved from refrigerators were contaminated and 64% of the refrigerators of infected cases contained at least one contaminated food item [112,134]. Table 1 demonstrates steps that can be taken by those at risk for listeriosis that help prevent the disease, based on foods known to be associated with infection or past outbreaks.

MICROBIOLOGY

Although, Hayem (France, 1891) and Henle (Germany, 1893) observed gram positive rods in the tissue sections from patients who died of a disease that, in retrospect, was almost certainly listeriosis, Hülphers (Sweden, 1911) appears to be the first to have recorded an encounter with the organism now known as *Listeria monocytogenes* [49]. Reflecting the association with liver abscesses and monocytosis in laboratory animals [26,103] (which is not a feature of disease in humans), the organism has been known as *Bacillus hepatitis* (1911) [66], *Bacterium monocytogenes* (1926) [33], *Listerella hepatolytica* (1927) [113], and following the first report in humans, *Bacterium monocytogenes hominis* (1929) [107]. The current name, *Listeria monocytogenes* [57], was finally agreed upon in 1940 in honour of Lord Joseph Lister, the “father of antisepsis” [45]. Of the seven species of *Listeria*, only four infect humans. Infection with *Listeria ivanii* is rare, and most cases, are due to three serotypes of *Listeria monocytogenes* (1/2a, 1/2b, and 4b), the last of which is responsible for the vast majority of listeriosis outbreaks [14,96,110,112].

L. monocytogenes is a small, facultatively anaerobic, Gram-positive, flagellated, linear motile rod, which is non-spore forming and microscopically difficult to distinguish from commensal *Diphtheroids* [17,106]. Therefore, it was initially thought to be of the genus *Corynebacterium* [80]. For an historical perspective on the microbiology of *Listeria*, the interested reader is referred to a comprehensive review by Farber and Peterkin [39].

Listeria grows well in broth, blood agar and most routine culture media [63]. Nevertheless, selective media, enrichment, and subculture may be necessary to avoid false-negatives from overgrowth of Gram-negative organisms from contaminated sites like the vagina and rectum [30,75,82]. Human pathogenic species produce a narrow zone of β -hemolysis on sheep blood agar [38,47,147], but while suspected to be a virulence factor, the presence of listeriolysin does not correlate with the severity of disease [44,54,78,79,109,140].

L. monocytogenes tolerates high and low temperatures as well as high salt concentrations. As a result, it replicates well in soil, dust, water, sewage, manure, silage and animal feeds [1,40,77,156,157]. *Listeria* grow at refrigerator temperatures (therefore contaminate

refrigerated foods and raw meat), and can be detected in the feces of animals and humans [144]. Although it can survive for many months in soil [156], pasteurization and most disinfecting agents eliminate the organism [115]. While *Listeria* can be isolated from stool cultures of 5% of healthy individuals [12,52,53,76,85,129,134] and fecal carriage of *Listeria* occurs in 1-15% of the population, vaginal carriage is lower [85]. Pathogenic strains have been detected in the gastrointestinal (GI) tract of asymptomatic individuals, including those at risk (such as pregnant women or immunocompromised patients) [76,82,137].

COLONIZATION WITH LISTERIA

Exposure to and transient colonization of the GI tract by *L. monocytogenes* appears to be common, but invasive disease is rare. Using repeated sampling, *Listeria* can be detected in the feces of nearly 70% of healthy non-pregnant individuals and 44% of pregnant women [75]. Single samples of feces during pregnancy (with careful attention to culture techniques) have a 12% culture positive rate [75], yet others have reported lower recovery rates of 1.1-3.9% [48,82]. Pregnancy does not appear to affect the fecal, cervico-vaginal or oropharyngeal carriage rate of *Listeria*, but such carriage has been reported as a possible predisposing factor for perinatal listeriosis [82]. A study of 750 healthy pregnant women reported that 3.9% had a positive rectal culture while only one woman (0.13%) had a positive vaginal culture [85]. Four studies comprising more than 1,000 healthy pregnant women overall found positive cervicovaginal cultures of *L. monocytogenes* to be very rare [65,82,85,116].

PATHOGENESIS OF INFECTION

L. monocytogenes is an intracellular pathogen which, following ingestion of contaminated food, is phagocytosed and internalised in epithelial cells by the interaction between the bacterial surface protein, internalin, and its receptor on the epithelial surface, E-cadherin. The organism is vacuolated by macrophages, polymorphonuclear leucocytes and other plasma cells, and escapes the vacuole through the action of listeriolysin, finally entering the cytoplasm where proliferation occurs [88]. In the cytoplasm, Act A, another virulence factor, induces actin polymerization to form filaments which become a propulsion system for the organism [114]. The organism then migrates to the cell membrane and produces pseudopodia-like structures which extrude from the host cell, are recognized by adjacent cells and phagocytosed, so that the intracellular cycle continues [144]. This cell-to-cell spread accounts for the fact that antibodies, complement and neutrophils have little or no effect in host protection [63,88,114,144]. Since *L. monocytogenes* can enter the host without disrupting the integrity of the GI tract [144], it has been suggested that GI symptoms may reflect a non-specific response to a systematic infection. Alternatively, GI symptoms may represent a concurrent illness through other organisms (such as *Shigella*) in the GI tract, which disrupt mucosal integrity and facilitate invasion of *L. monocytogenes* [12,88,132,135]. Otherwise, GI symptoms may simply reflect the spectrum of illness of listeriosis itself.

Listeria activates T-cell mediated immunity which, under the influence of cytokines, attracts macrophages that produce inflammatory granulomata where bacteria are destroyed [91]. Memory T-cells provide an acquired resistance to *Listeria* infection, and this might explain why listeriosis is linked with malignancy, immunosuppressive therapy, AIDS, pregnancy and the neonate. This can also account for the observation that neutropenia and disorders of complement or immunoglobulin synthesis are not associated with excessive prevalence of the disease [8,13,32,69,91]. Apart from the immune status of the host, other factors which influence whether or not invasive disease occurs include the virulence of the infecting strain and the size of the inoculum [68,118,131]. The infective dose is unknown, but is estimated

to be between 10^4 - 10^6 organisms/g of ingested product, although this estimate might be lower in groups at risk [61,101,128]. The duration of the incubation period is also unknown, but is likely to be about three weeks [112,134].

LISTERIOSIS IN PREGNANCY

Epidemiology

Listeriosis is 18 times more common in pregnancy (12/100,000) than in the non-pregnant population (0.7/100,000) [144], and 16-27% of all infections with *Listeria* occur in pregnant women [70,72,154]. Sporadic cases of perinatal listeriosis, as well as epidemic cases, have been reported with a prevalence that varies between 8.6 and 17.4/100,000 of live births [144,148]. The largest reported epidemic occurred in 1985 in Los Angeles, California, USA. Of 142 reported cases of listeriosis, 65.5% occurred in pregnant women [87]. The source was contaminated Mexican-style cheese made with unpasteurized milk, which is a well-known risk factor for the acquisition of listeriosis. Indeed, Hispanic (compared to non-Hispanic) women are at a significantly greater risk of pregnancy-associated listeriosis due to their consumption of food known to be at higher risk of contamination with *L. monocytogenes* [154].

Underlying maternal risk factors are uncommon (4.5%), and serious maternal disease is rare when other risk factors are absent [22,45]. However, one case of maternal meningo-encephalitis in a series of 248 cases of *Listeria* in pregnancy has been reported [95], as well as a case of maternal listerial endocarditis [64]. In addition, an analysis of mother-infant pairs with *Listeria* showed that one-third of mothers had *Listeria* isolated from blood cultures, and nearly half were hospitalized [70]. Listeriosis occurs mainly in the third trimester, perhaps due to deficient cell mediated immunity [14]; yet, cases have been observed at earlier gestational ages [48,111]. The incidence at lower gestational ages may be underestimated due to reluctance to culture aborted fetal tissue or products of conception.

Clinical Features

The presentation of listeriosis during pregnancy includes mild flu-like symptoms [10]. In a series of 191 cases of listeriosis in pregnancy, 32% of women had symptoms of a flu-like illness, 65% had a fever, and other symptoms included backache (21.5%) (which may be mistaken for a urinary tract infection), headache (10.5%), vomiting/diarrhea (7%), muscle pains (4%) and sore throat (4%). Approximately 29% of the women were asymptomatic [105].

Fetal transmission from maternal listeriosis is not invariable [67], and discordant infection *in utero* has been observed in a set of twins [142]. Once the placenta is infected, it becomes a reservoir for re-infection [6], and placental micro-abscesses may be histologically evident [31,152].

Listeriosis during pregnancy carries a poorer prognosis for fetuses who are affected at early gestations [105] as opposed to later gestations, and commonly results in spontaneous abortion or stillbirth [73,121]. The association with recurrent abortion has been reported in a series of cases in which *Listeria* was cultured from the cervix [119], though this has been questioned [4,92,93] mainly because of the inability to substantiate the presence of the organism microbiologically [2,50,86,125,126,150].

Preterm birth is common [9], and the highest mortality rate is predictably among infants born at the earliest gestations [95]. Two studies have been reported [95,105], one with 222 pregnant women comprising 11 culture-confirmed cases and 211 cases reported in the literature [105], and the other with 722 pregnant women who were either culture-positive or

had clinical suspicion of listeriosis [95]. Spontaneous abortion occurred in 10-20% of cases, approximately 50% delivered preterm, intrauterine fetal death occurred in approximately 11% [95], 34% demonstrated fetal distress, and 75% developed meconium staining of the amniotic fluid particularly at early gestation [95,105].

In another study, tissue from preterm births or spontaneous abortions (which occurred around 3-7 days post infection) tested positive for *Listeria* in 1.6% of cases [46]. Of 34 women with a history of recurrent abortions and 87 controls, positive cultures were found in 73.5% and 0% respectively [119]. Other studies [4,84,92,93] failed to confirm these findings; therefore, routine genital tract culture and treatment are not recommended [138]. Nonetheless, although rare and because of the potentially serious consequences, it is important that practicing obstetricians are familiar with the diagnosis, treatment, and prevention of *Listeria* infection [124].

NEONATAL LISTERIOSIS

While maternal illness “due to listeriosis” may be mild, neonatal illness is often severe and may be fatal. Neonatal listeriosis may occur by vertical transmission of *L. monocytogenes* from mother to fetus, either by inhalation of infected amniotic fluid, transplacentally from the maternal circulation [9] or by ascending colonization from the vagina. Though vaginal colonization is rare, nearly half of asymptomatic mothers of infants born with neonatal listeriosis will have a positive vaginal culture for *L. monocytogenes* [85]. This observation is consistent with other perinatal infections like group B *Streptococcus* (GBS). Similar to GBS, *Listeria* is likely to result from spread to the vagina from the lower GI tract [5,15,28]. Furthermore, ascending vaginal colonization is supported by the finding of listerial colonization solely in the firstborn of a set of twins [56]. However, haematogenous spread may be more likely, and documented maternal bacteriuria and *in utero* fetal infection with positive amniotic fluid cultures prior to labor have been recorded [94]. In addition, nosocomial cross-infection has been reported [37,115,133] 55,115,116].

Neonatal listeriosis occurs in approximately 8.6/100,000 of live births [20], and is one of the most common causes of neonatal meningitis. The clinical manifestations of neonatal listeriosis are similar to GBS, and there is a high mortality rate (20-60%) [2,87,95,105,136]. One study reported that 68% of newborns whose mothers were diagnosed with listeriosis as a result of positive cultures from placenta, maternal blood or cervix developed neonatal infection; of those 68.2% made a complete recovery, 12.7% developed long-term neurological sequelae, and the infant mortality rate was 24.5% [105]. Like GBS, there are two forms of the disease in the neonate, early- and late-onset, suggesting different modes of transmission (vertical and nosocomial) [150].

Early Onset

Symptoms of early-onset listeriosis occur at a mean of 36 hours after birth [123]. In 50-74% of cases, the mother is likely to have had a flu-like illness with symptoms of fever, headache and myalgia [56,87,95,123], and isolation of *L. monocytogenes* from maternal blood or genital tract is common (44-89%) [87,95]. Neonates with early-onset infection are more frequently born preterm and associated with chorioamnionitis. The fetus presents with clinical features like septicaemia (81-88%), respiratory distress or pneumonia (38%), meningitis (24%) [87,95] and occasionally disseminated inflammatory granulomata (Granulomatosis infantisepticum) [7,17,65,73,119,120], from which pure culture of *L. monocytogenes* can be obtained [121,122]. Granulomatosis Infantisepticum was reported for the first time in 1893 (at which time, the disease was named pseudotuberculosis) [60], and is a pathognomonic feature of listeriosis [80].

Late Onset

In contrast, late-onset neonatal listeriosis (commonly due to serotype 4b) [3], tends to occur between five days and two or more weeks postpartum, typically in term neonates [2,95]. Also typical of these cases is that the neonate is born to asymptomatic mothers with failure to isolate *L. monocytogenes* from maternal cultures [87,95]. The clinical features may be non-specific, but septicaemia (17-95%) and meningitis (67-93%) are common [115,141]. Neonatal listeriosis is one of the few congenital infections in which antibiotic therapy can improve outcome [23,42,105,159].

DIAGNOSIS

Diagnosis of maternal listerial infection may be difficult due to the lack of GI symptoms normally associated with food-borne pathogens and non-specific symptoms of malaise and influenza. Hematologically, a leucocytosis occurs rather than a monocytosis as the species name suggests. On wet mount microscopy, the organisms may be seen in characteristic “tumbling” motion, or may be evident on Gram stain, particularly if done on neonatal meconium before polymicrobial colonisation of the gut occurs. Gram stain is only useful in a minority of cases because, as an intracellular organism, it is easily missed and microscopically resembles many other bacteria commonly found in the vagina. This being the case, warning the microbiological microscopist of the possibility of listerial infection may improve the diagnosis using Gram stain [72]. Isolation is usually by culture from blood (maternal or neonatal), neonatal cerebrospinal fluid (CSF), amniotic fluid, the uterine cavity or the placenta [70,105]. The organism grows well from blood or CSF without the need for enrichment or selective media, and post-partum, histological evidence of micro-abscesses in the placenta can be found [31].

Using the ability of *Listeria* to grow at extreme temperatures, “cold enrichment” involving culture at 4°C (which discourages the growth of other organisms) may help identification [51]. However, this is time-consuming and not as accurate as selective media to isolate the organism from contaminated sites such as food or stools, which contain multiple species of *Listeria* [58]. Rapid detection, at least to the genus level, is possible using monoclonal antibody tests or DNA hybridization [25,29,45,143,155]. Serological testing for anti-listeriolysin may assist in diagnosing both invasive and non-invasive listeriosis [11,24,36], and identification to the species level is possible using specific primers for *L. monocytogenes*. Recently, a 16S rDNA assay with specific primers for the four major food-borne pathogens (*E. coli*, *Salmonella*, *Campylobacter* and *Listeria*) has been successfully tested [146].

PREVENTION OF LISTERIOSIS

In the USA, *Listeria* became nationally notifiable in 1999, and every occurrence should be reported to local health departments. Currently no vaccine for *Listeria* exists. Most cases of listeriosis are sporadic and not associated with an outbreak [105]. Most sporadic cases, as well as all large outbreaks, are due to manufactured foods [16], particularly ready-to-eat meats (especially hot dogs [97]), [71,98,105,138,144], and dairy products such as soft cheeses [21,87,141,148]. Most dairy-associated outbreaks are due to inadequate pasteurization, unpasteurized dairy products [87], or contamination post pasteurization [21].

The Centers for Disease Control and Prevention (CDC) have issued general recommendations for the prevention of listeriosis (Table 1). CDC recommendations state that all deli meats can be consumed safely if reheated until steaming hot. Prevention or avoidance of cross-contamination is an important preventative measure, and all utensils and surfaces should be cleaned after preparing meat dishes or cutting unprepared foods.

Following the 1985 listeriosis outbreak [87], the CDC began monitoring dairy products and sporadic cases of listeriosis, including the voluntary recall of products and has maintained a zero tolerance policy for contaminated ready to eat products. This has resulted in a 44% reduction in the incidence of perinatal listeriosis in the USA (from 17.4/100,000 to 8.6/100,000 live births) [148].

Patient awareness

Despite the CDC recommendations, in a US national survey of 403 pregnant women, only 18% were familiar with listeriosis and less than 30% knew that it could be prevented by avoiding delicatessen meats, soft cheeses and unpasteurized dairy products. In focus groups, many had heard of *E. coli* and *Salmonella*, but few knew that pregnancy increased the risk for development of listeriosis, none had made any change to their food-handling habits, and all had continued to eat hot dogs and delicatessen meats during pregnancy [108]. Both commercially and in the home, individuals at low risk for listeriosis (who may therefore not be aware of the precautions needed to reduce the risk of listeriosis) often prepare food for pregnant women who are at higher risk. For these individuals, multimedia education has been recommended [19].

TREATMENT OF LISTERIOSIS

Treatment of listeriosis in pregnancy requires collaboration with an infectious diseases specialist, and should also involve pediatricians. Irrespective of the gestational age at which infection occurs in pregnancy, treatment is directed toward improving neonatal outcome. Early recognition and intervention are associated with improved outcome, and some physicians have advocated the use of antibiotics for the suspicion of *Listeria* infection for high risk cases after samples for culture have been obtained [138].

In contrast to other causes of chorioamnionitis in which induction of labor is the standard of care [138], treatment of maternal listeriosis in pregnancy and delay of delivery can result in the birth of a healthy infant at term [23,42,67,74,105,159]. Amniotic fluid cultures have been observed to change from positive to negative with appropriate antibiotic therapy [27]. Ideally, successful reports of *in utero* therapy should be accompanied by documentation of resolution by demonstrating negative amniotic fluid cultures.

Treatment challenges

No randomized controlled trials to assess type of antibiotic or duration of treatment have been performed. Therapeutic considerations include the fact that: 1) host susceptibility is linked to the atypical onset of the disease. The non-specific nature of the normal symptomatology may also make diagnosis and treatment difficult; and 2) the intracellular nature of the organism presents therapeutic challenges. Both of these factors may delay diagnosis and treatment. Bacteriostatic concentrations *in vivo* may be insufficient to be effective intracellularly. As a result, treatment recommendations are based mainly upon *in vitro* susceptibility testing or animal-model experimentation or are experience-based, following trials with small numbers of historical controls rather than being based on a foundation of robust medical evidence [149].

Antibiotics

The antibiotic of choice should penetrate host cells, distribute within the cells and remain stable in the intracellular environment [115,138,144]. As a result of these unusual antibiotic requirements, antibiotics may be ineffective in up to 70% of listeriosis cases [62]. Most of the experience of treating *L. monocytogenes* is with the use of ampicillin, penicillin and amoxicillin [149], and to date, no bacterial resistance to penicillin has been detected [63]. *In*

vitro, *L. monocytogenes* has a wide range of antibiotic sensitivities but is resistant to cephalosporins, clindamycin and chloramphenicol [34,105]. Chloramphenicol should not be used because of unacceptable rates of failure and relapse [21,69,88], and quinolones do not have adequate *in vitro* activity [35]. Cephalosporins are ineffective against *Listeria* because they do not bind to bacterial penicillin-bound protein-3 (PBP3) [144,153].

Though resistance to ampicillin has been reported [90], it is preferred over penicillin [10,105]. *L. monocytogenes* responds poorly to bacteriostatic antibiotics due to the intracellular nature of the pathogen, but ampicillin provides good host cell penetration without any significant change in pH or concentration which might compromise efficiency. Ampicillin also crosses the placenta in adequate amounts and binds to PBP3, causing bacterial death [63]. Gentamicin and tobramycin have been reported to have greater *in vitro* activity than streptomycin, kanamycin and amikacin [3,102,127]. Listerial resistance to vancomycin is rare [63], and its use (though limited) has been employed in endocarditis, as well as in listerial meningitis (in which cerebral intraventricular injections were administered) [104].

Synergy of ampicillin with gentamicin is claimed, but this has only been demonstrated *in vitro* [21,34,102,127,144]. Nevertheless, these two antibiotics are often used in combination in cases of meningitis despite concerns with respect to toxicity [149]. In patients with an allergy to ampicillin and/or gentamicin, erythromycin may be used [149], but transplacental passage of erythromycin provides sub-therapeutic concentrations in both amniotic fluid and fetal serum [59]. Alternatively, the combination of trimethoprim and sulfamethoxazole (which is bactericidal and achieves good serum and CSF concentrations) may be used [18,34,55,72,89,100,106,144,145,158]. However, there are concerns regarding the use of sulphonamides due to sulphur allergy and the risk of kernicterus (possible displacement of bilirubin from albumin) from use immediately prior to delivery. There are also concerns about possible teratogenic effects, such as neural tube defects due to the folate antagonism of trimethoprim during pregnancy, as well as hemolytic anemia in male infants due to glucose-6-phosphate deficiency. The preference for ampicillin over other options is such that it has been proposed to test sensitivity for suspected allergy before recourse to alternative therapy [72].

During pregnancy, the current recommendation for first line therapy is IV ampicillin 6g/day [149] and erythromycin 4g/day as second line IV. The duration for both regimes is 14 days, though the length of course may be longer in different clinical syndromes [45] and if the pregnant woman is immunosuppressed [72,149].

CONCLUSION

L. monocytogenes is an unusual pathogen due to the intracellular nature of its life cycle, which may explain the ready ease of transplacental passage. With an incidence of 12/100,000 pregnancies, listeriosis in pregnancy is still 18 times more common in pregnant women than in the general population. Pregnant women with co-morbidities are at increased risk due to compromised cell-mediated immunity. Maternal illness is usually mild, but neonatal disease carries a 20-30% mortality rate (although fetal and neonatal mortality are potentially preventable with early diagnosis and intervention). Ready-to-eat meats and dairy products made with unpasteurized milk should be avoided, and care should be taken to prevent cross contamination of foods by ensuring that preparation and cooking utensils and food preparation surfaces are clean. High dose ampicillin (6g/day) for a prolonged period of 14 days is recommended for treatment in pregnancy.

Acknowledgments

This research was supported in part by the Perinatology Research Branch, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NIH, DHHS.

Reference List

1. Al-Ghazali MR, Al-Azawi SK. Detection and enumeration of *Listeria monocytogenes* in a sewage treatment plant in Iraq. *J Appl Bacteriol.* 1986; 60:251–4. [PubMed: 3086271]
2. Albritton WL, Cochi SL, Feeley JC. Overview of neonatal listeriosis. *Clin Invest Med.* 1984; 7:311–4. [PubMed: 6398177]
3. Albritton WL, Wiggins GL, Feeley JC. Neonatal listeriosis: distribution of serotypes in relation to age at onset of disease. *J Pediatr.* 1976; 88:481–3. [PubMed: 812974]
4. Ansbacher R, Borchardt KA, Hannegan MW, Boyson WA. Clinical investigation of *Listeria monocytogenes* as a possible cause of human fetal wastage. *Am J Obstet Gynecol.* 1966; 94:386–90. [PubMed: 4955545]
5. Badri MS, Zawaneh S, Cruz AC, Mantilla G, Baer H, Spellacy WN, et al. Rectal colonization with group B streptococcus: relation to vaginal colonization of pregnant women. *J Infect Dis.* 1977; 135:308–12. [PubMed: 320278]
6. Bakardjiev AI, Theriot JA, Portnoy DA. *Listeria monocytogenes* traffics from maternal organs to the placenta and back. *PLoS Pathog.* 2006; 2:e66. [PubMed: 16846254]
7. Barber M, Okubadejo OA. Maternal and neonatal listeriosis: report of case and brief review of literature of listeriosis in man. *Br Med J.* 1965; 2:735–8. [PubMed: 5317930]
8. Beckerman KP, Rogers HW, Corbett JA, Schreiber RD, McDaniel ML, Unanue ER. Release of nitric oxide during the T cell-independent pathway of macrophage activation. Its role in resistance to *Listeria monocytogenes*. *J Immunol.* 1993; 150:888–95. [PubMed: 7678626]
9. Becroft DM, Farmer K, Seddon RJ, Sowden R, Stewart JH, Vines A, et al. Epidemic listeriosis in the newborn. *Br Med J.* 1971; 3:747–51. [PubMed: 5097970]
10. Benshushan A, Tsafir A, Arbel R, Rahav G, Ariel I, Rojansky N. *Listeria* infection during pregnancy: a 10 year experience. *Isr Med Assoc J.* 2002; 4:776–80. [PubMed: 12389339]
11. Berche P. Bacteremia is required for invasion of the murine central nervous system by *Listeria monocytogenes*. *Microb Pathog.* 1995; 18:323–36. [PubMed: 7476097]
12. Bojsen-Moller J. Human listeriosis. Diagnostic, epidemiological and clinical studies. *Acta Pathol Microbiol Scand B Microbiol Immunol.* 1972; 229:1–157. [PubMed: 4624477]
13. Bortolussi R, Issekutz T, Burbridge S, Schellekens H. Neonatal host defense mechanisms against *Listeria monocytogenes* infection: the role of lipopolysaccharides and interferons. *Pediatr Res.* 1989; 25:311–5. [PubMed: 2468128]
14. Bortolussi R, McGregor DD, Kongshavn PA, Galsworthy S, Albritton W, Davies JW, et al. Host defense mechanisms to perinatal and neonatal *Listeria monocytogenes* infection. *Surv Synth Pathol Res.* 1984; 3:311–32. [PubMed: 6390622]
15. Boyer KM, Gadzala CA, Burd LI, Fisher DE, Paton JB, Gotoff SP. Selective intrapartum chemoprophylaxis of neonatal group B streptococcal early-onset disease. I. Epidemiologic rationale. *J Infect Dis.* 1983; 148:795–801. [PubMed: 6355316]
16. Braden CR. Listeriosis. *Pediatr Infect Dis J.* 2003; 22:745–6. [PubMed: 12913780]
17. Buchner LH, Schneierson S. Clinical and laboratory aspects of *Listeria monocytogenes* infections. With a report of ten cases. *Am J Med.* 1968; 45:904–21. [PubMed: 4972645]
18. Bula CJ, Bille J, Glauser MP. An epidemic of food-borne listeriosis in western Switzerland: description of 57 cases involving adults. *Clin Infect Dis.* 1995; 20:66–72. [PubMed: 7727673]
19. Cates SC, Carter-Young HL, Conley S, O'Brien B. Pregnant women and listeriosis: preferred educational messages and delivery mechanisms. *J Nutr Educ Behav.* 2004; 36:121–7. [PubMed: 15202987]
20. Centers for Disease Control and Prevention. Preliminary FoodNet data on the incidence of infection with pathogens transmitted commonly through food--10 states, 2007. *MMWR Morb Mortal Wkly Rep.* 2008; 57:366–70. [PubMed: 18401330]

21. Cherubin CE, Appleman MD, Heseltine PN, Khayr W, Stratton CW. Epidemiological spectrum and current treatment of listeriosis. *Rev Infect Dis.* 1991; 13:1108–14. [PubMed: 1775844]
22. Ciesielski CA, Hightower AW, Parsons SK, Broome CV. Listeriosis in the United States: 1980-1982. *Arch Intern Med.* 1988; 148:1416–9. [PubMed: 3377624]
23. Cruikshank DP, Warenski JC. First-trimester maternal *Listeria monocytogenes* sepsis and chorioamnionitis with normal neonatal outcome. *Obstet Gynecol.* 1989; 73:469–71. [PubMed: 2783771]
24. Dalton CB, Austin CC, Sobel J, Hayes PS, Bibb WF, Graves LM, et al. An outbreak of gastroenteritis and fever due to *Listeria monocytogenes* in milk. *N Engl J Med.* 1997; 336:100–5. [PubMed: 8988887]
25. Datta AR, Wentz BA, Hill WE. Detection of hemolytic *Listeria monocytogenes* by using DNA colony hybridization. *Appl Environ Microbiol.* 1987; 53:2256–9. [PubMed: 2823710]
26. Dawson KG, Emerson JC, Burns JL. Fifteen years of experience with bacterial meningitis. *Pediatr Infect Dis J.* 1999; 18:816–22. [PubMed: 10493344]
27. Delgado AR. Listeriosis in pregnancy. *J Midwifery Womens Health.* 2008; 53:255–9. [PubMed: 18455101]
28. Dillon HC Jr, Gray E, Pass MA, Gray BM. Anorectal and vaginal carriage of group B streptococci during pregnancy. *J Infect Dis.* 1982; 145:794–9. [PubMed: 7045248]
29. Donnelly CW, Baigent GJ. Method for flow cytometric detection of *Listeria monocytogenes* in milk. *Appl Environ Microbiol.* 1986; 52:689–95. [PubMed: 3096202]
30. Doyle MP, Schoeni JL. Selective-enrichment procedure for isolation of *Listeria monocytogenes* from fecal and biologic specimens. *Appl Environ Microbiol.* 1986; 51:1127–9. [PubMed: 3089147]
31. DRISCOLL SG, GORBACH A, FELDMAN D. Congenital listeriosis: diagnosis from placental studies. *Oncologia.* 1962; 20:216–20. [PubMed: 13887884]
32. Dunn PL, North RJ. Resolution of primary murine listeriosis and acquired resistance to lethal secondary infection can be mediated predominantly by Thy-1+ CD4- CD8- cells. *J Infect Dis.* 1991; 164:869–77. [PubMed: 1682394]
33. Murray EGD, Webb RA, Swann MBR. A disease of rabbits characterised by a large mononuclear leucocytosis, caused by a hitherto undescribed bacillus *Bacterium monocytogenes* (n.sp.). *The Journal of Pathology and Bacteriology.* 1926; 29:407–39.
34. Espaze EP, Reynaud AE. Antibiotic susceptibilities of *Listeria*: in vitro studies. *Infection.* 1988; 16(Suppl 2):S160–S164. [PubMed: 3047066]
35. Espaze EP, Roubex YG, Le Berre JY, Courtieu AI. In vitro susceptibility of *Listeria monocytogenes* to some antibiotics and their combinations. *Zentralbl Bakteriol Orig A.* 1978; 240:76–85. [PubMed: 416625]
36. Facinelli B, Giovanetti E, Varaldo PE, Casolari P, Fabio U. Antibiotic resistance in foodborne *Listeria*. *Lancet.* 1991; 338:1272. [PubMed: 1682666]
37. Facinelli B, Varaldo PE, Casolari C, Fabio U. Cross-infection with *Listeria monocytogenes* confirmed by DNA fingerprinting. *Lancet.* 1988; 2:1247–8. [PubMed: 2903973]
38. Farber JM, Losos JZ. *Listeria monocytogenes*: a foodborne pathogen. *CMAJ.* 1988; 138:413–8. [PubMed: 3124948]
39. Farber JM, Peterkin PI. *Listeria monocytogenes*, a food-borne pathogen. *Microbiol Rev.* 1991; 55:476–511. [PubMed: 1943998]
40. Fenlon DR. Rapid quantitative assessment of the distribution of *Listeria* in silage implicated in a suspected outbreak of listeriosis in calves. *Vet Rec.* 1986; 118:240–2. [PubMed: 3085322]
41. Filice GA, Cantrell HF, Smith AB, Hayes PS, Feeley JC, Fraser DW. *Listeria monocytogenes* infection in neonates: Investigation of an epidemic. *J Infect Dis.* 1978; 138:17–23. [PubMed: 98594]
42. Fleming AD, Ehrlich DW, Miller NA, Monif GR. Successful treatment of maternal septicemia due to *Listeria monocytogenes* at 26 weeks' gestation. *Obstet Gynecol.* 1985; 66:52S–3S. [PubMed: 4022517]

43. Fleming DW, Cochi SL, MacDonald KL, Brondum J, Hayes PS, Plikaytis BD, et al. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. *N Engl J Med*. 1985; 312:404–7. [PubMed: 3918263]
44. Gaillard JL, Berche P, Sansonetti P. Transposon mutagenesis as a tool to study the role of hemolysin in the virulence of *Listeria monocytogenes*. *Infect Immun*. 1986; 52:50–5. [PubMed: 3007363]
45. Gellin BG, Broome CV. Listeriosis. *JAMA*. 1989; 261:1313–20. [PubMed: 2492614]
46. Giraud JR, Denis F, Gargot F, Fizazi T, Babin P, Rautlin de la Roy YD, et al. [Listeriosis. Occurrence in spontaneous interruptions of pregnancy]. *Nouv Presse Med*. 1973; 2:215–8. [PubMed: 4631152]
47. Gray DI, Kroll RG. Polymerase chain reaction amplification of the *flaA* gene for the rapid identification of *Listeria* spp. *Lett Appl Microbiol*. 1995; 20:65–8. [PubMed: 7765872]
48. Gray JW, Barrett JF, Pedler SJ, Lind T. Faecal carriage of listeria during pregnancy. *Br J Obstet Gynaecol*. 1993; 100:873–4. [PubMed: 8218017]
49. GRAY ML, Killinger AH. *Listeria monocytogenes* and listeric infections 1. *Bacteriol Rev*. 1966; 30:309–82. [PubMed: 4956900]
50. GRAY ML, Seeliger HP, POTEL J. PERINATAL INFECTIONS DUE TO *LISTERIA MONOCYTOGENES*. DO THESE AFFECT SUBSEQUENT PREGNANCIES? *Clin Pediatr (Phila)*. 1963; 2:614–23. [PubMed: 14079241]
51. GRAY ML, STAFSETH HJ, Thorp F, Sholl LB, Riley WF. A New Technique for Isolating *Listerellae* from the Bovine Brain. *J Bacteriol*. 1948; 55:471–6.
52. Grif K, Hein I, Wagner M, Brandl E, Mpamugo O, McLauchlin J, et al. Prevalence and characterization of *Listeria monocytogenes* in the feces of healthy Austrians. *Wien Klin Wochenschr*. 2001; 113:737–42. [PubMed: 11715752]
53. Grif K, Patscheider G, Dierich MP, Allerberger F. Incidence of fecal carriage of *Listeria monocytogenes* in three healthy volunteers: a one-year prospective stool survey. *Eur J Clin Microbiol Infect Dis*. 2003; 22:16–20. [PubMed: 12582739]
54. Groves RD, Welshimer HJ. Separation of pathogenic from apathogenic *Listeria monocytogenes* by three in vitro reactions. *J Clin Microbiol*. 1977; 5:559–63. [PubMed: 407244]
55. Gunther G, Philipson A. Oral trimethoprim as follow-up treatment of meningitis caused by *Listeria monocytogenes*. *Rev Infect Dis*. 1988; 10:53–5. [PubMed: 3258438]
56. Halliday HL, Hirata T. Perinatal listeriosis-a review of twelve patients. *Am J Obstet Gynecol*. 1979; 133:405–10. [PubMed: 434005]
57. Harvey Pirie JH. *Listeria*: Change of Name for a Genus Bacteria: Abstract: *Nature*. 1940; 145:264.
58. Hayes PS, Graves LM, Ajello GW, Swaminathan B, Weaver RE, Wenger JD, et al. Comparison of cold enrichment and U.S. Department of Agriculture methods for isolating *Listeria monocytogenes* from naturally contaminated foods. The *Listeria* Study Group. *Appl Environ Microbiol*. 1991; 57:2109–13. [PubMed: 1768082]
59. Heikkinen T, Laine K, Neuvonen PJ, Ekblad U. The transplacental transfer of the macrolide antibiotics erythromycin, roxithromycin and azithromycin. *BJOG*. 2000; 107:770–5. [PubMed: 10847234]
60. Henle A. Pseudotuberkulose bei neugeborenen zwillingen. *Arb Pathol Inst Gottingen*. 1893
61. Ho JL, Shands KN, Friedland G, Eckind P, Fraser DW. An outbreak of type 4b *Listeria monocytogenes* infection involving patients from eight Boston hospitals. *Arch Intern Med*. 1986; 146:520–4. [PubMed: 3954524]
62. Hof H. An update on the medical management of listeriosis. *Expert Opin Pharmacother*. 2004; 5:1727–35. [PubMed: 15264987]
63. Hof H, Nichterlein T, Kretschmar M. Management of listeriosis. *Clin Microbiol Rev*. 1997; 10:345–57. [PubMed: 9105758]
64. Holshouser CA, Ansbacher R, McNitt T, Steele R. Bacterial endocarditis due to *Listeria monocytogenes* in a pregnant diabetic. *Obstet Gynecol*. 1978; 51:9s–10s. [PubMed: 618485]

65. HOOD M. Listeriosis as an infection of pregnancy manifested in the newborn. *Pediatrics*. 1961; 27:390–6. [PubMed: 13715686]
66. Hulpfers G. Lefvernekos hos kanin orsakad af en ej forut beskrifven bakterie. *Svenska Vet Tidskr*. 1911; 2:265–8.
67. Hume OS. Maternal *Listeria monocytogenes* septicemia with sparing of the fetus. *Obstet Gynecol*. 1976; 48:33S–4S. [PubMed: 821025]
68. Ireton K. Entry of the bacterial pathogen *Listeria monocytogenes* into mammalian cells. *Cell Microbiol*. 2007; 9:1365–75. [PubMed: 17419717]
69. Issekutz TB, Evans J, Bortolussi R. The immune response of human neonates to *Listeria monocytogenes* infection. *Clin Invest Med*. 1984; 7:281–6. [PubMed: 6532634]
70. Jackson KA, Iwamoto M, Swerdlow D. Pregnancy-associated listeriosis. *Epidemiol Infect*. 2010; 138:1503–9. [PubMed: 20158931]
71. Jacquet C, Catimel B, Brosch R, Buchrieser C, Dehaumont P, Goulet V, et al. Investigations related to the epidemic strain involved in the French listeriosis outbreak in 1992. *Appl Environ Microbiol*. 1995; 61:2242–6. [PubMed: 7793944]
72. Janakiraman V. Listeriosis in pregnancy: diagnosis, treatment, and prevention. *Rev Obstet Gynecol*. 2008; 1:179–85. [PubMed: 19173022]
73. JOHNSTON WH, MORTON SA, WONG MH, ROY TE. Septicaemia of the newborn due to *Listeria monocytogenes*. *Can Med Assoc J*. 1955; 73:402–5. [PubMed: 13250472]
74. Kalstone C. Successful antepartum treatment of listeriosis. *Am J Obstet Gynecol*. 1991; 164:57–8. [PubMed: 1986626]
75. Kampelmacher EH, Huysinga WT, van Noorle Jansen LM. The presence of *Listeria monocytogenes* in feces of pregnant women and neonates. *Zentralbl Bakteriol Orig A*. 1972; 222:258–62. [PubMed: 4144801]
76. Kampelmacher EH, van Noorle Jansen LM. Isolation of *Listeria monocytogenes* from faeces of clinically healthy humans and animals. *Zentralbl Bakteriol Orig*. 1969; 211:353–9. [PubMed: 4989347]
77. Kampelmacher EH, van Noorle Jansen LM. Listeriosis in humans and animals in the Netherlands (1958–1977). *Zentralbl Bakteriol A*. 1980; 246:211–27. [PubMed: 6775441]
78. Kathariou S, Metz P, Hof H, Goebel W. Tn916-induced mutations in the hemolysin determinant affecting virulence of *Listeria monocytogenes*. *J Bacteriol*. 1987; 169:1291–7. [PubMed: 3029033]
79. Kathariou S, Rocourt J, Hof H, Goebel W. Levels of *Listeria monocytogenes* hemolysin are not directly proportional to virulence in experimental infections of mice. *Infect Immun*. 1988; 56:534–6. [PubMed: 3123396]
80. Kelly CS, Gibson JL. Listeriosis as a cause of fetal wastage. *Obstet Gynecol*. 1972; 40:91–7. [PubMed: 4625596]
81. Kessler SL, Dajani AS. Listeria meningitis in infants and children. *Pediatr Infect Dis J*. 1990; 9:61–3. [PubMed: 2105480]
82. Lamont RJ, Postlethwaite R. Carriage of *Listeria monocytogenes* and related species in pregnant and non-pregnant women in Aberdeen, Scotland. *J Infect*. 1986; 13:187–93. [PubMed: 3093591]
83. Larsson S, Cederberg A, Ivarsson S, Svanberg L, Cronberg S. *Listeria monocytogenes* causing hospital-acquired enterocolitis and meningitis in newborn infants. *Br Med J*. 1978; 2:473–4. [PubMed: 678926]
84. LAWLER FC, WOOD WS, KING S, METZGER WI. LISTERIA MONOCYTOGENES AS A CAUSE OF FETAL LOSS. *Am J Obstet Gynecol*. 1964; 89:915–23. [PubMed: 14207559]
85. Lennon D, Lewis B, Mantell C, Becroft D, Dove B, Farmer K, et al. Epidemic perinatal listeriosis. *Pediatr Infect Dis*. 1984; 3:30–4. [PubMed: 6701102]
86. Levy E, NASSAU E. Experience with listeriosis in the new-born. An account of a small epidemic in a nursery ward. *Ann Paediatr*. 1960; 194:321–30. [PubMed: 14416386]
87. Linnan MJ, Mascola L, Lou XD, Goulet V, May S, Salminen C, et al. Epidemic listeriosis associated with Mexican-style cheese. *N Engl J Med*. 1988; 319:823–8. [PubMed: 3137471]
88. Lorber B. Listeriosis. *Clin Infect Dis*. 1997; 24:1–9. [PubMed: 8994747]

89. Louria DB, Blevins A, Armstrong D. Listeria infections. *Ann N Y Acad Sci.* 1970; 174:545–51. [PubMed: 4993531]
90. Lyn PC, Lim TH. Listeria meningitis resistant to ampicillin 1. *J Singapore Paediatr Soc.* 1986; 28:247–9. [PubMed: 3110504]
91. Mackaness GB. The immunology of antituberculous immunity. *Am Rev Respir Dis.* 1968; 97:337–44. [PubMed: 4966267]
92. MACNAUGHTON MC. Listeria monocytogenes in abortion. *Lancet.* 1962; 2:484. [PubMed: 14468045]
93. Manganiello PD, Yearke RR. A 10-year prospective study of women with a history of recurrent fetal losses fails to identify Listeria monocytogenes in the genital tract. *Fertil Steril.* 1991; 56:781–2. [PubMed: 1915959]
94. Mazor M, Froimovich M, Lazer S, Maymon E, Glezerman M. Listeria monocytogenes. The role of transabdominal amniocentesis in febrile patients with preterm labor. *Arch Gynecol Obstet.* 1992; 252:109–12. [PubMed: 1471911]
95. McLauchlin J. Human listeriosis in Britain, 1967–85, a summary of 722 cases. 1. Listeriosis during pregnancy and in the newborn. *Epidemiol Infect.* 1990; 104:181–9. [PubMed: 2108869]
96. McLauchlin J. Human listeriosis in Britain, 1967–85, a summary of 722 cases. 2. Listeriosis in non-pregnant individuals, a changing pattern of infection and seasonal incidence. *Epidemiol Infect.* 1990; 104:191–201. [PubMed: 2108870]
97. McLauchlin J. Listeriosis. *BMJ.* 1992; 304:1583–4. [PubMed: 1628083]
98. McLauchlin J, Hall SM, Velani SK, Gilbert RJ. Human listeriosis and pate: a possible association. *BMJ.* 1991; 303:773–5. [PubMed: 1932944]
99. Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, et al. Food-related illness and death in the United States. *Emerg Infect Dis.* 1999; 5:607–25. [PubMed: 10511517]
100. Meyer RD, Liu S. Determination of the effect of antimicrobics in combination against Listeria monocytogenes. *Diagn Microbiol Infect Dis.* 1987; 6:199–206. [PubMed: 3105948]
101. Miller JK. Listeriosis in man. *Med Times.* 1967; 95:754–8. [PubMed: 6068726]
102. Moellering RC Jr. Medoff G, Leech I, Wennersten C, Kunz LJ. Antibiotic synergism against Listeria monocytogenes. *Antimicrob Agents Chemother.* 1972; 1:30–4. [PubMed: 4207757]
103. Murray EG, Ghosh BK, Murray RG. The interaction of guanofuracin and Listeria monocytogenes. *Can J Microbiol.* 1966; 12:285–97. [PubMed: 4959244]
104. Mylonakis E, Hohmann EL, Calderwood SB. Central nervous system infection with Listeria monocytogenes. 33 years' experience at a general hospital and review of 776 episodes from the literature. *Medicine (Baltimore).* 1998; 77:313–36. [PubMed: 9772921]
105. Mylonakis E, Paliou M, Hohmann EL, Calderwood SB, Wing EJ. Listeriosis during pregnancy: a case series and review of 222 cases. *Medicine (Baltimore).* 2002; 81:260–9. [PubMed: 12169881]
106. Nieman RE, Lorber B. Listeriosis in adults: a changing pattern. Report of eight cases and review of the literature, 1968–1978. *Rev Infect Dis.* 1980; 2:207–27. [PubMed: 6771866]
107. Nyfeldt A. Etiologie de la mononucleose infectieuse. *Compt rend Soc Biol.* 1929; 101:590–6.
108. Ogunmodede F, Jones JL, Scheftel J, Kirkland E, Schulkin J, Lynfield R. Listeriosis prevention knowledge among pregnant women in the USA. *Infect Dis Obstet Gynecol.* 2005; 13:11–5. [PubMed: 16040322]
109. Parrisius J, Bhakdi S, Roth M, Tranum-Jensen J, Goebel W, Seeliger HP. Production of listeriolysin by beta-hemolytic strains of Listeria monocytogenes. *Infect Immun.* 1986; 51:314–9. [PubMed: 3079734]
110. Paterson Y, Ikonomidis G. Recombinant Listeria monocytogenes cancer vaccines. *Curr Opin Immunol.* 1996; 8:664–9. [PubMed: 8902392]
111. Pezeshkian R, Fernando N, Carne CA, Simanowitz MD. Listeriosis in mother and fetus during the first trimester of pregnancy. Case report. *Br J Obstet Gynaecol.* 1984; 91:85–6. [PubMed: 6691949]

112. Pinner RW, Schuchat A, Swaminathan B, Hayes PS, Deaver KA, Weaver RE, et al. Role of foods in sporadic listeriosis. II. Microbiologic and epidemiologic investigation. The Listeria Study Group. *JAMA*. 1992; 267:2046–50. [PubMed: 1552640]
113. Pirie JHH. A new disease of wild rodents “Tiger River disease”. Publications of the South African Institute for Medical Research. 1927; 3:163–86.
114. Portnoy DA, Chakraborty T, Goebel W, Cossart P. Molecular determinants of *Listeria monocytogenes* pathogenesis. *Infect Immun*. 1992; 60:1263–7. [PubMed: 1312514]
115. Posfay-Barbe KM, Wald ER. Listeriosis. *Pediatr Rev*. 2004; 25:151–9. [PubMed: 15121906]
116. Quarles JM Jr, Pittman B. Unsuccessful attempt to detect *Listeria monocytogenes* in healthy pregnant women. *J Bacteriol*. 1966; 91:2112–3. [PubMed: 4957031]
117. Ralovich BS, Shahamat M, Woodbine M. Further data on the characters of *Listeria* strains. *Med Microbiol Immunol*. 1977; 163:125–39. [PubMed: 895679]
118. Ramaswamy V, Cresence VM, Rejitha JS, Lekshmi MU, Dharsana KS, Prasad SP, et al. Listeria--review of epidemiology and pathogenesis. *J Microbiol Immunol Infect*. 2007; 40:4–13. [PubMed: 17332901]
119. RAPPAPORT F, RABINOVITZ M, TOAFF R, KROCHIK N. Genital listeriosis as a cause of repeated abortion. *Lancet*. 1960; 1:1273–5. [PubMed: 14436286]
120. Ray CG, WEDGWOOD RJ. NEONATAL LISTERIOSIS. SIX CASE REPORTS AND A REVIEW OF THE LITERATURE. *Pediatrics*. 1964; 34:378–92. [PubMed: 14211104]
121. REISS HJ, KREBS A. Septic granulomatosis in infants, a fetal sepsis caused by a specific pathogen. *Klin Wochenschr*. 1951; 29:29. [PubMed: 14825743]
122. REISS HJ, POTEL J, KREBS A. [Granulomatosis infantiseptica; a general infection of infants and the newborn characterized by the presence of miliary granuloma.]. *Z Gesamte Inn Med*. 1951; 6:451–7. [PubMed: 14901913]
123. Relier JP. Perinatal and neonatal infections: listeriosis. *J Antimicrob Chemother*. 1979; 5(Suppl A):51–7. [PubMed: 112093]
124. Ross DS, Rasmussen SA, Cannon MJ, Anderson B, Kilker K, Tumpey A, et al. Obstetrician/gynecologists’ knowledge, attitudes, and practices regarding prevention of infections in pregnancy. *J Womens Health (Larchmt)*. 2009; 18:1187–93. [PubMed: 19670963]
125. RUFFOLO EH, WILSON RB, WEED LA. *Listeria monocytogenes* as a cause of pregnancy wastage. *Obstet Gynecol*. 1962; 19:533–6. [PubMed: 14495173]
126. Saxbe WB Jr. *Listeria monocytogenes* and Queen Anne. *Pediatrics*. 1972; 49:97–101. [PubMed: 4621771]
127. Scheld WM. Evaluation of rifampin and other antibiotics against *Listeria monocytogenes* in vitro and in vivo. *Rev Infect Dis*. 1983; 5(Suppl 3):S593–S599. [PubMed: 6415782]
128. Schlech WF III. *Listeria gastroenteritis*--old syndrome, new pathogen. *N Engl J Med*. 1997; 336:130–2. [PubMed: 8988894]
129. Schlech WF III. Foodborne listeriosis. *Clin Infect Dis*. 2000; 31:770–5. [PubMed: 11017828]
130. Schlech WF III, Lavigne PM, Bortolussi RA, Allen AC, Haldane EV, Wort AJ, et al. Epidemic listeriosis--evidence for transmission by food. *N Engl J Med*. 1983; 308:203–6. [PubMed: 6401354]
131. Scholing M, Schneeberger PM, van den DP, Drenth JP. Clinical features of liver involvement in adult patients with listeriosis. Review of the literature. *Infection*. 2007; 35:212–8. [PubMed: 17646920]
132. Schroter GP, Weil R III. *Listeria monocytogenes* infection after renal transplantation. *Arch Intern Med*. 1977; 137:1395–9. [PubMed: 411435]
133. Schuchat A, Lizano C, Broome CV, Swaminathan B, Kim C, Winn K. Outbreak of neonatal listeriosis associated with mineral oil. *Pediatr Infect Dis J*. 1991; 10:183–9. [PubMed: 2041663]
134. Schuchat A, Swaminathan B, Broome CV. Epidemiology of human listeriosis. *Clin Microbiol Rev*. 1991; 4:169–83. [PubMed: 1906370]
135. Schwartz B, Hexter D, Broome CV, Hightower AW, Hirschhorn RB, Porter JD, et al. Investigation of an outbreak of listeriosis: new hypotheses for the etiology of epidemic *Listeria monocytogenes* infections. *J Infect Dis*. 1989; 159:680–5. [PubMed: 2494267]

136. Schwarze R, Bauermeister CD, Ortel S, Wichmann G. Perinatal listeriosis in Dresden 1981-1986: clinical and microbiological findings in 18 cases. *Infection*. 1989; 17:131-8. [PubMed: 2737754]
137. Senecal J, Fabiani G, Cormier M, Le MB. [Detection of *Listeria* infection in premature babies]. *Nouv Presse Med*. 1973; 2:1074. [PubMed: 4122420]
138. Silver HM. Listeriosis during pregnancy. *Obstet Gynecol Surv*. 1998; 53:737-40. [PubMed: 9870235]
139. Simmons MD, Cockcroft PM, Okubadejo OA. Neonatal listeriosis due to cross-infection in an obstetric theatre. *J Infect*. 1986; 13:235-9. [PubMed: 3098856]
140. Skalka B, Smola J, Elischerova K. Routine test for in vitro differentiation of pathogenic and apathogenic *Listeria monocytogenes* strains. *J Clin Microbiol*. 1982; 15:503-7. [PubMed: 6804488]
141. Skidmore AG. Listeriosis at Vancouver General Hospital, 1965-79. *Can Med Assoc J*. 1981; 125:1217-21. [PubMed: 6800624]
142. Smith AR, Lieberman BA, Allen L, Barson AJ. Listeriosis and pregnancy. *Lancet*. 1983; 2:1364. [PubMed: 6139696]
143. Sontakke S, Farber JM. The use of PCR ribotyping for typing strains of *Listeria* spp. *Eur J Epidemiol*. 1995; 11:665-73. [PubMed: 8861851]
144. Southwick FS, Purich DL. Intracellular pathogenesis of listeriosis. *N Engl J Med*. 1996; 334:770-6. [PubMed: 8592552]
145. Spitzer PG, Hammer SM, Karchmer AW. Treatment of *Listeria monocytogenes* infection with trimethoprim-sulfamethoxazole: case report and review of the literature. *Rev Infect Dis*. 1986; 8:427-30. [PubMed: 3523699]
146. Suo B, He Y, Tu SI, Shi X. A Multiplex Real-Time Polymerase Chain Reaction for Simultaneous Detection of *Salmonella* spp., *Escherichia coli* O157, and *Listeria monocytogenes* in Meat Products. *Foodborne Pathog Dis*. 2010
147. Swaminathan B, Hayes PS, Przybyszewski VA, Plikaytis BD. Evaluation of enrichment and plating media for isolating *Listeria monocytogenes*. *J Assoc Off Anal Chem*. 1988; 71:664-8. [PubMed: 3134340]
148. Tappero JW, Schuchat A, Deaver KA, Mascola L, Wenger JD. Reduction in the incidence of human listeriosis in the United States. Effectiveness of prevention efforts? The Listeriosis Study Group. *JAMA*. 1995; 273:1118-22. [PubMed: 7707600]
149. Temple ME, Nahata MC. Treatment of listeriosis. *Ann Pharmacother*. 2000; 34:656-61. [PubMed: 10852095]
150. Tessier F, Bouillie J, Daguet GL, Barrat J. [Listeriosis in obstetrical environment. Report on 10-year experience in a maternity hospital in Paris]. *J Gynecol Obstet Biol Reprod (Paris)*. 1986; 15:305-13. [PubMed: 3734345]
151. Tim MW, Jackson MA, Shannon K, Cohen B, McCracken GH Jr. Non-neonatal infection due to *Listeria monocytogenes*. *Pediatr Infect Dis*. 1984; 3:213-7. [PubMed: 6739324]
152. Topalovski M, Yang SS, Boonpasat Y. Listeriosis of the placenta: clinicopathologic study of seven cases. *Am J Obstet Gynecol*. 1993; 169:616-20. [PubMed: 8372871]
153. Vicente MF, Perez-Daz JC, Baquero F, Angel de PM, Berenguer J. Penicillin-binding protein 3 of *Listeria monocytogenes* as the primary lethal target for beta-lactams. *Antimicrob Agents Chemother*. 1990; 34:539-42. [PubMed: 2111657]
154. Voetsch AC, Angulo FJ, Jones TF, Moore MR, Nadon C, McCarthy P, et al. Reduction in the incidence of invasive listeriosis in foodborne diseases active surveillance network sites, 1996-2003. *Clin Infect Dis*. 2007; 44:513-20. [PubMed: 17243053]
155. Vogt RL, Donnelly C, Gellin B, Bibb W, Swaminathan B. Linking environmental and human strains of *Listeria monocytogenes* with isoenzyme and ribosomal RNA typing. *Eur J Epidemiol*. 1990; 6:229-30. [PubMed: 2113872]
156. Watkins J, Sleath KP. Isolation and enumeration of *Listeria monocytogenes* from Sewage, Sewage Sludge and River Water. *J Appl Bacteriol*. 1981; 50:1-9. [PubMed: 6262293]
157. Welshimer HJ, Donker-Voet J. *Listeria monocytogenes* in nature. *Appl Microbiol*. 1971; 21:516-9. [PubMed: 4994904]

158. Winslow DL, Pankey GA. In vitro activities of trimethoprim and sulfamethoxazole against *Listeria monocytogenes*. *Antimicrob Agents Chemother*. 1982; 22:51–4. [PubMed: 6812496]
159. Zervoudakis IA, Cederqvist LL. Effect of *Listeria monocytogenes* septicemia during pregnancy on the offspring. *Am J Obstet Gynecol*. 1977; 129:465–7. [PubMed: 910831]

Table 1CDC Recommendations for the Prevention of *Listeriosis***For the general public:**

- Thoroughly cook raw food from animal sources, such as beef, pork, or poultry
- Wash raw vegetables thoroughly before eating
- Keep uncooked meats separate from vegetables and from cooked foods and ready-to-eat foods
- Avoid unpasteurized (raw) milk or foods made from unpasteurized milk
- Wash hands, knives, and cutting boards after the handling of uncooked foods
- Consume perishable and ready-to-eat foods as soon as possible

For persons at high-risk, such as pregnant women and persons with weakened immune systems, in addition to the recommendations listed above:

- Do not eat hot dogs, luncheon meats, or deli meats, unless they are reheated until steaming hot
- Avoid getting fluid from hot dog packages on other foods, utensils, and food preparation surfaces, and wash hands after handling hot dogs, luncheon meats, and deli meats
- Do not eat soft cheeses, such as feta, brie, and camembert; blue-veined cheeses; or Mexican-style cheeses, such as *queso blanco*, *queso fresco*, and *Panela*, unless they have labels that clearly state they are made from pasteurized milk
- Do not eat refrigerated pâtés or meat spreads. Canned or shelf-stable pâtés and meat spreads may be eaten.
- Do not eat refrigerated smoked seafood, unless it is contained in a cooked dish, such as a casserole. Refrigerated smoked seafood, such as salmon, trout, whitefish, cod, tuna or mackerel, is most often labeled as “nova-style,” “lox,” “kippered,” “smoked,” or “jerky.” The fish is found in the refrigerator section or sold at deli counters of grocery stores and delicatessens. Canned or shelf-stable smoked seafood may be eaten.

Source: Centers for Disease Control and Prevention Centers for Disease Control and Prevention Web site, Listeriosis technical information and FAQs www.cdc.gov/ncidod/dbmd/diseaseinfo/listeriosis_t.htm [Accessed April 7th 2010]