Published in final edited form as:

Physiol Behav. 2014 January 17; 123: . doi:10.1016/j.physbeh.2012.07.014.

Maternal High Fat Diet Consumption during the Perinatal Period Programs Offspring Behavior

Elinor L. Sullivan^{a,b}, Libby Nousen^a, and Katie Chamblou^a

^aDepartment of Biology, University of Portland, Portland, OR, USA

^bDivision of Neuroscience, Oregon National Primate Research Center, Beaverton, OR, USA

Abstract

The environment that developing offspring experience during the perinatal period is markedly influenced by maternal health and diet composition. Evidence from both epidemiological studies and animal models indicates that maternal diet and metabolic status play a critical role in programming the neural circuitry that regulates behavior, resulting in long-term consequences for offspring behavior. Maternal diet and metabolic state influence the behavior of offspring directly by impacting the intrauterine environment and indirectly by modulating maternal behavior. The mechanisms by which maternal diet and metabolic profile shape the perinatal environment remain largely unknown, but recent research has found that increases in inflammatory cytokines, nutrients (glucose and fatty acids), and hormones (insulin and leptin) affect the environment of the developing offspring. Offspring exposed to maternal obesity and high fat diet consumption during development are more susceptible to developing mental health and behavioral disorders such as anxiety, depression, attention deficit hyperactivity disorder, and autism spectrum disorders. Recent evidence suggests that this increased risk for behavioral disorders is driven by modifications in the development of neural pathways involved in behavioral regulation. In particular, research indicates that the development of the serotonergic system is impacted by exposure to maternal obesity and high fat diet consumption, and this disruption may underlie many of the behavioral disturbances observed in these offspring. Given the high rates of obesity and high fat diet consumption in pregnant women, it is vital to examine the influence that maternal nutrition and metabolic profile have on the developing offspring.

Keywords

maternal high fat diet; obesity; anxiety; depression; inflammation; autism spectrum disorders; attention deficit hyperactivity disorder

1.0 Introduction

The majority of infants born in the United States and other western countries are exposed to maternal obese or overweight metabolic status and high fat diet (HFD) consumption during perinatal development. In the U.S., data from latest National Health and Nutrition

© 2012 Elsevier Inc. All rights reserved.

Corresponding author: Elinor L. Sullivan, PhD, University of Portland, 5000 N. Willamette Blvd., Portland, OR 97203, sullivan@up.edu, (503) 943-8861 (phone), (503) 943-7784 (fax).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Examination Survey (compiled from 1999–2010) indicates that 69% of adults are overweight or obese [1]. This increased risk of obesity is thought to be due to hyperphagia [2] and decreased physical activity [3]. At this time, about one third of pregnant women in the U.S. are obese [4], and this metabolic state has persistent implications for both the women and their children. Mammals encounter two environments during neural development: the intrauterine and early postnatal environments. Both of these are highly influenced by maternal diet and metabolic status. Animal models provide strong evidence that perinatal nutrition has an enduring impact on numerous aspects of offspring physiology and behavior, including a higher likelihood of developing mental health disorders [5, 6], impairments in social behaviors [7], decreased cognitive abilities [6, 8, 9], enhanced response to stress [10, 11], and altered reward-based behaviors [12]. The metabolic profile of the mother has an additional impact on the developing offspring. Factors associated with maternal obesity, such as inflammation [13], hyperlipidemia, lipotoxicity, hyperglycemia, and insulin resistance, have each been demonstrated to have a long-term effect on the developing offspring and are associated with increased risk of mental health disorders including anxiety and depression [5, 6, 14], attention deficit hyperactivity disorder (ADHD) [15], and autism spectrum disorders (ASD) [16–18].

With the current prevalence of obesity in western nations, it is vital to examine the impact that this may have on the behavior and physiology of future generations. In animal models, a HFD is commonly used to promote maternal obesity. In the majority of these studies, most of the females consuming the HFD become obese, and so it is difficult to determine which outcomes are due to factors associated with maternal obesity (i.e. hyperlipidemia, hyperglycemia, hyperinsulinemia) versus the consumption of a HFD. Furthermore, maternal HFD consumption and obesity often co-occur in humans, making it difficult to discriminate the effects of the HFD itself from the maternal metabolic profile. Our group has developed a nonhuman primate (NHP) model of maternal obesity, and an important advantage of this model is that approximately one third of the females consuming the HFD are resistant to weight gain in response to the diet. These animals have body weights similar to control animals, and this allows us to distinguish the effects of HFD intake from maternal obesity. As the majority of studies examine maternal obesity induced by HFD consumption, this review will therefore discuss the effects of both factors and will assume that consumption of a diet high in fat results in obesity. In addition, we will highlight studies that are able to distinguish between the effects of maternal HFD consumption and maternal obesity on offspring.

2.0 Co-morbidity Between Obesity and Mental Health Disorders

In humans, the obese state itself is associated with increased risk of developing mental health disorders, including anxiety [19], depression [19], and ADHD [20]. Obesity also increases susceptibility to neurodegenerative disorders such as Alzheimer's [21–23] and Parkinson's disease [24, 25]. When combined with obesity, factors like socioeconomic status, ethnicity, and maternal attitudes further increase the propensity for children to develop problems with internalization and social interaction [26]. Conversely, anxiety and depression are known to influence human obesity risk by impacting ingestive behavior, food preference, and physical activity level. Mood disorders like anxiety and depression are associated with enhanced desire for palatable energy-dense foods [27] and decreased levels of physical activity [28]. Reward-based eating has been shown to decrease feelings of stress in human patients [29], and has been described as an addiction or a type of self-medication with comfort food [30]. This effect is supported by animal studies that show that consumption of a palatable diet decreases anxiety in the short-term [31] and decreases anxiety and depression-like symptoms in rats exposed to perinatal stress [32]. Evidence from rodent studies indicates that the observed reduction in anxiety-like behaviors is linked to an

increase in glucocorticoid receptor mRNA expression in the hippocampus, providing support to the idea that palatable diet consumption influences the endocrine system to alleviate anxiety [32]. Perinatal exposure to a HFD has been demonstrated to alter the development of key pathways implicated in regulating mood and behavior such as the serotonergic [33] and dopaminergic [2, 34] systems.

3.0 HFD Consumption Impacts Offspring Behavior by Modifying Maternal Behavior

When diet affects the behavior of mothers, it indirectly results in changes in the behavior of the offspring. In rodent models, naturally occurring individual differences in maternal care during early postnatal life are associated with the programming of differences in offspring behavior and stress response [35–37]. Mothers that display decreased grooming and a lack of attentive behavior towards their offspring have offspring with increased anxiety-like behavior, and mothers who are attentive have offspring who are less anxious and better suited to handle stressful situations [38, 39]. Offspring from attentive mothers have improved hippocampal plasticity during stress and enhanced contextual fear conditioning [37]. In addition, male offspring of rat dams that exhibited a high frequency of licking and grooming behavior towards their pups display decreased aggression towards their peers [35]. All of these measures indicate that the nature of the interaction between mother and offspring influences the behavior of the offspring.

There is also evidence that experiencing maternal rejection or separation has a long-term impact on offspring behavior. NHP offspring experiencing maternal rejection are reported to be more prone to anxiety-like behaviors [40]. Female rodent offspring exposed to maternal separation displayed increased anxiety-like behavior in social situations [41]. However, other studies find that early maternal rejection increases offspring independence and decreases infant stress behaviors [42]. Therefore, the behavioral outcome appears to be influenced by the time period when the offspring experiences the maternal separation or rejection. For example, monkeys separated from their mother at one week of age showed increased self-comforting behaviors (such as thumb sucking), whereas those separated from their mother at one month sought more social comfort [43]. These studies provide strong evidence that maternal behavior (or lack thereof) has an important effect on the behavior of offspring.

Several human studies further support the impact that maternal behavior and stress have on offspring behavior. For example, postpartum depression (PPD) is known to influence the behavior of both mother and child [44]. Children of mothers who experienced PPD exhibit an increase in both violent and internalizing behaviors (seen in withdrawal or depression) [44]. Adolescent daughters of mothers with major depression are also more likely to develop psychiatric disorders: 30 months after recruitment, 45% of these daughters had a psychiatric disorder while none of the daughters of control mothers exhibited any type of psychopathology [45]. Human and animal offspring exposed to maternal stress display elevated anxiety and an increase in negative interactions with peers [46–48]. In addition, the quality of parental emotional care is inversely associated with the offspring's risk of developing coronary heart disease [49]. It is hypothesized that these effects of maternal stress seen in the infant may be due to changes in the placental environment or epigenetics of the offspring [46]. However, the factors that affect this maternal environment remain largely unknown, and knowledge about the mechanisms by which maternal behavior influences the behavior of the next generation is limited.

Data from several studies indicates that maternal diet impacts maternal behavior and parenting style and therefore indirectly affects offspring behavior. Research from rodent

models reveals that consumption of a HFD increases nursing [50, 51] and grooming [50]. Higher grooming rates have been documented in rodent models to have a beneficial effect on offspring behavior [38, 52], yet the consequences of increased nursing have not been well characterized. Furthermore, evidence from overfeeding studies suggests that higher rates of nursing will lead to hyperphagia, increased body weight, and reprogramming of the hypothalamic pathways that regulate energy balance [53]. Another way that diet may affect maternal behavior is demonstrated in a rodent model in which consumption of a high energy diet disrupted maternal hippocampal function and, consequently, learning and memory [54]. This was hypothesized to be due to the observed decrease in mRNA expression of tight junction proteins in the choroid plexus and blood-brain barrier. Human studies also provide preliminary evidence that diet influences maternal behavior. Consumption of a healthy diet has been found to reduce risk for PPD and improve symptoms while reduced access to healthy food increases PPD symptomology [55]. These relationships exemplify the interconnectivity of maternal diet and behavior as well as demonstrating pathways in humans through which maternal diet affects maternal behavior and therefore indirectly influences the behavior of the next generation.

4.0 Evidence from Animal Studies that Maternal HFD Impacts Offspring Behavior

4.1 Maternal HFD Programming of Anxiety

Intake of a HFD during gestation has been well documented in animal studies to alter offspring behavior. Maternal HFD consumption is associated with heightened anxiety in NHP [5] and rodent offspring [6]. These studies find that maternal HFD consumption exposes offspring to increased inflammatory cytokines that impact neural development [6]. In the rat model, mothers consuming a diet high in saturated or trans fat during pregnancy and lactation produced male pups that were more anxious as adults. The male rats on a trans fat diet swam faster in a swim test, indicating a higher level of anxiety and a desire to escape. The female rats did not exhibit this increase in anxiety, but this may have been because their baseline anxiety level was already elevated. In contrast, our NHP model found that exposure to a perinatal HFD resulted in increased anxiety in female offspring, which was associated with a reduction in neural serotonin [5]. Greater occurrence of anxiety in these female NHP offspring corresponds with human studies, which report that females have an increased susceptibility to anxiety and have a stronger association between obesity and anxiety [56].

It is clear that the timing of introduction to a diet high in fat during the perinatal period is critical. Offspring from mothers fed a HFD during gestation display increased anxiety [6], but exposure to a similar diet during lactation shows contradictory results. A cafeteria diet, which consists of chow supplemented with highly palatable food items such as cakes, meat pies, and cookies, resulted in reduced levels of anxiety when provided during lactation [57]. Additional support for this idea comes from findings that the anxiety and depression-like behaviors seen in offspring who experienced gestational stress are reduced by cafeteria diet consumption during the early postnatal period [32]. This data indicates that the behavioral outcome of offspring is differentially dependent on the timing of exposure to the HFD.

4.2 Maternal HFD Programming of Social Behaviors

The capacity to interact with conspecifics is essential for establishing stable social networks, for reproduction, and for survival, and these communicative behaviors are influenced by maternal diet. Several studies indicate that exposure to maternal HFD during the perinatal period impacts the social behavior of offspring. For example, maternal consumption of a diet high in polyunsaturated fatty acids (PUFA) led to offspring that displayed increased

aggressiveness to intruders, hyperlocomotion, and decreased immobility in a swim test [7]. This study examined several different strains of mice, and females in all three strains showed elevated rates of aggression. The increases in aggressiveness and locomotion and the decrease in immobility in the swim test were hypothesized to be due to an observed elevation in hypothalamic protein kinase C (PKC) activity. PKC was permanently down-regulated in most of the brain except for the hypothalamus, and it was proposed that PKC could serve as a biological mediator of the effect of maternal HFD diet on offspring behavior. PKC is also involved in the regulation of memory, manic-depressive illnesses, and alcohol-related behaviors [58,59], so the altered levels of PKC from a high PUFA maternal diet suggest a relationship between human mental health disorders and diet.

4.3 Maternal HFD Consumption Impacts Cognition

Evidence also demonstrates that maternal HFD is linked with differences in the cognitive abilities of offspring. Adult male rats from mothers exposed to a diet high in saturated or trans fat during gestation and lactation displayed deficits in spatial cognitive function [6]. Offspring of obese mothers had decreased hippocampal brain-derived neurotrophic factor (BDNF) production that resulted in decreased neurogenesis in this area [9]. These deficits in BDNF were linked to a delay in spatial learning in mouse offspring, but this cognitive impairment did not persist into adulthood, which indicates that the effects of maternal HFD diet on exploratory behavior may be most prominent in early life stages [9]. However, another study reported that maternal consumption of a diet high in lard impacted the spatial memory and learning ability of adult offspring [60]. HFD offspring also exhibited increased hippocampal microglial activation at birth that continued into adulthood [6], revealing the presence of inflammation in areas of the brain responsible for cognitive function. Thus, there is preliminary evidence that maternal diet and metabolic profile may negatively affect offspring cognitive abilities, and it will be important for future studies to clarify the role that maternal diet and obesity play in offspring cognition.

4.4 Maternal HFD Consumption Impacts Reward Pathways

Reward-based behaviors are also altered in the offspring of mothers consuming a HFD [61–64]. Preliminary findings from our NHP model indicate that the offspring exposed to maternal HFD consumption have heightened preference for diets with a high sugar and fat content (Sullivan and Grove, unpublished observation). Increased desire for fat, sugar, and salt was also documented in adult rat offspring exposed to junk food during either gestation or lactation [61–63]. These preferences create a positive feedback cycle where maternal HFD consumption programs cravings for palatable food in the offspring, which further increases HFD consumption. These results are supported by evidence that maternal HFD intake causes perturbations in the dopaminergic system of adult rodents in areas associated with the rewarding value of food, such as the nucleus accumbens and ventral tegmental area [12]. This alteration of the dopaminergic system in the nucleus accumbens also leads to decreased behavioral sensitization to amphetamine due to a reduction in negative feedback [65]. Thus, perinatal nutrition appears to have a persistent impact on reward-seeking behaviors like palatable food consumption and desensitization to illicit substances.

5.0 Maternal Obesity is Linked to Elevated Risk of Mental Health Disorders

Evidence from animal models indicates that maternal obesity increases the risk for offspring to develop neurological and psychological dysfunction. Results from human studies provide further support that maternal obesity or overnutrition is linked to mental health disorders in children. Pre-pregnancy maternal obesity predicted a two-fold increase in the likelihood of the child having difficulties with emotional intensity and the regulation of sadness, fear, and anger [66]. It is also twice as likely for the mother of a child with ADHD to be obese

compared to the mother of a child without ADHD [15]. Obesity prior to pregnancy also doubles the risk of the child developing ADHD symptoms compared to the children of women with a healthy weight status during pregnancy [67]. Genetic predispositions for being overweight and having ADHD did not explain this relationship because it remained after adjusting for ADHD symptoms in the parent [66]. A proposed mechanism behind this correlation is dysfunction in dopaminergic and serotonergic systems that leads to ADHD symptoms [67]. These serotonergic disruptions, as previously mentioned, have been found in a primate model in response to maternal HFD exposure, and future studies are examining ADHD symptomology in these animals.

Diabetes is another condition associated with maternal obesity and HFD consumption, and correlational studies in humans have found associations between maternal diabetes and psychological dysfunction in children. Children of diabetic mothers have higher rates of ADHD symptoms and hyperactivity, similar to the offspring of overweight or obese mothers [68]. Children of diabetic mothers are also at increased risk for disturbances like anxiety, depression, and trouble with social situations [26]. The children of diabetic mothers display deficits in fine and gross motor function, but their IQ score does not differ from control children [68] despite deficits in expressive verbal skills [69]. These findings further support the presence of a link between maternal metabolic state and the mental and social functioning of offspring.

Recent evidence indicates that exposure to maternal obesity and HFD consumption during fetal development may also increase the risk of developing ASD. Rising rates of ASD are matched with the high frequency of adult obesity in North America [70]. Moreover, maternal metabolic states such as obesity, diabetes, and hypertension have been recently shown to be associated with greater risk of ASD in offspring [69]. A proposed mechanism is the role of leptin: leptin is released in proportion to body fat, and high levels of leptin observed in obesity are associated with placental dysfunction that disrupts normal neurological development [71, 72]. When compared with controls, children with ASD have higher plasma levels of leptin [18]. Further research will need to examine the strength of the relationship between these factors.

The role that placental dysfunction plays in the development of ASD is relevant because of the physiological similarity between maternal HFD consumption and infections or illness, all of which have been found to increase ASD risk when they occur during pregnancy [73]. Infections or illness during pregnancy provoke an inflammatory response and cause the transfer of elevated inflammatory cytokines to the developing fetus, similar to the effects of HFD consumption and gestational obesity [74]. This increased exposure to cytokines causes an inflammatory reaction in the fetal brain during crucial developmental periods [16]. The placental dysfunction that is seen with hyperleptinemia has also been documented with high levels of inflammatory cytokines [75, 76], making offspring of obese mothers particularly susceptible to placental dysfunction. Numerous studies have demonstrated that environmental factors that increase exposure of the developing offspring to inflammatory cytokines result in behavioral abnormalities consistent with ASD [73, 77, 78] and ADHD [78]. Obesity is considered an inflammatory and immune disease, and it also results in an increase in a host of inflammatory cytokines circulating through the body and numerous organs, including the brain [13]. Maternal obesity and gestational diabetes have been associated with the activation of inflammatory responses in the placenta [79–82], and NHP [74] and ovine [83] models show a link between HFD consumption and placental dysfunction. Importantly, gestational consumption of a HFD stimulates inflammatory cytokines such as interleukin (IL)-4, IL-5, monocyte chemoattractant protein-1, RANTES, and granulocyte macrophage colony-stimulating factor, and these cytokines have been associated with ASD in humans [84–86]. Furthermore, these HFD offspring have a suppression of the

adaptive immune response, which is also a marker of the ASD profile in humans [73]. The inflammatory cytokines passed from an obese or diabetic mother to the fetus could therefore initiate the physiological and behavioral responses observed in children with ASD whose mothers experienced infections during pregnancy. Additionally, inflammation from poor maternal metabolic health and diet differs from infection-induced inflammation in that it results in prolonged exposure to inflammatory cytokines, and therefore it has a long-term impact on fetal development.

6.0 Mechanisms for HFD Programming of Behavior

The field is at an early stage of defining the mechanisms by which maternal HFD consumption and obesity influence the development of the complex neural circuitry that regulates behavior. There is evidence that neurotransmitter signaling pathways such as the serotonergic [5] and dopaminergic [34] systems are modified by maternal HFD consumption. The development of these pathways is influenced by exposure to increased levels of circulating inflammatory cytokines, hormones (leptin, insulin), and nutrients (fatty acids, triglycerides, and glucose). It is also possible (as discussed in sub section 3.0) that maternal HFD consumption impacts offspring neural development indirectly by modifying maternal behavior toward the infant, which has also been shown to induce changes in neural pathways critical in regulating behavior: the serotonergic [5], dopaminergic [2], and melanocortinergic [87] systems.

6.1 Inflammation-Induced Programming

Enduring inflammation occurs in the obese state because greater adiposity results in an elevation in many circulating markers of inflammation, including c reactive protein, IL-6, IL-1 β , and tumor necrosis factor- α [13]. Risk for a number of metabolic diseases such as cardiovascular disease, heart disease, insulin resistance, type II diabetes mellitus, and hypertension increases with the exposure to elevated levels of these inflammatory markers [13]. The relationship between obesity and higher levels of inflammatory cytokines has been observed in pregnant women; obese pregnant women have increased levels of inflammatory cytokines, which leads to endothelial [75] and placental dysfunction [88, 89]. There is strong evidence that exposure to higher levels of circulating cytokines during the neonatal period impacts neural development and thus may be a potential mechanism by which maternal HFD consumption programs the brain. The development of neurotransmitter systems that are important for behavioral regulation, such as the serotonergic system, dopaminergic system, and melanocortinergic system, have all been shown to be sensitive to circulating cytokine levels [13]. Fetal NHP offspring from mothers consuming a HFD exhibited elevated systemic and hypothalamic cytokines early in the third trimester [87], which may have led to the observed disruptions in the melanocortinergic [87] and serotonergic systems [5].

Further evidence for this hypothesis is supported by human studies, which find that emotional and behavioral disorders including anxiety [90, 91], depression [90, 92], ASD [84–86], and ADHD [93, 94] are associated with increased inflammatory cytokines. As exposure to inflammation during development causes a nonspecific response that impacts many neurotransmitter systems, it is important for future research to directly examine the influence of maternal obesity and HFD consumption-induced inflammation on each neural pathway important in behavioral regulation. A step in this direction is seen in the research on ursolic acid, which improved cognitive function when given to mice consuming a HFD [60]. As ursolic acid inhibits inflammatory signaling, the observed improvements suggest that anti-inflammatory compounds may be therapeutic for treating cognitive deficits due to HFD consumption [60]. An animal model has recently demonstrated that anti-inflammatory

compounds may be beneficial for treating Parkinson's disease as they block the neurodegenerative pathways stimulated by pro-inflammatory cytokines [24].

6.2 Programming by Excess Hormones and Nutrients

Maternal obesity has clear associations with gestational diabetes, maternal hyperglycemia, and hyperinsulinemia [95]. Glucose, but not insulin, can pass through the blood-placenta barrier and enter the fetal environment [96], and consequently the fetal pancreas must compensate for the resultant hyperglycemia by enhancing insulin release. Insulin is an important neural growth factor [97], and exposure to high levels of insulin during critical periods of brain development causes perturbations in the neural circuitry critical in regulating energy balance. For example, insulin administration during the last trimester of gestation results in offspring obesity [98, 99], and direct administration of insulin to the hypothalamus when the projections between the arcuate nucleus and the paraventricular nucleus are forming results in rodent offspring that have elevated body weight, hyperinsulinemia, impaired glucose tolerance, and elevated diabetes risk [100].

The exposure of offspring of obese mothers to high levels of leptin during development is also postulated to influence circuitry in the brain that regulates behavior and physiology. The concentration of leptin experienced during fetal development is critical in regulating the wiring of brain areas such as the hypothalamus [97], thus an elevation in fetal leptin concentration may be a critical factor in programming behavior and physiology. Rodent models indicate that postnatal leptin plays an important role in the development of hypothalamic neural circuitry [101, 102]. A role for leptin is further confirmed by studies in humans that demonstrated increased leptin levels in obese and diabetic mothers [103, 104] and lower leptin in infants that experienced intrauterine growth restriction at term [105]. It is important to note that circulating leptin levels do not rise until after completion of the development of the majority of neural circuitry in the hypothalamus in human and NHP gestation [105, 106]. Though leptin has been shown to be critical in rodent brain development, its role in primate brain development is uncertain [107]. However, high levels of leptin are associated with dysfunction of the placenta [81, 108], and so it is possible that leptin impacts primate brain development indirectly by acting on the placenta. Leptin's influence on the development of brain regions and pathways that regulate behavior has been largely unexplored. Given recent studies which indicate an important role of maternal metabolic state on offspring behavior, it is imperative that the role that leptin plays in programming behavior is further explored.

6.3 The Serotonin System is Suppressed by Maternal HFD Consumption

Numerous aspects of brain development including neuronal migration, neurogenesis, and synaptogenesis are regulated by the serotonin (5-HT) system [109, 110]. In addition, metabolism of 5-HT's precursor tryptophan (TRP) by the kynurenine (KYN) pathway plays an important role in modulating immune function during pregnancy by preventing allogeneic rejection of the fetus [111] and regulating blood flow between the fetus and placenta. The induction of maternal inflammation in animal models produces an increase in KYN metabolites [112]. As the KYN pathway competes with 5-HT for TRP, an up-regulation of this pathway decreases TRP availability for 5-HT synthesis. This reduction in 5-HT synthesis may therefore impact immune function and placental blood flow, which matches the results seen in our NHP model. As described earlier in this review, chronic consumption of a HFD during pregnancy in NHPs reduces placental blood flow, and this effect is amplified if the animals are obese and insulin resistant. Reduction in neural 5-HT synthesis is associated with mental health and behavioral disorders in humans, including anxiety [33], depression [113], ADHD [114], and ASD [81, 115], and so this suppression of

the serotonergic system increases the risk of offspring developing a number of behavioral disorders.

7.0 Conclusion

The mechanisms by which maternal obesity and HFD consumption may impact neural development and the regulation of offspring behavior include increased exposure to inflammatory cytokines, nutrients, and hormones in offspring of obese mothers in addition to the consequences of placental dysfunction. A primary potential mediator of maternal HFD-induced behavioral dysregulation is the serotonergic system because suppression of this system in response to maternal HFD consumption has been documented in a number of animal models. Furthermore, this suppression of the serotonin system results in the increased occurrence of behavioral abnormalities that are documented across animal models of maternal HFD-induced obesity. These behavioral irregularities are consistent with psychopathologies linked to maternal obesity in humans. This body of research shows that the current frequency of maternal obesity and HFD consumption will put future generations at greater risk of developing behavioral and mental health disorders. In response to the present staggering rates of maternal obesity, future studies need to investigate therapies that prevent and reduce the impact that maternal HFD consumption has on offspring behavioral regulation.

References

1. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA : the journal of the American Medical Association*. 2012; 307:491–497. [PubMed: 22253363]
2. Vucetic Z, Reyes TM. Central dopaminergic circuitry controlling food intake and reward: implications for the regulation of obesity. *Wiley interdisciplinary reviews. Systems biology and medicine*. 2010; 2:577–593. [PubMed: 20836049]
3. Scheers T, Philippaerts R, Lefevre J. Patterns of physical activity and sedentary behavior in normal-weight, overweight and obese adults, as measured with a portable armband device and an electronic diary. *Clin Nutr*. 2012
4. King JC. Maternal obesity, metabolism, and pregnancy outcomes. *Annual review of nutrition*. 2006; 26:271–291.
5. Sullivan EL, Grayson B, Takahashi D, Robertson N, Maier A, Bethea CL, Smith MS, Coleman K, Grove KL. Chronic consumption of a high-fat diet during pregnancy causes perturbations in the serotonergic system and increased anxiety-like behavior in nonhuman primate offspring. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2010; 30:3826–3830. [PubMed: 20220017]
6. Bilbo SD, Tsang V. Enduring consequences of maternal obesity for brain inflammation and behavior of offspring. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2010; 24:2104–2115. [PubMed: 20124437]
7. Raygada M, Cho E, Hilakivi-Clarke L. High maternal intake of polyunsaturated fatty acids during pregnancy in mice alters offsprings' aggressive behavior, immobility in the swim test, locomotor activity and brain protein kinase C activity. *The Journal of nutrition*. 1998; 128:2505–2511. [PubMed: 9868200]
8. Yu H, Bi Y, Ma W, He L, Yuan L, Feng J, Xiao R. Long-term effects of high lipid and high energy diet on serum lipid, brain fatty acid composition, and memory and learning ability in mice. *International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience*. 2010; 28:271–276. [PubMed: 20015474]
9. Tozuka Y, Kumon M, Wada E, Onodera M, Mochizuki H, Wada K. Maternal obesity impairs hippocampal BDNF production and spatial learning performance in young mouse offspring. *Neurochemistry international*. 2010; 57:235–247. [PubMed: 20538025]

10. D'Asti E, Long H, Tremblay-Mercier J, Grajzer M, Cunnane SC, Di Marzo V, Walker CD. Maternal dietary fat determines metabolic profile and the magnitude of endocannabinoid inhibition of the stress response in neonatal rat offspring. *Endocrinology*. 2010; 151:1685–1694. [PubMed: 20160134]
11. Augustyniak RA, Singh K, Zeldes D, Singh M, Rossi NF. Maternal protein restriction leads to hyperresponsiveness to stress and salt-sensitive hypertension in male offspring. *American journal of physiology. Regulatory, integrative and comparative physiology*. 2010; 298:R1375–R1382.
12. Naef L, Srivastava L, Gratton A, Hendrickson H, Owens SM, Walker CD. Maternal high fat diet during the perinatal period alters mesocorticolimbic dopamine in the adult rat offspring: reduction in the behavioral responses to repeated amphetamine administration. *Psychopharmacology*. 2008; 197:83–94. [PubMed: 18004547]
13. Das UN. Is obesity an inflammatory condition? *Nutrition*. 2001; 17:953–966. [PubMed: 11744348]
14. Vickers MH, Breier BH, Cutfield WS, Hofman PL, Gluckman PD. Fetal origins of hyperphagia, obesity, and hypertension and postnatal amplification by hypercaloric nutrition. *American journal of physiology. Endocrinology and metabolism*. 2000; 279:E83–E87. [PubMed: 10893326]
15. Ray GT, Croen LA, Habel LA. Mothers of children diagnosed with attention-deficit/hyperactivity disorder: health conditions and medical care utilization in periods before and after birth of the child. *Medical care*. 2009; 47:105–114. [PubMed: 19106738]
16. Patterson PH. Maternal infection and immune involvement in autism. *Trends in molecular medicine*. 2011; 17:389–394. [PubMed: 21482187]
17. Patterson PH. Maternal infection and autism. *Brain, behavior, and immunity*. 2011
18. Ashwood P, Kwong C, Hansen R, Hertz-Picciotto I, Croen L, Krakowiak P, Walker W, Pessah IN, Van de Water J. Brief report: plasma leptin levels are elevated in autism: association with early onset phenotype? *Journal of autism and developmental disorders*. 2008; 38:169–175. [PubMed: 17347881]
19. Rofey DL, Kolko RP, Iosif AM, Silk JS, Bost JE, Feng W, Szigethy EM, Noll RB, Ryan ND, Dahl RE. A Longitudinal Study of Childhood Depression and Anxiety in Relation to Weight Gain. *Child Psychiatry Hum Dev*. 2009
20. Waring ME, Lapane KL. Overweight in children and adolescents in relation to attention-deficit/hyperactivity disorder: results from a national sample. *Pediatrics*. 2008; 122:e1–e6. [PubMed: 18595954]
21. Pasinetti GM, Eberstein JA. Metabolic syndrome and the role of dietary lifestyles in Alzheimer's disease. *Journal of neurochemistry*. 2008; 106:1503–1514. [PubMed: 18466323]
22. Grant WB, Campbell A, Itzhaki RF, Savory J. The significance of environmental factors in the etiology of Alzheimer's disease. *Journal of Alzheimer's disease : JAD*. 2002; 4:179–189.
23. Ross MG, Desai M, Khorram O, McKnight RA, Lane RH, Torday J. Gestational programming of offspring obesity: a potential contributor to Alzheimer's disease. *Current Alzheimer research*. 2007; 4:213–217. [PubMed: 17430249]
24. Peterson LJ, Flood PM. Oxidative stress and microglial cells in Parkinson's disease. *Mediators of inflammation*. 2012; 2012:401264. [PubMed: 22544998]
25. Priyadarshini M, Kamal MA, Greig NH, Realef M, Abuzenadah AM, Chaudhary AG, Damanhour GA. Alzheimer's disease and type 2 diabetes: exploring the association to obesity and tyrosine hydroxylase. *CNS & neurological disorders drug targets*. 2012; 11:482–489. [PubMed: 22583431]
26. Rizzo TA, Silverman BL, Metzger BE, Cho NH. Behavioral adjustment in children of diabetic mothers. *Acta Paediatr*. 1997; 86:969–974. [PubMed: 9343277]
27. Kelley AE, Schiltz CA, Landry CF. Neural systems recruited by drug- and food-related cues: studies of gene activation in corticolimbic regions. *Physiol Behav*. 2005; 86:11–14. [PubMed: 16139315]
28. Teychenne M, Ball K, Salmon J. Physical activity, sedentary behavior and depression among disadvantaged women. *Health Educ Res*. 2010
29. Adam TC, Epel ES. Stress, eating and the reward system. *Physiology & behavior*. 2007; 91:449–458. [PubMed: 17543357]
30. Dallman MF, Pecoraro NC, la Fleur SE. Chronic stress and comfort foods: self-medication and abdominal obesity. *Brain, behavior, and immunity*. 2005; 19:275–280.

31. Prasad A, Prasad C. Short-term consumption of a diet rich in fat decreases anxiety response in adult male rats. *Physiology & behavior*. 1996; 60:1039–1042. [PubMed: 8873290]
32. Maniam J, Morris MJ. Palatable cafeteria diet ameliorates anxiety and depression-like symptoms following an adverse early environment. *Psychoneuroendocrinology*. 2010; 35:717–728. [PubMed: 19939573]
33. Kiyohara C, Yoshimasu K. Molecular epidemiology of major depressive disorder. *Environ Health Prev Med*. 2009; 14:71–87. [PubMed: 19568851]
34. Vucetic Z, Kimmel J, Totoki K, Hollenbeck E, Reyes TM. Maternal high-fat diet alters methylation and gene expression of dopamine and opioid-related genes. *Endocrinology*. 2010; 151:4756–4764. [PubMed: 20685869]
35. Parent CI, Meaney MJ. The influence of natural variations in maternal care on play fighting in the rat. *Developmental psychobiology*. 2008; 50:767–776. [PubMed: 18846499]
36. Bagot RC, Meaney MJ. Epigenetics and the biological basis of gene x environment interactions. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010; 49:752–771. [PubMed: 20643310]
37. Bagot RC, van Hasselt FN, Champagne DL, Meaney MJ, Krugers HJ, Joels M. Maternal care determines rapid effects of stress mediators on synaptic plasticity in adult rat hippocampal dentate gyrus. *Neurobiology of learning and memory*. 2009; 92:292–300. [PubMed: 19292996]
38. Weaver IC, Meaney MJ, Szyf M. Maternal care effects on the hippocampal transcriptome and anxiety-mediated behaviors in the offspring that are reversible in adulthood. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103:3480–3485. [PubMed: 16484373]
39. Walker CD. Maternal touch and feed as critical regulators of behavioral and stress responses in the offspring. *Developmental psychobiology*. 2010; 52:638–650. [PubMed: 20862707]
40. Maestripieri D, McCormack K, Lindell SG, Higley JD, Sanchez MM. Influence of parenting style on the offspring's behaviour and CSF monoamine metabolite levels in crossfostered and noncrossfostered female rhesus macaques. *Behavioural brain research*. 2006; 175:90–95. [PubMed: 16971003]
41. Tsuda MC, Ogawa S. Long-lasting consequences of neonatal maternal separation on social behaviors in ovariectomized female mice. *PloS one*. 2012; 7:e33028. [PubMed: 22412981]
42. Schino G, Speranza L, Troisi A. Early maternal rejection and later social anxiety in juvenile and adult Japanese macaques. *Developmental psychobiology*. 2001; 38:186–190. [PubMed: 11279595]
43. Sabatini MJ, Ebert P, Lewis DA, Levitt P, Cameron JL, Mirnics K. Amygdala gene expression correlates of social behavior in monkeys experiencing maternal separation. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2007; 27:3295–3304. [PubMed: 17376990]
44. Verbeek T, Bockting CL, van Pampus MG, Ormel J, Meijer JL, Hartman CA, Burger H. Postpartum depression predicts offspring mental health problems in adolescence independently of parental lifetime psychopathology. *Journal of affective disorders*. 2012; 136:948–954. [PubMed: 21930302]
45. Gershon A, Hayward C, Schraedley-Desmond P, Rudolph KD, Booster GD, Gotlib IH. Life stress and first onset of psychiatric disorders in daughters of depressed mothers. *Journal of psychiatric research*. 2011; 45:855–862. [PubMed: 21524424]
46. Buss C, Davis EP, Hobel CJ, Sandman CA. Maternal pregnancy-specific anxiety is associated with child executive function at 6–9 years age. *Stress*. 2011; 14:665–676. [PubMed: 21995526]
47. Kersten-Alvarez LE, Hosman CM, Riksen-Walraven JM, van Doesum KT, Smeekens S, Hoefnagels C. Early school outcomes for children of postpartum depressed mothers: comparison with a community sample. *Child psychiatry and human development*. 2012; 43:201–218. [PubMed: 22011810]
48. Bardi M, Bode AE, Ramirez SM, Brent LY. Maternal care and development of stress responses in baboons. *American journal of primatology*. 2005; 66:263–278. [PubMed: 16015661]
49. Almeida ND, Loucks EB, Kubzansky L, Pruessner J, Maselko J, Meaney MJ, Buka SL. Quality of parental emotional care and calculated risk for coronary heart disease. *Psychosomatic medicine*. 2010; 72:148–155. [PubMed: 20064905]

50. Bertino M. Effects of high fat protein supplemented diets on maternal behavior in rats. *Physiology & behavior*. 1982; 29:999–1005. [PubMed: 7163403]
51. Purcell RH, Sun B, Pass LL, Power ML, Moran TH, Tamashiro KL. Maternal stress and high-fat diet effect on maternal behavior, milk composition, and pup ingestive behavior. *Physiology & behavior*. 2011; 104:474–479. [PubMed: 21605577]
52. Francis D, Diorio J, Liu D, Meaney MJ. Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science*. 1999; 286:1155–1158. [PubMed: 10550053]
53. Glavas MM, Kirigiti MA, Xiao XQ, Enriori PJ, Fisher SK, Evans AE, Grayson BE, Cowley MA, Smith MS, Grove KL. Early overnutrition results in early-onset arcuate leptin resistance and increased sensitivity to high-fat diet. *Endocrinology*. 2010; 151:1598–1610. [PubMed: 20194730]
54. Kanoski SE, Zhang Y, Zheng W, Davidson TL. The effects of a high-energy diet on hippocampal function and blood-brain barrier integrity in the rat. *Journal of Alzheimer's disease : JAD*. 2010; 21:207–219.
55. Okubo H, Miyake Y, Sasaki S, Tanaka K, Murakami K, Hirota Y. Dietary patterns during pregnancy and the risk of postpartum depression in Japan: the Osaka Maternal and Child Health Study. *The British journal of nutrition*. 2011; 105:1251–1257. [PubMed: 21144112]
56. Desai RA, Manley M, Desai MM, Potenza MN. Gender differences in the association between body mass index and psychopathology. *CNS Spectr*. 2009; 14:372–383. [PubMed: 19773713]
57. Wright T, Langley-Evans SC, Voigt JP. The impact of maternal cafeteria diet on anxiety-related behaviour and exploration in the offspring. *Physiology & behavior*. 2011; 103:164–172. [PubMed: 21241725]
58. Manji HK, Etcheberrigaray R, Chen G, Olds JL. Lithium decreases membrane-associated protein kinase C in hippocampus: selectivity for the alpha isozyme. *Journal of neurochemistry*. 1993; 61:2303–2310. [PubMed: 8245981]
59. Serrano PA, Rodriguez WA, Pope B, Bennett EL, Rosenzweig MR. Protein kinase C inhibitor chelerythrine disrupts memory formation in chicks. *Behavioral neuroscience*. 1995; 109:278–284. [PubMed: 7619317]
60. Lu J, Wu DM, Zheng YL, Hu B, Cheng W, Zhang ZF, Shan Q. Ursolic acid improves high fat diet-induced cognitive impairments by blocking endoplasmic reticulum stress and IkappaB kinase beta/nuclear factor-kappaB-mediated inflammatory pathways in mice. *Brain, behavior, and immunity*. 2011; 25:1658–1667.
61. Bayol SA, Farrington SJ, Stickland NC. A maternal 'junk food' diet in pregnancy and lactation promotes an exacerbated taste for 'junk food' and a greater propensity for obesity in rat offspring. *The British journal of nutrition*. 2007; 98:843–851. [PubMed: 17697422]
62. Walker CD, Naef L, d'Asti E, Long H, Xu Z, Moreau A, Azeddine B. Perinatal maternal fat intake affects metabolism and hippocampal function in the offspring: a potential role for leptin. *Ann N Y Acad Sci*. 2008; 1144:189–202. [PubMed: 19076377]
63. Nakashima Y. Fish-oil high-fat diet intake of dams after day 5 of pregnancy and during lactation guards against excessive fat consumption of their weaning pups. *J Nutr Sci Vitaminol (Tokyo)*. 2008; 54:46–53. [PubMed: 18388407]
64. Ong ZY, Muhlhauser BS. Maternal "junk-food" feeding of rat dams alters food choices and development of the mesolimbic reward pathway in the offspring. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2011; 25:2167–2179. [PubMed: 21427213]
65. Naef L, Moquin L, Dal Bo G, Giros B, Gratton A, Walker CD. Maternal high-fat intake alters presynaptic regulation of dopamine in the nucleus accumbens and increases motivation for fat rewards in the offspring. *Neuroscience*. 2011; 176:225–236. [PubMed: 21187125]
66. Rodriguez A. Maternal pre-pregnancy obesity and risk for inattention and negative emotionality in children. *Journal of child psychology and psychiatry, and allied disciplines*. 2010; 51:134–143.
67. Rodriguez A, Miettunen J, Henriksen TB, Olsen J, Obel C, Taanila A, Ebeling H, Linnet KM, Moilanen I, Jarvelin MR. Maternal adiposity prior to pregnancy is associated with ADHD symptoms in offspring: evidence from three prospective pregnancy cohorts. *Int J Obes (Lond)*. 2008; 32:550–557. [PubMed: 17938639]

68. Ornoy A. Growth and neurodevelopmental outcome of children born to mothers with pregestational and gestational diabetes. *Pediatric endocrinology reviews : PER*. 2005; 3:104–113. [PubMed: 16361984]
69. Krakowiak P, Walker CK, Bremer AA, Baker AS, Ozonoff S, Hansen RL, Hertz-Picciotto I. Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders. *Pediatrics*. 2012; 129:e1121–1128. [PubMed: 22492772]
70. Dodds L, Fell DB, Shea S, Armson BA, Allen AC, Bryson S. The role of prenatal, obstetric and neonatal factors in the development of autism. *Journal of autism and developmental disorders*. 2011; 41:891–902. [PubMed: 20922473]
71. Chappell LC, Seed PT, Briley A, Kelly FJ, Hunt BJ, Charnock-Jones DS, Mallet AI, Poston L. A longitudinal study of biochemical variables in women at risk of preeclampsia. *American journal of obstetrics and gynecology*. 2002; 187:127–136. [PubMed: 12114900]
72. Hauguel-de Mouzon S, Lepercq J, Catalano P. The known and unknown of leptin in pregnancy. *American journal of obstetrics and gynecology*. 2006; 194:1537–1545. [PubMed: 16731069]
73. Onore C, Careaga M, Ashwood P. The role of immune dysfunction in the pathophysiology of autism. *Brain, behavior, and immunity*. 2011
74. Frias AE, Morgan TK, Evans AE, Rasanen J, Oh KY, Thornburg KL, Grove KL. Maternal high-fat diet disturbs uteroplacental hemodynamics and increases the frequency of stillbirth in a nonhuman primate model of excess nutrition. *Endocrinology*. 2011; 152:2456–2464. [PubMed: 21447636]
75. Stewart FM, Freeman DJ, Ramsay JE, Greer IA, Caslake M, Ferrell WR. Longitudinal assessment of maternal endothelial function and markers of inflammation and placental function throughout pregnancy in lean and obese mothers. *J Clin Endocrinol Metab*. 2007; 92:969–975. [PubMed: 17192290]
76. Park CW, Moon KC, Park JS, Jun JK, Yoon BH. The frequency and clinical significance of intra-uterine infection and inflammation in patients with placenta previa and preterm labor and intact membranes. *Placenta*. 2009; 30:613–618. [PubMed: 19447490]
77. Buehler MR. A proposed mechanism for autism: an aberrant neuroimmune response manifested as a psychiatric disorder. *Medical hypotheses*. 2011; 76:863–870. [PubMed: 21421290]
78. Tohmi M, Tsuda N, Watanabe Y, Kakita A, Nawa H. Perinatal inflammatory cytokine challenge results in distinct neurobehavioral alterations in rats: implication in psychiatric disorders of developmental origin. *Neuroscience research*. 2004; 50:67–75. [PubMed: 15288500]
79. Roberts KA, Riley SC, Reynolds RM, Barr S, Evans M, Statham A, Hor K, Jabbour HN, Norman JE, Denison FC. Placental structure and inflammation in pregnancies associated with obesity. *Placenta*. 2011; 32:247–254. [PubMed: 21232790]
80. Basu S, Haghiac M, Surace P, Challier JC, Guerre-Millo M, Singh K, Waters T, Minium J, Presley L, Catalano PM, Hauguel-de Mouzon S. Pregravid obesity associates with increased maternal endotoxemia and metabolic inflammation. *Obesity (Silver Spring)*. 2011; 19:476–482. [PubMed: 20930711]
81. Challier JC, Basu S, Bintein T, Minium J, Hotmire K, Catalano PM, Hauguel-de Mouzon S. Obesity in pregnancy stimulates macrophage accumulation and inflammation in the placenta. *Placenta*. 2008; 29:274–281. [PubMed: 18262644]
82. Radaelli T, Varastehpour A, Catalano P, Hauguel-de Mouzon S. Gestational diabetes induces placental genes for chronic stress and inflammatory pathways. *Diabetes*. 2003; 52:2951–2958. [PubMed: 14633856]
83. Wallace JM, Milne JS, Matsuzaki M, Aitken RP. Serial measurement of uterine blood flow from mid to late gestation in growth restricted pregnancies induced by overnourishing adolescent sheep dams. *Placenta*. 2008; 29:718–724. [PubMed: 18579200]
84. Goines PE, Croen LA, Braunschweig D, Yoshida CK, Grether J, Hansen R, Kharrazi M, Ashwood P, Van de Water J. Increased midgestational IFN-gamma, IL-4 and IL-5 in women bearing a child with autism: A case-control study. *Molecular autism*. 2011; 2:13. [PubMed: 21810230]
85. Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah IN, Van de Water J. Altered T cell responses in children with autism. *Brain, behavior, and immunity*. 2011; 25:840–849.

86. Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah IN, Van de Water J. Associations of impaired behaviors with elevated plasma chemokines in autism spectrum disorders. *Journal of neuroimmunology*. 2011; 232:196–199. [PubMed: 21095018]
87. Grayson BE, Levasseur PR, Williams SM, Smith MS, Marks DL, Grove KL. Changes in melanocortin expression and inflammatory pathways in fetal offspring of nonhuman primates fed a high-fat diet. *Endocrinology*. 2010; 151:1622–1632. [PubMed: 20176722]
88. Leung P, Bryant RA. Autobiographical memory in diabetes mellitus patients. *Journal of psychosomatic research*. 2000; 49:435–438. [PubMed: 11182437]
89. Nordahl CW, Simon TJ, Zierhut C, Solomon M, Rogers SJ, Amaral DG. Brief report: methods for acquiring structural MRI data in very young children with autism without the use of sedation. *Journal of autism and developmental disorders*. 2008; 38:1581–1590. [PubMed: 18157624]
90. Terness P, Bauer TM, Rose L, Dufter C, Watzlik A, Simon H, Opelz G. Inhibition of allogeneic T cell proliferation by indoleamine 2,3-dioxygenase-expressing dendritic cells: mediation of suppression by tryptophan metabolites. *The Journal of experimental medicine*. 2002; 196:447–457. [PubMed: 12186837]
91. Maes M, Verkerk R, Bonaccorso S, Ombelet W, Bosmans E, Scharpe S. Depressive and anxiety symptoms in the early puerperium are related to increased degradation of tryptophan into kynurenine, a phenomenon which is related to immune activation. *Life sciences*. 2002; 71:1837–1848. [PubMed: 12175700]
92. Zanolli P, Cannazza G, Baraldi M. Prenatal exposure to methyl mercury in rats: focus on changes in kynurenine pathway. *Brain research bulletin*. 2001; 55:235–238. [PubMed: 11470321]
93. Oades RD. An exploration of the associations of pregnancy and perinatal features with cytokines and tryptophan/kynurenine metabolism in children with attention-deficit hyperactivity disorder (ADHD). *Attention deficit and hyperactivity disorders*. 2011; 3:301–318. [PubMed: 21785943]
94. Oades RD, Myint AM, Dauvermann MR, Schimmelmann BG, Schwarz MJ. Attention-deficit hyperactivity disorder (ADHD) and glial integrity: an exploration of associations of cytokines and kynurenine metabolites with symptoms and attention. *Behavioral and brain functions : BBF*. 2010; 6:32. [PubMed: 20534153]
95. Leung TW, Lao TT. Placental size and large-for-gestational-age infants in women with abnormal glucose tolerance in pregnancy. *Diabet Med*. 2000; 17:48–52. [PubMed: 10691159]
96. Oken E, Gillman MW. Fetal origins of obesity. *Obes Res*. 2003; 11:496–506. [PubMed: 12690076]
97. Simerly RB. Hypothalamic substrates of metabolic imprinting. *Physiol Behav*. 2008; 94:79–89. [PubMed: 18262209]
98. Jones AP, Pothos EN, Rada P, Olster DH, Hoebel BG. Maternal hormonal manipulations in rats cause obesity and increase medial hypothalamic norepinephrine release in male offspring. *Brain research. Developmental brain research*. 1995; 88:127–131. [PubMed: 8665659]
99. Jones AP, Dayries M. Maternal hormone manipulations and the development of obesity in rats. *Physiology & behavior*. 1990; 47:1107–1110. [PubMed: 2204072]
100. Plagemann A, Heidrich I, Gotz F, Rohde W, Dorner G. Lifelong enhanced diabetes susceptibility and obesity after temporary intrahypothalamic hyperinsulinism during brain organization. *Exp Clin Endocrinol*. 1992; 99:91–95. [PubMed: 1639125]
101. Djiane J, Attig L. Role of leptin during perinatal metabolic programming and obesity. *J Physiol Pharmacol*. 2008; 59(Suppl 1):55–63. [PubMed: 18802216]
102. Bouret SG. Development of hypothalamic neural networks controlling appetite. *Forum Nutr*. 2010; 63:84–93. [PubMed: 19955776]
103. Lepercq J, Hauguel-De Mouzon S, Timsit J, Catalano PM. Fetal macrosomia and maternal weight gain during pregnancy. *Diabetes Metab*. 2002; 28:323–328. [PubMed: 12442070]
104. Hauguel-de Mouzon S, Shafrir E. Carbohydrate and fat metabolism and related hormonal regulation in normal and diabetic placenta. *Placenta*. 2001; 22:619–627. [PubMed: 11504530]
105. Davidowa H, Plagemann A. Decreased inhibition by leptin of hypothalamic arcuate neurons in neonatally overfed young rats. *Neuroreport*. 2000; 11:2795–2798. [PubMed: 10976965]
106. Grayson BE, Allen SE, Billes SK, Williams SM, Smith MS, Grove KL. Prenatal development of hypothalamic neuropeptide systems in the nonhuman primate. *Neuroscience*. 2006; 143:975–986. [PubMed: 17029798]

107. Grayson BE, Kievit P, Smith MS, Grove KL. Critical determinants of hypothalamic appetitive neuropeptide development and expression: species considerations. *Frontiers in neuroendocrinology*. 2010; 31:16–31. [PubMed: 19822169]
108. Uzelac PS, Li X, Lin J, Neese LD, Lin L, Nakajima ST, Bohler H, Lei Z. Dysregulation of leptin and testosterone production and their receptor expression in the human placenta with gestational diabetes mellitus. *Placenta*. 2010; 31:581–588. [PubMed: 20421132]
109. Kannan S, Saadani-Makki F, Balakrishnan B, Dai H, Chakraborty PK, Janisse J, Muzik O, Romero R, Chugani DC. Decreased cortical serotonin in neonatal rabbits exposed to endotoxin in utero. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2011; 31:738–749. [PubMed: 20827261]
110. Daws LC, Gould GG. Ontogeny and regulation of the serotonin transporter: providing insights into human disorders. *Pharmacology & therapeutics*. 2011; 131:61–79. [PubMed: 21447358]
111. Munn DH, Zhou M, Attwood JT, Bondarev I, Conway SJ, Marshall B, Brown C, Mellor AL. Prevention of allogeneic fetal rejection by tryptophan catabolism. *Science*. 1998; 281:1191–1193. [PubMed: 9712583]
112. Pfaff AW, Mousli M, Senegas A, Marcellin L, Takikawa O, Klein JP, Candolfi E. Impact of foetus and mother on IFN-gamma-induced indoleamine 2,3-dioxygenase and inducible nitric oxide synthase expression in murine placenta following *Toxoplasma gondii* infection. *Int J Parasitol*. 2008; 38:249–258. [PubMed: 17822706]
113. Sullivan GM, Mann JJ, Oquendo MA, Lo ES, Cooper TB, Gorman JM. Low cerebrospinal fluid transthyretin levels in depression: correlations with suicidal ideation and low serotonin function. *Biological psychiatry*. 2006; 60:500–506. [PubMed: 16487493]
114. Oades RD, Lasky-Su J, Christiansen H, Faraone SV, Sonuga-Barke EJ, Banaschewski T, Chen W, Anney RJ, Buitelaar JK, Ebstein RP, Franke B, Gill M, Miranda A, Roeyers H, Rothenberger A, Sergeant JA, Steinhausen HC, Taylor EA, Thompson M, Asherson P. The influence of serotonin- and other genes on impulsive behavioral aggression and cognitive impulsivity in children with attention-deficit/hyperactivity disorder (ADHD): Findings from a family-based association test (FBAT) analysis. *Behav Brain Funct*. 2008; 4:48. [PubMed: 18937842]
115. Chugani DC, Muzik O, Behen M, Rothermel R, Janisse JJ, Lee J, Chugani HT. Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Ann Neurol*. 1999; 45:287–295. [PubMed: 10072042]

Highlights

- Maternal diet and metabolic status have a long-term impact on offspring behavior.
- Obesity is associated with increased inflammatory cytokines, nutrients, and hormones.
- Increases in circulating factors impact the environment of the developing offspring.
- Offspring exposed to maternal obesity during development are at increased risk for mental health disorders.