Published in final edited form as:

J Am Chem Soc. 2012 October 17; 134(41): 16933–16936. doi:10.1021/ja3070528.

A Transition-State Interaction Shifts Nucleobase Ionization Toward Neutrality to Facilitate Small Ribozyme Catalysis

Joseph A. Liberman^{†,‡}, Man Guo^{§,‡,⊥}, Jermaine L. Jenkins[†], Jolanta Krucinska^{†,||}, Yuanyuan Chen^{§,#}, Paul R. Carey^{§,*}, and Joseph E. Wedekind^{†,*}

[†]Department of Biochemistry and Biophysics, Center for RNA Biology, University of Rochester School of Medicine & Dentistry, 601 Elmwood Avenue, Box 712, Rochester, New York 14642 USA

[§]Department of Biochemistry, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio 44106 USA

Abstract

One mechanism by which ribozymes can accelerate biological reactions is by adopting folds that favorably perturb nucleobase ionization. Herein we used Raman crystallography to directly measure pK_a values for the Ade38 N1-imino group of a hairpin ribozyme in distinct conformational states. A transition-state analogue gave a pK_a value of 6.27 ± 0.05 , which agrees strikingly well with values measured by pH-rate analyses. To identify the chemical attributes that contribute to the shifted pK_a we determined crystal structures of hairpin ribozyme variants containing single-atom substitutions at the active site and measured their respective Ade38 N1 pK_a values. This approach led to the identification of a single interaction in the transition-state conformation that elevates the base $pK_a > 0.8$ log units relative to the precatalytic state. The agreement of the microscopic and macroscopic pK_a values and the accompanying structural analysis support a mechanism in which Ade38 N1(H)⁺ functions as a general acid in phosphodiester bond cleavage. Overall the results quantify the contribution of a single electrostatic interaction to base ionization, which has broad relevance for understanding how RNA structure can control chemical reactivity.

RNA plays a central role in several key biocatalytic reactions including non-coding RNA processing, protein translation, and RNA splicing^{1–6}. Distinct strategies identified in the protein world have been posited to account for this catalysis, including electrostatic complementation and general acid-base catalysis^{7–10}. These strategies require base ionization, which necessitates the generation of an imine with a pK_a near neutrality¹⁰. However, unlike proteins, there are no chemical groups in RNA that ionize near

*Corresponding Author: prc5@case.edu; joseph.wedekind@rochester.edu.

||Present Addresses

Brae Burn, Glastonbury CT 06033

⊥Bayer Health Care, Warrendale PA 15086

#Case Western Reserve University, Dept. Pharmacology, Cleveland OH USA 44106

‡Author Contributions

These authors contributed equally.

ASSOCIATED CONTENT

Supporting Information. Methods, the diffraction table, and figures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

The authors declare no competing financial interest.

Dedication

This work is dedicated to Professor David B. McKay in honor of his 67th birthday.

physiological pH, requiring the evolution of microenvironments that favorably perturb pK_a values¹¹. Although methods have been developed for measuring nucleotide base pK_a values in a ribozyme, no study has quantified how specific stereochemical interactions modulate pK_a shifting¹². These knowledge gaps hinder efforts to model RNA structure-function relationships, including rigorous computational treatment of folding and catalysis¹³. Here, we used Raman crystallography, X-ray diffraction, and mutagenesis to pinpoint how specific functional groups influence the pK_a of an essential imino group in the active site of an RNA enzyme known as the hairpin ribozyme (HPRZ), which is a model system for studying the architecture and function of RNA-mediated phosphoryl transfer.

The HPRZ is a member of the autolytic, small-ribozyme class⁸. It is derived from the negative-polarity strand of the satellite tobacco ringspot virus where it processes concatenated transcripts to unit length¹⁴. Like other small ribozymes, its chemical mechanism involves nucleophilic attack of the O2' group from position -1 upon the scissile phosphorus at position +1 (Figure 1A). Unlike larger selfsplicing ribozymes, the HPRZ does not require divalent ions for activity⁸, making it ideal for investigating nucleobase-assisted catalysis. Indeed, the HPRZ pH-rate profile shows an apparent pK_a of 6.2 ± 0.2 , consistent with a single ionizable group^{8,15}.

A synthetic variant of the HPRZ was developed in our lab for single-atom probing and structure-function investigations (Figures 1B and S1, Supporting Information). Crystallographic analyses of this minimal construct in the context of precatalytic and transition-state analogues revealed Gua8 and Ade38 adjacent to the scissile bond (Figures 1B and 1C)¹⁶. A functional analysis in which Gua8 was made abasic caused an 850-fold reduction in cleavage without altering the pH-rate profile¹⁷. By contrast, abasic replacement of Ade38 caused a 14,000-fold activity loss and a three log unit shift of the apparent pK_a toward the basic¹⁵. The subtle exchange of Ade38 for N1-deaza-Ade also resulted in complete loss of detectable activity¹⁸, whereas similar active-site modifications at nearby Ade9 and Ade10 had modest effects¹⁸. These data indicate that the N1 imino of Ade38 has a critical role in catalysis, and an elevated N1 pK_a was inferred.

Because the Ade38 protonation state figures prominently in any proposed HPRZ mechanism, we previously used Raman crystallography to measure directly the N1 pK_a of Ade38 in the context of an inert precatalytic analog (PCA) (Figure 1C). The results revealed a pK_a of 5.46 ± 0.04 ¹⁹. Although this value is elevated relative to AMP control measurements of 3.68 ± 0.06 , it did not agree with the apparent reaction pK_a of 6.2 ± 0.2 , which reports on transition-state proton transfer⁹. We hypothesized that an active-site conformation closer to the transition state would fine tune the Ade38 N1 pK_a and bring the microscopic pK_a into better agreement with the apparent pK_a of the reaction. To test this possibility, we utilized a transition-state analogue (TSA) to restrain the non-bridging oxygens of the scissile bond into a geometry consistent with the proposed transition state^{20,21} (Figure 1D). Using Raman crystallography we measured a pK_a of 6.27 ± 0.05 for N1 of Ade38 in the context of the TSA, which represents a 0.81 log unit increase over the PCA value (Figure 1E and Table 1). This microscopic pK_a is identical to the apparent pK_a of the reaction¹⁵, thereby providing direct evidence that the N1 imino of Ade38 is the source of rate-limiting proton transfer.

To understand the molecular basis for the ~1 log unit difference between the PCA and TSA ionization constants, we compared their corresponding structures, which were derived previously from crystals of the same space group as those used for Raman measurements in Figure 1E. Two main differences were apparent in the active site. Whereas the TSA exhibits hydrogen bonds between the *pro*-R_p oxygen equivalent and the N6 amine of Ade38, as well as the O5' leaving group and the Ade38 N1 imine, these interactions are missing in the PCA

(Figure 1D versus 1C). By contrast, both active sites showed interactions between one non-bridging oxygen of the scissile bond and the N2 amino group of Gua8. To determine the importance of hydrogen bonding between the *pro*-R_p oxygen of the scissile bond and the N6 group of Ade38 for p*K*_a shifting, we substituted a purine nucleotide at position 38 that is devoid of the exocyclic amine, and the N1 p*K*_a was measured. The results revealed a PCA with a p*K*_a value of 5.00 ± 0.06 (Table 1 and Figure S2A). By contrast, the control p*K*_a for 2'-deoxynebularine in which N6 is a hydrogen, was 2.92 ± 0.04 (Figure S3), indicating a 2.08 log unit shift toward the basic in the context of the HPRZ active site. Application of the latter shift to AMP gave a p*K*_a of 5.76, which is comparable to the wildtype PCA p*K*_a (Table 1). To assess how the Ade38Pur substitution influences the active site geometry, we determined its crystal structure (Table S1), which shows an active site conformation similar to wildtype (Figure 2A). The phosphate group at the scissile bond adopted two conformations; one is similar to wildtype, but the other is rotated such that the *pro*-R_p oxygen cannot hydrogen bond with N2 of Gua8. Neither of these conformations supports a hydrogen bond between the O3' and N1 of position 38, as seen for the wildtype.

We then asked how loss of the N6 amine at position 38 would affect its N1 p*K*_a in the context of the TSA. The Ade38Pur variant produced a p*K*_a of 4.88 ± 0.06 (Table 1 and Figure S2B), representing a 2.0 log unit shift toward the basic relative to wildtype. When this shift is applied relative to the controls, the Ade38 N1 p*K*_a is 5.66, which is significantly less than the p*K*_a of 6.27 ± 0.05 measured for the wildtype TSA (Table 1). To understand how the Ade38Pur variant influences the HPRZ active site in the context of the TSA, we determined its crystal structure (Table S1). The results showed that the *pro*-R_p oxygen of the scissile bond alters its conformation to be within hydrogen bonding distance of the N1 and O6 groups of Gua8 (Figure 2B). This represents a significant loss of scissile bond localization, and documents the extent to which the interaction between the *pro*-R_p oxygen and the N6 amine of Ade38 contributes to imine p*K*_a shifting. Our p*K*_a measurements and structural results are consistent with reports of Ade38Pur activity loss ranging from 10² to 10³-fold^{15,22}.

We next asked whether the N2 group of Gua8 influences Ade38 ionization. Structures of the wildtype HPRZ indicated that hydrogen bonding occurs between the exocyclic amine of Gua8 and the scissile bond in both PCA and TSA conformations (Figures 1C and 1D). To assess the importance of these interactions we replaced Gua8 with inosine, which lacks an exocyclic amine. In the context of the PCA, the N1 p*K*_a for Ade38 was 5.36 ± 0.05 (Figure S2C), which is nearly identical to wildtype (Table 1). A prior PCA structural comparison of the Gua8Ino variant to wildtype revealed only minor structural changes localized to the scissile bond¹⁶ (Figure 2C) even though the hydrogen bond is absent between the *pro*-R_p oxygen and the N2 amine of Gua8.

By contrast, the Gua8Ino substitution in the context of the TSA had a significant impact on Ade38 ionization via an indirect effect that repositioned the scissile phosphate. Our results revealed a p*K*_a of 5.45 ± 0.09 , which represents a 0.82 log unit decrease compared to wildtype (Figure S2D and Table 1). We determined the crystal structure of this variant in the context of the TSA (Table S1) and compared it to wildtype. The superposition indicates very subtle changes in the overall structure but a significant change in the hydrogen-bonding pattern to the scissile bond. In particular, the *pro*-S_p oxygen – unable to form a hydrogen bond to N2 of Ino8 – forms a new hydrogen bond to N1 (Figure 2D). This change shifts the *pro*-R_p oxygen location so that it is unable to hydrogen bond with N6 of Ade38. Consequently, the O5' leaving group is misoriented so that it does not interact readily with the Ade38 imine, consistent with the >10-fold loss in *k*_{cat} for the Gua8Ino variant²³.

Here we have shown that the microscopic pK_a of Ade38 N1, in the context of a transition state analogue, is identical to the apparent pK_a measured for the cleavage reaction for the most active, four-way-helical junction form of the HPRZ¹⁵. This agreement provides strong evidence that the N1 imine of Ade38 is involved in rate-limiting proton transfer. Independent evidence that the protonated form of Ade38 is operative in catalysis is supported by substitution of the base with 2-fluoro adenosine ($pK_a < 1$), which significantly reduces activity²⁴. Simulated pH-rate profiles that assumed a pK_a -shifted Ade38 and an unperturbed Gua8 pK_a appeared similar to experimental pH-rate profiles, supporting the plausibility of general acid/base catalysis¹⁰. Herein, we also visualized HPRZ active sites that retained functional characteristics of the overall reaction, thereby allowing us to draw stereochemical inferences about the mechanism. Significantly, TSA crystal structures containing either vanadate^{20,21} or a 2',5' linkage at the scissile bond²⁰ – identical to that employed here – do not show water localized at the N1 imine of Ade38, as proposed in a prior N1-imino pK_a analysis²⁵. As such, our observations strongly support a general acid role for Ade38. Similar roles have been ascribed previously to base Cyt75 of the hepatitis delta virus ribozyme²⁶ and Ade756 of the Varkud satellite ribozyme²⁷. We anticipate that when TSA structures become available for other small ribozymes, pK_a -shifting interactions will be exhibited that are similar to those observed here. Our experimental analysis also provides a quantitative result that should be useful for computational biologists seeking ways to benchmark pK_a prediction algorithms.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank R. Spitale, D. Perrin, N. Walter, P. Bevilacqua and D. Herschlag for helpful discussions. We thank the staff of SSRL for assistance with data collection. Portions of this research were carried out at the Stanford Synchrotron Radiation Laboratory (SSRL), a facility operated by Stanford University on behalf of the U.S. DOE. The SSRL Structural Molecular Biology Program is supported by the DOE, Office of Biological and Environmental Research, and by the NIH/NCRR, Biomedical Technology Program, and NIGMS.

Funding Sources

Financial support was provided in part by grants from the NIH to PRC (GM84120) and to JEW (GM63162).

REFERENCES

1. Reiner R, Ben-Asouli Y, Krilovetzky I, Jarrous N. *Genes Dev.* 2006; 20:1621. [PubMed: 16778078]
2. Guerrier-Takada C, Gardiner K, Marsh T, Pace N, Altman S. *Cell.* 1983; 35:849. [PubMed: 6197186]
3. Steitz TA, Moore PB. *Trends Biochem Sci.* 2003; 28:411. [PubMed: 12932729]
4. Noller HF, Hoffarth V, Zimniak L. *Science.* 1992; 256:1416. [PubMed: 1604315]
5. Valadkhan S, Mohammadi A, Jaladat Y, Geisler S. *Proc Natl Acad Sci U S A.* 2009; 106:11901. [PubMed: 19549866]
6. Toor N, Keating KS, Taylor SD, Pyle AM. *Science.* 2008; 320:77. [PubMed: 18388288]
7. Fedor MJ, Williamson JR. *Nat Rev Mol Cell Biol.* 2005; 6:399. [PubMed: 15956979]
8. Fedor MJ. *Annu Rev Biophys.* 2009; 38:271. [PubMed: 19416070]
9. Doudna JA, Lorsch JR. *Nat Struct Mol Biol.* 2005; 12:395. [PubMed: 15870731]
10. Bevilacqua PC. *Biochemistry.* 2003; 42:2259. [PubMed: 12600192]
11. Bevilacqua PC, Brown TS, Nakano S, Yajima R. *Biopolymers.* 2004; 73:90. [PubMed: 14691943]
12. Liberman JA, Wedekind JE. *Curr Opin Struct Biol.* 2011; 21:327. [PubMed: 21530235]
13. Moser A, Range K, York DM. *J Phys Chem B.* 2010; 114:13911. [PubMed: 20942500]

14. Forster AC, Symons RH. *Cell*. 1987; 49:211. [PubMed: 2436805]
15. Kuzmin YI, Da Costa CP, Cottrell JW, Fedor MJ. *J Mol Biol*. 2005; 349:989. [PubMed: 15907933]
16. Salter J, Krucinska J, Alam S, Grum-Tokars V, Wedekind JE. *Biochemistry*. 2006; 45:686. [PubMed: 16411744]
17. Kuzmin YI, Da Costa CP, Fedor MJ. *J Mol Biol*. 2004; 340:233. [PubMed: 15201049]
18. Spitale RC, Volpini R, Heller MG, Krucinska J, Cristalli G, Wedekind JE. *J Am Chem Soc*. 2009; 131:6093. [PubMed: 19354216]
19. Guo M, Spitale RC, Volpini R, Krucinska J, Cristalli G, Carey PR, Wedekind JE. *J Am Chem Soc*. 2009; 131:12908. [PubMed: 19702306]
20. Torelli AT, Krucinska J, Wedekind JE. *RNA*. 2007; 13:1052. [PubMed: 17488874]
21. Rupert PB, Massey AP, Sigurdsson ST, Ferre-D'Amare AR. *Science*. 2002; 298:1421. [PubMed: 12376595]
22. Grasby JA, Mersmann K, Singh M, Gait MJ. *Biochemistry*. 1995; 34:4068. [PubMed: 7535099]
23. Pinard R, Hampel KJ, Heckman JE, Lambert D, Chan PA, Major F, Burke JM. *EMBO J*. 2001; 20:6434. [PubMed: 11707414]
24. Suydam IT, Levandoski SD, Strobel SA. *Biochemistry*. 2010; 49:3723. [PubMed: 20373826]
25. Cottrell JW, Scott LG, Fedor MJ. *J Biol Chem*. 2011; 286:17658. [PubMed: 21454684]
26. Gong B, Chen JH, Chase E, Chadalavada DM, Yajima R, Golden BL, Bevilacqua PC, Carey PR. *J Am Chem Soc*. 2007; 129:13335. [PubMed: 17924627]
27. Wilson TJ, Li NS, Lu J, Frederiksen JK, Piccirilli JA, Lilley DM. *Proc Natl Acad Sci U S A*. 2010; 107:11751. [PubMed: 20547881]

Figure 1.

The chemical reaction of small ribozymes, HPRZ active site organization, and directly measured microscopic pK_a values for the Ade38 N1 imine. **(A)** The HPRZ reaction entails deprotonation of the Ade-1 2'-OH and attack on the neighboring phosphate of Gua+1. The reaction proceeds via a phosphorane transition state that is favored by protonation of the O5' leaving group and electrostatic stabilization of the negatively charged phosphorane. The products are a cyclic-2',3' phosphodiester and a free 5'-hydroxyl. **(B)** The three dimensional structure of a 'minimal' synthetic hairpin ribozyme (PDB: 2OUE) with Ade-1 and Gua+1 in red; Ade38 is teal and Gua8 is maroon. **(C)** Ball-and-stick diagram of the active site of the HPRZ precatalytic analogue (PCA) reported previously in the presence of a 2'-O-Me cap that blocks activity¹⁶ (PDB ID 2OUE). Here and elsewhere a black arrow indicates the scissile bond. **(D)** The active site of a transition-state analogue (TSA) of the HPRZ reported previously with a 3'-deoxy-2',5'-phosphodiester linkage between Ade-1 and Gua+1²⁰ (PDB ID 2P7F). **(E)** Raman crystallography results showing the area under the peak at 734 cm^{-1} , which reports on protonation of Ade38 N1, between pH 4.5 and 8.5. The red line shows the titration of a PCA HPRZ described previously¹⁹; the blue line shows the TSA Raman crystallography titration from the investigation herein.

Figure 2. Superpositions of HPRZ active sites comparing wildtype structures to Ade38 or Gua8 variants in the context of PCA or TSA conformations. Structures of the base variants are depicted in red, green and purple with the wildtype structure in blue; for clarity, the atom present in the wildtype but absent in the base variant is highlighted in red. Hydrogen bonds (dashed lines) are indicated only for the Ade38 and Gua8 variants. **(a)** The PCA with an Ade38Pur substitution exhibits two scissile bond conformations. **(b)** The TSA with an Ade38Pur substitution. **(c)** The PCA with a Gua8Ino substitution reported previously¹⁶ (PDB ID IZFT). **(d)** The TSA with a Gua8Ino substitution.

Table 1Direct Measurements of Ade38 N1 pK_a Values

Sample	PCA	TSA
wildtype	5.46 ± 0.05	6.27 ± 0.05
Ade38Pur	5.00 ± 0.06 (measured) ^a	4.88 ± 0.06 (measured) ^a
	5.76 (corrected) ^b	5.66 (corrected) ^b
Gua8Ino	5.36 ± 0.08	5.45 ± 0.09

^a Measured values are derived directly from Raman experiments.

^b The corrected value is obtained by adding 0.78 log units to the pK_a values of the respective Ade38Pur PCA and TSA variants. The value of 0.78 represents the difference between the solution pK_a values measured by Raman spectroscopy for the respective AMP control, with a value of 3.68 ± 0.06 ¹⁹, and the 2'-deoxynebularine control with a value of 2.92 ± 0.04 (Figure S3, Supporting Information).