Published in final edited form as:

Nature. ; 486(7401): 113–117. doi:10.1038/nature10958.

Preferential electrical coupling regulates lineage-dependent microcircuit assembly in the neocortex

Yong-Chun Yu^{1,6}, Shuijin He^{2,6}, She Chen², Yinghui Fu¹, Keith N. Brown^{2,3}, Xing-Hua Yao¹, Jian Ma¹, Peng Gao^{2,3}, Gina E. Sosinsky⁴, Kun Huang⁵, and Song-Hai Shi^{2,3}

¹Institute of Neurobiology, Institutes of Brain Science and State Key Laboratory of Medical Neurobiology, Fudan University, 138 Yixueyuan Road, Shanghai, 200032, China

²Developmental Biology Program, Memorial Sloan-Kettering Cancer Centre, 1275 York Avenue, New York, NY 10065

³Neuroscience Graduate Program, Weill Cornell Medical College, 1230 York Avenue, New York, NY 10065

⁴National Centre for Microscopy and Imaging Research and Department of Neurosciences, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0608

⁵Department of Biomedical Informatics, Comprehensive Cancer Center Biomedical Informatics Shared Resource, The Ohio State University, 333 West 10th Avenue, Columbus, OH 43210, USA

Abstract

Radial glial cells are the primary neural progenitor cells in the developing neocortex ¹. Consecutive asymmetric divisions of individual radial glial progenitor cells produce a number of sister excitatory neurons that migrate along the elongated radial glial fibre, resulting in the formation of ontogenetic columns ^{2–4}. Moreover, sister excitatory neurons in ontogenetic columns preferentially develop specific chemical synapses with each other rather than with nearby non-siblings ⁵. While these findings provide crucial insights into the emergence of functional columns in the neocortex, little is known about the basis for this lineage-dependent assembly of excitatory neuron microcircuits with single-cell resolution. Here we show that transient electrical coupling between radially aligned sister excitatory neurons regulates the subsequent formation of specific chemical synapses in the neocortex. Multiple-electrode whole-cell recordings revealed that sister excitatory neurons preferentially form strong electrical coupling with each other rather than with adjacent non-sister excitatory neurons during early postnatal stages. This coupling allows selective electrical communication between sister excitatory neurons, promoting their action potential generation and synchronous firing. Interestingly, while this electrical communication largely disappears prior to the appearance of chemical synapses, its blockade impairs the subsequent formation of specific chemical synapses between sister excitatory neurons in ontogenetic columns. These results suggest a strong link between a lineage-dependent transient electrical coupling and the assembly of precise excitatory neuron microcircuits in the neocortex.

Sister excitatory neurons in individual ontogenetic columns preferentially develop specific chemical synapses with each other instead of with adjacent non-sister excitatory neurons ⁵. Given the near complete overlap of the dendritic fields of neighbouring excitatory neurons, how this lineage-dependent assembly of precise columnar microcircuits at the individual cell level is controlled remains an outstanding question. Gap junction-mediated neuronal

*Correspondence and request for materials should be addressed to Y.-C.Y. (ycyu@fudan.edu.cn), S. H. (hes@mskcc.org) or S.-H.S. (shis@mskcc.org).

⁶These authors contributed equally to this work

communication has been suggested to be involved in the formation of local connectivity in the developing neocortex^{6–8}, even though direct evidence of electrically coupled neocortical excitatory neurons at early developmental stages is lacking⁹. In this study, we investigated whether gap junction-mediated electrical coupling exists between sister excitatory neurons in ontogenetic columns and if so, whether it regulates the preferential formation of chemical synapses between them.

Gap junctions are small cell-to-cell contacts containing membrane-embedded channels that mediate intercellular adhesion and exchange of small molecules (typically less than 1 kDa), including low molecular weight dyes and ions that can be detected experimentally^{10, 11}. Previous dye injection experiments have suggested the presence of gap junctions between progenitor cells in the embryonic neocortex^{2, 12–14} and between neurons in the neonatal neocortex^{7, 15}. While these studies have provided important insights, the accuracy of dye coupling in revealing gap junctions has been debated^{16–18}. To circumvent this issue and to quantitatively examine gap junction channel activity, we performed whole-cell patch clamp recording experiments to study gap junctions between sister excitatory neurons in ontogenetic columns. We prepared acute neocortical slices from postnatal mice (postnatal day 1 to 21, P1–P21) that received *in utero* intraventricular injection of low titre EGFP-expressing retrovirus at embryonic day 12 to 13 (E12–E13). As shown in previous studies^{2, 3, 5}, these injections led to reliable labelling of isolated ontogenetic columns of sister excitatory neurons in the developing neocortex (Fig. 1a, top).

Guided by infra-red differential interference contrast (IR-DIC) and epi-fluorescence illumination, we simultaneously recorded from two radially aligned EGFP-expressing sister excitatory neurons (Fig. 1a, green). We identified excitatory neurons on the basis of their morphological characteristics including a pyramid-shaped cell body and a long apical dendrite. After the recordings were established, we tested their electrical coupling under current-clamp (Fig. 1a, bottom) or voltage-clamp (Supplementary Fig. 1a) conditions. Under current-clamp conditions, hyperpolarization (V_1) of one of the neurons (neuron 1, driver) by current injection (I_1) produced a simultaneous hyperpolarization (V_2) of the non-injected neuron (neuron 2, receiver) (Fig. 1a, bottom). Similarly, depolarizing driver neuron 1 produced a simultaneous depolarization of receiver neuron 2. As expected for electrotonic propagation, voltage deflections recorded in the non-injected receiver neuron (red, Fig. 1a, bottom) had a smaller amplitude and slower time course than those in the injected driver neuron (black, Fig. 1a, bottom). In all cases, electrical transmission between the two neurons was found to be reciprocal (Fig. 1a, bottom). Similar bidirectional electrotonic propagation was recorded under voltage-clamp conditions (Supplementary Fig. 1a). To confirm that the electrical coupling between sister excitatory neurons is mediated by gap junctions, we exposed electrically coupled sister excitatory neuron pairs to the gap junction blocker meclofenamic acid (MFA, 100 μ M), or glutamate receptor antagonists 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, 10 μ M) and 2-carboxypiperazin-4-propyl-1phosphonic acid (CPP, 20 μ M), or γ -aminobutyric acid (GABA)-A receptor antagonist bicuculline methiodide (BMI, 10 μ M). While CNQX, CPP and BMI had no effect (Supplementary Fig. 1b), MFA completely eliminated electrical transmission in all cases ($n=6$). Moreover, electrical transmission recovered partially after washout of MFA (Fig. 1b). Similar results were obtained with another gap junction blocker, 18- α -Glycyrrhetinic acid (100 μ M, data not shown).

The frequency of observing gap junction-mediated electrical coupling between sister excitatory neurons in ontogenetic columns at P1 and P2 was 38.9% (14 out of 36 pairs tested) and 35.3% (12 out of 34 pairs tested), respectively (Fig. 1c). As development proceeded, this coupling frequency progressively decreased to 20–25% at P3–P4 (9 out of 39 pairs tested at P3, and 9 out of 35 pairs tested at P4) and to ~10% at P5–P6 (5 out of 48

pairs tested) (Fig. 1c). After P6, electrical coupling between sister excitatory neurons in ontogenetic columns was rare; only 3 out of 193 pairs were coupled (Fig. 1c), consistent with prior observations of very sparse electrical coupling among mature excitatory neurons in the neocortex^{18–21}. It is worth noting that no chemical synapses were detected between sister excitatory neurons at early postnatal stages (P1–P6), as previously reported⁵.

To test whether sister excitatory neurons preferentially form gap junctions with each other, we performed quadruple whole-cell recordings on two EGFP-expressing sister excitatory neurons in individual ontogenetic columns and two non-EGFP-expressing excitatory neurons adjacent to the EGFP-expressing neurons on the same side in the developing neocortex at P1–P6, when electrical coupling is prominent (Fig. 2a). We targeted non-EGFP-expressing neurons with radially aligned pyramid-shaped cell bodies because they were most likely to be excitatory neurons and the progeny of different radial glial progenitor cells. Thereby, they served as adjacent non-sibling excitatory neuron controls. Once all four recordings were established, hyperpolarizing and depolarizing currents were injected sequentially into one of the four neurons and the voltage changes were monitored in all four neurons to probe gap junction-mediated electrical coupling. As shown in Figure 2b, when EGFP-expressing neuron 1 was hyperpolarized or depolarized, only its sister neuron 3 exhibited simultaneous hyperpolarization or depolarization, despite the near complete overlap of the neurite arbours of adjacent neurons 3 and 4 (Fig. 2c). Similarly, when EGFP-expressing neuron 3 was hyperpolarized or depolarized, only its sister neuron 1 showed simultaneous hyperpolarization or depolarization (Fig. 2b). Hyperpolarizing or depolarizing neurons 2 or 4 failed to trigger simultaneous voltage deflection in all three remaining neurons (Fig. 2b).

We analyzed a total of 174 pairs of radially aligned EGFP-expressing sister excitatory neurons and their neighbouring non-sibling excitatory neurons at P1–P6 (Fig. 2d). Of these, 28.2% (49 out of 174 pairs) of sister excitatory neuron pairs in an ontogenetic column were electrically coupled. In contrast, only 2.6% (8 out of 303 pairs) of radially situated non-sister excitatory neuron pairs (one EGFP-expressing and one non-EGFP-expressing) were electrically coupled (Fig. 2d). In addition, only 1.6% (2 out of 129 pairs) of similarly situated non-EGFP expressing excitatory neuron pairs and only 3.0% (9 out of 303) of nearby EGFP-expressing and non-EGFP-expressing excitatory neuron pairs were coupled (Fig. 2d). These results clearly demonstrate that sister excitatory neurons in ontogenetic columns have a strong preference to form gap junction-mediated electrical coupling with each other instead of with adjacent non-sister excitatory neurons, and that the frequency of coupling between non-sister excitatory neurons is low.

We also compared the coupling coefficient between coupled sister excitatory neuron pairs and rarely coupled non-sister excitatory neuron pairs. The coupling coefficient, estimated as the ratio of the amplitude of the low-frequency voltage change in the receiver neuron to that in the driver neuron, reflects the strength of the electrical coupling. We found that the coupling coefficient between coupled sister excitatory neurons in ontogenetic columns was substantially higher than that between coupled non-sister excitatory neurons (sisters: $5.7 \pm 0.7\%$, $n=74$; non-sisters: $1.2 \pm 0.3\%$; $n=24$; $p < 1e-8$; Fig. 2e), suggesting that the electrical coupling between sister excitatory neurons is much stronger than that between non-sisters. Furthermore, we observed a progressive decrease in the coupling coefficient of coupled sister excitatory neuron pairs as development proceeded (P1–P2, $7.4 \pm 1.2\%$, $n=36$; P3–P4, $4.4 \pm 0.7\%$, $n=32$; P5–P6, $2.3 \pm 0.6\%$, $n=6$; $p < 0.01$), while the coupling coefficient of rarely coupled non-sister pairs did not change significantly (P1–P2, $1.1 \pm 0.2\%$, $n=11$; P3–P4, $1.5 \pm 0.8\%$, $n=7$; P5–P6, $1.1 \pm 0.5\%$, $n=6$; Fig. 2f).

Having found that sister excitatory neurons in ontogenetic columns preferentially form strong gap junction-mediated electrical coupling with each other, we then examined the properties of this electrical transmission and tested whether this selective electrical coupling modulates the neuronal activity of sister excitatory neurons at neonatal stages. We injected a series of sinusoidal current waveforms of the same amplitude but different frequencies into one excitatory neuron and measured the response in its coupled sister excitatory neuron (Supplementary Fig. 2). We found that responses to low-frequency sine waves showed higher coupling coefficients and smaller phase lags than those triggered by high-frequency sine waves (Supplementary Fig. 2). These results suggest that the efficacy of signal transmission through electrical coupling between sister excitatory neurons in ontogenetic columns is frequency-dependent, which is similar to previous observations of electrical coupling between inhibitory interneurons in the more mature neocortex^{19, 20}

To determine whether this selective electrical coupling modulates neuronal activity, we injected sub-threshold depolarizing current pulses at the same or different times into two electrically coupled sister excitatory neurons in ontogenetic columns labelled by EGFP (Fig. 3a). In most cases, the asynchronous pulses did not generate an action potential in either neuron (Fig. 3a, arrowheads, and Fig. 3b, open bars; neuron 1, 0.1 ± 0.1 spike/pulse; neuron 2, 0.3 ± 0.2 spike/pulse). However, when the same current pulses were synchronously injected into both neurons, the two sister neurons reached action potential threshold and generated spikes (Fig. 3a, arrows, and Fig. 3b, filled bars; neuron 1, 1.5 ± 0.3 spike/pulse; neuron 2, 1.8 ± 0.4 spike/pulse). Similar observations were made in seven pairs of electrically coupled sister excitatory neurons, but not in non-coupled sister pairs (Fig. 3c, $p < 0.001$). These results showed that this selective electrical coupling can strongly facilitate the generation of action potentials in coupled sister excitatory neurons in ontogenetic columns.

We next determined whether this selective electrical coupling could facilitate synchronous spiking of radially aligned sister excitatory neurons in response to natural stimuli. First, we simulated natural activity by injecting two electrically coupled or non-coupled sister excitatory neurons with uncorrelated random current signals obtained by convolving Poisson trains of recorded spontaneous excitatory postsynaptic current waveforms^{20, 22} (I_1 and I_2 , Fig. 3d). Second, we directly recorded spontaneous sub-threshold activity of neurons in the neonatal neocortex and then injected two electrically coupled or non-coupled sister excitatory neurons with the recorded uncorrelated native neuronal activity (I_1 and I_2 , Fig. 3f). We analyzed the cross-correlogram of firing evoked in these sister excitatory neurons and found that there was a significant increase in firing centred at 0 milliseconds (ms) in coupled pairs compared with non-coupled ones (Fig. 3e, coupled, $n=4$; non-coupled, $n=5$; $p < 0.005$; and Fig. 3g, coupled, $n=10$; non-coupled, $n=9$; $p < 0.05$). These results suggest that gap-junction mediated electrical transmission between sister excitatory neurons in ontogenetic columns can effectively promote precise (within 1 ms) synchronous firing in response to uncorrelated neuronal activity.

It has long been postulated that correlated neuronal activity facilitates chemical synapse formation and neuronal circuit assembly^{23, 24}. Given that sister excitatory neurons in ontogenetic columns preferentially form strong gap junction-mediated electrical coupling, which promotes action potential generation and synchronous firing, this could be a mechanism that leads to preferential formation of chemical synapses between sister excitatory neurons in ontogenetic columns. We therefore tested whether the selective electrical coupling between sister excitatory neurons is required for preferential formation of chemical synapses within individual ontogenetic columns.

Gap junctions are composed of two juxtaposed hexameric hemi-channels consisting of Connexin proteins on two adjacent cells²⁵. There are ~20 genes encoding Connexin

proteins in rodents and they are often abbreviated as Cx plus the molecular mass of the protein²⁶. Among them, Cx26 and Cx43 have been shown to be abundantly expressed in the developing neocortex at embryonic and neonatal stages^{27, 28}. Consistent with this, we found that developing neurons in the neonatal neocortex expressed Cx26 (Supplementary Fig. 3a). Moreover, Cx26-positive puncta were present at the dendrodendritic and dendrosomatic contacts of radially aligned sister excitatory neurons (Supplementary Fig. 3 b–f and Movie 1). These results suggest that Cx26 is a major Connexin isoform that mediates neonatal electrical coupling between sister excitatory neurons in ontogenetic columns.

Previous studies have shown that mutation of a conserved Threonine (T135) in the third transmembrane helix of Cx26 to Alanine (A) (Cx26(T135A)) creates a dominant-negative closed gap junction channel without affecting the synthesis, assembly or trafficking of the channel²⁹. Mutant Cx26 also exerts a trans-dominant-negative effect on other Connexins³⁰, providing a useful tool for eliminating broad gap junction channel functionality²⁹. Notably, this point mutation should spare the gap junction-mediated adhesion required for proper radial migration of newly-generated excitatory neurons in the developing neocortex^{27, 31}. We engineered retroviruses expressing either wild type Cx26 or the dominant-negative closed channel mutant Cx26(T135A) together with EGFP using an internal ribosomal entry site (IRES) sequence and performed *in utero* intraventricular injection of retroviruses at E12–E13. We found that expression of wild type Cx26 had no discernible effect, but expression of Cx26(T135A) largely eliminated the electrical coupling between sister excitatory neurons at P1–P5 (Cx26: 26.4%, 14 out of 53 pairs; Cx26(T135A): 9.8%, 4 out of 41 pairs; Supplementary Fig. 4). More importantly, when we examined the chemical synapses formed between sister excitatory neurons at P10–P21, we found that only ~8.2% (4 out of 49) of sister pairs expressing the dominant-negative closed channel mutant Cx26(T135A) were connected with chemical synapses (Fig. 4, d–g), compared to 30–35% of sister pairs expressing either EGFP alone (12 out of 33 pairs tested) or wild type Cx26 (22 out of 63 pairs tested) (Fig. 4, a–c and g). Expression of Cx26(T135A) did not affect neocortical excitatory neuron migration²⁷ (Supplementary Fig. 5) or maturation (Supplementary Fig. 6). Taken together, these results suggest that the blockade of electrical communication between sister excitatory neurons impairs subsequent chemical synapse formation between them.

Transient electrical coupling occurs before the establishment of mature patterns of synaptic connectivity in many developing nervous systems, and in some cases, is crucial for the development of chemical synapses^{32–34}. Previously, dye coupling was observed in the developing neocortex^{2, 7, 12, 15}. In addition, it has been suggested that the formation of electrically coupled neuronal domains might help guide the emergence of chemically transmitting neuronal circuits^{6, 8}. However, electrically coupled neuronal pairs in the neocortex at early developmental stages have not been reported⁹. By performing dual and quadruple whole-cell recording experiments, which allow the detection of gap junction-mediated electrical coupling with high sensitivity and spatial precision^{10, 18}, we directly demonstrated the electrical coupling of excitatory neurons in the early postnatal neocortex. Moreover, we revealed that neocortical excitatory neurons show a high preference for forming strong electrical coupling with their sister excitatory neurons, but not with nearby non-sister excitatory neurons.

Furthermore, we found that strong electrical coupling between sister excitatory neurons in ontogenetic columns promotes their action potential generation and synchronous firing. Although previous studies suggested that electrical coupling is critical for the robust, spontaneous and synchronous electrical activity in the neonatal neocortex^{6, 35–38}, the precise function of electric coupling remains elusive. Our results showed that the electrical transmission between sister excitatory neurons in ontogenetic columns is required for the

development of precise chemical synapses between them. These findings provide clear evidence on the role of gap junctions in regulating precise neuronal circuit assembly in the neocortex. It will be interesting in the future to determine how electrical coupling regulates the formation of specific chemical synapses between sister excitatory neurons in ontogenetic columns at the molecular and cellular level.

Methods

Retrovirus production and *in utero* intraventricular injection

Replication-incompetent EGFP-expressing Moloney murine leukemia virus was produced from a stably transfected packaging cell line (293gp NIT-GFP; a kind gift from Dr. F. H. Gage, The Salk Institute) as previously published². Animals were maintained according to protocols approved by the Institutional Animal Care and Use Committee at the Memorial Sloan-Kettering Cancer Centre and Fudan University. *In utero* intraventricular injection was performed as previously described⁵. In brief, uterine horns of E12–E13 gestation stage pregnant CD-1 mice (Charles River Laboratories, MA) were exposed in a clean environment. Retrovirus (~1.0 μ L) with Fast green (2.5 mg/ml, Sigma) was injected into the embryonic cerebral ventricle through a bevelled, calibrated glass micropipette (Drummond Scientific, Broomall, PA). After injection, the peritoneal cavity was lavaged with ~10 mL warm phosphate buffered saline (PBS, pH 7.4) containing antibiotics, the uterine horns were replaced, and the wound was closed. Avian RCAS (Replication-Competent Δ SLV long terminal repeat with a Splice acceptor) retroviruses expressing EGFP, Cx26-IRES-EGFP or Cx26(T135A)-IRES-EGFP were generated as previously described³⁹. Similar *in utero* intraventricular injection of RCAS retroviruses was performed on the embryos of Nestin-TVA transgenic mice generated previously⁴⁰.

Slice preparation, recording and data analysis

Embryos that received retroviral injections were delivered naturally. Brains were removed at different postnatal days and acute cortical slices (200–300 μ m thick) were prepared in artificial cerebrospinal fluid (ACSF) containing (in mM) 126 NaCl, 3 KCl, 1.25 KH_2PO_4 ; 1.3 MgSO_4 , 3.2 CaCl_2 , 26 NaHCO_3 , and 10 glucose, bubbled with 95% O_2 /5% CO_2) with a Vibratome (Leica Microsystems) at 4°C. Slices were allowed to recover in an interface chamber at 35°C for at least one hour and then kept at room temperature before being transferred to a recording chamber containing ACSF at 34°C. An infrared-differential interference contrast (IR-DIC) microscope (Olympus) equipped with epi-fluorescence illumination, a CCD camera, and two water immersion lenses (10X and 60X) were used to visualize and target recording electrodes to EGFP-expressing sister cells in ontogenetic columns and their nearby control cells. Glass recording electrodes (20–30 M Ω resistance) were filled with an intracellular solution consisting of (in mM) 130 K-gluconate, 6 KCl, 2 MgCl_2 , 0.2 EGTA, 10 HEPES, 2.5 Na_2ATP , 0.5 Na_2GTP , 10 K-phosphocreatine and 0.3% Alexa Fluor 568 hydrazide (Invitrogen) (pH 7.25 and 295 mOsm/kg). In all dual and quadruple recordings, electrical coupling was assessed by injecting currents to trigger hyperpolarization or depolarization under current-clamp mode or by voltage steps under voltage-clamp mode. In some experiments, meclofenamic acid (100 μ M) or 18- α -Glycyrrhetic acid (100 μ M) (Sigma) were added to the bath to block gap junctions, and BMI (10 μ M), CPP (20 μ M), and CNQX (10 μ M) (Tocris Biosciences) were used to block GABA-A, NMDA, and AMPA receptors, respectively. Native neuronal activity stimuli were simulated as previously described²².

Chemical connections between neuron pairs were assessed by injecting current to evoke action potentials in one of the neurons kept in current-clamp while testing for postsynaptic responses in other neurons during voltage-clamp recording at –70 mV. Glass recording

electrodes (8–12 M Ω resistance) were filled with an intracellular solution consisting of (in mM) 130 K-gluconate, 6 KCl, 2 MgCl₂, 0.2 EGTA, 10 HEPES, 2.5 Na₂ATP, 0.5 Na₂GTP, 10 K-phosphocreatine and 0.5% Neurobiotin. For every possible pair, connections were tested in both directions in at least 20 trials, with both single and trains of action potentials being generated in each presynaptic neuron. Recordings were collected and analyzed using Axopatch 700B amplifier and pCLAMP10 software (Molecular Devices). In whole-cell patch clamp recording experiments, slices were fixed in 4% paraformaldehyde (PFA) in PBS (pH 7.4) after the recordings were completed, the morphology of recorded neurons loaded with Alexa Fluor 568 hydrazide through the recording pipette was visualized using an Olympus FV1000 confocal laser scanning microscope. Z-series images were taken at 1- μ m steps and analyzed using FluoView (Olympus), Neurolucida (MicroBrightField Inc.) and Photoshop (Adobe Systems). In some recording experiments, Neurobiotin was later visualized with Alexa Fluor 647- or Alexa Fluor 568- conjugated streptavidin (Invitrogen).

For Cx26 and TuJ1 immunohistochemistry, after intracardial perfusion with cold PBS (pH 7.4) and 4% PFA in PBS (pH 7.4), the brains were removed and embedded in tissue freezing medium O.C.T. (Electron Microscopy Sciences) after sucrose treatment. Coronal sections (20 μ m) were prepared using a cryostat (Leica Microsystems) and incubated for 2 hours at room temperature in a blocking solution (10% normal goat serum and 0.1% Triton X-100 in PBS), followed by incubation with the primary antibodies including rabbit anti-Cx26 antibody (Invitrogen) and mouse anti-TuJ1 antibody (Covance) for 2 days at 4°C. Sections were then washed in 0.1% Triton X-100 in PBS and incubated with the appropriate secondary antibodies overnight at 4°C. For three-dimensional reconstruction, Z-series images were taken at 0.1- μ m steps using 100X objective lens (Olympus) in an Olympus confocal laser scanning microscope and analyzed using Neurolucida (MicroBrightField Inc.) and Imaris (Andor Technology).

Normalized cross-correlogram of firing patterns was analyzed as previously described⁴¹. In brief, the number of neuron 1 that fired within a time interval $[n\Delta t, (n+1)\Delta t]$ from spikes fired by neuron 2 was calculated (y_n = counts/bin; bin width, Δt = 1 msec). The cross-correlogram y_n was normalized in standard scores $[Z = (y_n - \gamma_E)/s_y]$, with $\gamma_E = f_1 * f_2 * T * \Delta t$ ($f_{1,2}$, average firing rate of neurons 1 and 2; T , recording time) and s_y , standard deviation of y_n . Data were presented as mean $\pm$ s.e.m. and statistical differences were determined using nonparametric Mann-Whitney-Wilcoxon and Kruskal-Wallis tests. Peaks in the cross-correlogram were considered significant if individual bins exceeded expected value by 3 standard deviations (Z score > 3).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Drs. C.I. Bargmann, Y. Dan, and A.L. Joyner for comments on the manuscript, Dr. F.H. Gage for 293gp NIT-GFP package cell line, Dr. E.C. Holland for Nestin-TVA transgenic mice, and members of the Shi and Yu laboratories for valuable input. This work was supported by grants from the National Institutes of Health (R01DA024681 and R21NS072483 to S.-H.S, R21MH083624 to S.-H.S and K.H., and R01GM065947 to G.E.S.), the McKnight Foundation, the March of Dimes Foundation (to S.-H.S), the Pujiang Talent Project of the Shanghai Science and Technology Committee (10PJ1400700), the Natural Science Foundation of China (31070947 and 30821002), the Foundation of Ministry of Education of China (20100071120061) and the National Key Scientific Program of China (2012CB966303) (to Y.-C.Y.).

References

1. Kriegstein A, Alvarez-Buylla A. The glial nature of embryonic and adult neural stem cells. *Annu Rev Neurosci.* 2009; 32:149–84. [PubMed: 19555289]
2. Noctor SC, Flint AC, Weissman TA, Dammerman RS, Kriegstein AR. Neurons derived from radial glial cells establish radial units in neocortex. *Nature.* 2001; 409:714–20. [PubMed: 11217860]
3. Noctor SC, Martinez-Cerdeno V, Ivic L, Kriegstein AR. Cortical neurons arise in symmetric and asymmetric division zones and migrate through specific phases. *Nat Neurosci.* 2004; 7:136–44. [PubMed: 14703572]
4. Rakic P. Specification of cerebral cortical areas. *Science.* 1988; 241:170–6. [PubMed: 3291116]
5. Yu YC, Bultje RS, Wang X, Shi SH. Specific synapses develop preferentially among sister excitatory neurons in the neocortex. *Nature.* 2009; 458:501–4. [PubMed: 19204731]
6. Yuste R, Nelson DA, Rubin WW, Katz LC. Neuronal domains in developing neocortex: mechanisms of coactivation. *Neuron.* 1995; 14:7–17. [PubMed: 7826643]
7. Peinado A, Yuste R, Katz LC. Extensive dye coupling between rat neocortical neurons during the period of circuit formation. *Neuron.* 1993; 10:103–14. [PubMed: 8427699]
8. Yuste R, Peinado A, Katz LC. Neuronal domains in developing neocortex. *Science.* 1992; 257:665–9. [PubMed: 1496379]
9. Montoro RJ, Yuste R. Gap junctions in developing neocortex: a review. *Brain Res Brain Res Rev.* 2004; 47:216–26. [PubMed: 15572173]
10. Connors BW, Long MA. Electrical synapses in the mammalian brain. *Annu Rev Neurosci.* 2004; 27:393–418. [PubMed: 15217338]
11. Bennett MV, Zukin RS. Electrical coupling and neuronal synchronization in the Mammalian brain. *Neuron.* 2004; 41:495–511. [PubMed: 14980200]
12. Lo Turco JJ, Kriegstein AR. Clusters of coupled neuroblasts in embryonic neocortex. *Science.* 1991; 252:563–6. [PubMed: 1850552]
13. Bittman K, Owens DF, Kriegstein AR, LoTurco JJ. Cell coupling and uncoupling in the ventricular zone of developing neocortex. *J Neurosci.* 1997; 17:7037–44. [PubMed: 9278539]
14. Duval N, et al. Cell coupling and Cx43 expression in embryonic mouse neural progenitor cells. *J Cell Sci.* 2002; 115:3241–51. [PubMed: 12140256]
15. Connors BW, Benardo LS, Prince DA. Coupling between neurons of the developing rat neocortex. *J Neurosci.* 1983; 3:773–82. [PubMed: 6187902]
16. Gutnick MJ, Lobel-Yaakov R, Rimon G. Incidence of neuronal dye-coupling in neocortical slices depends on the plane of section. *Neuroscience.* 1985; 15:659–66. [PubMed: 3906427]
17. Knowles WD, Funch PG, Schwartzkroin PA. Electrotonic and dye coupling in hippocampal CA1 pyramidal cells in vitro. *Neuroscience.* 1982; 7:1713–22. [PubMed: 6289179]
18. Wang Y, Barakat A, Zhou H. Electrotonic coupling between pyramidal neurons in the neocortex. *PLoS One.* 2010; 5:e10253. [PubMed: 20436674]
19. Gibson JR, Beierlein M, Connors BW. Two networks of electrically coupled inhibitory neurons in neocortex. *Nature.* 1999; 402:75–9. [PubMed: 10573419]
20. Galarreta M, Hestrin S. A network of fast-spiking cells in the neocortex connected by electrical synapses. *Nature.* 1999; 402:72–5. [PubMed: 10573418]
21. Thomson AM, Deuchars J. Synaptic interactions in neocortical local circuits: dual intracellular recordings in vitro. *Cereb Cortex.* 1997; 7:510–22. [PubMed: 9276176]
22. Stevens CF, Zador AM. Input synchrony and the irregular firing of cortical neurons. *Nat Neurosci.* 1998; 1:210–7. [PubMed: 10195145]
23. Hebb, DO. The organization of behavior. Wiley & Sons; New York: 1949.
24. Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. *Science.* 1996; 274:1133–8. [PubMed: 8895456]
25. Sosinsky GE, Nicholson BJ. Structural organization of gap junction channels. *Biochim Biophys Acta.* 2005; 1711:99–125. [PubMed: 15925321]
26. Willecke K, et al. Structural and functional diversity of connexin genes in the mouse and human genome. *Biol Chem.* 2002; 383:725–37. [PubMed: 12108537]

27. Elias LA, Wang DD, Kriegstein AR. Gap junction adhesion is necessary for radial migration in the neocortex. *Nature*. 2007; 448:901–7. [PubMed: 17713529]
28. Nadarajah B, Jones AM, Evans WH, Parnavelas JG. Differential expression of connexins during neocortical development and neuronal circuit formation. *J Neurosci*. 1997; 17:3096–111. [PubMed: 9096144]
29. Beahm DL, et al. Mutation of a conserved threonine in the third transmembrane helix of alpha- and beta-connexins creates a dominant-negative closed gap junction channel. *J Biol Chem*. 2006; 281:7994–8009. [PubMed: 16407179]
30. Rouan F, et al. trans-dominant inhibition of connexin-43 by mutant connexin-26: implications for dominant connexin disorders affecting epidermal differentiation. *J Cell Sci*. 2001; 114:2105–13. [PubMed: 11493646]
31. Fushiki S, et al. Changes in neuronal migration in neocortex of connexin43 null mutant mice. *J Neuropathol Exp Neurol*. 2003; 62:304–14. [PubMed: 12638734]
32. Personius KE, Balice-Gordon RJ. Loss of correlated motor neuron activity during synaptic competition at developing neuromuscular synapses. *Neuron*. 2001; 31:395–408. [PubMed: 11516397]
33. Chuang CF, Vanhoven MK, Fetter RD, Verselis VK, Bargmann CI. An innexin-dependent cell network establishes left-right neuronal asymmetry in *C. elegans*. *Cell*. 2007; 129:787–99.
34. Curtin KD, Zhang Z, Wyman RJ. Gap junction proteins expressed during development are required for adult neural function in the *Drosophila* optic lamina. *J Neurosci*. 2002; 22:7088–96. [PubMed: 12177205]
35. Corlew R, Bosma MM, Moody WJ. Spontaneous, synchronous electrical activity in neonatal mouse cortical neurones. *J Physiol*. 2004; 560:377–90. [PubMed: 15297578]
36. Dupont E, Hanganu IL, Kilb W, Hirsch S, Luhmann HJ. Rapid developmental switch in the mechanisms driving early cortical columnar networks. *Nature*. 2006; 439:79–83. [PubMed: 16327778]
37. Yang JW, Hanganu-Opatz IL, Sun JJ, Luhmann HJ. Three patterns of oscillatory activity differentially synchronize developing neocortical networks in vivo. *J Neurosci*. 2009; 29:9011–25. [PubMed: 19605639]
38. Kandler K, Katz LC. Coordination of neuronal activity in developing visual cortex by gap junction-mediated biochemical communication. *J Neurosci*. 1998; 18:1419–27. [PubMed: 9454851]
39. Du Z, et al. Introduction of oncogenes into mammary glands in vivo with an avian retroviral vector initiates and promotes carcinogenesis in mouse models. *Proc Natl Acad Sci U S A*. 2006; 103:17396–401. [PubMed: 17090666]
40. Holland EC, Hively WP, DePinho RA, Varmus HE. A constitutively active epidermal growth factor receptor cooperates with disruption of G1 cell-cycle arrest pathways to induce glioma-like lesions in mice. *Genes Dev*. 1998; 12:3675–85. [PubMed: 9851974]
41. Vos BP, Maex R, Volny-Luraghi A, De Schutter E. Parallel fibers synchronize spontaneous activity in cerebellar Golgi cells. *J Neurosci*. 1999; 19:RC6. [PubMed: 10341267]

Figure 1. Electrical coupling between sister excitatory neurons in neocortical ontogenetic columns

(a) Dual whole-cell recordings of sister excitatory neurons in ontogenetic columns at different neonatal stages. (top) Images of sister excitatory neurons in ontogenetic columns expressing EGFP (green) filled with Alexa Fluor 568 hydrazide (red) through the recording pipette and stained with DNA dye 4',6-diamidino-2-phenylindole (DAPI, blue). Scale bars: 50 μ m. (bottom) Simultaneous depolarization or hyper-polarization of two sister excitatory neurons in an ontogenetic column (V_1 , V_2) when a positive or negative current (+100 pA or -100 pA) was injected into only one of the neurons (I_1 or I_2) under current-clamp conditions, indicating electrical coupling. Subsequent figures use a similar panel layout. Scale bars: 50 mV (black); 10 mV (red); 200 msec (blue) (P1, P2 and P4); 50 mV (black); 2 mV (red); 200 msec (blue) (P6). (b) Blockade of electrical coupling by the gap junction blocker meclofenamic acid (MFA, 100 μ M). Scale bars: 25 mV; 5 mV; 200 msec. (c) Summary of the frequency of electrical coupling between sister excitatory neurons at different postnatal stages.

Figure 2. Preferential formation of strong electrical coupling between sister excitatory neurons in neocortical ontogenetic columns

(a) Image of a quadruple whole-cell recording of two EGFP-expressing sister excitatory neurons in an ontogenetic column (green, neurons 1 and 3) and two non-EGFP-expressing excitatory neurons (neurons 2 and 4) adjacent to the sisters on the same side, filled with Alexa Fluor 568 hydrazide (red) and stained with DAPI (blue). Scale bar: 20 μ m. (b) Sample traces of voltage changes in the four neurons in response to sequential current injection into one of the four neurons. Green circles indicate EGFP-expressing sister excitatory neurons and white circles indicate non-EGFP-expressing neighbouring excitatory neurons. The average traces are shown in the rectangle. Scale bars: 50 mV; 5 mV; 200 msec. (c) Morphological reconstruction of the four neurons in a quadruple recording. The wavy line and arrows indicate reciprocal electrical coupling between sister excitatory neurons 1 and 3. (d) Summary of the frequency of electrical coupling observed between sister excitatory neurons in ontogenetic columns and their adjacent non-sister excitatory neurons at P1–P6. (e) Summary of the coupling coefficient observed between sister excitatory neurons in ontogenetic columns and their adjacent non-sister excitatory neurons at P1–P6. ***, $p < 1e-8$. (f) Progressive decrease in coupling coefficient between sister excitatory neurons in ontogenetic columns as development proceeds. *, $p < 0.01$.

Figure 3. Electrical coupling promotes action potential generation and synchronous firing of sister excitatory neurons in neocortical ontogenetic columns

(a–c) Electrical coupling facilitates action potential generation in sister excitatory neurons. (a) Sample traces of synchronous (1+2) or asynchronous (1 or 2) injection of sub-threshold current pulses into electrically coupled sister excitatory neurons in ontogenetic columns. Arrowheads indicate voltage deflection due to the electrical transmission. Note that synchronous but not asynchronous injection results in action potential generation (arrows). Scale bars: 50 pA; 20 mV; 100 msec; (b) Summary of the firing rate of two sister excitatory neurons in *c* responding to 50 pA current injection (**, $p < 0.001$). Filled bars correspond to synchronous injection; open bars correspond to asynchronous injection. (c) Summary of the firing rate in electrically coupled (closed circles) or non-coupled (open circles) sister excitatory neurons responding to simultaneous current injections ($n=7$; **, $p < 0.001$). (d–g) Electrical coupling promoting synchronous firing of sister excitatory neurons in ontogenetic columns in response to uncorrelated simulated current injection (d, e) or uncorrelated native neuronal activity (f, g). (d, f) Sample traces of voltage changes in electrically coupled sister

excitatory neurons. Arrows indicate the spikes occurring in both neurons within a 1-msec window. Scale bars: 200 pA; 30 mV; 100 msec (d); 100 pA; 25 mV; 100 msec (f). (e, g) Normalized cross-correlogram analysis. Bin size is 1 msec. Note that the firing frequency is significantly increased near 0 msec (Z score is larger than 3) for coupled (red) but not for non-coupled (black), sister excitatory neuron pairs, indicating synchronous firing.

Figure 4. Preferential electrical coupling is required for chemical synapse formation between sister excitatory neurons in neocortical ontogenetic columns
(a–f) Quadruple whole-cell recordings of two sister excitatory neurons (2 and 4) in ontogenetic columns expressing Cx26-IRES-EGFP (a–c) or Cx26(T135A)-IRES-EGFP (d–f) and two adjacent non-sibling excitatory neurons (1 and 3). (a, d) DIC and fluorescence images and (b, e) morphological reconstruction of the respective quadruple whole-cell recording. Scale bars: 100 μ m. (c, f) Summary of the chemical synaptic connection detected in the quadruple recordings. Scale bars: 10 pA; 200 msec. **(g)** Summary of the frequency of chemical synapse formation between sister excitatory neurons in ontogenetic columns expressing EGFP, Cx26-IRES-EGFP or Cx26(T135A)-IRES-EGFP, and their adjacent non-sister excitatory neurons.