Published in final edited form as:

Mol Oral Microbiol. 2013 April ; 28(2): 83–101. doi:10.1111/omi.12012.

Microbial interactions in building of communities

Christopher J. Wright^{1,*}, Logan H. Burns^{1,*}, Alison A. Jack², Catherine R. Back², Lindsay C. Dutton², Angela H. Nobbs², Richard J. Lamont¹, and Howard F. Jenkinson²

¹Department of Oral Health and Systemic Disease, University of Louisville, 570 South Preston Street, Louisville, Kentucky, 40202, USA

²School of Oral and Dental Sciences, University of Bristol, Lower Maudlin Street, Bristol BS12LY, UK

SUMMARY

Establishment of a community is considered to be essential for microbial growth and survival in the human oral cavity. Biofilm communities have increased resilience to physical forces, antimicrobial agents, and nutritional variations. Specific cell-to-cell adherence processes, mediated by adhesin-receptor pairings on respective microbial surfaces, are able to direct community development. These interactions co-localize species in mutually beneficial relationships, such as streptococci, veillonellae, *Porphyromonas gingivalis* and *Candida albicans*. In transition from the planktonic mode of growth to a biofilm community, microorganisms undergo major transcriptional and proteomic changes. These occur in response to sensing of diffusible signals, such as autoinducer molecules, and to contact with host tissues or other microbial cells. Underpinning many of these processes are intracellular phosphorylation events that regulate a large number of microbial interactions relevant to community formation and development.

Keywords

biofilm; *Streptococcus*; *Porphyromonas*; *Candida*; phosphorylation; signaling

INTRODUCTION

In the natural world, microorganisms are mostly organized into communities, and these in turn are found assembled upon abiotic or living substrates as biofilms. A typical biofilm forms at an interface of two phases and comprises microbial cells enclosed within a matrix consisting of polysaccharides, proteins, nucleic acid and lipids (Flemming & Wingender, 2010), derived from microbe and environmental sources. Mono-species biofilm formation often proceeds through distinct developmental stages, as exemplified by *Pseudomonas aeruginosa* (Sauer *et al.*, 2002) and by *Candida albicans* (Chandra *et al.*, 2001). The process is initiated through low affinity attachment of planktonic cells to a substrate, followed by high affinity adhesion mediated by specific receptors. Microcolonies develop upon growth and division of attached cells, sometimes referred to as a linking film. Subsequently, recruitment of planktonic cells (from the fluid phase) leads to the further development of the community. Moreover, recruitment of heterotypic bacterial species, and/or initial adhesion

Correspondence: Richard J. Lamont, Department of Oral Health and Systemic Disease, University of Louisville, 570 South Preston Street, Louisville, Kentucky, 40202, USA. Tel: +502 852 2112; Fax: +502 852 6394; rich.lamont@louisville.edu or Howard F. Jenkinson, School of Oral and Dental Sciences, University of Bristol, Lower Maudlin Street, Bristol BS12LY, UK. Tel: +44117 3424423; Fax: +44117 342 4313; howard.jenkinson@bristol.ac.uk.

*Contributed equally

by multiple species in close proximity leads to the formation of multi-species (polymicrobial) communities, which represents the most common situation in nature. The integrity of the biofilm community is maintained by inter-microbial adhesion, cell signaling by means of cell-to-cell contact, and metabolic communication including quorum sensing (Swift et al., 2001; Blango & Mulvey, 2009).

The advantages for microorganisms growing in biofilm communities over remaining in planktonic conditions are numerous. The inherent matrix of the biofilm, such as extracellular polymeric substances (EPS), and the presence of persister cells surviving at low metabolic rates, contribute to the widely described phenomenon of reduced sensitivity to antimicrobial agents (Hoyle & Costerton, 1991). Biofilms are also more resilient to mechanical removal and to killing by the host immune system (Leid *et al.*, 2005). More recently, extracellular DNA (eDNA) has been shown to play an important structural role in stabilizing biofilms (Barnes *et al.*, 2012). In addition, this eDNA may be a source for potential transfer of antibiotic resistance or virulence genes between species within the communities (Roberts & Mullany, 2010).

GENERAL CHARACTERISTICS OF ORAL BIOFILMS

The organization of oral microorganisms into dental plaque biofilms plays an essential role in their survival (Jakubovics & Kolenbrander, 2010). The microorganisms are continually subjected to environmental challenges in the oral cavity including variations in oxygen and nutrient availability, pH fluctuations and the antimicrobial properties of saliva (Abiko & Saitoh, 2007). Microorganisms that fail to attach to host surfaces, or to adhered antecedent organisms, are unable to participate in community development and are subjected to eventual displacement due to the flow of saliva and other mechanical shearing forces (Scannapieco, 1994).

Oral bacteria bind to accessible host or bacterial surfaces and form complex communities in an orderly fashion. Mature dental plaque on teeth contains about 10^9 bacteria per gram and anywhere up to about 200 microbial species or phylotypes (Dewhirst et al., 2010). Certain species initiate community formation by interacting directly with the salivary pellicle that is deposited on newly available tooth surfaces. Notable pioneer organisms include many species of oral streptococci (Nyvad & Kilian, 1990), *Actinomyces* spp. *Granulicatella adiacens*, *Abiotrophia defectiva*, *Gemella* spp. and *Rothia* (Jenkinson, 2011). These early colonizers are all components of the natural microbiota (Aas *et al.*, 2005), and few are known to be directly responsible for the development of a diseased state. However, early colonizers such as streptococci (Palmer *et al.*, 2001) can alter the pathogenic potential of the oral biofilm through both their influence on how the biofilm community develops and by elevating the pathogenic potential of other bacteria (Whitmore & Lamont, 2011).

***Streptococcus* adhesion molecules**

Streptococci are facultatively anaerobic and adhere to an array of salivary molecules including mucins, proline-rich proteins, statherin, salivary agglutinin (gp-340), and α -amylase (Nobbs *et al.*, 2009). They also bind a wide variety of oral microorganisms, leading to development of complex microbial networks that stabilize communities. Accordingly, streptococci express a diversity of cell surface molecules that enable adherence to host or bacterial receptors. For example, long thread-like structures termed pili are produced by *Streptococcus sanguinis*, and by other *Streptococcus* species, and are composed of polymers of three different protein subunits (PilA, PilB, PilC). These promote attachment to host receptors (Okahashi *et al.*, 2010) and in *Streptococcus pneumoniae* are required for full virulence (Barocchi *et al.*, 2006). Pili and many other surface proteins are found covalently-linked to the cell wall peptidoglycan through their C-terminal anchorage

sequences (Nobbs *et al.*, 2009). Other cell-wall anchored polypeptides identified in oral streptococci functioning as adhesins include: CshA (and CshB) which forms surface fibrils that interact with fibronectin and other oral microbes (Holmes *et al.*, 1996; McNab *et al.*, 1999); Hsa (and GspB) that interact with salivary pellicle, epithelial cells and blood platelets (Kerrigan *et al.*, 2007); PadA which interacts with blood platelets (Petersen *et al.*, 2010) and salivary pellicle components; Fap1 which binds salivary pellicle and mediates biofilm formation (Ramboarina *et al.*, 2010); BapA1, which represents a new family of streptococcal adhesins involved in biofilm formation (Liang *et al.*, 2011); AbpA which binds α -amylase (Nikitkova *et al.*, 2012); and glucan-binding proteins (GbpB, GbpC) that promote adhesion of bacteria to polysaccharide matrix (Mattos-Graner *et al.*, 2006; Biswas *et al.*, 2007). Genomic sequencing has revealed that some streptococcal strains may carry up to 30 or more genes encoding proteins with predicted cell-wall anchorage (Nobbs *et al.*, 2009). Therefore it is likely that in future more functional adhesins will be characterized that play a role in streptococcus colonization and biofilm development.

Antigen I/II (AgI/II) family protein adhesins are produced by most oral streptococci. The sequences and structures are well-conserved, but they have a diverse range of functions in mediating adhesion to host surfaces, and coaggregation with other oral microorganisms (Brady *et al.*, 2010). The AgI/II protein expressed by *S. mutans* (variously designated SpaP, Pac or AgB) is responsible for adhesion of bacteria to salivary pellicle. *S. gordonii* produces two AgI/II family proteins termed SspA (172 kDa) and SspB (164 kDa). These polypeptides mediate coaggregation of streptococci with *Actinomyces oris* (Egland *et al.*, 2001) but with strain specificities (Jakubovics *et al.*, 2005). SspB also interacts directly with *Porphyromona gingivalis* through a C-terminal region designated BAR that is recognized by the shorter fimbriae on the surface of *P. gingivalis* (Daep *et al.*, 2008).

Microbial interactions between earlier and later colonizers

Streptococci and other precursor organisms provide unique receptor sites for later, more pathogenic colonizers such as *Fusobacterium nucleatum* (He *et al.*, 2012), *Tannerella forsythia*, *Treponema denticola*, and *Porphyromonas gingivalis* (Fig. 1) (Kuboniwa & Lamont, 2010; Periasamy & Kolenbrander, 2010), which are closely associated with the development of periodontitis (Haffajee & Socransky, 1994; van Winkelhoff *et al.*, 2002). Adherence of *P. gingivalis* to antecedent bacteria promotes initial colonization, and ultimately facilitates periodontal destruction (Slots & Gibbons, 1978). A well understood interspecies interaction is between that of *P. gingivalis* and the oral commensal *Streptococcus gordonii*. This occurs through two sets of adhesin-receptor pairs (Fig. 2). *P. gingivalis* cells display preferential binding to oral surfaces coated with certain streptococci, such as *S. gordonii* and other members of the oralis group (Lamont *et al.*, 1992). It is likely that this interaction begins primarily on the supragingival tooth surface (Ximenez-Fyvie *et al.*, 2000; Mayanagi *et al.*, 2004; Haffajee *et al.*, 2008). From here, *P. gingivalis* may spread laterally to the subgingival region via an increase in biomass or by cell dispersal as the result of active cellular release or passive mechanical shearing of the supragingival biofilm. *P. gingivalis*, and some other secondary colonizers such as *Fusobacterium nucleatum* can also provide bridging functions by expressing multiple adhesins that bind other later colonizers (Kolenbrander *et al.*, 2002). Both attachment-based and physiological interactions between late colonizers and compatible precursor organisms can promote progression of the plaque biofilm towards a more pathogenic state (Kuboniwa & Lamont, 2010; Whitmore & Lamont, 2011).

Interspecies recognition and co-adhesion contributes to community formation and to the success of the participating bacteria. Bacterial growth within a community can bring metabolic advantages and access to nutrients that would be unavailable to planktonic organisms. As certain species flourish within the community they release metabolites that

can be utilized by other community inhabitants. In such a heterotypic biofilm, bacteria will often co-localize with other constituents that are metabolically compatible (Jenkinson & Lamont, 2005). This metabolic synergy within the community can allow the development of a more complex microbiota. One example of metabolic synergy occurs between *T. denticola* and *P. gingivalis*. When grown in a dual species biofilm, these organisms produce a significantly larger biomass than the total of the individual monospecies biofilms (Grenier, 1992). *P. gingivalis* produces isobutyric acid which stimulates growth of *T. denticola*, while *T. denticola* produces succinic acid that enhances growth of *P. gingivalis* (Fig. 1). The chymotrypsin-like proteinase produced by *T. denticola* also stimulates formation of a dual species biofilm with *P. gingivalis* (Cogoni *et al.*, 2012). The co-operation of *P. gingivalis* with other oral species such as *F. nucleatum* has also been demonstrated. The ability of some *F. nucleatum* strains to tolerate higher oxygen concentrations than *P. gingivalis* means that *F. nucleatum* facilitates generation of reduced oxygen conditions that promote growth and survival of *P. gingivalis* (Bradshaw *et al.*, 1998; Diaz *et al.*, 2002). This modification of the microenvironment by *F. nucleatum* may allow growth of other strictly anaerobic oral species (Kolenbrander *et al.*, 1995). *F. nucleatum* can also elevate the pH of its environment through the generation of ammonia, thus neutralizing acid produced by fermenting microorganisms and creating a more favorable environment for *P. gingivalis* and other acid-sensitive organisms (Takahashi, 2003).

Bacterial interactions are often established by pairings of adhesin (protein) and receptor (saccharide) components found on the surfaces of the associated bacteria (Kolenbrander *et al.*, 2006). An example of such interaction involves the type 2 fimbriae on the surface of *A. oris* (*naeslundii*) that recognize a (GalNAc β 1 $\rightarrow$ 3Gal) linkage present within cell wall polysaccharide on *Streptococcus oralis* thus allowing the cells to co-aggregate (Palmer *et al.*, 2003; Yoshida *et al.*, 2006). In the case of interaction of *P. gingivalis* with *F. nucleatum* (Rosen & Sela, 2006), the latter expresses a lectin adhesin that specifically recognizes galactose, which is present in the capsule and lipopolysaccharide of *P. gingivalis*. Similar galactose-containing receptors are found on *Aggregatibacter actinomycetemcomitans* (Rupani *et al.*, 2008) and *T. denticola* (Rosen *et al.*, 2008), in the form of the serotype-specific O polysaccharide and outer membrane carbohydrate groups, respectively. Thus, strains of *F. nucleatum* actively bind these different organisms, both earlier and later colonizers. Also, *T. denticola* and *T. forsythia* each express leucine-rich repeat proteins that mediate mutual attachment and facilitate binding to *F. nucleatum*, further adding to the developing bacterial network (Ikegami *et al.*, 2004; Sharma *et al.*, 2005) (Fig. 1).

METABOLIC NETWORKS

Many bacteria rely on metabolic cooperation provided by close proximity of cells to grow and become incorporated within oral microbial communities. For example, *Veillonella* species are Gram-negative, anaerobic cocci, which occur in plaque in high abundance (Bik *et al.*, 2010) and are part of the pioneer oral community after birth (Cephas *et al.*, 2011). Growth of streptococci leads to the formation of lactic acid, which is a favored substrate of *Veillonella atypica*. This in turn accelerates the glycolysis rate in streptococci by removing the end-product (lactate) inhibition (Fig. 1).

When *S. gordonii* and *V. atypica* are grown in co-culture a diffusible signal produced leads to up-regulation of the *S. gordonii* amylase gene, *amyB*. Increased amylase activity on a starch substrate would produce more fermentable glucose, generating further lactic acid and more favorable conditions for *V. atypica* (Egland *et al.*, 2004). On the other hand, *S. gordonii* appears to benefit from interaction with *A. naeslundii* (Egland *et al.*, 2001). When co-cultured, a number of genes involved in arginine biosynthesis are differentially expressed in *S. gordonii* (Jakubovics *et al.*, 2008) potentially increasing the efficiency of arginine

biosynthesis in *S. gordonii*. An observation that highlights the benefits of interspecies cooperation is in the degradation of salivary mucins. Individually, *Streptococcus* species do not necessarily produce all of the required enzymes for mucin hydrolysis, while cooperatively they are able to more efficiently utilize the mucin oligosaccharides for growth (Byers *et al.*, 1999).

The examples above begin to paint a picture of a web of metabolic exchanges that occur in the oral cavity (Jenkinson, 2011). But more simply, colonization by early pioneering colonizers e.g. streptococci can enhance the growth and virulence of potentially pathogenic bacteria such as *P. gingivalis* and *T. denticola*. This has led to the mitis-group streptococci e.g. *S. gordonii*, *S. oralis* etc. being termed accessory pathogens in the oral cavity (Whitmore & Lamont 2011).

Antagonism

Microorganisms are not always greeted into a community with open arms. A number of bacterial species have evolved specific mechanisms to inhibit growth and attachment of competing organisms. Hydrogen peroxide produced by some of the oral streptococci provides one mechanism of bacterial antagonism (Holmberg & Hallander, 1973). However, interspecies interactions are often multi-threaded. Kreth *et al.* (2005) observed two separate mechanisms by which *S. sanguinis* and *S. mutans* are mutually antagonistic, based upon hydrogen peroxide (H₂O₂) production by *S. sanguinis* and bacteriocin production by *S. mutans*. When grown simultaneously, both species proliferate; however, prior establishment of one of the species prevents the other from occupying the same niche (Kreth *et al.*, 2005). Further work demonstrated that *S. gordonii* also inhibits growth of *S. mutans*, and that this is promoted under aerobic conditions which led to elevated H₂O₂ levels. Interestingly, H₂O₂ production by bacteria may have also co-evolved to act as a signaling molecule for the fungus *Candida albicans* to undergo filamentation (Srinivasa *et al.*, 2012). More aerobic conditions appear to stimulate production of bacteriocins by *S. mutans*, through activation of the Gram-positive competence-stimulating peptide (CSP) signaling system encoded by the *com* genes (Kreth *et al.*, 2008). As covered in more detail below, *S. mutans* expresses a eukaryotic serine/threonine type kinase known as STPK which contributes to resistance to peroxide (Zhu & Kreth, 2010). This thrust and counter-thrust, driven by co-evolution, continues through strategies to subvert the production of antagonistic elements. An example of this is a gene *sgc* in *S. gordonii* encoding a protease capable of interfering with bacteriocin production in *S. mutans* (Wang & Kuramitsu, 2005).

Certain oral streptococci have been shown to have a negative impact on biofilm formation by *P. gingivalis*. Contact with the later streptococcal colonizer *S. cristatus* has been shown to down-regulate expression of *fimA*, which encodes the major fimbrial adhesin of *P. gingivalis*, and thus prevent *P. gingivalis* accumulation on *S. cristatus*-rich substrata (Wang *et al.*, 2009). Arginine deaminase (ArcA) in *S. cristatus* provides the communication signal responsible for the down-regulation of *fimA* in *P. gingivalis*, although enzymatic function of ArcA is not essential for signaling activity (Xie *et al.*, 2007). *S. intermedius* also produces arginine deaminase that can repress the expression of both FimA and Mfa1 (minor fimbria) in *P. gingivalis* (Christopher *et al.*, 2010). Although *S. gordonii* produces ArcA, *cis* catabolite response elements function to repress expression in *S. gordonii* in comparison to *S. cristatus* (Lin *et al.*, 2008). This antagonistic interaction has been shown to have biological consequences. In a mouse model colonization of the oral cavity by ArcA-expressing *S. cristatus* followed by *P. gingivalis* infection reduces the levels of *P. gingivalis* colonization and subsequent bone loss (Xie *et al.*, 2012).

Detachment of microorganisms from the biofilm may occur as a consequence of antagonism, or exclusion. Release of cells from the biofilm is anyway an important

mechanism for dispersal, and may be passive or active. The main mechanism is shear force, such as salivary flow or external applications such as tooth-brushing. However, some microbial species are known to actively disperse from the biofilm. Following *P. aeruginosa* biofilm formation, cells on the outer layer remain as a stationary biofilm phenotype, while cells on the inside of the biofilm become motile (planktonic phenotype) and can swim out of the biofilm, leaving a hollow mound (Sauer *et al.*, 2002). The dental pathogen *A. actinomycetemcomitans* is not a motile species, but does have the ability to become released from the biofilm, utilizing dispersin B, a biofilm-releasing β -hexosaminidase (Manuel *et al.*, 2007). Salivary flow then seeds other areas of the mouth with the released *A. actinomycetemcomitans* cells (Kaplan & Fine 2002). *Candida albicans* biofilms are composed mainly of a network of hyphae (filaments) that provide the biofilm structure. A transcriptional regulator Nrg1p blocks yeast to hyphae transition and controls dispersion of yeast morphology cells from the biofilm (Uppuluri *et al.*, 2010).

INTERACTIONS OF STREPTOCOCCI WITH PORPHYROMONAS AND CANDIDA

The AgI/II adhesins of oral streptococci

The streptococcal AgI/II family of proteins are multifunctional adhesins with the ability to bind a variety of host components such as collagen, laminin, and salivary substrates, as well as other microorganisms (Brady *et al.*, 2010). Ag I/II family protein functions vary according to streptococcal species and strains in which they are expressed. For example, *S. mutans* strain NG5 expresses a single Ag I/II polypeptide, SpaP, whereas *S. gordonii* strains express two AgI/II family proteins (SspA and SspB). The production of these proteins is affected by environmental factors including salivary proteins, variations in pH, osmolarity and temperature. SspA and SspB expression levels increase under elevated temperature and acidic pH, while SspB expression is reduced under lower NaCl concentration (El-Sabaeny *et al.*, 2000). Also, SspA levels are generally higher than those of SspB, and the SspA polypeptide positively regulates *sspB* by binding its promoter region (El-Sabaeny *et al.*, 2001). One major receptor for AgI/II is salivary glycoprotein gp340, also found in most if not all mucosal secretions. This protein participates in innate immunity by promoting microbial cell agglutination and clearance (Prakobphol *et al.*, 2000). However, when gp340 is adsorbed onto an oral cavity surface it provides sites for streptococcal binding (Lamont *et al.*, 1991). The AgI/II proteins play a critical role in the association of *P. gingivalis* with *S. gordonii* as described in more detail below.

The SspA and SspB proteins also appear to mediate interactions of *S. gordonii* with hyphal filaments of *C. albicans*, thus promoting co-colonization by these microorganisms (Bamford *et al.*, 2009) (Fig. 1). The receptor on *C. albicans* that interacts with SspB is a protein expressed on the hyphal cell surface, designated Als3 (Silverman *et al.*, 2010). This is one of a group of proteins expressed by the agglutinin-like sequence (ALS) family of genes (Hoyer *et al.*, 1995). They encode surface glycoprotein adhesins involved in host-pathogen interactions and other adhesive functions (Hoyer *et al.*, 2008). There are eight Als proteins (Als1-8p), but the largest decrease in *C. albicans* adhesion is observed by deletion of both *als3* alleles (*C. albicans* is generally diploid) (Zhao *et al.*, 2004). Als3p is hypha-specific (Murciano *et al.*, 2012) and is in all probability involved in early establishment of biofilms in addition to interacting with oral streptococci (Silverman *et al.*, 2010).

P. gingivalis fimbriae

Numerous peritrichous fimbriae protrude from the cellular envelope of most strains of *P. gingivalis* (Listgarten & Lai, 1979; Handley & Tipler, 1986). Binding of *P. gingivalis* to sites in the oral cavity and to other bacteria is dependent, at least in part, upon these fimbrial

structures (Slots & Gibbons, 1978) which are of two kinds. The major fimbriae are longer, with lengths 0.3–1.6 μm and width approximately 5 nm (Yoshimura *et al.*, 1984), while the minor fimbriae are 80–120 nm in length and 3.5 – 6.5 nm in diameter (Park *et al.*, 2005). The major fimbriae bind to salivary proteins and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) found on the surface of *S. oralis* (Maeda *et al.*, 2004b) and other streptococci. The primary unit of the major fimbria is fimbrillin (FimA). There are at least six different variants of *fimA* among strains of *P. gingivalis*, and binding differences among these fimbrillin types could impact the likelihood of periodontal disease development (Amano *et al.*, 2004). The *fimA* locus is flanked by genes that are involved in transcriptional regulation or encode proteins that contribute to the structure of the fully mature fimbriae. The two genes upstream of *fimA* are involved in regulation via a response regulator, FimR (Nishikawa *et al.*, 2004), part of the FimS/FimR two-component signal transduction system which governs transcriptional levels of *fimA* (Hayashi *et al.*, 2000). Environmental cues detected by *P. gingivalis* that influence *fimA* expression include changes in temperature and hemin concentration (Amano *et al.*, 1994; Xie *et al.*, 1997). The arginine and lysine-specific gingipains produced by *P. gingivalis* also regulate the amount of FimA on the bacterial surface, potentially affecting adhesion and colonization (Xie *et al.*, 2000). The genes downstream of *fimA* encode the products FimC - FimE, which are minor components of the mature fimbriae. FimE is responsible for assembly of FimC and FimD proteins onto the fimbrial fiber and for maintaining a stable attachment to the bacterial surface (Nishiyama *et al.*, 2007). The absence of any of these three accessory proteins manifests as a significant reduction of FimA binding to GAPDH, which in turn affects initial *P. gingivalis* binding to oral streptococci (Maeda *et al.*, 2004a).

The minor fimbriae bind to components of the streptococcal surface. The primary subunit of the minor fimbriae, Mfa1 (67 kDa) has a role in inflammatory processes and is involved in induction of several cytokines such as IL-1 α , IL-1 β , and TNF- α in peritoneal macrophages, in response to *P. gingivalis* (Lin *et al.*, 2006). Additional genes encode accessory proteins that associate or co-operate with the main protein subunit. Downstream of *mfa1* is a co-transcribed gene encoding the protein Mfa2. This protein is known to have a role in the regulation of the length of the minor fimbriae and is required for their attachment to the cell envelope (Hasegawa *et al.*, 2009). *P. gingivalis* cells that do not express Mfa2, but still produce Mfa1, have abnormally lengthened minor fimbrial extensions and these fimbriae are only weakly bound to the cells. There are three additional, less characterized products encoded by the genes situated downstream of *mfa2* which are predicted to be accessory proteins that interact directly with the polymerized Mfa1 since they are co-purified with the filaments (Hasegawa *et al.*, 2009).

Interaction of SspA and SspB with Mfa1

P. gingivalis binding to *S. gordonii* SspA and SspB proteins is important for development of dual species biofilm communities (Lamont *et al.*, 2002). *P. gingivalis* does not interact with all AgI/II family members, despite conservation in primary amino acid sequences and secondary structure (Brady *et al.*, 2010). For example, *P. gingivalis* does not recognize SpaP of *S. mutans* (Brooks *et al.*, 1997). The precise region of SspA/B involved in binding *P. gingivalis* was determined by examining a series of truncated SspB polypeptides and chimeric proteins consisting of portions of SspB and SpaP. The adherence characteristics of the chimeras, composed of different sections of SspB and SpaP sequences, pointed toward a region (aa residues 1167 – 1250 in SspB) as being necessary for *P. gingivalis* binding. This region was designated BAR (SspB adherence region) and shown to have a significantly different secondary structure from the corresponding region in SpaP (Forsgren *et al.*, 2010). A critical region of 26 amino acid residues within BAR contains the necessary motifs recognized by Mfa and a synthetic BAR peptide can adhere to *P. gingivalis*, whereas the

corresponding SpaP peptide does not. It was also shown that BAR peptides with specific mutations confer conformational changes that ablate binding to *P. gingivalis* cells (Demuth *et al.*, 2001).

The NITVK motif within BAR helps define the binding specificity for AgI/II protein members as NITVK is only found in the AgI/II proteins of the oralis group of streptococci, which includes *S. gordonii*, *S. oralis* and *S. sanguinis*. The AgI/II proteins expressed by other streptococci have Gly in place of Asn¹¹⁸², Pro in place of Val¹¹⁸⁵, or potentially both of these substitutions. Both Gly and Pro are known to terminate α helices in proteins, and these particular substitutions inhibit *P. gingivalis* binding. A VQDLL motif upstream of NITVK is also conserved in the oralis group and resembles a nuclear receptor box domain of eukaryotic proteins that is involved in protein-protein interactions (Daep *et al.*, 2008). This motif is also flanked by lysine residues, and the positive charge of these residues could participate in electrostatic interactions that stabilize the complex of SspB with Mfa1. The VQDLL motif is in an α -helical region while NITVK is in a predicted β sheet, and this region extends outside of the SspB core, making it accessible to Mfa1 (Forsgren *et al.*, 2010). The importance of the BAR region for co-colonization with *P. gingivalis* has been established in a mouse model. Peptides derived from BAR inhibited *P. gingivalis* colonization and disease in mice pre-infected with *S. gordonii* (Daep *et al.*, 2011). The efficacy and low toxicity to the host of BAR derivatives suggest that they could be developed as a therapeutic or prophylactic agent in periodontal disease (Daep *et al.*, 2008).

GENE REGULATION WITH THE COMMUNITY ENVIRONMENT

Transcriptional studies on the differences between planktonic versus the sessile (biofilm) state have highlighted the fundamental shift an organism undertakes as it becomes part of the biofilm community (O'Toole & Kolter, 1998). A dramatic response to monospecies biofilm formation is exhibited by *P. gingivalis*, with 18% of the genome differentially regulated compared to planktonic organisms. Many of the regulated genes are associated with cell envelope biogenesis, DNA replication and metabolism, supporting the concept that cells in the transition from planktonic to biofilm state exhibit a lower rate of growth and cell metabolism (Lo *et al.*, 2009). Genes involved in adhesion and early biofilm formation, *fimA* and *mfa1*, were up-regulated in early biofilms, while *fimA* was down-regulated in the later stages (Yamamoto *et al.*, 2011). In a community with *S. gordonii*, thirty-three *P. gingivalis* genes showed up- or down-regulation by microarray analysis (Simionato *et al.*, 2006), one of which was *ltp1*, a tyrosine phosphatase.

Prevailing environmental conditions, influenced by diet, can have an important role in regulating gene activity within the oral microbial community (Bradshaw *et al.*, 1989; Percival *et al.*, 1991). Different sugars or complex carbohydrates influence expression of specific metabolic pathways (Klein *et al.*, 2010) and impact on mechanisms controlling cell integrity and secretion of extracellular biofilm matrix. Carbohydrate metabolism is integrated with cell-cell signaling systems, such as the autoinducer-2 (LuxS/AI-2, see below) pathway. A LuxS deficient strain of *S. mutans* was affected in expression of genes involved in carbohydrate metabolism, DNA repair, amino acid and protein synthesis and stress tolerance (Wen *et al.*, 2011). A phenotypic outcome of loss of LuxS is a fundamental difference in biofilm architecture.

The correlation between mRNA levels and protein amounts is not always strong (Nie *et al.*, 2006) and thus it is also important to understand community adaptation at the proteome level. In *T. forsythia*, 44 proteins were found to be altered between planktonic and biofilm cultures (Pham *et al.*, 2010). Many of these proteins were associated with the outer membrane and transport systems, and effects were observed in amount of S-layer produced

by the cells. Oxidative stress response proteins were up-regulated and the resulting biofilm cells were 10- to 20-fold more resistant to oxidative stress as compared to their planktonic counterparts. This could enhance survival of *T. forsythia* in the presence of H₂O₂-producing streptococci (Pham *et al.*, 2010).

P. gingivalis displays differential abundance of 47 proteins when grown in planktonic versus biofilm conditions (Ang *et al.*, 2008). A high percentage of these changes were associated with the cell envelope. Increased presence of proteins associated with hemin transport and metabolism indicated that *P. gingivalis* cells were entering a starved state (Ang *et al.*, 2008). In a three-species community with *S. gordonii* and *F. nucleatum*, levels of *P. gingivalis* proteins involved in cell envelope structure and DNA repair were decreased (Kuboniwa *et al.*, 2009), indicating that the multi-species community environment was less stressful to *P. gingivalis*.

Signaling within a biofilm setting

Communication is an important part of any society, including bacterial communities, and intra- and inter-species communication facilitates community development. The first example of bacterial signaling was described in the marine organism *Vibrio fischeri* where a diffusible signal N-acyl homoserine lactone (AHL) was responsible for the induction of bioluminescence (Nealson & Hastings, 1979). The same AHL molecule is utilized by both *P. aeruginosa* and *Burkholderia cepacia* in co-ordination of virulence genes and biofilm formation in cystic fibrosis (Eberl & Tumbler, 2004). AHL is not commonly utilized by oral bacterial species for communication (Kolenbrander *et al.*, 2002); rather oral bacteria rely on two distinct signaling systems. The first, restricted to Gram-positive organisms such as the early colonizing streptococci, utilizes short peptides termed Competence Signaling Peptides (CSP) (Suntharalingam & Cvitkovitch, 2005) or other small peptides (Son *et al.*, 2012). These have been described in a number of streptococci, including *S. mutans* and *S. gordonii*, where they play a role in genetic exchange and virulence. The second signaling system involves autoinducer-2 (AI-2), a family of signaling molecules produced by the action of the LuxS enzyme on S-ribosyl-homocysteine (SRH), generating 4,5-dihydroxyl-2,3-pentanedione or DPD, which breaks down to produce AI-2 (Sun *et al.*, 2004). AI-2 was originally described in the marine organism *Vibrio harveyi* (Bassler *et al.*, 1993) and is now recognized as a species-independent signal that is widespread in oral bacteria including *P. gingivalis*, *A. actinomycetemcomitans* and also oral streptococci such as *S. gordonii* and *S. mutans*.

There are a number of studies on the role of AI-2 and community development within the oral cavity. For example, AI-2 is required for biofilm growth of *A. actinomycetemcomitans* (Shao *et al.*, 2007). AI-2 is linked to the two-component system QseBC, which is induced by AI-2 through uptake of AI-2 into the cell. This uptake is reliant on two AI-2 receptors, linked to ABC transporters, termed LsrB and RbsB. Deletion of either of these elements reduces *A. actinomycetemcomitans*-induced alveolar bone resorption in animal models. Loss of QseC also diminishes biofilm formation consistent with a role for this system in colonization and virulence (Novak *et al.*, 2010)(see Fig. 3). Other oral species are also dependent on AI-2 signaling for biofilm formation. *A. oris* can utilize AI-2 produced by *S. oralis* for biofilm growth and development. Interestingly, the concentration of signal is an important factor in dual species biofilm formation, and at higher concentrations of AI-2 there are significantly lower levels of biofilm formation (Rickard *et al.*, 2006). Moreover, the AI-2 concentration in *A. oris*-*S. oralis* biofilms decreases over time, possibly contributing to the persistence of the dual species communities (Rickard *et al.*, 2008).

S. gordonii also exhibits altered biofilm development in the absence of AI-2, and AI-2 is required for community development between *S. gordonii* and *P. gingivalis* (McNab *et al.*,

2003). Interkingdom effects of AI-2 have been seen between *C. albicans* and *S. gordonii*, where a *luxS* knockout of *S. gordonii* is substantially affected in dual-species biofilm formation with *C. albicans* (Bamford *et al* 2009). It is suggested that *luxS* mutation affects the ability of *S. gordonii* to promote hyphal growth in *C. albicans*. This might occur by suppressing effects of the quorum-sensing molecule farnesol, which normally inhibits filamentation (Hornby *et al.*, 2001) (Fig. 4).

A range of signaling molecules have been identified produced by bacteria that affect *C. albicans* biofilm formation or morphogenesis. These include lactic acid, H₂O₂, CO₂, and bacterial peptidoglycan (Xu *et al.*, 2008) all of which appear to promote filamentation, while HSLs inhibit filamentation (Hall *et al.*, 2011). By way of return, fatty acids, carboxylic acids and glycans produced by *C. albicans* are able to promote growth of bacteria, while farnesol inhibits bacterial biofilm formation (Pammi *et al.*, 2011). Thus filamentation of *C. albicans* and mixed-species biofilm formation are regulated by recognition of a complex array of self- or non-self signaling molecules (Fig. 4).

A number of genes are regulated by LuxS in *S. gordonii* including those involved in carbohydrate synthesis (McNab *et al.*, 2003). The pathways in *P. gingivalis* involved in signal transduction, including AI-2 dependent signaling, following contact with *S. gordonii* were identified by Chawla *et al.* (2010). A LuxR family orphan transcriptional regulator designated CdhR was shown to constrain development of *S. gordonii* and *P. gingivalis* communities. The community function of this regulator was attributed to two genes that are under its control. The first is *mfaI*, which as mentioned previously encodes the minor fimbrial antigen in *P. gingivalis* and is responsible for primary interactions with AgI/II polypeptides on the surface of *S. gordonii*. CdhR was also shown to regulate LuxS which directly affects the amount of AI-2 thus ultimately affecting dual species development (Chawla *et al.*, 2010). CdhR was also shown to be part of the same regulatory circuit as Ltp1, a tyrosine phosphatase.

PHOSPHORYLATION AND BIOFILM COMMUNITY DEVELOPMENT

The importance of post translational modification on serine, threonine and tyrosine residues has long been known in eukaryotic systems, and is gaining increasing significance in bacterial systems. Bacteria and some plant systems also possess another type of phosphorylation system based on histidine and aspartate. Two component signal transduction systems (TCSs) comprise a sensory kinase and response regulator and are the most common examples of bacterial regulatory systems involving phospho-transfer. The sensor kinase responds to external signals resulting in the autophosphorylation of a histidine residue. The transfer of the phosphate group to the response regulator results in the phosphorylation of an aspartate residue and a downstream effect on gene transcription (Gao & Stock, 2009). TCSs have been shown to be involved in biofilm development in a number of bacteria (Kolar *et al.*, 2011; Zhang *et al.*, 2009).

Serine/threonine protein kinases (STPKs)

A number of STPKs now described in bacteria are of the Hanks-type kinase, and show homology to eukaryotic kinases. This allows intracellular bacteria to utilize STPKs and the corresponding phosphatases to subvert host signal transduction via phosphorylation and de-phosphorylation of signaling components within the host (Kobir *et al.*, 2011). As well as having a role in virulence and host subversion, STPKs have been implicated in biofilm formation e.g. PrkC of *Bacillus subtilis* (Madec *et al.*, 2002) and Stk in *S. epidermidis* (Liu *et al.*, 2011). *S. mutans* possesses a homologue of the eukaryotic-like STPKs, PknB. A strain deficient in PknB exhibited reduced biofilm formation on hydroxyapatite discs, and an inability to tolerate acid conditions (Hussain *et al.*, 2006). Further characterization of PknB,

and the corresponding phosphatase PppL, showed that both enzymes were important for biofilm formation, cell shape, acid tolerance, genetic transformation and cariogenicity in a rat model (Banu *et al.*, 2010). PknB is also involved in controlling bacteriocin production, possibly through modulating the activity of a TCS in *S. mutans*. Additionally, PknB participates in oxidative stress tolerance, and a decreased fitness was observed in a *pknB* deficient strain of *S. mutans* when grown with *S. sanguinis* (Zhu & Kreth, 2010). Taken together, the results indicate that the STPK *pknB* in *S. mutans* provides an important link in the establishment of a *S. mutans* community and persistence within the oral cavity.

Tyrosine kinases

Over recent years a number of tyrosine kinases and phosphatases have been described in prokaryotes leading to the realization that phosphorylation on tyrosine is not limited to occurring in eukaryotic systems, as was once believed. Significant structural differences are apparent between Gram-positive and Gram-negative bacterial tyrosine kinases (BY-kinases). In Gram-negative species, the BY-kinase comprises a single polypeptide. A short N-terminal region is usually present within the cytoplasm, followed by a transmembrane stretch and a region occupying the periplasmic space. A second transmembrane stretch brings the polypeptide back into the cytoplasm where the C-terminal region contains the enzymatically active Walker (ATP binding) domains. In Gram-positive bacteria the kinase domain and transmembrane regions are encoded by neighboring genes on the bacterial chromosome (Lee & Jia, 2009). Initially thought of as purely autophosphorylating peptides, the discovery of phosphorylation of specific substrates has led to the appreciation that BY-kinases play a critical role in many aspects of virulence. One of the first substrates of BY-kinases to be recognized was a UDP-glucose dehydrogenase in *E. coli* (Grangeasse *et al.*, 2003). Phosphorylation of this substrate increases its activity in generating precursors for polysaccharide synthesis. BY-kinases have since been shown to play a significant role in the transport and synthesis of cellular polysaccharide (Fig. 3), and are thus likely to impact community development and biofilm formation. Tyrosine kinase activity has been found to be important for biofilm formation by *Bacillus subtilis* (Kiley & Stanley-Wall, 2010); however, the function of tyrosine kinase in oral bacteria has yet to be investigated.

Tyrosine phosphatases

Tyrosine kinases generally have partner phosphatases, such that reversible phosphorylation of substrates allows for regulation of cellular processes. Bacterial protein tyrosine phosphatases (PTPs) possess similar structures to those typically found in eukaryotes. Bacterial PTPs fall into 3 classes: the conventional PTP; dual specificity phosphatase (DSP); and a low molecular weight (LMW) PTP class. The LMW-PTP class of phosphatases has been found to be important in virulence and other physiologically important cellular events (Grangeasse *et al.*, 2007). In Gram-positive species, LMW-PTP genes are found upstream of their corresponding kinase genes, whereas in Gram-negative species, the genes are at distinct locations on the chromosome.

In *P. aeruginosa*, deletion of *tpbA* encoding a tyrosine phosphatase resulted in >100-fold increase in biofilm formation over an 8 h period. This dramatic effect was attributed to increase in initial attachment levels, decrease in swimming activity and loss of swarming motility (Ueda & Wood, 2009). TpbA also appeared to constrain pellicle and EPS formation. Two potential mechanisms account for the observed phenotype. The first is increased transcription of the *pel* locus responsible for production of EPS. The second is regulation of cyclic di-GMP, also shown to contribute to biofilm formation in *P. aeruginosa* (Ueda & Wood, 2009). The *tpbA* gene is regulated by AHL signaling in *P. aeruginosa*, indicating cross-talk between quorum sensing and tyrosine phosphatase activity (Ueda & Wood, 2009). Another aspect of the functionality of TpbA is the ability to regulate the

amount of extracellular DNA released from *P. aeruginosa* cells (Ueda & Wood, 2010). Extracellular DNA is a major component of the biofilm matrix and can provide a structural role in biofilm development (Whitchurch *et al.*, 2002; Martins *et al.*, 2010).

A tyrosine phosphatase has also been characterized in *P. gingivalis* (Maeda *et al.*, 2008). Ltp1, a LMW-PTP, influences a number of cellular processes and is a key component of a regulatory pathway that constrains heterotypic community development between *P. gingivalis* and *S. gordonii*. Ltp1 activity restricts EPS production at the transcriptional level and also negatively regulates expression of *luxS*. By contrast, Ltp1 through modulating the activity of the transcriptional regulator CdhR (see above), positively regulates *hmu* (hemin uptake operon), thus increasing hemin/iron uptake by the organism (Fig. 3). Secretion of the RgpA/B gingipain (proteinase) is reduced in a *ltp1* mutant while increase in the cell-associated Kgp gingipain is observed (Maeda *et al.*, 2008). It is interesting to note that deletion of the genes encoding the Kgp and RgpA/B gingipains markedly alters the mono-species biofilm phenotype (Kuboniwa *et al.*, 2009a). Thus, kinase-phosphatase networks have pleiotropic effects on the formation and stability of oral microbial biofilm communities.

CONCLUSIONS

Biofilm development and maintenance is an essential factor for microbial survival and growth, both in the environmental setting and the host. Benefits of the biofilm lifestyle include reduced sensitivity to mechanical shearing and to the actions of antimicrobial agents, and enhanced nutritional flexibility. Cells undergo a multitude of changes in the transition from planktonic to biofilm mode of growth. These begin with the sensing of diffusible signals that are secreted by one set of microbial cells and recognized by others. Quorum sensing along with contact dependent sensing instigates changes in gene and protein expression. Specific cell-to-cell adherence as dictated by adhesin-receptor pairings on respective bacterial surfaces can also direct community development. Interbacterial binding helps optimize co-localization of species that can coexist in a mutually beneficial relationship. Underpinning many of these processes are the intracellular phosphorylation events that regulate a large number of bacterial cell processes relevant to community formation and development.

Acknowledgments

Research of the authors is currently funded by the National Institutes of Health (NIDCR) (DE11111, DE12505, DE16690, DE17921, and DE22867) and by the Wellcome Trust (097285). Owing to space limitations it has sometimes not been possible to cite all previous publications related to a finding or topic, and for this we apologize.

REFERENCES

- Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol.* 2005; 43:5721–5732. [PubMed: 16272510]
- Abiko Y, Saitoh M. Salivary defensins and their importance in oral health and disease. *Curr Pharm Des.* 2007; 13:3065–3072. [PubMed: 17979749]
- Amano A, Nakagawa I, Okahashi N, Hamada N. Variations of *Porphyromonas gingivalis* fimbriae in relation to microbial pathogenesis. *J Periodontal Res.* 2004; 39:136–142. [PubMed: 15009522]
- Amano A, Sharma A, Sojar HT, Kuramitsu HK, Genco RJ. Effects of temperature stress on expression of fimbriae and superoxide dismutase by *Porphyromonas gingivalis*. *Infect Immun.* 1994; 62:4682–4685. [PubMed: 7927742]
- Ang CS, Veith PD, Dashper SG, Reynolds EC. Application of 16O/18O reverse proteolytic labeling to determine the effect of biofilm culture on the cell envelope proteome of *Porphyromonas gingivalis* W50. *Proteomics.* 2008; 8:1645–1660. [PubMed: 18409167]

- Bamford CV, d'Mello A, Nobbs AH, Dutton LC, Vickerman MM, Jenkinson HF. *Streptococcus gordonii* modulates *Candida albicans* biofilm formation through intergeneric communication. *Infect Immun*. 2009; 77:3696–3704. [PubMed: 19528215]
- Banu LD, Conrads G, Rehrauer H, Hussain H, Allan E, van der Ploeg JR. The *Streptococcus mutans* serine/threonine kinase, PknB, regulates competence development, bacteriocin production, and cell wall metabolism. *Infect Immun*. 2010; 78:2209–2220. [PubMed: 20231406]
- Barnes AM, Ballering KS, Leibman RS, Wells CL, Dunne GM. *Enterococcus faecalis* produces abundant extracellular structures containing DNA in the absence of cell lysis during early biofilm formation. *MBio*. 2012; 3:e00193-12.
- Barocchi MA, Ries J, Zogai X, et al. A pneumococcal pilus influences virulence and host inflammatory responses. *Proc Natl Acad Sci USA*. 2006; 103:2857–2862. [PubMed: 16481624]
- Bassler BL, Wright M, Showalter RE, Silverman MR. Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence. *Mol Microbiol*. 1993; 9:773–786. [PubMed: 8231809]
- Bik EM, Long CD, Armitage GC, et al. Bacterial diversity in the oral cavity of 10 healthy individuals. *ISME J*. 2010; 4:962–974. [PubMed: 20336157]
- Biswas I, Drake L, Biswas S. Regulation of *gbpC* expression in *Streptococcus mutans*. *J Bacteriol*. 2007; 189:6521–6531.
- Blango MG, Mulvey MA. Bacterial landlines: contact-dependent signaling in bacterial populations. *Curr Opin Microbiol*. 2009; 12:177–181. [PubMed: 19246237]
- Bradshaw DJ, McKee AS, Marsh PD. Effects of carbohydrate pulses and pH on population shifts within oral microbial communities in vitro. *J Dent Res*. 1989; 68:1298–1302. [PubMed: 2674233]
- Bradshaw DJ, Marsh PD, Watson GK, Allison C. Role of *Fusobacterium nucleatum* and coaggregation in anaerobe survival in planktonic and biofilm oral microbial communities during aeration. *Infect Immun*. 1998; 66:4729–4732. [PubMed: 9746571]
- Brady LJ, Maddocks SE, Larson MR, et al. The changing faces of *Streptococcus* antigen I/II polypeptide family adhesions. *Mol Microbiol*. 2010; 77:276–286. [PubMed: 20497507]
- Brooks W, Demuth DR, Gil S, Lamont RJ. Identification of a *Streptococcus gordonii* SspB domain that mediates adhesion to *Porphyromonas gingivalis*. *Infect Immun*. 1997; 65:3753–3758. [PubMed: 9284148]
- Byers HL, Tarelli E, Homer KA, Hambley H, Beighton D. Growth of viridians streptococci on human serum α 1-acid glycoprotein. *J Dent Res*. 1999; 78:1370–1380. [PubMed: 10403465]
- Cephas KD, Kim J, Mathai RA, et al. Comparative analysis of salivary bacterial microbiome diversity in edentulous infants and their mothers or primary care givers using pyrosequencing. *PLoS One*. 2011; 6:e23503. [PubMed: 21853142]
- Chandra J, Kuhn DM, Mukherjee PK, Hoyer LL, McCormick T, Ghannoum MA. Biofilm formation by the fungal pathogen *Candida albicans*: development, architecture, and drug resistance. *J Bacteriol*. 2001; 183:5385–5394. [PubMed: 11514524]
- Chawla A, Hirano T, Bainbridge BW, Demuth DR, Xie H, Lamont RJ. Community signalling between *Streptococcus gordonii* and *Porphyromonas gingivalis* is controlled by the transcriptional regulator CdhR. *Mol Microbiol*. 2010; 78:1510–1522. [PubMed: 21143321]
- Christopher AB, Arndt A, Cugini C, Davey ME. A streptococcal effector protein that inhibits *Porphyromonas gingivalis* biofilm development. *Microbiology*. 2010; 156:3469–3477. [PubMed: 20705665]
- Cogoni V, Morgan-Smith A, Fenno JC, Jenkinson HF, Dymock D. *Treponema denticola* chymotrypsin-like proteinase (CTLP) integrates spirochaetes within oral microbial communities. *Microbiology*. 2012; 158:759–770. [PubMed: 22313692]
- Daep CA, Lamont RJ, Demuth DR. Interaction of *Porphyromonas gingivalis* with oral streptococci requires a motif that resembles the eukaryotic nuclear receptor box protein-protein interaction domain. *Infect Immun*. 2008; 76:3273–3280. [PubMed: 18474648]
- Daep CA, Novak EA, Lamont RJ, Demuth DR. Structural dissection and *in vivo* effectiveness of a peptide inhibitor of *Porphyromonas gingivalis* adherence to *Streptococcus gordonii*. *Infect Immun*. 2011; 79:67–74. [PubMed: 21041492]

- Demuth DR, Irvine DC, Costerton JW, Cook GS, Lamont RJ. Discrete protein determinant directs the species-specific adherence of *Porphyromonas gingivalis* to oral streptococci. *Infect Immun*. 2001; 69:5736–5741. [PubMed: 11500450]
- Dewhirst FE, Chen T, Izard J, et al. The human oral microbiome. *J Bacteriol*. 2010; 192:5002–5017. [PubMed: 20656903]
- Diaz PI, Zilm PS, Rogers AH. *Fusobacterium nucleatum* supports the growth of *Porphyromonas gingivalis* in oxygenated and carbon-dioxide-depleted environments. *Microbiology*. 2002; 148:467–472. [PubMed: 11832510]
- Eberl L, Tummlier B. *Pseudomonas aeruginosa* and *Burkholderia cepacia* in cystic fibrosis: genome evolution, interactions and adaptation. *Int J Med Microbiol*. 2004; 294:123–131. [PubMed: 15493822]
- Egland PG, Du LD, Kolenbrander PE. Identification of independent *Streptococcus gordonii* SspA and SspB functions in coaggregation with *Actinomyces naeslundii*. *Infect Immun*. 2001; 69:7512–7516. [PubMed: 11705927]
- Egland PG, Palmer RJ Jr, Kolenbrander PE. Interspecies communication in *Streptococcus gordonii*-*Veillonella atypica* biofilms: signaling in flow conditions requires juxtaposition. *Proc Natl Acad Sci USA*. 2004; 101:16917–16922. [PubMed: 15546975]
- El-Sabaeny A, Demuth DR, Lamont RJ. Regulation of *Streptococcus gordonii* *sspB* by the *sspA* gene product. *Infect Immun*. 2001; 69:6520–6522. [PubMed: 11553598]
- El-Sabaeny A, Demuth DR, Park Y, Lamont RJ. Environmental conditions modulate the expression of the *sspA* and *sspB* genes in *Streptococcus gordonii*. *Microb Pathog*. 2000; 29:101–113. [PubMed: 10906265]
- Flemming HC, Wingender J. The biofilm matrix. *Nat Rev Microbiol*. 2010; 8:623–633. [PubMed: 20676145]
- Forsgren N, Lamont RJ, Persson K. Two intramolecular isopeptide bonds are identified in the crystal structure of the *Streptococcus gordonii* SspB C-terminal domain. *J Mol Biol*. 2010; 397:740–751. [PubMed: 20138058]
- Gao R, Stock AM. Biological insights from structures of two-component proteins. *Annu Rev Microbiol*. 2009; 63:133–154. [PubMed: 19575571]
- Grangeasse C, Cozzzone AJ, Deutscher J, Mijakovic I. Tyrosine phosphorylation: an emerging regulatory device of bacterial physiology. *Trends Biochem Sci*. 2007; 32:86–94. [PubMed: 17208443]
- Grangeasse C, Obadia B, Mijakovic I, Deutscher J, Cozzzone AJ, Doublet P. Autophosphorylation of the *Escherichia coli* protein kinase Wzc regulates tyrosine phosphorylation of Ugd, a UDP-glucose dehydrogenase. *J Biol Chem*. 2003; 278:39323–39329. [PubMed: 12851388]
- Grenier D. Nutritional interactions between two suspected periodontopathogens, *Treponema denticola* and *Porphyromonas gingivalis*. *Infect Immun*. 1992; 60:5298–5301. [PubMed: 1333450]
- Haffajee AD, Socransky SS. Microbial etiological agents of destructive periodontal diseases. *Periodontol*. 1994; 5:78–111. [PubMed: 9673164]
- Haffajee AD, Socransky SS, Patel MR, Song X. Microbial complexes in supragingival plaque. *Oral Microbiol Immunol*. 2008; 23:196–205. [PubMed: 18402605]
- Hall RA, Turner KJ, Chaloupka J, et al. The quorum-sensing molecules farnesol/homoserine lactone and dodecanol operate via distinct modes of action in *Candida albicans*. *Eukaryot Cell*. 2011; 10:1034–1042. [PubMed: 21666074]
- Handley PS, Tipler LS. An electron microscope survey of the surface structures and hydrophobicity of oral and non-oral species of the bacterial genus *Bacteroides*. *Arch Oral Biol*. 1986; 31:325–335. [PubMed: 2875705]
- Hasegawa Y, J. Iwami J, K. Sato K, et al. Anchoring and length regulation of *Porphyromonas gingivalis* Mfa1 fimbriae by the downstream gene product Mfa2. *Microbiology*. 2009; 155:3333–3347. [PubMed: 19589838]
- Hayashi J, K. Nishikawa K, Hirano K, Noguchi T, Yoshimura F. Identification of a two-component signal transduction system involved in fimbriation of *Porphyromonas gingivalis*. *Microbiol Immunol*. 2000; 44:279–282. [PubMed: 10832973]

- He X, Hu W, Kaplan CW, Guo L, Shi W, Lux R. Adherence to streptococci facilitates *Fusobacterium nucleatum* integration into an oral microbial community. *Microb Ecol.* 2012; 63:532–542. [PubMed: 22202886]
- Holmberg K, Hallander HO. Production of bactericidal concentrations of hydrogen peroxide by *Streptococcus sanguis*. *Arch Oral Biol.* 1973; 18:423–434. [PubMed: 4515970]
- Holmes AR, McNab R, Jenkinson HF. *Candida albicans* binding to the oral bacterium *Streptococcus gordonii* involves multiple adhesin-receptor interactions. *Infect Immun.* 1996; 64:4680–4685. [PubMed: 8890225]
- Hornby JM, Jensen EC, Lisec AD, et al. Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. *Appl Environ Microbiol.* 2001; 67:2982–2992. [PubMed: 11425711]
- Hoyer LL, Green CB, Oh SH, Zhao X. Discovering the secrets of the *Candida albicans* agglutinin-like sequence (ALS) gene family – a sticky pursuit. *Med Mycol.* 2008; 46:1–15. [PubMed: 17852717]
- Hoyer LL, Scherer S, Shatzman AR, Livi GP. *Candida albicans* ALS1: domains related to *Saccharomyces cerevisiae* sexual agglutinin separated by a repeating motif. *Mol Microbiol.* 1995; 15:39–54. [PubMed: 7752895]
- Hoyle BD, Costerton JW. Bacterial resistance to antibiotics: the role of biofilms. *Prog Drug Res.* 1991; 37:91–105. [PubMed: 1763187]
- Hussain H, Branny P, Allan E. A eukaryotic-type serine/threonine protein kinase is required for biofilm formation, genetic competence, and acid resistance in *Streptococcus mutans*. *J Bacteriol.* 2006; 188:1628–1632. [PubMed: 16452447]
- Ikegami A, Honma K, Sharma A, H.K. Kuramitsu HK. Multiple functions of the leucine-rich repeat protein LrrA of *Treponema denticola*. *Infect Immun.* 2004; 72:4619–4627. [PubMed: 15271922]
- Jakubovics NS, Gill S, Iobst, Vickerman MM, Kolenbrander PE. Regulation of gene expression in a mixed-genus community: stabilized arginine biosynthesis in *Streptococcus gordonii* by coaggregation with *Actinomyces naeslundii*. *J Bacteriol.* 2008; 190:3646–3657. [PubMed: 18359813]
- Jakubovics NS, Kolenbrander PE. The road to ruin: the formation of disease-associated oral biofilms. *Oral Diseases.* 2010; 16:729–739. [PubMed: 20646235]
- Jakubovics NS, Stromberg N, van Dolleweerd CJ, Kelly CG, Jenkinson HF. Differential binding specificities of oral streptococcal antigen I/II family adhesins for human or bacterial ligands. *Mol Microbiol.* 2005; 55:1591–1605. [PubMed: 15720563]
- Jenkinson HF. Beyond the oral microbiome. *Environ Microbiol.* 2011; 13:3077–3087. [PubMed: 21906224]
- Jenkinson HF, Lamont RJ. Oral microbial communities in sickness and in health. *Trends Microbiol.* 2005; 13:589–595. [PubMed: 16214341]
- Kaplan JB, Fine DH. Biofilm dispersal of *Neisseria subflava* and other phylogenetically diverse oral bacteria. *Appl Environ Microbiol.* 2002; 68:4943–4950. [PubMed: 12324342]
- Kerrigan SW, Jakubovics NS, Keane C, et al. Role of *Streptococcus gordonii* surface proteins SspA/SspB and Hsa in platelet function. *Infect Immun.* 2007; 75:5740–5747. [PubMed: 17893126]
- Kiley TB, Stanley-Wall NR. Post-translational control of *Bacillus subtilis* biofilm formation mediated by tyrosine phosphorylation. *Mol Microbiol.* 2010; 78:947–963. [PubMed: 20815827]
- Klein MI, DeBaz L, Agidi, et al. Dynamics of *Streptococcus mutans* transcriptome in response to starch and sucrose during biofilm development. *PLoS One.* 2010; 5:e13478. [PubMed: 20976057]
- Kobir A, Shi L, Boskovic A, Grangeasse C, Franjevic D, Mijakovic I. Protein phosphorylation in bacterial signal transduction. *Biochim Biophys Acta.* 2011; 1810:989–994. [PubMed: 21266190]
- Kolenbrander PE, Andersen RN, Blehert DS, Eglund PG, Foster JS, Palmer RJ Jr. Communication among oral bacteria. *Microbiol Mol Biol Rev.* 2002; 66:486–505. [PubMed: 12209001]
- Kolar SL, Nagarajan V, Oszmiana A, et al. NsaRS is a cell-envelope-stress-sensing two-component system of *Staphylococcus aureus*. *Microbiology.* 2011; 157:2206–2219. [PubMed: 21565927]
- Kolenbrander PE, Palmer RJ Jr, Rickard AH, Jakubovics NS, Chalmers NI, Diaz PI. Bacterial interactions and successions during plaque development. *Periodontol 2000.* 2006; 42:47–79. [PubMed: 16930306]

- Kolenbrander PE, Parrish KD, Andersen RN, Greenberg EP. Intergeneric coaggregation of oral *Treponema* spp. with *Fusobacterium* spp. and intrageneric coaggregation among *Fusobacterium* spp. *Infect Immun*. 1995; 63:4584–4588. [PubMed: 7591109]
- Kreth J, Merritt J, Shi W, Qi F. Competition and coexistence between *Streptococcus mutans* and *Streptococcus sanguinis* in the dental biofilm. *J Bacteriol*. 2005; 187:7193–7203. [PubMed: 16237003]
- Kreth J, Zhang Y, Herzberg MC. Streptococcal antagonism in oral biofilms: *Streptococcus sanguinis* and *Streptococcus gordonii* interference with *Streptococcus mutans*. *J Bacteriol*. 2008; 190:4632–4640. [PubMed: 18441055]
- Kuboniwa M, Amano A, Hashino E, et al. Distinct roles of long/short fimbriae and gingipains in homotypic biofilm development by *Porphyromonas gingivalis*. *BMC Microbiol*. 2009a; 9:105. [PubMed: 19470157]
- Kuboniwa M, Hendrickson EL, Xia Q, et al. Proteomics of *Porphyromonas gingivalis* within a model oral microbial community. *BMC Microbiol*. 2009b; 9:98. [PubMed: 19454014]
- Kuboniwa M, Lamont RJ. Subgingival biofilm formation. *Periodontol* 2000. 2010; 52:38–52. [PubMed: 20017794]
- Lamont RJ, Demuth DR, Davis CA, Malamud D, Rosan B. Salivary-agglutinin-mediated adherence of *Streptococcus mutans* to early plaque bacteria. *Infect Immun*. 1991; 59:3446–3450. [PubMed: 1894355]
- Lamont RJ, Hershey SG, Rosan B. Characterization of the adherence of *Porphyromonas gingivalis* to oral streptococci. *Oral Microbiol Immunol*. 1992; 7:193–197. [PubMed: 1328995]
- Lamont RJ, El-Sabaeny A, Park Y, Cook GS, Costerton JW, Demuth DR. Role of the *Streptococcus gordonii* SspB protein in the development of *Porphyromonas gingivalis* biofilms on streptococcal substrates. *Microbiology*. 2002; 148:1627–1636. [PubMed: 12055284]
- Lee DC, Jia Z. Emerging structural insights into bacterial tyrosine kinases. *Trends Biochem Sci*. 2009; 34:351–357. [PubMed: 19525115]
- Leid JG, Willson CJ, Shirliff ME, Hassett DJ, Parsek MR, Jeffers AK. The exopolysaccharide alginate protects *Pseudomonas aeruginosa* biofilm bacteria from IFN- γ -mediated macrophage killing. *J Immunol*. 2005; 175:7512–7518. [PubMed: 16301659]
- Liang X, Chen YY, Ruiz T, Wu H. New cell surface protein involved in biofilm formation by *Streptococcus parasanguinis*. *Infect Immun*. 2011; 79:3239–3248. [PubMed: 21576336]
- Lin X, Lamont RJ, Wu J, Xie H. Role of differential expression of streptococcal arginine deiminase in inhibition of *fimA* expression in *Porphyromonas gingivalis*. *J Bacteriol*. 2008; 190:4367–4371. [PubMed: 18408031]
- Lin X, Wu J, Xie H. *Porphyromonas gingivalis* minor fimbriae are required for cell-cell interactions. *Infect Immun*. 2006; 74:6011–6015. [PubMed: 16988281]
- Listgarten MA, Lai CH. Comparative ultrastructure of *Bacteroides melaninogenicus* subspecies. *J Periodontal Res*. 1979; 14:332–340. [PubMed: 44319]
- Liu Q, Fan J, Niu C, et al. The eukaryotic-type serine/threonine protein kinase Stk is required for biofilm formation and virulence in *Staphylococcus epidermidis*. *PLoS One*. 2011; 6:e25380. [PubMed: 21966513]
- Lo AW, Seers CA, Boyce JD, et al. Comparative transcriptomic analysis of *Porphyromonas gingivalis* biofilm and planktonic cells. *BMC Microbiol*. 2009; 9:18. [PubMed: 19175941]
- Madec E, Laszkiewicz A, Iwanicki A, Obuchowski M, Seror S. Characterization of a membrane-linked Ser/Thr protein kinase in *Bacillus subtilis* implicated in developmental processes. *Mol Microbiol*. 2002; 46:571–586. [PubMed: 12406230]
- Maeda K, Nagata H, Kuboniwa M, et al. Characterization of binding of *Streptococcus oralis* glyceraldehyde-3-phosphate dehydrogenase to *Porphyromonas gingivalis* major fimbriae. *Infect Immun*. 2004a; 72:5475–5477. [PubMed: 15322048]
- Maeda K, Nagata H, Yamamoto Y, et al. Glyceraldehyde-3-phosphate dehydrogenase of *Streptococcus oralis* functions as a coadhesin for *Porphyromonas gingivalis* major fimbriae. *Infect Immun*. 2004b; 72:1341–1348. [PubMed: 14977937]

- Maeda K, Tribble GD, Tucker CM, et al. A *Porphyromonas gingivalis* tyrosine phosphatase is a multifunctional regulator of virulence attributes. *Mol Microbiol.* 2008; 69:1153–1164. [PubMed: 18573179]
- Manuel SG, Ragunath C, Sait HB, Izano EA, Kaplan JB, Ramasubbu N. Role of active-site residues of dispersin B, a biofilm-releasing β -hexosaminidase from a periodontal pathogen, in substrate hydrolysis. *FEBS J.* 2007; 274:5987–5999. [PubMed: 17949435]
- Martins M, Uppuluri P, Thomas DP, et al. Presence of extracellular DNA in the *Candida albicans* biofilm matrix and its contribution to biofilms. *Mycopathologia.* 2010; 169:323–331. [PubMed: 20012895]
- Mattos-Graner RO, Porter KA, Smith DJ, Hosogi Y, Duncan MJ. Functional analysis of glucan binding protein B from *Streptococcus mutans*. *J Bacteriol.* 2006; 188:3813–3925. [PubMed: 16707674]
- Mayanagi G, Sato T, Shimauchi H, Takahashi N. Detection frequency of periodontitis-associated bacteria by polymerase chain reaction in subgingival and supragingival plaque of periodontitis and healthy subjects. *Oral Microbiol Immunol.* 2004; 19:379–385. [PubMed: 15491463]
- McNab R, Ford SK, El-Sabaeny A, Barbieri B, Cook GS, Lamont RJ. LuxS-based signaling in *Streptococcus gordonii*: autoinducer 2 controls carbohydrate metabolism and biofilm formation with *Porphyromonas gingivalis*. *J Bacteriol.* 2003; 185:274–284. [PubMed: 12486064]
- McNab R, Forbes H, Handley PS, Loach DM, Tannock GW, Jenkinson HF. Cell wall-anchored CshA polypeptide (259 kilodaltons) in *Streptococcus gordonii* forms surface fibrils that confer hydrophobic and adhesive properties. *J Bacteriol.* 1999; 181:3087–3095. [PubMed: 10322009]
- Murciano C, Moyes DL, Runglall M, et al. Evaluation of the role of *Candida albicans* agglutinin-like sequence (Als) proteins in human oral epithelial cell interactions. *PLoS One.* 2012; 7:e33362. [PubMed: 22428031]
- Nealson KH, Hastings JW. Bacterial bioluminescence: its control and ecological significance. *Microbiol Rev.* 1979; 43:496–518. [PubMed: 396467]
- Nie L, Wu G, Zhang W. Correlation between mRNA and protein abundance in *Desulfovibrio vulgaris*: a multiple regression to identify sources of variations. *Biochem Biophys Res Commun.* 2006; 339:603–610. [PubMed: 16310166]
- Nikitkova AE, Haase EM, Scannapieco FA. Effect of starch and amylase on the expression of amylase-binding protein A in *Streptococcus gordonii*. *Mol Oral Microbiol.* 2012; 27:284–294. [PubMed: 22759313]
- Nishikawa K, F. Yoshimura F, Duncan MJ. A regulation cascade controls expression of *Porphyromonas gingivalis* fimbriae via the FimR response regulator. *Mol Microbiol.* 2004; 54:546–560. [PubMed: 15469523]
- Nishiyama S, Murakami Y, Nagata H, Shizukuishi S, Kawagishi I, Yoshimura F. Involvement of minor components associated with the FimA fimbriae of *Porphyromonas gingivalis* in adhesive functions. *Microbiology.* 2007; 153:1916–1925. [PubMed: 17526848]
- Nobbs AH, Lamont RJ, Jenkinson HF. *Streptococcus* adherence and colonization. *Microbiol Mol Biol Rev.* 2009; 73:407–450. [PubMed: 19721085]
- Novak EA, Shao H, Daep CA, Demuth DR. Autoinducer-2 and QseC control biofilm formation and in vivo virulence of *Aggregatibacter actinomycetemcomitans*. *Infect Immun.* 2010; 78:2919–2926. [PubMed: 20404080]
- Nyvad B, Kilian M. Comparison of the initial streptococcal microflora on dental enamel in caries-active and in caries-inactive individuals. *Caries Res.* 1990; 24:267–272. [PubMed: 2276164]
- Okahashi N, Nakata M, Sakurai A, et al. Pili of oral *Streptococcus sanguinis* bind to fibronectin and contribute to cell adhesion. *Biochem Biophys Res Commun.* 2010; 391:1192–1196. [PubMed: 20004645]
- O'Toole GA, Kolter R. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol.* 1998; 28:449–461. [PubMed: 9632250]
- Palmer RJ Jr, Gordon SM, Cisar JO, Kolenbrander PE. Coaggregation-mediated interactions of streptococci and actinomyces detected in initial human dental plaque. *J Bacteriol.* 2003; 185:3400–3409. [PubMed: 12754239]

- Palmer RJ Jr, Kazmerzak K, Hansen MC, Kolenbrander PE. Mutualism versus independence: strategies of mixed-species oral biofilms in vitro using saliva as the sole nutrient source. *Infect Immun.* 2001; 69:5794–5804. [PubMed: 11500457]
- Pammi M, Liang R, Hicks JM, Barrish J, Versalovic J. Farnesol decreases biofilms of *Staphylococcus epidermidis* and exhibits synergy with nafcillin and vancomycin. *Pediatr Res.* 2011; 70:578–583. [PubMed: 21857375]
- Park Y, Simionato MR, Sekiya K, et al. Short fimbriae of *Porphyromonas gingivalis* and their role in coadhesion with *Streptococcus gordonii*. *Infect Immun.* 2005; 73:3983–3989. [PubMed: 15972485]
- Percival RS, Challacombe SJ, Marsh PD. Age-related microbiological changes in the salivary and plaque microflora of healthy adults. *J Med Microbiol.* 1991; 35:5–11. [PubMed: 2072378]
- Perisasamy S, Kolenbrander PE. Central role of the early colonizer *Veillonella* sp. in establishing multispecies biofilm communities with initial, middle, and late colonizers of enamel. *J Bacteriol.* 2010; 192:2965–2972. [PubMed: 20154130]
- Petersen HJ, Keane C, Jenkinson HF, et al. Human platelets recognize a novel surface protein, PadA, on *Streptococcus gordonii* through a unique interaction involving fibrinogen receptor GPIIb/IIIa. *Infect Immun.* 2010; 78:413–422. [PubMed: 19884334]
- Pham TK, Roy S, Noirel J, Douglas I, Wright PC, Stafford GP. A quantitative proteomic analysis of biofilm adaptation by the periodontal pathogen *Tannerella forsythia*. *Proteomics.* 2010; 10:3130–3141. [PubMed: 20806225]
- Prakobphol A, Xu F, Hoang VM, et al. Salivary agglutinin, which binds *Streptococcus mutans* and *Helicobacter pylori* is the lung scavenger receptor cysteine-rich protein gp-340. *J Biol Chem.* 2000; 275:39860–39866. [PubMed: 11007786]
- Ramboarina S, Garnett JA, Zhou M, et al. Structural insights into serine-rich fimbriae from Gram-positive bacteria. *J Biol Chem.* 2010; 285:32446–32457. [PubMed: 20584910]
- Rickard AH, Campagna SR, Kolenbrander PE. Autoinducer-2 is produced in saliva-fed flow conditions relevant to natural oral biofilms. *J Appl Microbiol.* 2008; 105:2096–2103. [PubMed: 19120655]
- Rickard AH, Palmer RJ Jr, Blehert DS, et al. Autoinducer 2: a concentration-dependent signal for mutualistic bacterial biofilm growth. *Mol Microbiol.* 2006; 60:1446–1456. [PubMed: 16796680]
- Roberts AP, Mullany P. Oral biofilms: a reservoir of transferable, bacterial, antimicrobial resistance. *Expert Rev Anti Infect Ther.* 2010; 8:1441–1450. [PubMed: 21133668]
- Rosen G, Genzler T, Sela MN. Coaggregation of *Treponema denticola* with *Porphyromonas gingivalis* and *Fusobacterium nucleatum* is mediated by the major outer sheath protein of *Treponema denticola*. *FEMS Microbiol Lett.* 2008; 289:59–66. [PubMed: 19054094]
- Rosen G, Sela MN. Coaggregation of *Porphyromonas gingivalis* and *Fusobacterium nucleatum* PK 1594 is mediated by capsular polysaccharide and lipopolysaccharide. *FEMS Microbiol Lett.* 2006; 256:304–310. [PubMed: 16499621]
- Rupani D, Izano EA, Schreiner HC, Fine DH, Kaplan JB. *Aggregatibacter actinomycetemcomitans* serotype f O-polysaccharide mediates coaggregation with *Fusobacterium nucleatum*. *Oral Microbiol Immunol.* 2008; 23:127–130. [PubMed: 18279180]
- Sauer K, Camper AK, Ehrlich GD, Costerton JW, Davies DG. *Pseudomonas aeruginosa* displays multiple phenotypes during development as a biofilm. *J Bacteriol.* 2002; 184:1140–1154. [PubMed: 11807075]
- Scannapieco FA. Saliva-bacterium interactions in oral microbial ecology. *Crit Rev Oral Biol Med.* 1994; 2:203–248. [PubMed: 7703323]
- Shao H, Lamont RJ, Demuth DR. Autoinducer 2 is required for biofilm growth of *Aggregatibacter (Actinobacillus) actinomycetemcomitans*. *Infect Immun.* 2007; 75:4211–4218. [PubMed: 17591788]
- Sharma A, Inagaki S, Sigurdson W, Kuramitsu HK. Synergy between *Tannerella forsythia* and *Fusobacterium nucleatum* in biofilm formation. *Oral Microbiol Immunol.* 2005; 20:39–42. [PubMed: 15612944]

- Silverman RJ, Nobbs AH, Vickerman MM, Barbour ME, Jenkinson HF. Interaction of *Candida albicans* cell wall Als3 protein with *Streptococcus gordonii* SspB adhesin promotes development of mixed-species communities. *Infect Immun*. 2010; 78:4644–4652. [PubMed: 20805332]
- Simionato MR, Tucker CM, Kuboniwa M, et al. *Porphyromonas gingivalis* genes involved in community development with *Streptococcus gordonii*. *Infect Immun*. 2006; 74:6419–6428. [PubMed: 16923784]
- Slots J, Gibbons RJ. Attachment of *Bacteroides melaninogenicus* subsp. asaccharolyticus to oral surfaces and its possible role in colonization of the mouth and of periodontal pockets. *Infect Immun*. 1978; 19:254–264. [PubMed: 24002]
- Son M, Ahn SJ, Guo Q, Burne RA, Hagen SJ. Microfluidic study of competence regulation in *Streptococcus mutans*: environmental inputs modulate bimodal and unimodal expression of *comX*. *Mol Microbiol*. 2012 [pages to be added in proof].
- Srinivasa K, Kim J, Yee S, Kim W, Choi W. A MAP kinase pathway is implicated in the pseudohyphal induction by hydrogen peroxide in *Candida albicans*. *Mol Cells*. 2012; 33:183–193. [PubMed: 22358510]
- Sun J, Daniel R, Wagner-Dobler I, Zeng AP. Is autoinducer-2 a universal signal for interspecies communication: a comparative genomic and phylogenetic analysis of the synthesis and signal transduction pathways. *BMC Evol Biol*. 2004; 4:36. [PubMed: 15456522]
- Suntharalingam P, Cvitkovitch DG. Quorum sensing in streptococcal biofilm formation. *Trends Microbiol*. 2005; 13:3–6. [PubMed: 15639624]
- Swift S, Downie JS, Whitehead NA, Barnard AM, Salmond GP, Williams P. Quorum sensing as a population-dependent determinant of bacterial physiology. *Adv Microbial Physiol*. 2001; 34:199–270.
- Takahashi N. Acid-neutralizing activity during amino acid fermentation by *Porphyromonas gingivalis* *Prevotella intermedia* and *Fusobacterium nucleatum*. *Oral Microbiol Immunol*. 2003; 18:109–113. [PubMed: 12654101]
- Ueda A, Wood TK. Connecting quorum sensing, c-di-GMP, pel polysaccharide, and biofilm formation in *Pseudomonas aeruginosa* through tyrosine phosphatase TpbA (PA3885). *PLoS Pathog*. 2009; 5:e1000483. [PubMed: 19543378]
- Ueda A, Wood TK. Tyrosine phosphatase TpbA of *Pseudomonas aeruginosa* controls extracellular DNA via cyclic diguanylic acid concentrations. *Environ Microbiol*. 2010; 2:449–455. [PubMed: 21552365]
- Uppuluri P, Pierce CG, Thomas DP, Bubeck SS, Saville SP, Lopez-Ribot JL. The transcriptional regulator Nrg1p controls *Candida albicans* biofilm formation and dispersion. *Eukaryot Cell*. 2010; 9:1531–1537. [PubMed: 20709787]
- van Winkelhoff AJ, Loos BG, van der Reijden WA, van der Velden U. *Porphyromonas gingivalis* *Bacteroides forsythus* and other putative periodontal pathogens in subjects with and without periodontal destruction. *J Clin Periodontol*. 2002; 29:1023–1028. [PubMed: 12472995]
- Wang BY, Kuramitsu HK. Interactions between oral bacteria: inhibition of *Streptococcus mutans* bacteriocin production by *Streptococcus gordonii*. *Appl Environ Microbiol*. 2005; 71:354–362. [PubMed: 15640209]
- Wang BY, Wu J, Lamont RJ, Lin X, Xie H. Negative correlation of distributions of *Streptococcus cristatus* and *Porphyromonas gingivalis* in subgingival plaque. *J Clin Microbiol*. 2009; 47:3902–3906. [PubMed: 19846640]
- Wen ZT, Nguyen AH, Bitoun JP, Abranches J, Baker HV, Burne RA. Transcriptome analysis of LuxS-deficient *Streptococcus mutans* grown in biofilms. *Mol Oral Microbiol*. 2011; 26:2–18. [PubMed: 21214869]
- Whitchurch CB, Tolker-Nielsen T, Ragas PC, Mattick JS. Extracellular DNA required for bacterial biofilm formation. *Science*. 2002; 295:1487. [PubMed: 11859186]
- Whitmore SE, Lamont RJ. The pathogenic persona of community-associated oral streptococci. *Mol Microbiol*. 2011; 81:305–314. [PubMed: 21635580]
- Xie H, Cai S, Lamont RJ. Environmental regulation of fimbrial gene expression in *Porphyromonas gingivalis*. *Infect Immun*. 1997; 65:2265–2271. [PubMed: 9169762]

- Xie H, Chung WO, Park Y, Lamont RJ. Regulation of the *Porphyromonas gingivalis* *fimA* (fimbriin) gene. *Infect Immun*. 2000; 68:6574–6579. [PubMed: 11083767]
- Xie H, Lin X, Wang BY, Wu J, Lamont RJ. Identification of a signalling molecule involved in bacterial intergeneric communication. *Microbiology*. 2007; 153:3228–3234. [PubMed: 17906122]
- Xie H, Hong J, Sharma A, Wang B-Y. *Streptococcus cristatus* ArcA interferes with *Porphyromonas gingivalis* pathogenicity in mice. *J Perio Res*. 2012; 47:578–583.
- Ximenez-Fyvie LA, Haffajee AD, Socransky SS. Comparison of the microbiota of supra- and subgingival plaque in health and periodontitis. *J Clin Periodontol*. 2000; 27:648–657. [PubMed: 10983598]
- Xu XL, Lee RT, Fang HM, et al. Bacterial peptidoglycan triggers *Candida albicans* hyphal growth by directly activating the adenyl cyclase Cyr1p. *Cell Host Microbe*. 2008; 4:28–39. [PubMed: 18621008]
- Yamamoto R, Noiri Y, Yamaguchi M, Asahi Y, Maezono H, Ebisu S. Time course of gene expression during *Porphyromonas gingivalis* strain ATCC 33277 biofilm formation. *Appl Environ Microbiol*. 2011; 77:6733–6736. [PubMed: 21803908]
- Yoshida Y, Palmer RJ, Yang J, Kolenbrander PE, Cisar JO. Streptococcal receptor polysaccharides: recognition molecules for oral biofilm formation. *BMC Oral Health*. 2006; 6(Suppl 1):S12. [PubMed: 16934113]
- Yoshimura F, Takahashi K, Nodasaka Y, Suzuki T. Purification and characterization of a novel type of fimbriae from the oral anaerobe. *Bacteroides gingivalis*. *J Bacteriol*. 1984; 160:949–957.
- Zhang Y, Whiteley M, Kreth J, et al. The two-component system BfrAB regulates expression of ABC transporters in *Streptococcus gordonii* and *Streptococcus sanguinis*. *Microbiology*. 2009; 155:165–173. [PubMed: 19118357]
- Zhao X, Oh SH, Cheng G, et al. ALS3 and ALS8 represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology*. 2004; 150:2415–2428. [PubMed: 15256583]
- Zhu L, Kreth J. Role of *Streptococcus mutans* eukaryotic-type serine/threonine protein kinase in interspecies interactions with *Streptococcus sanguinis*. *Arch Oral Biol*. 2010; 55:385–390. [PubMed: 20381010]

Fig. 1.

Bacterial cell-to-cell interactions in the dental plaque biofilm. One example of mutualism is displayed between *P. gingivalis* and *T. denticola*. Each releases a metabolic factor that the other can utilize, thus enhancing the growth of both species. Lines linking the microorganisms represent adhesive interactions. Aa = *Aggregatibacter actinomycetemcomitans*

Fig. 2.

The interaction of *P. gingivalis* with *S. gordonii* depends upon two sets of adhesin-receptor pairs. The major and minor fimbriae of *P. gingivalis* bind glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and SspA/B on the surface of *S. gordonii*, respectively. The domains of SspA/B that interact with the minor fimbriae subunit, Mfa1, are highlighted in the yellow ovals. These domains reside within the SspA/B adherence region (BAR) and are required to maintain the contacts between the two species in the oral cavity.

Fig. 3. Phosphorylation events within a bacterial cell are complex and dynamic. The two-component system in *A. actinomycetemcomitans* is induced by AI-2, with the loss of this TCS resulting in diminished biofilm formation. Tyrosine kinases in Gram-negative bacteria such as Wzc in *E. coli* have been shown to regulate a variety of targets including DNA-binding proteins, capsule synthesis genes and the heat shock response. The phosphatase Ltp1 in *P. gingivalis* has been shown to affect a number of cellular activities including dual-species community formation, reduced EPS synthesis and hemin uptake.

Communication Circuits (physical and chemical)

Fig. 4.

Communication circuits between *C. albicans* and oral streptococci. The diagram depicts self- or non-self control of the yeast to hypha transition, with bacterial products variously impacting on morphogenesis and biofilm formation, and *C. albicans* products positively or negatively influencing bacterial growth or biofilm formation (see text for discussion).