Published in final edited form as:

ACS Comb Sci. 2012 October 8; 14(10): 527–530. doi:10.1021/co300076k.

Resin-Supported Catalysts for CuAAC Click Reactions in Aqueous or Organic Solvents

Stanislav I. Presolski, Sreeman K. Mamidyala, Florian Manzenrieder, and M.G. Finn

Department of Chemistry, The Scripps Research Institute, 10550 N. Torrey Pines Rd., La Jolla, CA 92037, USA, mgfinn@scripps.edu

Abstract

The copper-catalyzed azide-alkyne cycloaddition click reaction is a valuable process for the synthesis of libraries of drug candidates, derivatized polymers and materials, and a wide variety of other functional molecules. In some circumstances, the removal of the copper catalyst is both necessary and inconvenient. We describe here two immobilized forms of a Cu-binding ligand that has been shown to accelerate triazole formation under many different conditions, using different resin supports that are appropriate for aqueous or organic solvents. Copper leaching from these resins was modest, allowing them to be reused in many reaction/filtration cycles without recharging with metal ion. The utility of this catalyst form was demonstrated in the convenient synthesis of 20 N-acetylgalactosamine derivatives for biological testing.

Introduction

The copper-catalyzed azide-alkyne cycloaddition (CuAAC) reaction^{1,2} is well suited to many solution-phase applications with the simple use of Cu(I) salts, either added directly or generated by reduction of Cu(II) or oxidation of Cu(0). This process is approximately one million times faster than the very sluggish uncatalyzed [3+2] cycloaddition with unactivated alkynes and azides.³ When even faster reaction rates are needed, certain heterocyclic chelating agents have been shown to provide strong ligand-accelerated catalysis.^{4–6} The intrinsically tolerant nature of Cu catalysts to variations in reaction conditions has allowed the CuAAC process to be applied in an extraordinarily wide range of situations.

Reliable methods for the removal of the catalyst from the reaction mixture are desirable. In the case of CuAAC, which is commonly applied to the synthesis of materials intended for biological application, this can be of great concern because of the perception that copper ions are toxic to cells and organisms. In most cases, we find that washing with or dialysis against solutions of EDTA removes even small traces of copper ions from CuAAC reaction mixtures and isolated products. However, this procedure is not always convenient, especially in parallel or combinatorial synthesis.

A common strategy is to immobilize the catalyst on a support that allows access to reactants and can be easily separated from the reaction mixture upon completion. A handful of reports have appeared of heterogeneous CuAAC catalysts, but this field is not yet fully developed.^{7–17} (See Supporting Information for an annotated list of these contributions.) We describe here the synthesis and reactivity of two versions of a system first described by Chan and Fokin, in which an accelerating chelating ligand is covalent attached to a resin material commonly used for solid-phase organic and peptide synthesis.⁸

Results and Discussion

We have described a highly versatile mono(benzimidazolymethyl)-bis(pyridylmethyl) amine ligand (designated **BimPy₂**) that performs well in both aqueous and coordinating organic solvents.⁶ Its ease of synthesis and readily functionalized single attachment point made it attractive for immobilization, which we performed on two different solid supports as shown in Figure 1. The undecyl alcohol BimPy₂ derivative **1** was attached to standard 2-chlorotrityl polystyrene resin (1% divinylbenzene crosslinked, available from several suppliers, 1.04 mmol/g in the exemplary case here, but a variety of loadings between 0.5 and 2 mmol/g work equally well), followed by capping with methanol, to provide the immobilized **PS-BimPy₂**. The mass increase of the resin beads indicated a loading of 290 μmol/g. Treatment of the resin with trifluoroacetic acid to cleave the trityl connector, followed by MS analysis, revealed that only the desired ligand had been attached. A negative control resin (**PS-OMe**) was prepared using methanol in place of **1**. To provide immobilized catalyst capability for polar and hydrophilic media, the BimPy₂ ligand was also attached to Nova-PEG-amine resin (Novabiochem #855126) via an 11-carbon aldehyde (**2**) by reductive amination, to give the **NovaPEG-BimPy₂** ligand. The unreacted primary amines and the secondary amines resulting from the coupling were capped with acetic anhydride. A loading of 230 μmol/g was indicated by the mass increase from the attachment reaction, representing occupancy of approximately 30% of the resin amine groups. Reductive amination with cinnamaldehyde provide the negative control material in this case (**NovaPEG-Cin**). The undecyl hydrocarbon tether was selected in order to allow the ligand enhanced mobility in the swollen resin (gel) state, but other chain lengths and types would likely also be effective. We also explored Sepharose-epoxide as a water-swelling solid support, but the attachment yields were low and the material had insufficient mechanical stability to stand up to the agitation required in repeat reactions.

The polystyrene-based catalysts were initially tested by adding CuSO₄, sodium ascorbate, and small amounts of resin (10 mg) to CuAAC reactions in 1.5 mL 95:5 DMF:H₂O, a solvent system that effectively swells the resin. Yields of triazole **3** were determined the following day by HPLC-MS against a quantitative calibration curve (Figure 2). The reactions were performed in fritted syringes, allowing for easy separation of the solution from the solid-phase material. For this length of reaction time, the ligand-free reaction produced relatively high yield but failed to reach completion, presumably because of competitive oxidative inactivation of the Cu(I) catalyst. In contrast, the ligand-bearing resin provided complete reaction, reflecting an accelerated reaction rate.

After removal of the reaction solvent, each of these resins was recharged with another batch of azide, alkyne, and sodium ascorbate, but no additional copper source. Only those resins bearing the BimPy₂ ligand produced triazole, showing that the ligand is required to retain the catalytic metal center. The **PS-BimPy₂** catalyst was subjected to 13 cycles of CuAAC reaction, filtration, and re-charge with azide, alkyne, and ascorbate. The Cu content of the filtered solutions was measured after each cycle by inductively coupled plasma atomic emission spectroscopy (ICP-AES). As shown in Figure 2A, all of these reactions went to completion, and an average of 3% of the Cu was lost in each cycle (approximately 3 μM of Cu in the solution), leaving more than 60% of the initial Cu retained on the resin after the 13th reaction. Note that low-micromolar concentrations of Cu(I) without accelerating ligand do not catalyze the CuAAC reaction under these conditions (data not shown).

When the solvent was changed from DMF to methanol, the overnight reactions catalyzed by the polystyrene-immobilized ligand did not reach completion, probably due to the poor swelling capacity of the solid support. The ligand-modified NovaPEG resin served well in this case, being swelled in methanol (and a wide variety of other solvents) and mediating

complete formation of triazole in two consecutive reactions (Figure 2B). In the three subsequent runs the yields were about 85%, after which performance declined to a greater degree without Cu recharging. Under these “synthetic” conditions (100 mM concentrations of substrates, organic solvent), the greatest amount of Cu contamination was observed, with approximately 400 μM Cu observed in solution after the first reaction. This is likely due in part to the high concentration of triazole formed in this reaction, which competes for Cu with the immobilized ligand. The **NovaPEG-Cin** resin initially induced modest levels of triazole formation (27% yield with 1 mol% Cu), presumably due to some acceleration provided by adventitious Cu ligands such as the amide groups installed on the resin. Following the course of the NovaPEG-based reactions more closely revealed supported BimPy₂ to be far more active than the control acylated cinnamoylamine-bearing resin (Figure 2C). When no resin was employed, the reaction gave only a 5% yield of triazole before exhaustion of the reducing agent needed to support CuAAC.

The NovaPEG resin has excellent swelling capacity in both dichloromethane and water. In the former solvent, the use of CuBr and CuI gave incomplete reactions that contained substantial quantities of copper, suggesting that counterion substitution is important in noncoordinating solvents.¹⁸ In aqueous reactions performed under typical bioconjugation conditions, however, 10 mg of **NovaPEG-BimPy₂** in 90% H₂O, 10% DMSO with 100 μM CuSO₄, and 5 mM NaAsc catalyzed the complete conversion of 100 μM BnN₃ to triazole, when 200 μM PhCCH were used. Low levels of Cu were observed in solution (approximately 5.4 μM , 0.34 ppm, in each cycle). Such minimal leaching makes the resin-immobilized catalyst an excellent choice for library syntheses of compounds that can be used in many biological tests without concern for significant copper contamination.

The **NovaPEG-BimPy₂** ligand was tested in preparatory-scale reactions involving glycan precursors **4** and **6** with ten azides and alkynes each, including aromatic, aliphatic, and benzylic examples (Figure 3).¹⁹ All triazole products were isolated in >90% yields after filtration and flash column chromatography, in purities >95% as judged by ¹H NMR.

Conclusions

We show that the resin immobilization of the BimPy₂ ligand provides readily accessible and useful catalysts for the CuAAC reaction in both organic and aqueous solvents. These systems show convenient properties of rate, metal retention, and separability from soluble reaction components. The previously reported and very useful resin of Chan and Fokin⁸ employs the tris(triazolylmethyl)amine ligand, which is a weaker binder of Cu^I than BimPy₂.⁶ For this reason, this ligand is not well suited to reactions in strongly coordinating solvents. If the user intends to perform reactions in DMSO, DMF, or similar solvents, **PS-BimPy₂** is an effective and inexpensive catalyst. When polar protic and aqueous solvent systems are desired, we recommend **NovaPEG-BimPy₂** due to its superior swelling properties. For all other solvents except acetonitrile (which gives poor results with this class of catalysts; see Figure 5 of reference 6), both ligand-functionalized resins should provide good results if the solvent swells the resin to a sufficient degree.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (R56 AI090832), The Skaggs Institute for Chemical Biology, and Pfizer Global R&D. We thank Dr. Vu Hong for helpful discussions.

References

1. Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. A Stepwise Huisgen Cycloaddition Process: Copper(I)-Catalyzed Regioselective Ligation of Azides and Terminal Alkynes. *Angew. Chem., Int. Ed.* 2002; 41:2596–2599.
2. Tornøe CW, Christensen C, Meldal M. Peptidotriazoles on Solid Phase: [1,2,3]-Triazoles by Regiospecific Copper(I)-Catalyzed 1,3-Dipolar Cycloadditions of Terminal Alkynes to Azides. *J. Org. Chem.* 2002; 67:3057–3062. [PubMed: 11975567]
3. Lewis WG, Green LG, Grynszpan F, Radic Z, Carlier PR, Taylor P, Finn MG, Sharpless KB. Click Chemistry *in situ*: Acetylcholinesterase as a Reaction Vessel for the Selective Assembly of a Femtomolar Inhibitor from an Array of Building Blocks. *Angew. Chem. Int. Ed.* 2002; 41:1053–1057.
4. Chan TR, Hilgraf R, Sharpless KB, Fokin VV. Polytriazoles as Copper(I)-Stabilizing Ligands in Catalysis. *Org. Lett.* 2004; 6:2853–2855. [PubMed: 15330631]
5. Rodionov VO, Presolski S, Díaz DD, Fokin VV, Finn MG. Ligand-Accelerated Cu-Catalyzed Azide-Alkyne Cycloaddition: a Mechanistic Report. *J. Am. Chem. Soc.* 2007; 129:12705–12712. and references therein. [PubMed: 17914817]
6. Presolski S, Hong V, Cho S-H, Finn MG. Tailored Ligand Acceleration of the Cu-Catalyzed Azide-Alkyne Cycloaddition Reaction. *J. Am. Chem. Soc.* 2010; 132:14570–14576. [PubMed: 20863116]
7. Lipshutz BH, Taft BR. Heterogeneous Copper-in-Charcoal-Catalyzed Click Chemistry. *Angew. Chem., Int. Ed.* 2006; 45:8235–8238.
8. Chan TR, Fokin VV. Polymer-Supported Copper(I) Catalysts for the Experimentally Simplified Azide-Alkyne Cycloaddition. *QSAR Comb. Sci.* 2007; 26:1274–1279.
9. Li P, Wang L, Zhang Y. SiO₂-NHC-Cu(I): an efficient and reusable catalyst for [3+2] cycloaddition of organic azides and terminal alkynes under solvent-free reaction conditions at room temperature. *Tetrahedron.* 2008; 64:10825–10830.
10. Dervaux B, Van Camp W, Du Prez FE. Cryogels as Novel Solid-Supported Copper Catalyst System for Click Chemistry and their use in Column Reactors. *Polym. Preprints.* 2008; 49:692–693.
11. Jialia I, Elamari H, Meganem F, Herscovici J, Girard C. Copper(I)-doped Wyoming's montmorillonite for the synthesis of disubstituted 1,2,3-triazoles. *Tetrahedron Lett.* 2008; 49:6756–6758.
12. Namitharan K, Kumarraja M, Pitchumani K. CuII-Hydrotalcite as an Efficient Heterogeneous Catalyst for Huisgen [3+2] Cycloaddition. *Chem. Eur. J.* 2009; 15:2755–2758. [PubMed: 19197920]
13. Chichigrovsky I, Primo A, Gonzalez P, Molvinger K, Robitzer M, Quignard F, Taran F. Functionalized Chitosan as a Green, Recyclable, Biopolymer-Supported Catalyst for the [3+2] Huisgen Cycloaddition. *Angew. Chem. Int. Ed.* 2009; 48:5916–5920.
14. Jialia I, Meganem F, Herscovici J, Girard C. "Flash" Solvent-free Synthesis of Triazoles Using a Supported Catalyst. *Molecules.* 2009; 14:528–539. [PubMed: 19169200]
15. Bonami L, Van Camp W, Van Rijckegem D, Du Prez FE. Facile Access to an Efficient Solid-Supported Click Catalyst System Based on Poly(ethyleneimine). *Macromol. Rapid Commun.* 2009; 30:34–38. [PubMed: 21706535]
16. Buckley BR, Dann SE, Harris DP, Heaney H, Stubbs EC. Alkynylcopper(I) polymers and their use in a mechanistic study of alkyne-azide click reactions. *Chem. Commun.* 2010; 46:2274–2276.
17. He Y, Bian Z, Kang C, Cheng Y, Gao L. Chiral binaphthylbisbipyridine-based copper(I) coordination polymer gels as supramolecular catalysts. *Chem. Commun.* 2010; 46:3532–3534.
18. Hein JE, Tripp JC, Krasnova LB, Sharpless KB, Fokin VV. Copper(I)-catalyzed cycloaddition of organic azides and 1-iodoalkynes. *Angew. Chem. Int. Ed.* 2009; 48:8018–8021.
19. Mamidyala SK, Dutta S, Chrnyk BA, Préville C, Wang C, Withka JM, McColl A, Subashi TA, Hawrylik SJ, Griffor MC, Kim S, Pfefferkorn JA, Price DA, Menhaji-Klotz E, Mascitti V, Finn MG. Glycomimetic Ligands for the Human Asialoglycoprotein Receptor. *J. Am. Chem. Soc.* 2012; 134:1978–1981. [PubMed: 22280495]

Figure 1.
Synthesis of resin-supported BimPy₂ CuAAC-accelerating ligands.

Figure 2.

Screening of activity of resin-bound ligands for the model CuAAC reaction shown. (A,B) Amounts of triazole **3** formed (solid circles), and amount of Cu remaining on the resin (open circles), in sequential cycles of CuAAC reaction using 10 mg (A) **PS-BimPy₂** or **PS-OMe** with an initial charge of 100 μM CuSO_4 , or (B) **NovaPEG-BimPy₂** or **NovaPEG-Cin** with initial charge of 1 mM CuSO_4 . Reaction conditions: (A) 1.25 mM BnN_3 , 2.5 mM PhCCH , 10 mM Na ascorbate, 95:5 DMF:H₂O, RT, 12 h. (B) 100 mM BnN_3 , 100 mM PhCCH , 20 mM Na ascorbate, 90:10 MeOH:H₂O, RT, 12 h. Each reaction was drained, washed 3 times with methanol, air dried, and recharged with azide, alkyne, and ascorbate to initiate the next cycle. (C) Time course for reactions mediated by 10 mg of the indicated NovaPEG derivatives; conditions as in the first reaction of plot (B), except for the omission of resin or CuSO_4 as noted.

Figure 3.
Parallel CuAAC reactions performed with **NovaPEG-BimPy₂** catalyst.