Published in final edited form as:

Cancer. 2012 August 1; 118(15): 3766–3775. doi:10.1002/cncr.26702.

Yoga for persistent fatigue in breast cancer survivors: A randomized controlled trial

Julienne E. Bower, Ph.D.^{1,2,3,4}, Deborah Garet, M.P.H.³, Beth Sternlieb⁵, Patricia A. Ganz, M.D.^{4,6}, Michael R. Irwin, M.D.^{1,2,3,4}, Richard Olmstead, Ph.D.^{2,3}, and Gail Greendale, M.D.⁷

¹ UCLA Department of Psychology, Geffen School of Medicine at UCLA

² Department of Psychiatry and Biobehavioral Sciences, Geffen School of Medicine at UCLA

³ Cousins Center for Psychoneuroimmunology, Semel Institute at UCLA

⁴ Division of Cancer Prevention and Control Research, Jonsson Comprehensive Cancer Center at UCLA

⁵ Pediatric Pain Program, Mattel Children's Hospital at UCLA

⁶ UCLA Schools of Medicine and Public Health

⁷ UCLA Department of Geriatrics

Abstract

Background—Cancer-related fatigue afflicts up to one-third of breast cancer survivors, yet there are no empirically-validated treatments for this symptom.

Methods—We performed a two-group RCT to determine the feasibility and efficacy of an Iyengar yoga intervention for breast cancer survivors with persistent post-treatment fatigue. Participants were breast cancer patients who had completed cancer treatments (other than endocrine therapy) at least 6 months prior to enrollment, reported significant cancer-related fatigue, and had no other medical conditions that would account for fatigue symptoms or interfere with yoga practice. Block randomization was used to assign participants to a 12-week Iyengar-based yoga intervention or to 12 weeks of health education (control). The primary outcome was change in fatigue measured at baseline, immediately post-treatment, and 3 months after treatment completion. Additional outcomes included changes in vigor, depressive symptoms, sleep, perceived stress, and physical performance. Intent to treat analyses were conducted with all randomized participants using linear mixed models.

Results—Thirty-one women were randomly assigned to yoga ($n = 16$) or health education ($n = 15$). Fatigue severity declined significantly from baseline to post-treatment and over a 3 month follow-up in the yoga group relative to controls ($P = .032$). In addition, the yoga group showed significant increases in vigor relative to controls ($P = .011$). Both groups showed positive changes in depressive symptoms and perceived stress ($P < .05$). No significant changes in sleep or physical performance were observed.

Conclusions—A targeted yoga intervention led to significant improvements in fatigue and vigor among breast cancer survivors with persistent fatigue symptoms.

INTRODUCTION

Approximately one-third of cancer survivors experience persistent fatigue of unknown origin, causing significant impairment in quality of life (1;2). Despite the prevalence and impact of this symptom, there are currently no empirically validated treatments for persistent cancer-related fatigue. To date, only two published behavioral intervention trials have specifically targeted cancer survivors with persistent fatigue (3;4). Other psychological (e.g., stress reduction) and activity-based (e.g., exercise) interventions have reduced fatigue in cancer populations (5-7); however, because these trials have not targeted fatigued patients, the feasibility and efficacy of these approaches for treating post-cancer fatigue is unclear. Indeed, fatigue is a significant barrier to participation in exercise programs for cancer survivors (8).

Mind-body interventions such as yoga are a promising approach for treating cancer-related fatigue. Yoga involves physical postures (*asanas*) that develop strength and flexibility, and promote relaxation. Yoga is also a meditative practice because the practitioner focuses on the body and breath in each pose (9). A growing body of research indicates that yoga has beneficial effects on physical and behavioral outcomes in cancer patients and survivors (10-12), including improvements in quality of life, mood, and fatigue (13-18). However, as with the behavioral interventions, none of the published yoga trials have targeted patients with fatigue. Moreover, very few of these trials have included an active control group to control for attention, group support, and other non-specific components of the treatment.

The primary goal of this randomized controlled trial (RCT) was to examine the feasibility and efficacy of an Iyengar yoga intervention for breast cancer survivors with persistent post-treatment fatigue relative to health education control. Based on promising results from a small, uncontrolled trial conducted by our group (19), we hypothesized that this intervention would lead to significant improvements in fatigue and vigor. Effects on secondary outcomes, including depressive symptoms, sleep disturbance, stress, and physical performance, were also assessed.

METHODS

Design

The study was a single-center, two armed RCT. The UCLA Institutional Review Board approved the study procedures, and written informed consent was obtained from all participants.

Participants

Breast cancer survivors with persistent post-treatment fatigue were recruited through multiple mechanisms, including tumor registry mailings, newspaper advertisements, and flyers distributed at cancer-related and community locations. Inclusion criteria were: 1) originally diagnosed with Stage 0 – II breast cancer; 2) completed local and/or adjuvant cancer therapy (with the exception of hormonal therapy) at least 6 months previously; 3) age 40 – 65; 4) post-menopausal; 5) no other cancer in last 5 years; and 6) experiencing persistent cancer-related fatigue. We focused on women with early-stage breast cancer to reduce heterogeneity in disease- and treatment-related factors impacting fatigue and its biological and psychological correlates. We focused on women at least 6 months post treatment completion to ensure that participants had persistent post-treatment fatigue and to reduce confounding residual effects of cancer treatment on biological parameters. The presence of cancer-related fatigue was indicated by scores of 50 or below on the SF-36 vitality scale, a reliable and valid measure of energy/fatigue in the past month (20), and self-report that fatigue was a consequence of cancer or cancer therapy. In our previous research,

breast cancer survivors who scored at or below 50 on the SF-36 vitality scale showed behavioral, immune, and neuroendocrine alterations relative to women who scored above 50, supporting this classification (1;21-26).

Exclusion criteria were: 1) chronic medical conditions or regular use of medications that are associated with fatigue (e.g., untreated hypothyroidism, diabetes, autoimmune disease, anemia (defined as hematocrit < 24), chronic fatigue syndrome); 2) evidence that fatigue was primarily driven by a medical or psychiatric disorder other than cancer (e.g., current major depression, insomnia, sleep apnea); 3) evidence that fatigue was primarily driven by other non-cancer related factors (e.g., shift work, recent change in activity or schedule); 4) physical problems or conditions that could make yoga unsafe (e.g., serious neck injuries, unstable joints); and 5) body mass index greater than 31 kg/m².

Potential participants were first screened by phone and then, if they met preliminary eligibility criteria, completed an in-person screening visit where they completed questionnaires and interviews about their medical history and were asked to perform simple movements (e.g., standing with feet touching for 30 seconds, lifting arms over head, moving from standing position to seated position on floor) to verify safety for yoga practice.

Randomization and blinding

Given class scheduling considerations, participants were randomized in blocks. Once a sufficient number of participants to comprise the yoga and health education groups had been screened as eligible (8-14 women), they were randomized (1:1) to the two treatments after completing baseline assessments. The allocation sequence was generated independently by the study statistician (RO) and concealed in opaque envelopes. Our recruitment documents, screening materials, and informed consent indicated that the purpose of this study was to compare two different treatments for post-cancer fatigue: yoga classes and a wellness seminar series. Thus, although participants were aware of the condition to which they were assigned, they did not know the study hypotheses. Outcomes assessors for the performance tasks were blinded to group assignment, and all were trained in standardized testing procedures.

Interventions

Iyengar yoga classes were conducted for 90 minutes twice a week for 12 weeks in groups of 4-6 women. Classes were taught by a certified Junior Intermediate Iyengar yoga instructor and an assistant under the guidance of a senior teacher. Iyengar yoga, a traditional form of Hatha yoga, prescribes specific therapeutic yoga practices for individuals with specific medical problems and conditions (27). This trial emphasized postures and breathing techniques believed to be effective for reducing fatigue among women with a history of breast cancer, with a focus on passive inversions (i.e., supported upside-down postures in which the head is lower than the heart) and passive backbends (i.e., supported spinal extensions). In supportive postures the shape of the pose is supported by props (e.g., blocks, bolsters, blankets, wall ropes, belts), rather than held by the strength of the body, so that participants can perform and maintain the postures without stress and tension. The postures were introduced using a standard progression from simpler to more challenging over the course of the intervention, and were adapted to suit individual needs. A complete list of study postures and the props used to support each is provided in Table 2. Of note, not all of these postures were included in each class. A typical sequence of postures in a mid-intervention class and the duration of each posture is as follows: 1) Supta Baddhakonasana (10 minutes), 2) Setubandha Sarvangasana on bolsters (5 minutes); 3) Adhomukha Svanasana (5 minutes); 4) Salamba Sirsasana (5 minutes); 5) Viparita Dandasana (5

minutes); 6) Setubandha Sarvangasana on a wooden bench (5 minutes); 7) Viparita Karani (10 minutes); and 8) Supported Savasana (10 minutes).

Health education classes were conducted for 120 minutes once a week for 12 weeks in groups of 4-7 women. Classes were led by a Ph.D. level psychologist with clinical experience in the treatment of breast cancer survivors. The classes were didactic in nature and consisted of lectures about topics of interest to breast cancer survivors followed by questions and discussion. The topics included 1) cancer-related fatigue, 2) introduction to cancer survivorship, 3) psychosocial issues in cancer survivorship, 4) weight and chronic disease management, 5) cancer genetic predisposition testing and counseling for breast/ovarian cancer syndromes, 6) stress and cancer, 7) diet, nutrition, and cancer survivorship, 8) sleep hygiene, 9) cognitive problems after cancer treatment, 10) osteoporosis and cancer survivorship, 11) body image and sexuality, 12) finding meaning and achieving goals. Our previous experience with this patient population suggested that women would be unlikely to travel to the study site twice a week for health education classes. Thus, we elected to have the health education group meet only once per week (vs. twice per week for yoga) for a longer duration (120 minutes/session vs. 90 minutes/session for yoga). As a result, the total number of class hours for yoga (36 hours) was higher than for health education (24 hours). Neither group was instructed to do home practice or reading.

Outcome Measures

Questionnaires and functional assessments were obtained at baseline, immediately post-intervention, and 3 months after the intervention was completed. The primary outcome of interest was subjective fatigue severity, assessed with the Fatigue Symptom Inventory (FSI), a reliable and valid measure of fatigue that was designed for use with cancer patients (28;29). A related outcome, vigor, was assessed using the vigor subscale of the Multidimensional Fatigue Symptom Inventory (30). Several secondary outcomes were also assessed. To determine whether intervention effects might extend to behavioral symptoms correlated with fatigue (1;31), we assessed depressive symptoms using the Beck Depression Inventory-II (BDI-II) (32) and subjective sleep quality using the Pittsburgh Sleep Quality Index (PSQI) (33). In addition, to examine more generalized intervention effects on feelings of stress, we administered the Perceived Stress Scale (PSS)(34). Physical performance tasks were administered to provide an objective measure of functional status. Timed chair stands were used to assess lower extremity strength and endurance (35), and the functional reach test was used to assess strength, flexibility and balance (36).

Self-reported demographic and disease-related variables were assessed at baseline, and expectations and beliefs about treatment efficacy were assessed at baseline and post-intervention. Self-efficacy for managing fatigue, a potentially important component of intervention effects (37), was assessed with the fatigue subscale of the HIV self-efficacy questionnaire (38) adapted for breast cancer. Fatigue interference with activities, mood, and enjoyment of life was assessed with the interference subscale of the FSI (28;29). Finally, blood and saliva samples were collected for assessment of biological functioning; results from these assays are still pending and will be reported separately.

Statistical analyses

Based on the large effect on subjective fatigue severity observed in our single arm pilot study of this yoga intervention ($d = 2.2$ at post-intervention, 1.2 at 3 month follow-up) (19), a sample of 30 participants would provide 90% or greater power to detect a between-group difference on the FSI. To account for the possibility of a smaller intervention effect than that observed in our single-arm pilot, our original enrollment target was 72 participants and assumed 20% loss to follow-up. Because of our stringent eligibility criteria, enrollment was

lower than planned ($n = 31$) but still adequate to detect a large intervention effect. All statistical analyses were performed on an intent-to-treat basis. Outcome measures were first tested for baseline equivalence using t-tests or chi-squared as appropriate. Primary analyses utilized mixed model ANOVAs to account for any incomplete data (i.e., lost to follow-up). Treatment (yoga, health education) by Time (baseline, post-treatment, 3 month post-treatment follow-up) were the independent factors, with the time factor covariances estimated individually (i.e., unstructured). Separate analyses were conducted for each of the outcome measures. We did not adjust for multiple testing. Exact p values are presented.

RESULTS

The study took place at the UCLA Medical Center, Westwood, CA between March 2007 and July 2010. We screened 255 women for eligibility; 188 (75%) were not eligible, and 36 (14%) were not interested in participating. The primary reasons for ineligibility were cancer-related, including diagnosis with Stage 3 or 4 breast cancer ($n = 23$), no previous breast cancer diagnosis ($n = 19$), less than 6 months post treatment completion ($n = 11$), and other cancer in last 5 years ($n = 3$). Medical exclusions were BMI greater than 31 ($n=23$), medical conditions that might interfere with safe yoga practice ($n = 17$), chronic medical conditions or medications that might impact fatigue ($n = 11$), and age and menopause-related ($n = 39$). Fatigue-related exclusions were not fatigued ($n = 30$) and fatigue not related to cancer ($n = 11$). We randomized 31 women to either a 12-week Iyengar yoga intervention ($n = 16$) or a 12-week health education program ($n = 15$). Twenty-eight women received the allocated intervention, which we defined as attending more than one class, and follow-up data was obtained on 29 participants (94%). A CONSORT flow diagram is provided in Figure 1.

Groups were balanced on demographic and disease-related characteristics at baseline, and both reported high levels of fatigue (Table 1). In the yoga group, the mean number of classes attended was 18.9 (78%) and the median was 22 (92%), out of a total of 24 classes. In the education group, the mean number of classes attended was 9.2 (77%) and the median was 11 (92%), out of a total of 12 classes. At the 3 month follow-up, 9 of the 14 women who attended the yoga classes (64%) were continuing to use techniques learned in class.

Intervention effects on fatigue and vigor

Yoga led to statistically significant improvements in fatigue severity (P for group $\times$ time interaction = .032; effect size for predicted change from baseline to 3 month follow-up in yoga vs. health education: $d = 1.5$). Participants in the yoga group reported steady declines in fatigue severity from baseline to post-treatment and over the 3 month follow-up, whereas women assigned to health education reported no change over this period (Figure 2). Similar effects were observed for vigor, with the yoga group showing a statistically significant increase over the assessment period compared to controls (P for group $\times$ time interaction = .011; effect size for predicted change from baseline to 3 month follow-up in yoga vs. health education: $d = 1.20$). We also explored intervention effects on fatigue interference, and found a significant effect of time ($P < .0001$) and a marginally significant group $\times$ time interaction ($P = .08$). Both groups reported decreases in fatigue interference over time, with a marginally larger decrease in the yoga group (mean change from baseline to 3 month follow-up = 15.8) than in health education (mean change from baseline to 3 month follow-up = 7.9). Yoga group participants also felt significantly more confident about their ability to manage fatigue and its impact on their lives than control group participants at post-treatment (yoga group mean = 7.9, education group mean = 6.1; $t(28) = -2.6$, $P = .017$).

To determine the clinical significance of these effects, we calculated the change in SF-36 vitality scale scores from baseline to 3 month follow-up in each group. Although this was not a primary outcome for the trial, there are published norms for determination of clinically

significant change on this measure (39). For women randomized to yoga, there was a mean increase in SF-36 vitality scores of 23.9 points, which exceeds the reliable change index of 22.7 for this scale and indicates a clinically significant improvement in fatigue. For women randomized to health education, the mean increase in SF-36 vitality scores was 7.7 points.

Intervention effects on depressive symptoms, stress, sleep, and physical performance

For depressive symptoms, there was a significant effect of time ($P < .0001$) and a significant group x time interaction ($P = .026$). Both groups reported reductions in depressive symptoms from baseline to post-treatment, although there was a greater decline in symptoms among women in the yoga group at this time point. By the 3 month follow-up, the yoga group had rebounded and the groups reported similar symptom levels. There was also a significant time effect for perceived stress ($P = .015$); feelings of stress decreased over the assessment period in both groups. There were no significant effects for subjective sleep quality. For the physical performance tasks, there was a significant time effect for chair stands ($P < .0001$) as both groups improved over time, but no group x time interaction for either task ($P > .60$). Table 3 presents mean scores for the primary and secondary outcomes at baseline, post-treatment, and 3 month follow-up as well as group differences in change scores.

Expectations and beliefs about treatment

We assessed expectations and beliefs about treatment efficacy to determine whether these might account for beneficial effects of yoga. After random assignment, participants in both groups believed that the intervention to which they had been assigned would be effective in improving their fatigue symptoms; on a 7-point Likert scale ranging from 0 “not at all effective” to 6 “very effective”, the mean score for the yoga group was 3.86 and for the health education group was 3.2 ($P = .33$). At post-treatment, participants in both groups reported that they had experienced “quite a bit” of benefit from attending the classes (mean score for the yoga group was 4.43 and for the health education group was 4.87 on a 7-point Likert scale ranging from 0 “no benefit” to 6 “very much benefit”; $P = .53$). Moreover, both groups reported positive changes in fatigue at post-intervention; on a 13-point Likert scale ranging from -6 “very much worse” to 6 “very much better”, the mean score for the yoga group was 1.93 and for the education group was 1.87 ($P = .94$). These findings suggest that the two groups had similar expectations and beliefs about treatment efficacy.

Safety

There was one adverse event related to the protocol; a participant with a history of back problems experienced a back spasm in yoga class. After evaluation by her physician, she was able to return to class and complete the intervention.

DISCUSSION

Results from this randomized, controlled trial indicate that a yoga intervention targeted at improving fatigue may be a feasible and effective treatment for breast cancer survivors with persistent cancer-related fatigue. Participants in the yoga group experienced significant reductions in fatigue and increases in vigor from pre to post-treatment that persisted over a three month follow-up, consistent with results from our single arm pilot study (19), whereas control group participants showed no change in these outcomes. Measures of effect size indicated that this was a large effect ($d = 1.5$ for fatigue severity), considerably larger than the small to moderate effect sizes observed in previous behavioral interventions for cancer-related fatigue (5;7). This change also appeared to be clinically significant. By the three month post-treatment follow up, scores on the Fatigue Symptom Inventory fell below the cutoff for clinically meaningful fatigue among women in the yoga group (40), and this group also showed a clinically significant increase on the SF-36 vitality scale (39).

Despite high levels of fatigue at study onset, adherence to the yoga intervention was excellent, with over 80% of participants attending at least 20 of the 24 yoga classes offered. This is notable because fatigue is typically a barrier to participation in behavioral interventions for cancer patients, including yoga programs not specifically designed to treat fatigue (18). We suspect that the careful selection of yoga poses for fatigued individuals and the use of props that enabled these poses to be performed without effort or strain played an important role in maintaining trial adherence.

Participants in the yoga and the health education interventions reported comparable declines in depressive symptoms and perceived stress by the three month follow-up. For the education group, these improvements may have resulted from the provision of information about breast cancer survivorship as well as non-specific aspects of the intervention, including attention and group support. Very few behavioral or mind-body intervention trials for cancer-related fatigue have included active control groups, relying instead on usual care or wait-list controls (5-7). Consistent with our findings, Irwin et al. found beneficial effects of health education on depressive symptoms in a study comparing Tai Chi to health education for insomnia (41). These results strongly support the inclusion of active control groups in mind-body and other behavioral intervention trials.

Our study had several limitations. First, we enrolled a relatively small number of participants, reflecting the challenges of recruiting fatigued breast cancer survivors with no co-morbid fatigue-related medical conditions or physical limitations who were willing to participate in a demanding intervention study. It will be important to replicate these findings in a larger trial, and to determine if effects are generalizable to a broader group of breast cancer survivors. We also restricted our sample to women who had been diagnosed with early stage disease, had completed cancer treatment, and had a BMI under 31. Thus, effects may not be generalizable to breast cancer survivors with more advanced disease, who are undergoing treatment, and/or who are obese, and the yoga intervention may require adaptation to meet the specific needs of these groups. In addition, although the majority of study participants (94%) were within 5 years post-diagnosis, we did enroll a few women who had been living with cancer-related fatigue for over 10 years. Future studies should examine whether the length of time since cancer treatment influences intervention efficacy, and possibly target either shorter-term or longer-term survivors. Next, it is not feasible to use a double-blind study design with this type of intervention; participants are unavoidably aware of the treatment that they receive. We attempted to minimize the impact of pre-existing beliefs and expectations on study outcomes by informing participants that we were testing two different treatments for cancer-related fatigue, each of which were presumably effective. This presentation appeared to have been successful, as both groups had similar positive expectations about the efficacy of the treatment to which they had been assigned. Finally, the yoga and health education conditions were not matched for class frequency or duration, and it is possible that benefits seen in the yoga group may be attributable in part to the higher number of intervention hours received. Future studies should match class frequency and duration across conditions. It will also be useful to compare yoga to a more physically-oriented control condition (e.g., stretching) or to relaxation, to probe the active components of this intervention.

These preliminary findings indicate that a specialized yoga intervention may have beneficial effects on cancer-related fatigue, even among survivors who have experienced fatigue for years post-treatment. Future investigation into this promising approach is warranted, including examination of mechanisms and duration of treatment effects. Yoga may have effects on the immune and neuroendocrine systems (42-46), which have been linked to post-treatment fatigue in previous research (21;23;24;26;47-49) and are currently under investigation in this sample.

Acknowledgments

Supported by NCCAM/NIH U01- AT003682 Iyengar Yoga for Breast Cancer Survivors with Persistent Fatigue. The authors report no financial disclosures

Reference List

1. Bower JE, Ganz PA, Desmond KA, Rowland JH, Meyerowitz BE, Belin TR. Fatigue in breast cancer survivors: occurrence, correlates, and impact on quality of life. *J.Clin.Oncol.* 2000; 18:743–53. [PubMed: 10673515]
2. Cella D, Davis K, Breitbart W, Curt G. Cancer-related fatigue: prevalence of proposed diagnostic criteria in a United States sample of cancer survivors. *J Clin Oncol.* 2001; 19:3385–91. [PubMed: 11454886]
3. Gielissen MF, Verhagen S, Witjes F, Bleijenberg G. Effects of cognitive behavior therapy in severely fatigued disease-free cancer patients compared with patients waiting for cognitive behavior therapy: a randomized controlled trial. *J Clin Oncol.* 2006; 24:4882–7. [PubMed: 17050873]
4. Jain S, Pavlik D, Distefan J, Bruyere RR, Acer J, Garcia R, et al. Complementary medicine for fatigue and cortisol variability in breast cancer survivors: A Randomized Controlled Trial. *Cancer.* 2011
5. Jacobsen PB, Donovan KA, Vadaparampil ST, Small BJ. Systematic review and meta-analysis of psychological and activity-based interventions for cancer-related fatigue. *Health Psychol.* 2007; 26:660–7. [PubMed: 18020836]
6. Schmitz KH, Holtzman J, Courneya KS, Masse LC, Duval S, Kane R. Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol.Biomarkers Prev.* 2005; 14:1588–95. [PubMed: 16030088]
7. Kangas M, Bovbjerg DH, Montgomery GH. Cancer-related fatigue: a systematic and meta-analytic review of non-pharmacological therapies for cancer patients. *Psychol.Bull.* 2008; 134:700–41. [PubMed: 18729569]
8. Courneya KS, Friedenreich CM, Quinney HA, Fields AL, Jones LW, Vallance JK, et al. A longitudinal study of exercise barriers in colorectal cancer survivors participating in a randomized controlled trial. *Ann.Behav.Med.* 2005; 29:147–53. [PubMed: 15823788]
9. Iyengar, BKS. *Light on Yoga.* Schocken; New York, NY: 1995.
10. Lin KY, Hu YT, Chang KJ, Lin HF, Tsao JY. Effects of yoga on psychological health, quality of life, and physical health of patients with cancer: a meta-analysis. *Evid.Based.Complement Alternat.Med.* 2011; 2011:659876. [PubMed: 21437197]
11. Bower JE, Woolery A, Sternlieb B, Garet D. Yoga for cancer patients and survivors. *Cancer Control.* 2005; 12:165–71. [PubMed: 16062164]
12. Smith KB, Pukall CF. An evidence-based review of yoga as a complementary intervention for patients with cancer. *Psychooncology.* 2009; 18:465–75. [PubMed: 18821529]
13. Culos-Reed SN, Carlson LE, Daroux LM, Hatley-Aldous S. A pilot study of yoga for breast cancer survivors: physical and psychological benefits. *Psychooncology.* 2006; 15:891–7. [PubMed: 16374892]
14. Danhauer SC, Mihalko SL, Russell GB, Campbell CR, Felder L, Daley K, et al. Restorative yoga for women with breast cancer: findings from a randomized pilot study. *Psychooncology.* 2009; 18:360–8. [PubMed: 19242916]
15. Rao MR, Raghuram N, Nagendra HR, Gopinath KS, Srinath BS, Diwakar RB, et al. Anxiolytic effects of a yoga program in early breast cancer patients undergoing conventional treatment: a randomized controlled trial. *Complement Ther.Med.* 2009; 17:1–8. [PubMed: 19114222]
16. Vadiraja HS, Rao MR, Nagarathna R, Nagendra HR, Rekha M, Vanitha N, et al. Effects of yoga program on quality of life and affect in early breast cancer patients undergoing adjuvant radiotherapy: a randomized controlled trial. *Complement Ther.Med.* 2009; 17:274–80. [PubMed: 19942107]

17. Chandwani KD, Thornton B, Perkins GH, Arun B, Raghuram NV, Nagendra HR, et al. Yoga improves quality of life and benefit finding in women undergoing radiotherapy for breast cancer. *J Soc.Integr.Oncol.* 2010; 8:43–55. [PubMed: 20388445]
18. Moadel AB, Shah C, Wylie-Rosett J, Harris MS, Patel SR, Hall CB, et al. Randomized controlled trial of yoga among a multiethnic sample of breast cancer patients: effects on quality of life. *J Clin Oncol.* 2007; 25:4387–95. [PubMed: 17785709]
19. Bower JE, Garet D, Sternlieb B. Yoga for persistent fatigue in breast cancer survivors: results of a pilot study. *Evid.Based.Complement Alternat.Med.* 2011; 2011:623168. [PubMed: 21274288]
20. Ware JE Jr. Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med.Care.* 1992; 30:473–83. [PubMed: 1593914]
21. Bower JE, Ganz PA, Aziz N, Fahey JL. Fatigue and proinflammatory cytokine activity in breast cancer survivors. *Psychosom.Med.* 2002; 64:604–11. [PubMed: 12140350]
22. Bower JE, Ganz PA, Aziz N, Fahey JL, Cole SW. T-cell homeostasis in breast cancer survivors with persistent fatigue. *J.Natl.Cancer Inst.* 2003; 95:1165–8. [PubMed: 12902446]
23. Bower JE, Ganz PA, Dickerson SS, Petersen L, Aziz N, Fahey JL. Diurnal cortisol rhythm and fatigue in breast cancer survivors. *Psychoneuroendocrinology.* 2005; 30:92–100. [PubMed: 15358446]
24. Bower JE, Ganz PA, Aziz N. Altered cortisol response to psychologic stress in breast cancer survivors with persistent fatigue. *Psychosom.Med.* 2005; 67:277–80. [PubMed: 15784794]
25. Bower JE, Ganz PA, Aziz N, Olmstead R, Irwin MR, Cole SW. Inflammatory responses to psychological stress in fatigued breast cancer survivors: relationship to glucocorticoids. *Brain Behav.Immun.* 2007; 21:251–8. [PubMed: 17008048]
26. Collado-Hidalgo A, Bower JE, Ganz PA, Cole SW, Irwin MR. Inflammatory biomarkers for persistent fatigue in breast cancer survivors. *Clin Cancer Res.* 2006; 12:2759–66. [PubMed: 16675568]
27. Iyengar, BKS. Yoga: The path to holistic health. Dorling Kindersley Ltd; London, England: 2001.
28. Donovan KA, Jacobsen PB. The Fatigue Symptom Inventory: a systematic review of its psychometric properties. *Support.Care Cancer.* 2010; 19:169–85. [PubMed: 20824482]
29. Hann DM, Jacobsen PB, Azzarello LM, Martin SC, Curran SL, Fields KK, et al. Measurement of fatigue in cancer patients: development and validation of the Fatigue Symptom Inventory. *Qual.Life Res.* 1998; 7:301–10. [PubMed: 9610214]
30. Stein KD, Martin SC, Hann DM, Jacobsen PB. A multidimensional measure of fatigue for use with cancer patients. *Cancer Pract.* 1998; 6:143–52. [PubMed: 9652245]
31. Bower JE. Behavioral symptoms in patients with breast cancer and survivors. *J Clin Oncol.* 2008; 26:768–77. [PubMed: 18258985]
32. Beck, AT.; Steer, RA.; Brown, GK. BDI-II Manual. 2 ed. The Psychological Corporation; San Antonio: 1996.
33. Buysse DJ, Reynolds CF III, Monk TH, Hoch CC, Yeager AL, Kupfer DJ. Quantification of subjective sleep quality in healthy elderly men and women using the Pittsburgh Sleep Quality Index (PSQI). *Sleep.* 1991; 14:331–8. [PubMed: 1947597]
34. Cohen, S.; Kessler, RC.; Gordon, LU. Measuring stress: A guide for health and social scientists. Oxford University Press; New York: 1997.
35. Csuka M, McCarty DJ. Simple method for measurement of lower extremity muscle strength. *Am.J Med.* 1985; 78:77–81. [PubMed: 3966492]
36. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: a new clinical measure of balance. *J Gerontol.* 1990; 45:M192–M197. [PubMed: 2229941]
37. Antoni MH, Lechner SC, Kazi A, Wimberly SR, Sifre T, Urcuyo KR, et al. How stress management improves quality of life after treatment for breast cancer. *J Consult Clin Psychol.* 2006; 74:1143–52. [PubMed: 17154743]
38. Shively M, Smith TL, Bormann J, Gifford AL. Evaluating Self-Efficacy for HIV Disease Management Skills. *AIDS and Behavior.* 2002; 6:371–9.
39. Ferguson RJ, Robinson AB, Splaine M. Use of the reliable change index to evaluate clinical significance in SF-36 outcomes. *Qual.Life Res.* 2002; 11:509–16. [PubMed: 12206571]

40. Donovan KA, Jacobsen PB, Small BJ, Munster PN, Andrykowski MA. Identifying clinically meaningful fatigue with the Fatigue Symptom Inventory. *J Pain Symptom Manage.* 2008; 36:480–7. [PubMed: 18495413]
41. Irwin MR, Olmstead R, Motivala SJ. Improving sleep quality in older adults with moderate sleep complaints: A randomized controlled trial of Tai Chi Chih. *Sleep.* 2008; 31:1001–8. [PubMed: 18652095]
42. Banasik J, Williams H, Haberman M, Blank SE, Bendel R. Effect of Iyengar yoga practice on fatigue and diurnal salivary cortisol concentration in breast cancer survivors. *J Am.Acad.Nurse Pract.* 2011; 23:135–42. [PubMed: 21355946]
43. Vadiraja HS, Raghavendra RM, Nagarathna R, Nagendra HR, Rekha M, Vanitha N, et al. Effects of a yoga program on cortisol rhythm and mood states in early breast cancer patients undergoing adjuvant radiotherapy: a randomized controlled trial. *Integr.Cancer Ther.* 2009; 8:37–46. [PubMed: 19190034]
44. Kiecolt-Glaser JK, Christian L, Preston H, Houts CR, Malarkey WB, Emery CF, et al. Stress, inflammation, and yoga practice. *Psychosom Med.* 2010; 72:113–21. [PubMed: 20064902]
45. Pullen PR, Nagamia SH, Mehta PK, Thompson WR, Benardot D, Hammoud R, et al. Effects of yoga on inflammation and exercise capacity in patients with chronic heart failure. *J Card Fail.* 2008; 14:407–13. [PubMed: 18514933]
46. West J, Otte C, Geher K, Johnson J, Mohr DC. Effects of Hatha yoga and African dance on perceived stress, affect, and salivary cortisol. *Ann.Behav.Med.* 2004; 28:114–8. [PubMed: 15454358]
47. Orre IJ, Murison R, Dahl AA, Ueland T, Aukrust P, Fossum SD. Levels of circulating interleukin-1 receptor antagonist and C-reactive protein in long-term survivors of testicular cancer with chronic cancer-related fatigue. *Brain, Behavior, and Immunity.* 2009; 23:868–74.
48. Alexander S, Minton O, Andrews P, Stone P. A comparison of the characteristics of disease-free breast cancer survivors with or without cancer-related fatigue syndrome. *European Journal of Cancer.* 2009; 45:384–92. [PubMed: 18977131]
49. Bower JE. Cancer-related fatigue: links with inflammation in cancer patients and survivors. *Brain Behav.Immun.* 2007; 21:863–71. [PubMed: 17543499]

Figure 1.
Consort diagram

Figure 2. Change in fatigue severity (Panel A) and vigor (Panel B) in yoga and health education groups. The yoga group showed significant declines in fatigue severity and significant improvements in vigor from baseline to post-treatment and over the 3-month post-treatment follow-up relative to health education controls.

Table 1

Demographic and treatment-related characteristics of study participants

Characteristic	Yoga (n = 16)	Health education (n = 15)
Age, mean (SD)	54.4 (5.7)	53.3 (4.9)
Race/ethnicity, # (%):		
White	15 (94)	12 (80)
Hispanic	0	2 (13)
Black	1 (6)	0
Other	0	1 (7)
Relationship status, # (%):		
Married/committed relationship	12 (75)	11 (73)
Education status, # (%):		
High school graduate/AA degree	6 (37.5)	7 (46)
College graduate	6 (37.5)	4 (27)
Graduate degree	4 (25)	4 (27)
Annual family income, # (%):		
< 45,000	3 (19)	1 (7)
45,000 – 75,000	5 (31)	4 (27)
75,000 – 100,000	4 (25)	3 (20)
> 100,000	3 (19)	7 (46)
Not reported	1 (6)	0
Body mass index, mean (SD)	24.0 (2.5)	25.3 (3.4)
Treated with radiation, # (%)	11 (69)	13 (87)
Treated with chemotherapy, # (%)	8 (50)	9 (60)
Tamoxifen/aromatase inhibitor, # (%)	12 (75)	10 (67)
Years since treatment completion, median (range)	1.7 (0.7 – 4.1)	1.7 (0.7 – 18.3)
SF-36 vitality score, mean (SD)	37.8 (16)	34.0 (16.3)

No differences are significant at $p < .20$

Table 2

List of yoga postures for persistent cancer-related fatigue

Sanskrit Name
Supta Baddhakonasana with bolster, strap, and blankets
Supta Svastikasana with bolster and blanket
Setubandha Sarvangasana with bolster, strap and blankets
Setubandha Sarvangasana on a wooden bench with a box and bolster
Purvottanasana with two chairs, bolsters, and blankets
Viparita Dandasana on two chairs with the head supported
Salamba Sarvangasana with a chair, bolster, sticky mat, and blanket
Salamba Sirsasana on ropes
Supta Konasana with legs apart on two chairs
Viparita Karani with two blocks, a wall, a bolster and blankets
Bharadvajasana on chair
Adhomukha Svanasana on ropes with chair
Urdhva Mukha Svanasana with chair
Urdhva Hastasana
Urdhva Baddhanguliyasana
Ropes 1
Savasana with bolster and blanket

A full description of each pose, including pictures, and a detailed intervention manual including class sequences is available from the first author.

Table 3

Mean Scores on Outcome Measures for Yoga and Health Education Groups

Outcome	Baseline Mean (SD)	Post-treatment Mean (SD)	3 month FU Mean (SD)	Group Difference in Change from Baseline- Post treatment Mean (95%CI)	Group Difference in Change from Baseline-3 month FU Mean (95%CI)
Primary Outcomes					
FSI fatigue severity (0-11; higher is worse)					
Iyengar Yoga	4.8 (1.1)	3.4 (1.8)	2.8 (2.3)	-1.24 (-0.04 to -2.45)*	-1.62 (-0.37 to -2.88)*
Health Education	5.1 (.95)	4.9 (1.3)	4.7 (1.5)		
MFSI vigor (0-24; higher is better)					
Iyengar Yoga	8.8 (3.2)	12.5 (3.8)	11.9 (4.7)	4.80 (1.86 to 7.74)*	4.25 (0.99 to 7.50)*
Health Education	9.1 (3.6)	8.4 (3.4)	8.6 (4.2)		
Secondary Outcomes					
BDI-II depressive symptoms (0-63; higher is worse)					
Iyengar yoga	15.5 (7.5)	7.7 (5.8)	9.9 (8.0)	-5.80 (-1.74 to -9.86)*	-3.06 (0.37 to -6.49)
Health education	14.3 (7.5)	11.6 (7.1)	10.5 (7.9)		
PSQI sleep disturbance (0-21; higher is worse)					
Iyengar yoga	9.2 (3.3)	8.1 (2.5)	7.6 (2.7)	0.20 (2.78 to -2.38)	-2.14 (1.06 to -5.39)
Health education	9.1 (3.5)	7.7 (2.6)	9.1 (3.3)		
Perceived stress scale (0-40; higher is worse)					
Iyengar yoga	26.6 (7.3)	23.5 (7.3)	23.6 (6.7)	-1.77 (1.71 to -5.26)	-1.51 (2.38 to -5.39)
Health education	26.8 (5.9)	25.4 (5.9)	24.2 (4.7)		
Chair stand time, sec (higher is worse)					
Iyengar yoga	13.4 (2.3)	12.3 (2.0)	11.1 (1.9)	1.31 (-5.00 to 2.38)	-0.74 (-3.74 to 2.26)
Health education	13.5 (2.6)	12.6 (3.2)	12.2 (2.8)		
Functional reach, cm (higher is better)					
Iyengar yoga	31.1 (5.7)	29.6 (6.2)	29.5 (5.1)	-2.00 (5.76 to -9.73)	-2.21 (5.56 to -9.98)
Health education	29.5 (6.7)	28.5 (7.2)	28.8 (8.0)		

*
p < .05