Published in final edited form as:

Am J Prev Med. 2013 April ; 44(4): 325–329. doi:10.1016/j.amepre.2012.11.032.

Promoting Colorectal Cancer Screening Discussion:

A Randomized Controlled Trial

Shannon M. Christy, MA, Susan M. Perkins, PhD, Yan Tong, PhD, Connie Krier, BS, Victoria L. Champion, PhD, RN, FAAN, Celette Sugg Skinner, PhD, Jeffrey K. Springston, PhD, Thomas F. Imperiale, MD, and Susan M. Rawl, PhD, RN, FAAN

Department of Psychology (Christy), School of Medicine (Perkins, Tong, Imperiale), School of Nursing (Krier, Champion, Rawl), Indiana University, Indianapolis, Indiana; Harold C. Simmons Cancer Center (Skinner), University of Texas Southwestern Medical Center, Dallas Texas; College of Journalism and Mass Communication (Springston), University of Georgia, Athens, Georgia

Abstract

Background—Provider recommendation is a predictor of colorectal cancer (CRC) screening.

Purpose—To compare the effects of two clinic-based interventions on patient–provider discussions about CRC screening.

Design—Two-group RCT with data collected at baseline and 1 week post-intervention.

Participants/setting—African-American patients that were non-adherent to CRC screening recommendations ($n=693$) with a primary care visit between 2008 and 2010 in one of 11 urban primary care clinics.

Intervention—Participants received either a computer-delivered tailored CRC screening intervention or a nontailored informational brochure about CRC screening immediately prior to their primary care visit.

Main outcome measures—Between-group differences in odds of having had a CRC screening discussion about a colon test, with and without adjusting for demographic, clinic, health literacy, health belief, and social support variables, were examined as predictors of a CRC screening discussion using logistic regression. Intervention effects on CRC screening test order by PCPs were examined using logistic regression. Analyses were conducted in 2011 and 2012.

Results—Compared to the brochure group, a greater proportions of those in the computer-delivered tailored intervention group reported having had a discussion with their provider about CRC screening (63% vs 48%, OR=1.81, $p<0.001$). Predictors of a discussion about CRC screening included computer group participation, younger age, reason for visit, being unmarried, colonoscopy self-efficacy, and family member/friend recommendation (all p -values <0.05).

Conclusions—The computer-delivered tailored intervention was more effective than a nontailored brochure at stimulating patient–provider discussions about CRC screening. Those who

© 2012 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

Address correspondence to: Susan M. Rawl, PhD, RN, FAAN, Indiana University School of Nursing, 1111 Middle Drive, NU 345E, Indianapolis IN 46202. srawl@iupui.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

No financial disclosures were reported by the authors of this paper.

received the computer-delivered intervention also were more likely to have a CRC screening test (fecal occult blood test or colonoscopy) ordered by their PCP.

Trial registration—This study is registered at www.clinicaltrials.gov NCT00672828.

Introduction

Tailored interventions are more effective than nontailored materials in promoting behavior change including cancer screening.^{1–5} The current RCT compared the efficacy of a clinic-based, computer-delivered tailored interactive program with a nontailored brochure to promote patient–provider discussions about colorectal cancer (CRC) screening among African-American patients. Demographic, clinic, and health belief variables were examined as predictors of a discussion. CRC screening test orders written during the visit also were examined. It was hypothesized that: (1) individuals who received the computer-delivered tailored intervention would be more likely to engage in CRC screening discussions with their primary care provider (PCP) than those in the brochure group; and (2) PCPs of individuals who received the computer-delivered tailored intervention would be more likely to write orders for colon tests than PCPs of those in the brochure group.

Methods

A total of 693 African-American patients of 118 PCPs were enrolled between 2008 and 2010. Patients were eligible if they were self-identified as black or African-American, aged 51–80 years, English-speaking, and currently non-adherent to CRC screening guidelines. Exclusion criteria were: personal history of CRC or adenomatous polyps requiring surveillance colonoscopy; medical condition precluding CRC screening; cognitive, speech, or hearing impairment; and current adherence to CRC screening guidelines. Sample sizes were determined based on detecting a difference in CRC screening at 15 months of 25% vs 15% for the computer-delivered tailored interactive program and nontailored brochure, respectively. Observed power for the patient–provider discussion outcome was 95%.

Procedure

The Indiana University IRB approved the study; all procedures were Health Insurance Portability and Accountability Act (HIPAA)–compliant. Patients were recruited from 11 Midwestern urban primary care clinics (five Veterans Affairs (VA) clinics and six non-VA). Potentially eligible participants were identified via clinic databases and approved for contact by their PCPs. Patients with upcoming PCP appointments were mailed an introductory letter, a study brochure, and an informed consent form.

Trained recruiters telephoned patients within 1 week of mailing letters to explain the study, answer questions, assess eligibility, and obtain verbal consent. After providing verbal consent, patients completed the baseline interview prior to their clinic visit. Randomization, stratified by site, gender, and age occurred following the baseline interview. Research staff met patients in the clinic 45 minutes before their scheduled PCP visit to obtain written consent and HIPAA authorization, assess health literacy, and deliver interventions.

Data Collection

Trained interviewers collected data using a computer-assisted telephone interview system. Baseline data were collected after verbal consent was obtained but prior to intervention. The second telephone interview was conducted 1 week following intervention delivery.

Interventions

Trained research staff delivered interventions in the clinic immediately prior to the PCP visit. Details of the intervention design and delivery have been published elsewhere.⁶ Briefly, the computer intervention delivered messages tailored to the patient's age; gender; objective CRC risk (family history); perceived CRC risk; and barriers to screening. The program produced a tailored printout that summarized the user's CRC risk factors and risk-based test recommendations, and encouraged them to discuss CRC screening with their PCP. The usual-care group received a nontailored CRC screening brochure.⁷

Measures

Demographic characteristics were collected during the baseline interview. CRC knowledge was measured using 11 items ($\alpha=0.63$). Perceived CRC risk was measured using three items ($\alpha=0.83$).^{8,9} Objective CRC risk was determined based on two items assessing strength of family history of CRC.¹⁰ Perceived barriers to fecal occult blood testing (FOBT) and colonoscopy were measured using nine items ($\alpha=0.82$) and 15 items ($\alpha=0.89$), respectively.¹¹ Perceived benefits of FOBT and colonoscopy were measured using three items ($\alpha=0.76$) and four items ($\alpha=0.67$), respectively.¹¹ Self-efficacy for FOBT and for colonoscopy was measured with eight items and 11 items, respectively (both $\alpha=0.87$).¹¹

Cancer fatalism was measured with 11 items ($\alpha=0.858$).¹²⁻¹⁴ Single items were used to assess whether a family member/friend had encouraged CRC screening and whether participants had ever received a PCP recommendation for CRC screening. Health literacy was measured using the shortened version of the Rapid Estimate of Adult Literacy in Medicine.^{15,16} At 1 week post-intervention, patients were asked whether they talked with their PCP about a colon test. Test order outcomes for FOBT and colonoscopy were collected from medical records.

Data Analysis

Analyses were performed using data from randomized patients who received the intervention and completed the 1 week post-intervention interview (95%); missing data were not imputed. Patients who were randomized but never received the interventions because they failed to attend their clinic visit ($n=124$) were excluded from analyses. Those who did not complete the 1 week post-intervention interview ($n=34$) also were excluded (Figure 1).

Demographics were examined using descriptive statistics. Logistic models were estimated using the generalized estimation equation methodology to account for clustering by PCP. Univariate logistic regression models determined predictors of patient-provider discussion about CRC screening. Potential predictors included: intervention group; age; gender; education; insurance status; marital status; employment status; site (VA vs non-VA); BMI; number of doctor visits in past year; reason for visit; objective CRC risk; health literacy; perceived CRC risk; FOBT benefits, barriers, and self-efficacy; colonoscopy benefits, barriers, and self-efficacy; cancer fatalism, CRC knowledge, PCP recommendation of FOBT, colonoscopy, and sigmoidoscopy; and family member/friend encouragement of CRC testing.

Regression analyses of each predictors' univariate effect on the outcome variable having a p -value <0.20 were entered into the final multivariable logistic regression model. In addition, univariate logistic regression was used to assess intervention effect on PCP orders of CRC screening tests. All data were analyzed using SAS 9.3 in 2011 and 2012.

Results

Baseline demographic data are listed in Table 1. Of the 693 primary care patients who received interventions (319 in the computer group, 340 in the brochure group), 659 (95%) completed the 1 week post-intervention interview. Univariate analysis of intervention effects on patient–provider discussions is presented in Table 2. Compared to those who received the nontailored brochure, participants who received the computer-delivered tailored intervention were more likely to report having engaged in a CRC screening discussion with their PCPs (63% vs 49%, OR=1.81, $p<0.001$).

Results from the final multivariable model also are summarized in Table 2 and show that the intervention effect held after controlling for all other variables. The computer group had higher odds of having a discussion about a colon test with their PCP than the brochure group ($p=0.003$). Participants who were older ($p=0.003$) and were married or living with a partner ($p=0.045$) had lower odds of having a discussion about a colon test with their PCP. Individuals had higher odds of having such a discussion if they were being seen for a preventive health visit ($p=0.011$); had higher colonoscopy self-efficacy scores ($p=0.002$); and had a family member/friend encourage CRC screening ($p=0.002$). PCPs of those who received the computer-delivered tailored intervention were more likely to write orders for a CRC screening test (OR=1.48; 95% CI=[1.11, 1.96]; p -value=0.007).

Discussion

This study compared efficacy of two clinic-based interventions to stimulate patient-reported CRC screening discussions between African-American primary care patients and their PCPs. Individuals who received the computer-delivered tailored intervention had higher odds of reporting a colon test discussion with their PCP and were more likely to have a CRC screening test ordered during the visit. This study is novel due to its focus on evaluating efficacy of an interactive CRC screening intervention to stimulate patient–provider discussions about CRC screening among African-American primary care patients as well as the test orders that resulted.^{17,18}

Strengths and Limitations

Strengths of the study include the large sample size, the RCT design, and use of multiple recruitment sites. Limitations include the fact that patient–provider discussion data were based on patient self-report. However, test order outcomes were collected, thereby confirming that a discussion occurred. In addition, data were not collected regarding whether the patient or the PCP initiated the discussion. Finally, results may not generalize to populations dissimilar to participants in the current study.

Future Directions

Future research is needed to investigate whether patient–provider discussions about CRC screening include risk-based recommendations for CRC testing and/or ultimately lead to higher CRC screening test completion rates among African-American patients.^{19–22} Older age was associated with lower likelihood of having a patient–provider discussion. It is unclear if this is due to older individuals being less likely to discuss CRC screening with their physicians or whether prior discussions have already occurred. Also, it may be that with increased age, there is a greater focus on other health concerns due to increased comorbidities and less focus on CRC. Given health disparities experienced by African Americans, it is imperative that CRC screening be promoted in this population.^{23,24} CRC interventions aimed at this underserved population have the potential to save lives if they

can promote patient–provider CRC screening discussions and prompt patients to complete CRC screening.

Conclusion

Clinic-based computer-delivered tailored interventions can successfully promote patient–provider discussions about CRC screening and subsequent screening test orders. Future analyses from this trial will examine relationships between CRC screening discussions and test completion.

Acknowledgments

The intervention trial was funded by a National Cancer Institute grant awarded to Susan M. Rawl, PhD, RN, FAAN (R01 CA115983; PI: Rawl).

The work of SMC was funded by the Training in Research for Behavioral Oncology and Cancer Control Program—R25 (R25 CA117865-06; PI: Champion). SMC is a predoctoral fellow funded by the Training in Research for Behavioral Oncology and Cancer Control Program—R25 (R25 CA117865-06; PI: Champion) and a doctoral student in the Department of Psychology in the Purdue School of Science at Indiana University–Purdue University Indianapolis.

SMP, YT, CK, VLC, CSS, and JKS were funded by a National Cancer Institute grant awarded to Susan M. Rawl, PhD, RN, FAAN (R01 CA115983; PI: Rawl).

SMP is an Associate Professor in the Indiana University School of Medicine and Indiana University Simon Cancer Center. YT is a Biostatistician in the Department of Biostatistics at Indiana University School of Medicine. CK is a Research Project Manager in the Indiana University School of Nursing. VLC is the Mary Margaret Walther Distinguished Professor, the Edward W. and Sarah Stam Cullipher Endowed Chair, and the Executive Associate Dean for Research Affairs at the Indiana University School of Nursing. She is also the Associate Director of Population Science and a Co-leader of the Cancer Control Program at the Indiana University Simon Cancer Center. CSS is Chief of the Division of Behavioral & Communication Sciences in the Department of Clinical Sciences and Associate Director of Population Research & Cancer Control at the Harold C. Simmons Cancer Center, University of Texas Southwestern Medical Center. JKS is a Professor and the Associate Dean for Research and Graduate Studies in the College of Journalism and Mass Communication at University of Georgia.

The work of TFI and SMR on this project was funded by a National Cancer Institute grant awarded to Susan M. Rawl, PhD, RN, FAAN (R01 CA115983; PI: Rawl).

TFI is a Professor in the Division of Gastroenterology and Hepatology, Department of Medicine, Indiana University School of Medicine, a Research Scientist at Regeneron Institute, Inc., and a Core Investigator in the Center for Implementation of Evidence-based Practices at Roudebush VA Medical Center. SMR is a Professor and Director of Research Training in Behavioral Nursing in the Indiana University School of Nursing. She is also a member of the Cancer Prevention and Control program at Indiana University Simon Cancer Center.

References

1. Champion VL, Skinner CS, Menon U, Seshadri R, Anzalone DC, Rawl SM. Comparisons of tailored mammography interventions at two months postintervention. *Ann Behav Med*. 2002; 24(3): 211–218. [PubMed: 12173678]
2. Skinner CS, Campbell MK, Rimer BK, Curry S, Prochaska JO. How effective is tailored print communication? *Ann Behav Med*. 1999; 21(4):290–298. [PubMed: 10721435]
3. Krebs P, Prochaska JO, Rossi JS. A meta-analysis of computer-tailored interventions for health behavior change. *Prev Med*. 2010; 51(3–4):214–221. [PubMed: 20558196]
4. de Vries H, Kremers SP, Smeets T, Brug J, Eijmael K. The effectiveness of tailored feedback and action plans in an intervention addressing multiple health behaviors. *Am J Health Promot*. 2008; 22(6):417–425. [PubMed: 18677882]
5. Christy SM, Mosher CE, Sloane R, Snyder DC, Lobach DF, Demark-Wahnefried W. Long-term dietary outcomes of the FRESH START intervention for breast and prostate cancer survivors. *J Am Diet Assoc*. 2011; 111(12):1844–1851. [PubMed: 22117660]

6. Rawl SM, Skinner CS, Perkins SM, et al. Computer-tailored Intervention Improves Colon Cancer Screening Knowledge and Health Beliefs of African Americans. *Health Educ Res.* 2012; 27(5):868–885. [PubMed: 22926008]
7. American Cancer Society. If you're 50 or older, you need to get tested for colon cancer. No matter how good your excuse is. In: American Cancer Society. , editor. Vol 2432.00. Atlanta, GA: American Cancer Society; 2005.
8. Champion VL. Instrument refinement for breast cancer screening behaviors. *Nurs Res.* 1993; 42(3): 139–143. [PubMed: 8506161]
9. Champion VL, Scott CR. Reliability and validity of breast cancer screening belief scales in African American women. *Nurs Res.* 1997; 46(6):331–337. [PubMed: 9422052]
10. Levin B, Lieberman DA, McFarland B, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: A joint guideline from the American Cancer Society, the U.S. Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *CA Cancer J Clin.* 2008; 58(3):130–160. [PubMed: 18322143]
11. Rawl S, Champion VL, Menon U, Loehrer P, Vance G, Skinner CS. Validation of scales to measure benefits and barriers to colorectal cancer screening. *J Psychosoc Oncol.* 2001; 19(3/4): 47–63.
12. Powe BD. Cancer fatalism among elderly Caucasians and African Americans. *Oncol Nurs Forum.* 1995; 22(9):1355–1359. [PubMed: 8539176]
13. Powe BD. Fatalism among elderly African Americans: Effects on colorectal cancer screening. *Cancer Nurs.* 1995; 18(5):385–392. [PubMed: 7585493]
14. Mayo RM, Ureda JR, Parker VG. Importance of fatalism in understanding mammography screening in rural elderly women. *J Women Aging.* 2001; 13(1):57–72. [PubMed: 11217186]
15. Davis TC, Long SW, Jackson RH, et al. Rapid estimate of adult literacy in medicine: a shortened screening instrument. *Fam Med.* 1993; 25(6):391–395. [PubMed: 8349060]
16. Davis TC, Michielutte R, Askov EN, Williams MV, Weiss BD. Practical assessment of adult literacy in health care. *Health Educ Behav.* 1998; 25(5):613–624. [PubMed: 9768381]
17. Steinwachs DAJ, Barlow WE, Duncan RP, et al. NIH Consensus State Sci Statements: NIH. 2010
18. Bazargan M, Ani C, Bazargan-Hejazi S, Baker RS, Bastani R. Colorectal cancer screening among underserved minority population: discrepancy between physicians' recommended, scheduled, and completed tests. *Patient Educ Couns.* 2009; 76(2):240–247. [PubMed: 19150198]
19. Zapka JG, Lemon SC, Puleo E, Estabrook B, Luckmann R, Erban S. Patient education for colon cancer screening: a randomized trial of a video mailed before a physical examination. *Ann Intern Med.* 2004; 141(9):683–692. [PubMed: 15520425]
20. Aragonés A, Schwartz MD, Shah NR, Gany FM. A randomized controlled trial of a multilevel intervention to increase colorectal cancer screening among Latino immigrants in a primary care facility. *J Gen Intern Med.* 2010; 25(6):564–567. [PubMed: 20213208]
21. Baron RC, Rimer BK, Breslow RA, et al. Client-directed interventions to increase community demand for breast, cervical, and colorectal cancer screening a systematic review. *Am J Prev Med.* 2008; 35(1 Suppl):S34–S55. [PubMed: 18541187]
22. Baron RC, Rimer BK, Coates RJ, et al. Client-directed interventions to increase community access to breast, cervical, and colorectal cancer screening a systematic review. *Am J Prev Med.* 2008; 35(1 Suppl):S56–S66. [PubMed: 18541188]
23. Ward E, Jemal A, Cokkinides V, et al. Cancer disparities by race/ethnicity and socioeconomic status. *CA Cancer J Clin.* 2004; 54(2):78–93. [PubMed: 15061598]
24. Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin.* 2011; 61(4): 212–236. [PubMed: 21685461]

Figure 1.

Study flowchart

Note: T1=initial interview; T2=follow-up interview 1 week after intervention T, timepoint

Table 1

Demographic characteristics of participants in intervention trial to evaluate CRC screening discussions, *n* (%) unless otherwise indicated

Variable	Computer Group (<i>n</i> =319)	Brochure Group (<i>n</i> =340)	<i>p</i> - value
Age (years), M (SD)	56.8 (6.0)	57.8 (6.4)	0.046
Years of education, M (SD)	12.2 (1.8)	12.3 (1.9)	0.683
Gender:			
Male	154 (48)	161 (47)	0.785
Female	165 (52)	180 (53)	
Married/partnered			
Yes	95 (30)	108 (32)	0.5814
No	224 (70)	232 (68)	
Employed			
Yes	71 (22)	73 (21)	0.792
No	248 (78)	268 (79)	
Insurance			
Yes	291 (91)	300 (88)	0.208
No	28 (9)	40 (12)	
Income (\$)			
<15,000	185 (60)	180 (55)	0.350
15,000–30,000	84 (27)	106 (32)	
>30,000	38 (12)	40 (12)	
Site			
VA	72 (23)	63 (18)	0.192
Non-VA	247 (77)	278 (82)	

Note: *t*-tests were performed for continuous variables (age, years of education); chi-square tests were performed for categorical variables (all others).

VA, Veterans Affairs

Table 2

Differences between groups for self-reported CRC screening discussions.

Predictor variables	Discussion of any colon test	
	OR (95% CI)	p-value
Univariate Model with Intervention:		
Intervention: computer vs brochure (ref)	1.81 (1.32, 2.47)	<0.001
Final Multivariable Model:		
Intervention: computer vs brochure (ref)	1.75 (1.21, 2.54)	0.003
Age (years)	0.95 (0.92, 0.98)	0.003
Gender: male vs female (ref)	1.20 (0.87, 1.66)	0.274
Education	1.05 (0.96, 1.16)	0.277
Married ^a	0.68 (0.47, 0.99)	0.045
Reason for visit: acute illness vs preventive care (ref)	0.62 (0.37, 1.04)	0.070
Reason for visit: not seen by a doctor vs preventive care (ref)	0.33 (0.14, 0.77)	0.011
Objective CRC risk	1.46 (0.80, 2.67)	0.219
Perceived CRC risk	1.19 (0.99, 1.42)	0.060
FOBT benefits	1.06 (0.78, 1.43)	0.729
FOBT self-efficacy	0.84 (0.59, 1.21)	0.356
COL benefits	1.21 (0.81, 1.82)	0.354
COL self-efficacy	1.73 (1.22, 2.44)	0.002
Cancer fatalism	0.82 (0.63, 1.06)	0.135
Knowledge	1.01 (0.93, 1.10)	0.823
Doctor recommendation of FOBT ^a	1.15 (0.81, 1.63)	0.420
Doctor recommendation of COL ^a	1.34 (0.93, 1.92)	0.114
Family/friend recommendation ^a	1.81 (1.24, 2.63)	0.002

Note: Boldface indicates significance.

^a Yes vs No (ref)

COL, colonoscopy; CRC, colorectal cancer; FOBT, fecal occult blood test