Figure 2 Illustration of a D-shaped incision with modified Burow's triangular excision for advancement (shaded). The D shape is skewed laterally.

Figure 3 The defect is closed, from medial to lateral. A skin hook marks the lateral extent of the 'D' to allow accurate apposition of the skin at the apex. The anterolateral part of the superior skin flap is actively advanced into a more anterior position.

my have included the fish-tail plasty method, which results in a Y-shaped closure.¹⁻³ However, the disadvantages of this method are that it involves additional scarring, potentially creates two smaller dog ears and can be complicated by skin necrosis at the Y confluence. Another alternative is a teardrop shape incision but this is less predictable in its outcome depending on the point at which the lateral apex of closure is commenced.⁴

Our technique results in consistently good cosmetic and functional outcomes. The triangular component of the technique may be regraded as a modified Burow's triangle excision.⁵ Excess tissue is removed laterally while avoiding buckling of the tissues that would create a dog-ear effect.

References

1. Farrar WB, Fanning WJ. Eliminating the dog-ear in modified radical mastec-

Figure 4 Final result

tomy. *Am J Surg* 1988; **156**: 401-402.

2. Nowacki MP, Towpik E, Tchórzewska H. Early experience with 'fish-shaped' incision for mastectomy. *Eur J Surg Oncol* 1991; **17**: 615-617.
3. Hussien M, Daltrey IR, Dutta S *et al*. Fish-tail plasty: a safe technique to improve cosmesis at the lateral end of mastectomy scars. *Breast* 2004; **13**: 206-209.
4. Mirza M, Sinha KS, Fortes-Mayer K. Tear-drop incision for mastectomy to avoid dog-ear deformity. *Ann R Coll Surg Engl* 2003; **85**: 131.
5. Burow KA. *Berliner Klinische Wochenschrift* 1870; 350-352.

Progressive dilatation of stenosed suprapubic tracts using Nelaton catheters in the emergency setting

A Shrestha, N Ravindranath, A Jain
Royal Oldham Hospital, Oldham, UK

CORRESPONDENCE TO

Anne Shrestha, E: shrestha.anne@doctors.net.uk

BACKGROUND

Suprapubic catheter (SPC) changes are generally managed in the community by district nurses. Failed reinsertion of SPCs is frequently encountered and usually dealt with by junior doctors in hospital in an emergency setting. An extensive literature review revealed only two related publications, both describing the use of an Otis urethrotome to dilate the old tract.^{1,2} There have been anecdotal references on the use of progressive dilatation.

TECHNIQUE

We present a technique using single use hydrophilic-coated female Nelaton catheters to progressively dilate the stenosed tract. Ultrasonography is performed to ensure a full bladder. After analgesia and local anaesthetic infiltration, starting with a well lubricated 8Fr catheter, the tract is dilated up to 18Fr. Successful insertion into the bladder each time is confirmed by efflux of urine. After dilatation, an 18Fr silicone long-term Foley catheter is inserted.

DISCUSSION

This is a simple procedure that can be carried out safely in an emer-

Figure 1 Hydrophilic-coated female Nelaton catheters

gency setting by junior doctors. It allows early discharge of the patient and saves additional invasive procedures. It is less traumatic compared to other methods as well as being cost effective. A silicone catheter is preferable to a latex catheter as it is more rigid and therefore easier to reinsert via the dilated tract. We have dilated five tracts in the last two months without any complications.

References

1. Thrasher JB, Kreder KJ. Suprapubic tube tract dilation using the Otis urethrotome. *Urology* 1993; **41**: 247–248.
2. McInnes BK. Use of Otis urethrotome to reopen stenotic suprapubic tract. *Urology* 1981; **17**: 373.

TECHNICAL TIPS

Aiding the insertion of polyurethane coated breast implants

ME Smith, AJ Durrani
Addenbrooke's Hospital, Cambridge, UK

CORRESPONDENCE TO

Matthew Smith, E: matthewsmith@addenbrookes.nhs.net

Polyurethane foam covered implants are a popular choice for breast augmentation but insertion can be more problematic than with standard implants due to the highly adherent textured surface. We utilise a simple method to facilitate implant insertion and adjustment, and simultaneously deliver perioperative analgesia. Prior to implant insertion, 11ml of Instillagel® (lidocaine hydrochloride 2.0%, chlorhexidine digluconate 0.25%; CliniMed Ltd, High Wycombe, HP10 9QY) is placed within the manufacturer's plastic insertion sleeve. The gel surrounds the implant, lubricating and delivering anaesthetic to the pocket. We routinely use this technique without complication or evidence of implant migration or rotation at follow up.

The modified recovery position for posterior leg surgery

JMD Williams, A Bebbington
Morriston Hospital, Swansea, UK

CORRESPONDENCE TO

John Williams, E: djwilliams1980@yahoo.co.uk

Conventionally used to approach the posterior leg, the prone position poses an anaesthetic challenge with changes in respiratory and cardiovascular physiology and an increased risk of pressure injuries and nerve damage.¹ To avoid prone positioning, for an approach to the left leg, the patient is placed in the left lateral position with hips and shoulders slightly rotated. With the right hip and knee flexed, the torso remains in the lateral position while the left leg is optimally placed for unilateral posterior leg surgery such as an Achilles' tendon repair.

Reference

1. Edcombe H, Carter K, Yarrow S. Anaesthesia in the prone position. *Br J Anaesth* 2008; **100**: 165–183.

Figure 1 Patient positioned to approach the posterior left leg