Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading doi:10.1016/S2221-1691(11)60211-6 © 2012 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Larvicidal and irritant activities of hexane leaf extracts of *Citrus sinensis* against dengue vector *Aedes aegypti* L.

Radhika Warikoo, Ankita Ray, Jasdeep Kaur Sandhu, Roopa Samal, Naim Wahab, Sarita Kumar*

Department of Zoology, Acharya Narendra Dev College, University of Delhi, New Delhi-110019, India

ARTICLE INFO

Article history:

Received 15 July 2011
 Received in revised form 1 August 2011
 Accepted 20 August 2011
 Available online 28 February 2012

Keywords:

Citrus
 Hexane
 Larvicide
 Irritancy
 Take-offs
Citrus sinensis
Aedes aegypti
 Larvicidal activity
 Irritant activity
 Dengue vector
 Mosquito control

ABSTRACT

Objective: To assess the larvicidal and irritant activities of the hexane extracts of leaves of *Citrus sinensis* (*C. sinensis*) against the early fourth instars and female adults of *Aedes aegypti* (*Ae. aegypti*). **Methods:** The larvicidal potential of the prepared leaf extract was evaluated against early fourth instar larvae of *Ae. aegypti* using WHO protocol. The mortality counts were made after 24 h and LC₅₀ and LC₉₀ values were calculated. The efficacy of extract as mosquito irritant was assessed by contact irritancy assays. Extract-impregnated paper was placed on a glass plate over which a perspex funnel with a hole on the top was kept inverted. Single female adult, 3-day old unfed/blood-fed, was released inside the funnel. After 3 min of acclimatization time, the time taken for the first take-off and total number of flights undertaken during 15 min were scored. **Results:** The citrus leaf extracts from hexane possessed moderate larvicidal efficiency against dengue vector. The bioassays resulted in an LC₅₀ and LC₉₀ value of 446.84 and 1 370.96 ppm, respectively after 24 h of exposure. However, the extracts were proved to be remarkable irritant against adults *Ae. aegypti*, more pronounced effects being observed on blood-fed females than unfed females. The extract-impregnated paper was thus proved to be 7–11 times more irritable as compared with the control paper. **Conclusions:** The hexane extracts from *C. sinensis* leaves are proved to be reasonably larvicidal but remarkably irritant against dengue vector. Further studies are needed to identify the possible role of extract as adulticide, oviposition deterrent and ovicidal agent. The isolation of active ingredient from the extract could help in formulating strategies for mosquito control.

1. Introduction

Mosquitoes being vector for many tropical and subtropical diseases are the most important single group of insects well-known for their public health importance. Despite progress in vaccine development, no effective and acceptable multi-valent vaccines are currently available against mosquito-borne diseases. The approach to combat these diseases largely relies on interruption of the disease transmission cycle by either destruction of the aquatic stages or by killing the adult mosquitoes using chemical insecticides[1,2].

The drastic effects of chemical insecticides-based intervention measures for the control of disease vectors have received wide public apprehension and have caused many problems like insecticide resistance, resurgence of pest

species, environmental pollution, toxic hazards to humans and other non-target organisms[3]. To alleviate these problems, major emphasis has been on the use of natural plant-based products as larvicides which can provide an alternate to synthetic insecticides[4]. Plants are rich sources of bioactive compounds that can be used to develop environmentally safe vector and pest-managing agents. A number of plants have been reported as selective with little or no harmful effect on non-target organisms and the environment. The botanical extracts from the plant leaves, roots, seeds, flowers and bark in their crude form have been used as conventional insecticides for centuries[5]. Plant products are important natural alternatives to insecticides, as phytochemicals extracted from the whole plant or specific part of the plant using different solvents may also act against mosquitoes as toxicant, growth regulators, repellents and ovipositional deterrent[6–8].

The preliminary screening is a good mean of evaluation of the potential larvicidal activity of plants popularly used for this purpose. *Citrus* plants have been known for their use as food, food-flavoring agents, and also for their mosquitocidal components. Different parts of the *Citrus* plant i.e. fruits, seeds, roots and leaves, have been tested

*Corresponding author: Sarita Kumar, Department of Zoology, Acharya Narendra Dev College, University of Delhi, New Delhi-110019, India.

Tel: +91-981-0309912

Fax: +91-11-26294540

E-mail: sarita.sanjay90@gmail.com

Foundation Project: This work was financially supported by University Grants Commission [grant No. F. 35-74/2009 (SR)].

for their use as mosquitocidal components^[9,10]. Amusan *et al*^[5] have tested the ethanolic extracts of the orange peel of *Citrus sinensis* (*C. sinensis*) for the toxicity effect on the larvae of the yellow fever mosquito *Aedes aegypti* (*Ae. aegypti*) and found them quite effective. Lee^[11] had tested essential oil of *Citrus bergamia* against larvae of *Ae. aegypti* and *Culex pipiens*, while Amer and Mehlhorn^[6] investigated essential oil from *Citrus limon* against *Ae. aegypti* and *Culex quinquefasciatus*, and *Anopheles stephensi*. The essential oils contained in the rind of the fruit and the leaves of bergamot from Greece (*C. aurantium* subsp. *bergamia*) have also been studied^[12]. All these reports suggest the toxic potential of *Citrus* plant against mosquitoes. Keeping in view this and that not much studies have been carried out on the *Citrus* leaves for the possible potential for dengue vector control, present investigations were performed to study the larvicidal and irritability effects of the hexane extract of *C. sinensis* leaves against dengue fever mosquito, *Ae. aegypti*. The *Citrus* leaves being commonly and widely available may prove to be an excellent alternative for mosquito control because of non-toxicity to environment and non-target organisms.

2. Materials and methods

2.1. Preparation of leaf extract

The fresh and tender leaves of *C. sinensis* were collected from the surrounding areas. These were thoroughly washed with tap water and kept under shade at room temperature [$(27 \pm 2)^\circ\text{C}$] for about 20 days till they dried completely. The dried leaves were mechanically grinded and sieved to get fine powder. The 200 g of powdered leaves was soaked in 1 L of hexane. The soaked material was left undisturbed for five days and the crude extract, thus formed, was concentrated using a vacuum evaporator at 45°C under low pressure. After complete evaporation of the solvent the concentrated extract was collected and stored in a refrigerator at 4°C as the stock solution of 1 000 ppm for further use.

2.2. Mosquitoes

The present investigations employed the dengue fever mosquito, *Ae. aegypti* originated from fields of Delhi and surrounding areas. The colony was maintained in an insectary at $(28 \pm 1)^\circ\text{C}$, $(80 \pm 5)\%$ relative humidity and 14/10 light:dark photoperiod^[13].

2.3. Larvicidal bioassays

The larvicidal bioassay was performed at $(28 \pm 1)^\circ\text{C}$ on the *Ae. aegypti* larvae in accordance with the procedure described by WHO with slight modifications^[14]. For experimental treatment, the graded series of the leaf extract was prepared using 5 parts of hexane and 95 parts of acetone as the diluting solvent. The early fourth instar larvae of *Ae. aegypti*, in batches of 25, were taken in plastic bowls containing 99 mL of distilled water and transferred to glass jar containing 150 mL of distilled water and 1 mL of the particular concentration of the leaf extract. Four replicates were carried out simultaneously for each dilution.

Controls were exposed to the 1 mL of the diluting solvent, *i.e.* 5 parts of hexane and 95 parts of acetone. During the treatment period, the larvae were not provided with any food. The dead and moribund larvae were recorded after 24 h as larval mortality.

2.4. Statistical analysis of data

The tests with more than 20% mortality in controls and pupae formed were discarded and repeated again. If the control mortality ranged between 5%–20%, it was corrected using Abbott's formula (Abbott 1925).

The data were subjected to regression analysis using computerized SPSS 11.5 Programme. The LC_{50} and LC_{90} values with 95% fiducial limits were calculated in each bioassay to measure difference between the test samples. The results obtained with different extracts were analyzed using Student's *t*-test with statistical significance considered for $P \leq 0.05$.

2.5. Contact irritancy assays

Whatman filter paper circles were impregnated with 100 ppm hexane leaf extracts, separately. These papers were completely dried and used afresh for contact irritancy assays. Each paper was placed on a glass plate and a perspex funnel with a hole on the top was kept inverted over them. Single female adult, 3-day old unfed, was released inside the funnel on the paper. The hole on the top of funnel was plugged with cotton. The adult was allowed to settle for 3 min after which the time taken for the first take-off was recorded. The experiment was continued for 15 min during which the total number of flights undertaken by the mosquito was scored.

Parallel control tests were performed with acetone-impregnated papers. Similar experiments were also carried out with freshly blood-fed females to compare the repellency effects of the *Citrus* leaves. Each treatment had 3 replicates. Data were analyzed and the relative irritability of the extract in each case was calculated with respect to control.

3. Results

The results of the larvicidal bioassay performed on the early fourth instars and contact irritancy carried out on the female adults of *Ae. aegypti* with the hexane leaf extract of *C. sinensis* were presented in Table 1 and Table 2. The results obtained showed the moderate larvicidal efficacy of the hexane leaf extract, but the extracts proved to be remarkably significant irritant against the dengue vector. The larvicidal bioassay resulted in the LC_{50} value of 446.84 ppm and LC_{90} value of 1 370.96 ppm (Table 1). The larvae remained alive developed into normal pupae and then adults within (60–72 h). No morphological alterations or the delayed mortality was noticed during the development negating the possibility of delayed toxicity of leaf extracts.

When the female adults of *Ae. aegypti* were subjected to contact irritancy assays, a significant behavioural response was observed. The most remarkable response on exposure to leaf extract was observed in blood-fed mosquitoes with

first flight only after 4 sec and an average total of 54.33 take-offs as against only 5.00 take-offs against control (Table 2). Against unfed females though the extract was relatively less effective yet caused a significant number of 34.33 take-offs in just 15 min. The extract-impregnated paper was thus proved to be 7–11 times more irritable as compared with the control paper.

Table 1

Larvicidal activity (in ppm) of hexane extract prepared from the leaves of *C. sinensis* against early fourth instars of *Ae. aegypti*.

Parameters	LC ₃₀	LC ₅₀	LC ₇₀	LC ₉₀	S.E.	χ^2 (df)	Regression coefficient
Value	282.44	446.84	706.92	1370.96	1.40	4.94	2.63
Upper FL	99.31	269.22	508.75	863.92	–	(4)	–
Lower FL	403.65	639.03	1451.36	7002.14	–	–	–

FL indicates the 95% lower and upper fiducial limits.

S.E. indicates standard error and *df* indicates degrees of freedom.

Table 2

Response of 3-day old adult females (blood-fed and non blood-fed) of *Ae. aegypti* to papers impregnated with hexane extracts of *C. sinensis* leaves in the contact irritancy assays (Mean $\pm$ SEM).

Test	Mean time lapse before first take- off (in min)	Mean number of take-offs for females (in 15 min)	Relative irritability
Control	0.00 $\pm$ 0.00 ^a (0.00)	5.00 $\pm$ 2.00 ^c	1.00 (1.15)
Non blood-fed	3.30 $\pm$ 0.00 ^a (1.00)	34.33 $\pm$ 8.69 ^d	6.86 (15.04)
Blood-fed	4.00 $\pm$ 0.58 ^b (1.00)	54.33 $\pm$ 21.60 ^d	10.86 (37.42)

Figures in parentheses indicate standard deviation value.

Figures in each column followed by the same letter are not significantly different at $P=0.05$ (Students' paired *t*-test).

4. Discussion

Natural pesticides derived from plants are a promising tool especially for targeting mosquitoes in the larval stage^[6]. Different parts of the *Citrus* plant *i.e.* fruits, seeds, roots and leaves have been tested for their use as mosquitocidal components^[9,10]. A number of studies have also been carried out on the larvicidal potential of essential oil extracted from the *Citrus* leaves and peels^[11,12]. However, very few reports are available regarding the larvicidal effects of crude extracts of *Citrus* leaves against *Ae. aegypti* whereas no work reports about their irritancy effects against the female adults.

Our investigations showed that 24 h exposure to early fourth instars of *Ae. aegypti* with hexane extracts of the leaves of *C. sinensis* resulted in 50% mortality at 446.84 ppm. This value is much higher than that obtained by Cavalcanti *et al*^[15] who reported LC₅₀ value of 69 ppm but with essential oil extracted from *Cymbopogon citratus* against early fourth instars of *Ae. aegypti*. Bagavan *et al*^[16] tested the crude extracts of *Citrus* peels against mosquito larvae and obtained the LC₅₀ value of 58.25 ppm and LC₉₀ value of 298.31 ppm with the chloroform extract of *C. sinensis* peels against *Anopheles subpictus* larvae; and the LC₅₀ value of 38.15 ppm and LC₉₀ value of 184.67 ppm with the peel methanolic extract against the larvae of *Culex tritaeniorhynchus*. Traboulsi *et al*^[9] have accounted the insecticidal activity of essential oil extracts from leaves, flowers and roots of *C. sinensis* Osbeck against fourth instar larvae of the mosquito *Culex pipiens molestus*

Forsk as 60.0 mg/L.

It has been reported that behavioural modifications of the mosquitoes through chemical actions by contact irritancy, are expected to have population effects that may ultimately reduce disease transmission^[17]. Several reports are available regarding the repellent properties of essential oils against adult mosquitoes^[18–20]; however very few works have been reported proving the irritability potential of leaf extracts and practically no work with *Citrus* leaf extracts. Our studies showed the significant irritancy potential of hexane extracts of *Citrus* leaves against the adults of *Ae. aegypti* as it caused 54.33 flights in blood-fed females and 34.33 flights in unfed females when exposed for just 15 min. The effect was, thus more pronounced in blood-fed females (10.86 irritability ratio) than unfed females (6.86 irritability ratio). This result may prove significant as the extract could be used as a repellent on the human skin surface to keep away the mosquitoes from biting. Similar significant repellency behaviour was reported in *Ae. aegypti* when the female adults were exposed to various extracts prepared from the leaves of *Parthenium hysterophorus*^[13] and to the crude leaf extracts of *Cassia fistula* prepared in acetone^[21]. The application of *Mentha piperita* oil on human skin had resulted in 100% protection till 150 min after which only 1–2 bites were reported for next 30 min^[22]. Essential oils from some Verbenaceae plants have shown repellent and/or insecticidal effects against mosquitoes^[7,23]. The five most effective oils reported were those of litsea (*Litsea cubeba*), cajeput (*Melaleuca leucadendron*), niaouli (*Melaleuca quinquenervia*), violet (*Viola odorata*), and catnip (*Nepeta cataria*), which induced a 100% repellency against *Ae. aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus*^[7]. Petroleum ether extracts of *Vitex negundo* leaves offered 8 h of bite protection (2 mg/cm²) by different mosquito species in the field^[24,25]. A patented extract of the seeds of *Vitex agnus castus* was shown effective against mosquitoes resulting in more than 6 h protection against *Culex quinquefasciatus* and *Anopheles stephensi*, and 3–4 h against *Ae. aegypti*^[26]. Nour *et al*^[27] suggested the use of the essential oils of *Ocimum basilicum* as promising new natural repellents at 0.1% concentration against *Anopheles* mosquito.

Mosquito control is vital for many countries and is still in a state of evolution. During the last decades, it depended upon synthetic organic insecticides, many of which have been removed from the arsenal of weapons^[28] and botanicals are the new weapons of mosquito control under exploration. The activity of crude plant extracts is often attributed to the complex mixture of active compounds. Natural pesticides derived from plants are a promising tool especially for targeting mosquitoes in the larval stage^[6], but there are some factors that need more investigation in order to extract useful conclusions: light and storing conditions^[29], enantioselectivity of major and minor ingredients^[30], harvesting time and plant material^[12], *etc.* Further studies are needed to identify the possible role of extract as adulticide, oviposition deterrent and ovicidal agent. The isolation of active ingredient from the extract could help in formulating strategies for mosquito control.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors are highly grateful to University Grants Commission for providing financial assistance to carry out the present investigations. Thanks are extended to Dr. Savithri Singh, Principal, Acharya Narendra Dev College for providing infrastructure and research facilities.

References

- [1] Joseph CC, Ndoile MM, Malima RC, Nkuniya MHM. Larvicidal and mosquitocidal extracts, a coumrin, isoflavonoids and pterocarpanes from *Neorautanenia mitis*. *Trans R Soc Trop Med Hyg* 2004; **98**: 451–455.
- [2] Grieco JP, Achee NL, Chareonviriyaphap T, Suwonkerd W, Chauhan KR, Sardelis M, et al. A new classification system for the actions of IRS chemicals traditionally used for malaria control. *PLoS ONE* 2007; **2**: e716.
- [3] Sarwar M, Ahmad N, Toufiq M. Host plant resistance relationships in chickpea (*Cicer arietinum* L.) against gram pod borer (*Helicoverpa armigera* Hubner). *Pak J Bot* 2009; **41**: 3047–3052.
- [4] Zhu J, Zeng X, Yanma, Liu T, Qian K, Han Y, et al. Adult repellency and larvicidal activity of five plant essential oils against mosquitoes. *J Am Mosq Control Assoc* 2006; **3**: 515–522.
- [5] Amusan AA, Idowu AB, Arowolo FS. Comparative toxicity effect of bush tea leaves (*Hyptis suaveolens*) and orange peel (*Citrus sinensis*) oil extract on larvae of the yellow fever mosquito *Aedes aegypti*. *Tanzan Health Res Bull* 2005; **7**: 174–178.
- [6] Amer A, Mehlhorn H. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitol Res* 2006a; **99**: 466–472.
- [7] Amer A, Mehlhorn H. Repellency effect of forty-one essential oils against *Aedes*, *Anopheles* and *Culex* mosquitoes. *Parasitol Res* 2006c; **99**: 478–490.
- [8] Rajkumar S, Jebanesan A. Larvicidal and oviposition activity of *Cassia obtusifolia* Linn (Family: Leguminosae) leaf extract against malarial vector, *Anopheles stephensi* Liston (Diptera: Culicidae). *Parasitol Res* 2009; **104**: 337–340.
- [9] Traboulsi AF, El-Haj S, Tuani M, Taoubi K, Nader NA, Mrad A. Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Manag Sci* 2005; **61**: 597–604.
- [10] Akram W, Khan HAA, Hafeez F, Bilal H, Kim YK, Lee JJ. Potential of citrus seed extracts against dengue fever mosquito, *Aedes albopictus* (Skuse) (Culicidae: Diptera). *Pak J Bot* 2010; **42**: 3343–3348.
- [11] Lee HS. Mosquito larvicidal activity of aromatic medicinal plant oils against *Aedes aegypti* and *Culex pipiens pallens*. *J Am Mosq Control Assoc* 2006; **22**: 292–295.
- [12] Melliou E, Michaelakis A, Koliopoulos G, Skaltsounis AL, Magiatis P. High quality bergamot oil from Greece: chemical analysis using enantiomeric GC–MS and larvicidal activity against the West Nile virus vector. *Molecules* 2009; **14**: 839–849.
- [13] Kumar S, Singh AP, Nair G, Batra S, Seth A, Wahab N, et al. Impact of *Parthenium hysterophorus* leaf extracts on the fecundity, fertility and behavioural response of *Aedes aegypti* L. *Parasitol Res* 2011a; **108**: 853–859.
- [14] WHO. *World malaria report*. WHO/UNICEF. Geneva: WHO; 2005.
- [15] Cavalcanti ES, Morais SM, Lima MA, Santana EW. Larvicidal activity of essential oils from Brazilian plants against *Aedes aegypti* L. *Mem Inst Oswaldo Cruz* 2004; **99**: 541–544.
- [16] Bagavan A, Kamaraj C, Rahuman AA, Elango G, Zahir AA, Pandiyan G. Evaluation of larvicidal and nymphicidal potential of plant extracts against *Anopheles subpictus* Grassi, *Culex tritaeniorhynchus* Giles and *Aphis gossypii* Glover. *Parasitol Res* 2008; **104**: 1109–1117.
- [17] Said SH, Grieco JP, Achee NL. Evaluation of contact irritant and spatial repellent behavioural responses of male *Aedes aegypti* to vector control compounds. *J Am Mosq Control Assoc* 2009; **25**: 436–441.
- [18] Rajkumar S, Jebanesan A. Repellent activities of selected plant essential oils against the malarial fever mosquito *Anopheles stephensi*. *Trop Biomed* 2007; **24**: 71–75.
- [19] Rajkumar S, Jebanesan A. Bioactivity of flavonoid compounds from *Poncirus trifoliata* L. (Family: Rutaceae) against the dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *J Parasitol Res* 2008; **104**: 19–25.
- [20] Oyewolo I, Ibiapo C, Moronkola D, Doula A, Adeoye G, Anyasor G, et al. Antimalarial and repellent activities of *Tithonia diversifolia* (Hemsl.) leaf extracts. *J Med Plant Res* 2008; **2**: 171–175.
- [21] Govindarajan M. Bioefficacy of *Cassia fistula* Linn. (Leguminosae) leaf extract against chikungunya vector, *Aedes aegypti* (Diptera: Culicidae). *Eur Rev Med Pharmacol Sci* 2009; **13**: 99–103.
- [22] Kumar S, Wahab N, Warikoo R. Bioefficacy of *Mentha piperita* essential oil against dengue fever mosquito, *Aedes aegypti* L. *Asian Pac J Trop Biomed* 2011; **1**: 90–93.
- [23] Gillij YG, Gleiser RM, Zygadlo JA. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. *Bioresour Technol* 2008; **99**: 2507–2515.
- [24] Karunamoorthi K, Mulelam A, Wassie F. Laboratory evaluation of traditional insect/mosquito repellent plants against *Anopheles arabiensis*, the predominant malaria vector in Ethiopia. *Parasitol Res* 2008a; **103**: 529–534.
- [25] Karunamoorthi K, Ramanujam S, Rathinasamy R. Evaluation of leaf extracts of *Vitex negundo* L. (Family: Verbenaceae) against larvae of *Culex tritaeniorhynchus* and repellent activity on adult vector mosquitoes. *Parasitol Res* 2008b; **103**: 545–550.
- [26] Mehlhorn H, Schmahl G, Schmidt J. Extract of the seeds of the plant *Vitex agnus castus* proven to be highly efficacious as a repellent against ticks, fleas, mosquitoes and biting flies. *Parasitol Res* 2005; **95**: 363–365.
- [27] Nour AH, Elhussien SA, Osman NA, Nour AH. Repellent activities of the essential oils of four Sudanese accessions of basil (*Ocimum basilicum* L.) against *Anopheles* mosquito. *J Appl Sci* 2009; **9**: 2645–2648.
- [28] Floore TG. Mosquito larval control practices: past and present. *J Am Mosq Control Assoc* 2006; **22**: 527–533.
- [29] Amer A, Mehlhorn H. Persistency of larvicidal effects of plant oil extracts under different storage conditions. *Parasitol Res* 2006b; **99**: 473–477.
- [30] Michaelakis A, Koliopoulos G, Stroggilos A, Bouzas E, Couladouros EA. Larvicidal activity of naturally occurring naphthoquinones and derivatives against the West Nile virus vector *Culex pipiens*. *Parasitol Res* 2009; **104**: 657–662.