ultrasound has been a part of the HNPCC/LS screening guidelines for years, but several studies have shown transvaginal ultrasound to be unhelpful for both endometrial and ovarian cancer early detection (12). Ovarian cancer screening remains problematic in all settings. However, we have been encouraged with results from a combined colon/endometrial screening program in which at-risk women underwent colonoscopy and, while under sedation from this examination, endometrial biopsy was performed (13). Although there were modest logistic considerations in coordinating schedules of endoscopists and gynecologists, the approach has been well-received by patients. It is too early to evaluate endometrial biopsy sensitivity and specificity, but some screening experience is being gained (14).

As clinicians from various fields become more familiar with HNPCC/LS, the opportunity to diagnose the condition based on a presentation of endometrial cancer will be possible. The study by Win et al. (8) was from a series of colorectal cancer patients worked up for evidence of MMR deficiency, but the same approach could have been taken with a series of endometrial cancer patients. Such efforts have been taken and largely parallel the experience when colorectal cancer is the starting point. Using the Bethesda Guidelines or modifications, a diagnostic algorithm can be followed (5,7,15). Selecting cases for microsatellite instability/immunohistochemistry according to various age cutoffs has been shown to have an acceptable sensitivity for detecting HNPCC/LS.

In conclusion, we have seen that patients whose sentinel HNPCC tumor is an endometrial cancer have a considerable risk of other cancer. The Win et al. study provides the most unbiased, and thus likely the most accurate, data to date on this issue. These data are highly consistent with other earlier reports and again support a modest increase in risk of breast cancer in HNPCC/LS. Consequently, it is time to redouble efforts to conduct reasonable screening trials to establish the screening benefit for those extracolonic tumors that have been relatively neglected to date. Substantial challenges exist in considering such screening. Although the relative risk of small bowel cancer is an extraordinary 63-fold after endometrial cancer, its absolute rarity still leaves the potential usefulness of screening, such as by capsule endoscopy, an open question. The standardized incidence ratio for pancreatic cancer of nearly 9 suggests a possible role for screening, but the costs (high) and yields (probably low) of tools such as endoscopic ultrasound and magnetic resonance imaging leave this an area of controversy as well. But at least now we have better measures of the actual risks.

References

- Hiripi E, et al. Survival from common and rare cancers in Germany in the early 21st century. *Ann Oncol*. 2012;23(2):472–479.
- Moreira L, et al. Identification of Lynch syndrome among patients with colorectal cancer. *JAMA*. 2012;308(15):1555–1565.
- Pinol V, et al. Accuracy of revised Bethesda guidelines, microsatellite instability, and immunohistochemistry for the identification of patients with hereditary nonpolyposis colorectal cancer. *JAMA*. 2005;293(16):1986–1994.
- Hampel H, et al. Screening for the Lynch syndrome (hereditary nonpolyposis colorectal cancer). *N Engl J Med*. 2005;352(18):1851–1860.
- Lu KH, et al. Prospective determination of prevalence of Lynch syndrome in young women with endometrial cancer. *J Clin Oncol*. 2007;25(33):5158–5164.
- Peterson LM, et al. Molecular characterization of endometrial cancer: a correlative study assessing microsatellite instability, *MLH1* hypermethylation, DNA mismatch repair protein expression, and *PTEN*, *PIK3CA*, *KRAS*, and *BRAF* mutation analysis. *Int J Gynecol Pathol*. 2012;31(3):195–205.
- Leenen CH, et al. Prospective evaluation of molecular screening for Lynch syndrome in patients with endometrial cancer ≤ 70 years. *Gynecol Oncol*. 2012;125(2):414–420.
- Win AK, Lindor NM, Winship I, et al. Risks of colorectal and other cancers following endometrial cancer for women with Lynch syndrome. *J Natl Cancer Inst*. 2012;105(4):274–279.
- Engel C, et al. Efficacy of annual colonoscopic surveillance in individuals with hereditary nonpolyposis colorectal cancer. *Clin Gastroenterol Hepatol*. 2010;8(2):174–182.
- Win AK, et al. Colorectal and other cancer risks for carriers and noncarriers from families with a DNA mismatch repair gene mutation: a prospective cohort study. *J Clin Oncol*. 2012;30(9):958–964.
- Lindor NM, et al. Recommendations for the care of individuals with an inherited predisposition to Lynch syndrome: a systematic review. *JAMA*. 2006;296(12):1507–1517.
- Auranen A, Joutsiniemi T. A systematic review of gynecological cancer surveillance in women belonging to hereditary nonpolyposis colorectal cancer (Lynch syndrome) families. *Acta Obstet Gynecol Scand*. 2011;90(5):437–444.
- Huang M, et al. Prospective study of combined colon and endometrial cancer screening in women with Lynch syndrome: a patient-centered approach. *J Oncol Pract*. 2011;7(1):43–47.
- Manchanda R, et al. Annual outpatient hysteroscopy and endometrial sampling (OHES) in HNPCC/Lynch syndrome (LS). *Arch Gynecol Obstet*. 2012;286(6):1555–1562.
- Walsh CS, et al. Lynch syndrome among gynecologic oncology patients meeting Bethesda guidelines for screening. *Gynecol Oncol*. 2010;116(3):516–521.

Notes

Affiliation of author: Department of Gastroenterology, Hepatology, Nutrition, University of Texas MD Anderson Cancer Center, Houston, TX (PML).

DOI:10.1093/jnci/djs641

Advance Access publication January 24, 2013

Published by Oxford University Press 2013.

Out of Pocket, Out of Sight? An Unmeasured Component of the Burden of Cancer

Matthew P. Banegas, K. Robin Yabroff

Correspondence to: Matthew P. Banegas, PhD, Health Services and Economics Branch Applied Research Program, Division of Cancer Control and Population Sciences, National Cancer Institute, 6130 Executive Blvd, Rm 4019, Bethesda, MD 20892–7344 (e-mail: banegasmp@mail.nih.gov).

While tremendous strides in prevention and treatment of cancer have been made in recent years, accumulating evidence shows that

the costs associated with these advances have led to an increasing financial burden on health-care systems, patients and their

families, and society overall (1,2). Medical care costs can often be documented from existing administrative health-care data systems, although other aspects of the financial burden of cancer—particularly, those aspects related to patient costs—are not measured systematically and, as a result, are rarely reported. In this issue of the Journal, Lauzier et al. (3) add valuable information to our understanding of the financial impact of cancer, describing out-of-pocket expenditures in the first year after treatment initiation among a cohort of women diagnosed with early-stage breast cancer and their spouses from Quebec, Canada. Using multiple measurement time points, the authors detail out-of-pocket costs related to cancer treatment, including those associated with transportation, domestic help, and other accommodations. Lauzier et al. report median net out-of-pocket costs of \$1,002 for patients, comprised predominantly of costs related to travel, accommodations and management of side effects, and \$111 for spouses (2003 Canadian dollars). Further, the authors underscore wage loss as the primary driver of patients' perceived change for the worse in their family's financial situation.

Although the authors conclude that out-of-pocket costs for the first year after treatment initiation are probably not unmanageable for most women with early-stage breast cancer, the financial consequences reported likely understate those for breast cancer patients diagnosed with distant metastases and other patients with more treatment-intensive cancers. Additionally, estimates from Canada may not reflect the out-of-pocket costs experienced by cancer patients in the United States or other countries because of fundamental differences in the organization and financing of health care. The magnitude of financial burden on cancer patients can vary widely, especially in the complex US health-care system, which is comprised of multiple payers, coverage policies, annual/lifetime benefit limits and insurance caps, coinsurance, copayments, and deductibles (4). In addition to costs for transportation and endocrine therapy faced by patients in Quebec, expenses for insured cancer patients in the United States would include deductibles and copayments for hospitalizations, doctor visits, surgical visits, and adjuvant treatments. For cancer patients without health insurance or those who have already reached a lifetime benefit limit, all costs of care would be considered out-of-pocket costs. It is not surprising, then, that a large proportion of US cancer patients and their families report hardship paying their cancer bills (5), with many families using all or most of their savings (6).

Moreover, because health insurance in the working-age population is predominantly employer-based in the United States, rather than coverage provided universally as in Canada, maintaining insurance coverage during treatment can be more challenging for patients and their families. Cancer treatment can lead to substantial work loss for both patients and their families, which is manifested as fewer hours worked, change in jobs, or job loss—all of which can lead to a change in or loss of health insurance (4,7,8). Time demands associated with cancer treatment may also lead to reduced work and wage loss for patients and caregivers, an indirect cost of cancer care that affects families' outlook about their financial well-being, as indicated by Lauzier et al. (3). In the United States, cancer patients may choose to forgo and/or delay recommended medical care, even potentially life-saving cancer treatment, because of an inability to pay or in an attempt to avoid further debt (4,6,9–11). Additionally, the decision

to forgo or delay care because of cost may be more likely among individuals who are uninsured, low-income, and/or racial/ethnic minorities (6,9), which further perpetuates disparities that lead to poor health outcomes among the most vulnerable populations.

As such, understanding the potential financial consequences of cancer care is increasingly important. Cancer accounts for approximately 5% of total health-care spending in the United States (12) and 5% to 7% of the health-care budget in Canada (13). Furthermore, the costs of cancer care in most developed countries are expected to rise because of population growth and aging alone (1,2). Much attention has recently focused on the soaring price of cancer drugs (14), some with price tags of more than \$125 000 per drug per year (15), often with modest clinical benefits. For US patients facing a coinsurance of 20% to 33% for “top-tier” drugs (16), this can translate to out-of-pocket costs of approximately \$41 000 a year. This phenomenon of patients being exposed to higher costs for expensive cancer medications is becoming more common in Canada as well (17). Therefore, for both US and Canadian oncologists, consideration of the patient costs associated with cancer treatment will be central to comprehensive discussions with patients and treatment recommendations.

Nevertheless, for providers to be able to have substantive conversations about the costs of cancer treatment with their patients, more studies like that from Lauzier et al. (3) are needed to provide detailed longitudinal data on patient out-of-pocket costs, including those associated with medical treatment and wage loss. National data, such as the Medical Expenditure Panel Survey *Experiences with Cancer Supplement*, will provide information about the impact of cancer and its treatment on financial burden, health-care use and expenditures, and the employment of cancer survivors as well as their caregivers (18). Overall, such evidence on the scope of financial consequences of cancer care for patients and their families, in addition to the clinical benefits and risks, will enhance discussions of the “value” of cancer treatment, which is something oncologists in both the United States and Canada acknowledge as a fundamental component of high-quality care (17,19).

References

1. Mariotto AB, Yabroff KR, Shao Y, Feuer EJ, Brown ML. Projections of the cost of cancer care in the United States: 2010–2020. *J Natl Cancer Inst*. 2011;103(2):117–128.
2. Sullivan R, Peppercorn J, Sikora K, et al. Delivering affordable cancer care in high-income countries. *Lancet Oncol*. 2011;12(10):933–980.
3. Lauzier S, Lévesque P, Mondor M, et al. Out-of-pocket costs in the year after early breast cancer among Canadian women and spouses. *J Natl Cancer Inst*. 2013;105(4):280–292.
4. Kaiser Family Foundation. *Spending To Survive: Cancer Patients Confront Holes in the Health Insurance System*. <http://www.kff.org/insurance/7851.cfm>. Accessed December 20, 2012.
5. Shankaran V, Jolly S, Blough D, Ramsey SD. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol*. 2012;30(14):1608–1614.
6. USA Today, Kaiser Family Foundation, Harvard School of Public Health. *National Survey of Households Affected by Cancer*. <http://www.kff.org/kaiserpolls/pomr112006pkg.cfm>. Accessed December 20, 2012.
7. Bradley CJ, Neumark D, Luo Z, Bednarek HL. Employment-contingent health insurance, illness, and labor supply of women: evidence from married women with breast cancer. *Health Econ*. 2007;16(7):719–737.

8. Bernard DS, Farr SL, Fang Z. National estimates of out-of-pocket health care expenditure burdens among nonelderly adults with cancer: 2001 to 2008. *J Clin Oncol*. 2011;29(20):2821–2826.
9. Weaver KE, Rowland JH, Bellizzi KM, Aziz NM. Forgoing medical care because of cost: assessing disparities in healthcare access among cancer survivors living in the United States. *Cancer*. 2010;116(14):3493–3504.
10. Kirchhoff AC, Lyles CR, Fluchel M, Wright J, Leisenring W. Limitations in health care access and utilization among long-term survivors of adolescent and young adult cancer. *Cancer*. 2012;118(23):5964–5972.
11. Streeter SB, Schwartzberg L, Husain N, Johnsrud M. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *J Oncol Pract*. 2011;7(3 Suppl):46s–51s.
12. Pauly MV. Is high and growing spending on cancer treatment and prevention harmful to the United States economy? *J Clin Oncol*. 2007;25(2):171–174.
13. Polder JJ, Meerding WJ, Bonneux L, van der Maas PJ. A cross-national perspective on cost of illness: a comparison of studies from the Netherlands, Australia, Canada, Germany, United Kingdom, and Sweden. *Eur J Health Econ*. 2005;6(3):223–232.
14. Bach PB. Limits on Medicare's ability to control rising spending on cancer drugs. *N Engl J Med*. 2009;360(6):626–633.
15. Faden RR, Chalkidou K, Appleby J, Waters HR, Leider JP. Expensive cancer drugs: a comparison between the United States and the United Kingdom. *Milbank Q*. 2009;87(4):789–819.
16. Lee TH, Emanuel EJ. Tier 4 drugs and the fraying of the social compact. *N Engl J Med*. 2008;359(4):333–335.
17. Berry SR, Bell CM, Ubel PA, et al. Continental divide? The attitudes of US and Canadian oncologists on the costs, cost-effectiveness, and health policies associated with new cancer drugs. *J Clin Oncol*. 2010;28(27):4149–4153.
18. Yabroff KR, Dowling E, Rodriguez J, et al. The Medical Expenditure Panel Survey (MEPS) experiences with cancer survivorship supplement. *J Cancer Surviv*. 2012;6(4):407–419.
19. Meropol NJ, Schrag D, Smith TJ, et al. American Society of Clinical Oncology guidance statement: the cost of cancer care. *J Clin Oncol*. 2009;27(23):3868–3874.

Funding

The authors are employees of the National Cancer Institute at the National Institutes of Health and no additional funding was provided specifically for this work.

Notes

The authors declare no conflicts of interest.

Affiliation of authors: Division of Cancer Control and Population Sciences, National Cancer Institute, National Institutes of Health, Bethesda, MD (MPB, KRY).

DOI:10.1093/jnci/djs651

Advance Access publication January 25, 2013

©The Author 2013. Published by Oxford University Press. All rights reserved.
For Permissions, please e-mail: journals.permissions@oup.com.

Detection of Mesothelioma Remains a Conundrum

Robert Kratzke

Correspondence to: Robert Kratzke, MD, University of Minnesota Medical School, 420 Delaware St SE, Minneapolis, MN 55455 (e-mail: kratz003@umn.edu).

Malignant pleural mesothelioma remains a deadly cancer. The frustrating aspect, as with every cancer of the thorax, is that we understand the population at risk, but collectively we have yet to identify adequate markers for the risk of development of the disease. Recently there have been exciting developments of circulating markers that may prove useful in monitoring exposed populations, but application has not yet universally occurred (1–3).

The article presented in this issue of the Journal by Pairon and colleagues (4) delivers a fascinating perspective to this conundrum. The standard belief has been that pleural plaques are not a signal of risk of mesothelioma. Pairon et al. present for the first time solid and useful evidence that the presence of pleural plaques can be an accurate indicator of mesothelioma risk. It needs to be understood that the plaques alone have not been proven to be premalignant lesions. For example, plaques are not shown to be positron emission tomography computed tomography avid and have largely not been useful as a good radiologic marker for a history of asbestos exposure. Pairon et al. (4) accessed a unique French population of the Asbestos Past Exposure survey (APEXS). Populations such as these have proven very useful for mesothelioma researchers and clinicians because it is well known at this time that asbestos exposure remains the major described risk factor for mesothelioma. As

noted by Pairon et al., the utility, as well as methods, of screening for the exposed population remains controversial. As noted above, it remains very frustrating to clinicians that, as with the population at risk for lung cancer after tobacco exposure, the group that will be at risk for development of mesothelioma is all too clear.

Without much doubt, the most uniformly characterized group of asbestos-exposed individuals are the workers and citizens that comprise the Wittenoom cohort that has been admirably characterized and monitored by Musk and colleagues from the University of Western Australia (5). Wittenoom was both a workplace and the adjacent city where asbestos was heavily mined and residentially used for 40 years. The mesothelioma rate among exposed workers in the asbestos mining operation appears to be 4.7% according to the data from the Perth research team. Although this rate continues to be unacceptably high for this fatal cancer, it is heartening to know that most of the highly exposed population of asbestos workers and local occupants did not develop mesothelioma. Indeed, Musk and colleagues continue to identify and explore preventive strategies for populations at risk for fatal asbestos-related disease, and we may all hope that patients with plaques will be included in future studies (6).

Going forward, if clinicians identify patients at risk for mesothelioma, either through circulating markers or by the identification