Published in final edited form as:

Curr Opin Chem Biol. 2010 October 1; 14(5): 623–633. doi:10.1016/j.cbpa.2010.08.015.

Nanoplasmonics-enabled On-Demand and Systematic Intracellular Gene Regulation for Quantitative Biology and Therapeutics

Somin Eunice Lee and Luke P. Lee*

Department of Bioengineering, University of California-Berkeley, UCSF/UCB Joint Graduate Group in Bioengineering, Berkeley Sensor & Actuator Center

Abstract

This review focuses on the recent developments in nanoplasmonic technologies for on-demand and systematic intracellular gene regulation. Types of nanoplasmonic carriers and DNA/RNA cargo are described. Strategies to liberate cargo from their carriers using light are reviewed. Nanoplasmonic technologies enable on-demand and systematic silencing of endogenous intracellular genes. In addition to inhibitory effects, exogenous foreign genes are also introduced and expressed on-demand using nanoplasmonic technologies. The magnitude and timing of genetic activities can therefore be systematically controlled on-demand. Equipped with new nanoplasmonic technologies to directly probe the intracellular space, quantitative approaches should capture many dynamic activities within living systems that were otherwise previously impossible to control using conventional methods.

Introduction

A single living cell is an amazingly dynamic system that is capable of constantly sensing and responding to its perpetually changing environment. As such, it is an integrated system consisting of extracellular input signals streaming in from the local external environment, interlinked signaling cascades of internal connections, and gene transcriptional networks that respond by producing the appropriate proteins to give rise to cell function (Fig. 1). Quantitatively understanding the inner workings of these systems is the quest set forth by quantitative cell biology and systems biology. By systematically perturbing with extracellular stimuli, specific internal connections can be mapped and the resulting cell function can be observed in response to the single environmental change. Spatial, temporal, and systematic variation of extracellular stimuli has been shown to be absolutely critical for developmental processes [1], growth [2], differentiation [3], apoptosis, and stem cell fate decisions [4].

In addition to extracellular control of environmental stimuli, intracellular control of the actual internal connections themselves can provide unparalleled insight into the inner workings of these systems. When a cell is treated as input-output “black box,” ideally, the

© 2010 Elsevier Ltd. All rights reserved.

*To whom correspondence should be addressed. Prof. Luke P. Lee, Department of Bioengineering, University of California-Berkeley, 408C Stanley Hall, Berkeley, CA 94720-1762, (510) 642-5855, lplee@berkeley.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

output response is directly correlated to the input signal. However, signal distortions - time delays, noise, and signal magnitude reductions - can confound this input-output relationship. Time delays, due to inherent time-dependent interactions, transcriptional rates, and translational rates, are associated with each interconnected step inside the “black box.” Noise, from stochastic fluctuations in proteins, is introduced at each stage inside the “black box.” Signal magnitude reductions, due to degradation, dilution and diffusion effects, are also inevitable. These signal distortions can have significant implications on the output response. Consider a simple transcriptional network, where gene X regulates gene Y, and gene Y in turn regulates the output. In the presence of its input signal, X becomes active. Threshold effects are known to govern many gene regulatory processes [5]. Thus, when the concentration of X is greater than the threshold concentration required for activating Y, then Y is produced. Subsequently, when the concentration of Y is greater than the threshold concentration required for activating the output, then the output is produced. Now, suppose that the input signal is externally introduced (Fig. 1a). As this extracellular input signal traverses through each step of the extracellular-to-intracellular cascade, signal distortions are introduced. If the signal is significantly confounded by noise or if the signal magnitude is significantly reduced such that the activity of Y no longer satisfies the threshold condition (Fig. 1b), the output is not produced (Fig. 1c). This result can give the misleading impression that no relationship exists between the input signal and Y, when in fact they are related. Internal connections can be incorrectly mapped. Suppose now that Y is directly controlled at the intracellular level (Fig. 1d). Steps in the extracellular-to-intracellular cascade are essentially bypassed, thereby minimizing signal distortions. As a result, the output is reliably produced (Fig. 1e). Internal connections can therefore be correctly mapped and modeled.

By delving directly into the intracellular “black box,” the magnitude and timing of intracellular processes can be precisely controlled. Interfering oligonucleotides, such as antisense DNA, single-stranded RNA, short hairpin RNA (shRNA), and small interfering RNA (siRNA), enable direct, sequence-specific control of intracellular genes, but alone, lack the temporal control necessary for precise manipulation. Recent advancements in chemical biology, nanotechnology, and plasmonics now enable new light-sensitive tools of sub-nanometer and nanometer size scales to directly interface with intracellular processes. For example, nanoplasmonic technologies can be used as carriers of oligonucleotide cargo (Fig. 1d). Initially, oligonucleotide functionality is inactivated. Using light illumination as a remote trigger to release free oligonucleotides and “activate” their functionality, endogenous intracellular genes can be silenced on-demand. In addition to the inhibitory effects of interfering oligonucleotides, exogenous foreign genes can be also introduced and expressed on-demand. In this way, the magnitude and timing of genetic activities can be systematically varied on-demand. Signal distortions can be minimized since light-sensitive tools are activated from within the intracellular space, and therefore, steps in the extracellular-to-intracellular cascade are bypassed at the time of activation. Equipped with new tools to directly probe the intracellular space, quantitative and systematic approaches should capture many dynamic activities within the living cell that were otherwise previously impossible to detect using conventional methods.

Caging Technologies

Caging is also an effective means to temporarily inactivate oligonucleotide functionality by incorporating photo-labile protective groups, otherwise known as caging groups, into the bases or the phosphate backbone [6]. Ultra-violet (UV) irradiation removes the caging groups and restores oligonucleotide functionality. UV-activated gene silencing using caged antisense DNA has been demonstrated in mouse NIH 3T3 fibroblast cells [7]. UV-activated gene silencing using caged siRNA has also been demonstrated in human HeLa cervical

carcinoma cells [8]. While caging enables excellent spatiotemporal control of intracellular genes, the particular use of UV wavelengths is of some concern since intracellular proteins and nucleic acids are widely known to absorb, crosslink, and mutagenize in the presence of UV irradiation. Therefore, activation using less biologically harmful wavelengths of light is highly desirable.

NIR Nanoplasmonic Technologies

Gold nanoplasmonic particles (GNPs), in the near-infrared (NIR) spectral region, are attractive candidates for intracellular control. The NIR wavelength regime is well suited for biological and biomedical applications since tissues and cells are essentially transparent between 700–1300 nm [9]. Due to their strong and sharp resonance peak in their optical properties, GNPs efficiently convert light energy into surface-localized heat, otherwise known as photothermal conversion [10–12], when the incident light is matched to their plasmon resonance wavelength. In the presence of this incident light, the conduction band electrons of the GNPs collectively oscillate in phase on resonance and subsequently make collisions with the metal lattice, thereby dissipating heat [12]. Heat transfer from the surface of GNPs to the surrounding cellular environment is highly localized, decaying exponentially within a few nanometers [10,13,14] and therefore is thought to have minimal adverse effects on cells. Among the GNPs, gold nanorods [13,15,16] and gold nanoshells [17,18], are of widespread interest and pervasive use due to their large absorption cross-section, facile tunability of their plasmon resonance wavelength based on geometry, and large-scale synthesis with uniform distribution (Fig. 2).

Because of their large surface-to-volume ratio, GNPs are ideal carriers of cargo, such as interfering oligonucleotides. While attached to their carriers, cargo is rendered inactive due to steric hindrances between the tightly-packed cargo. For 150 nm diameter gold nanoshells, for example, the surface coverage of dsDNA cargo was determined to be 6400 dsDNA molecules/nanoshell [17]. In the presence of light that is matched to their plasmon resonance wavelength, GNPs photothermally release their cargo to freely interact with the local environment. Several strategies, employing different carrier and cargo types, have been demonstrated (Table 1). For example, it has been shown that short single-stranded DNA, otherwise known as antisense DNA, can be hybridized to a thiolated complementary sense strand, bound to a gold nanorod's surface through the gold-thiol covalent bond, and photothermally dehybridized using continuous-wave incident light that is matched to the plasmon resonance wavelength of the gold nanorods [13] (Fig. 3a). Antisense DNA has also been photothermally dehybridized from gold nanoshells [17] (Fig. 3b) and gold nanoprisms [19] (Fig. 3c) using continuous-wave illumination. This strategy of photothermal dehybridization using continuous-wave illumination offers several notable advantages. Firstly, no chemical modifications are made to the antisense DNA strand itself since a thiolated complementary strand is used to directly conjugate to the GNP's surface. Because chemical modifications can interfere with nucleic acid functionality and gene silencing efficacy, unmodified antisense DNA is highly desirable. Secondly, gold-thiol covalent bonds are stable after illumination, such that the GNP's surface remains covered with the thiolated complementary sense strands. With respect to cytotoxicity, this surface coating of complementary strands after illumination is critical. While the gold core is widely accepted as being biocompatible, bare GNPs have been shown to interact with proteins and induce mis-folding at physiological conditions [20]. Maintaining surface coverage with complementary strands after illumination also prevents reattachment of antisense DNA strands back onto the GNP carrier since rehybridization events are thermodynamically unfavorable due to steric hindrances and electrostatic repulsive forces at the GNP's surface [21]. Finally, the structural integrity of GNPs is uncompromised after illumination. Maintaining structure after illumination is crucial for *in vivo* applications, where the size,

geometry, coating material, and core material of nanoparticles are precisely designed and carefully characterized for proper biodistribution [22]. Maintaining structure after illumination also allows unique nano-scale optical properties to be retained, thereby enabling the same incident light wavelength to be used. Repetitive or finely-graded release of cargo is conceivable for future applications requiring precise temporal patterns of cargo release. For example, the delivery of a drug at its most effective concentration profile is an on-going challenge. When a drug is introduced, its concentration steadily decreases, producing a large concentration swing between the intervals at which the drug is administered. Photothermal cargo release from GNPs can be used to temporally control drug release for delivering programmable drug concentration profiles that are tailored to specific patients and applications. In addition to temporal control, spatial control of photothermal cargo release from GNPs can also be utilized for localized gene therapy at the tumor location. Genetic abnormalities at the tumor location can be corrected without adversely affecting neighboring, functional organs.

Alternatively, thiol-modified antisense DNA can be covalently bound to gold nanorods directly. It has been demonstrated that pulsed incident light can photothermally melt gold nanorods into spheres, thereby destabilizing the gold-thiol covalent bond and releasing thiol-modified antisense DNA [23] (Fig. 4a). Photothermal melting has also been effective at releasing thiol-modified linearized plasmid DNA from gold nanorods [24] (Fig. 4b) and thiol-modified siRNA from gold hollow nanoshells [25,26] (Fig. 4d). Additionally, photothermal melting has been shown to destabilize electrostatically-attached circular plasmid DNA from gold nanorods [27,28] (Fig. 4c). This strategy of photothermal melting using pulsed illumination presents several unique functionalities. Firstly, photothermal melting ensures complete release of all cargo. Secondly, the resulting shape change from rods to spheres enables a distinct and defined shift in the plasmon resonance wavelength. A second light source can be easily employed to re-match to the plasmon resonance wavelength of these shape-transformed GNPs. These functionalities may be useful for future studies that involve multifunctional release and detection schemes. Conceivably, cargo can be completely release from gold nanorods using pulsed illumination, and after a shape transformation, binding activity can then be detected using the now bare gold spherical surface. In the case of gold nanoshells, their silica cores deform after pulsed illumination due to high temperature heating (lattice temperatures reaching $\sim 1064^{\circ}\text{C}$, the melting point of bulk gold) [29]. As a result, their plasmon resonance wavelength can also shift after illumination, but in a less distinct and controlled manner compared with photothermally melted gold nanorods. Re-matching the incident light wavelength can therefore be considerably more difficult if the plasmon resonance wavelength cannot be easily controlled after shape transformation.

Nanoplasmonics-enabled Intracellular Control

GNPs enable “nanoplasmonic control” of genetic activities with sequence-specificity and spatiotemporal resolution. GNPs carrying genetic cargo are internalized in living cells by endocytosis. While attached to their carriers, the cargo is temporarily “inactive”. The cargo is also protected from degradation by nucleases due to steric hindrances between the tightly-packed cargo at the GNPs’ surface [30]. Illumination matched to the plasmon resonance wavelength of GNPs is used to “activate” cargo by photothermally disrupting encapsulating endosomes [13,31–34] and photothermally releasing free cargo into the cytosol. In this way, exogenous foreign genes can be introduced and expressed on-demand. Endogenous intracellular genes can also be silenced on-demand.

Nanoplasmonic induction of exogenous EGFP expression has been demonstrated in human HeLa cervical carcinoma cells [24]. This mechanism is illustrated in Figure 5a. Thiol-

modified, linearized EGFP-N1 plasmid DNA is covalently bound to gold nanorods and internalized in HeLa cells. Pulsed illumination is used to photothermally melt gold nanorods into spheres, thereby destabilizing the thiol-gold bond and releasing the plasmid into the cytosol. The plasmid transports to the nucleus. Exogenous mRNA is then transcribed from the plasmid, exported out the nucleus, and translated into the corresponding EGFP protein. The nanoplasmonic gene induction of EGFP expression is shown in the fluorescent image in Figure 5b [24]. Thus, expression of exogenous foreign genes can be induced on demand using nanoplasmonic control.

In addition to inducing exogenous expression, endogenous intracellular genes can also be silenced on demand. Nanoplasmonic silencing of endogenous ERBB2 expression using antisense DNA has been demonstrated in human BT474 breast carcinoma cells [13]. As illustrated in Figure 6a, the duplex of thiol-modified sense and unmodified antisense DNA is covalently bound to gold nanorods and internalized in BT474 cells. In the presence of continuous-wave illumination, antisense DNA strands dehybridize and release into the cytosol, while the thiol-modified sense strands remain attached to the gold nanorods. The unbound antisense DNA then binds to a portion of the mature mRNA corresponding to ERBB2. Once the mRNA/antisense DNA heteroduplex is formed, it is recognized and degraded by RNase H enzymes in the cytosol, thereby silencing ERBB2. As seen in Figure 6b, nanoplasmonic silencing of ERBB2 expression is qualitatively demonstrated using immunofluorescence imaging and quantitatively shown using flow cytometry [13]. To ensure light illumination itself and photothermally generated heat had no adverse effects on cells, viability studies were conducted using a fluorescence assay [13]. Since RNase H, is ubiquitously present in both the cytosol and the nucleus [35,36], an alternative model [35] suggests that unbound antisense DNA transports to the nucleus. In the nucleus, the antisense DNA binds to the premature mRNA. Once this heteroduplex is formed, it is recognized and degraded by RNase H enzymes in the nucleus.

Intracellular genes can also be silenced on-demand using siRNA. siRNA also signals the degradation of specifically targeted mRNA; however, degradation is through a different mechanism compared to antisense DNA. Nanoplasmonic silencing of EGFP expression using siRNA has been shown in mouse endothelial C166 cells stably expressing EGFP [25]. In Figure 7a, thiol-modified siRNA targeting EGFP is covalently bound to gold hollow nanoshells and internalized in endothelial cells stably expressing EGFP. Pulsed illumination is used to photothermally melt gold hollow nanoshells, thereby destabilizing the thiol-gold bond and releasing siRNA into the cytosol. The unbound siRNA sequentially triggers cytosolic RNA-inducing silencing complex (RISC) to unwind the duplex, binds to complementary mature mRNA, and silences EGFP expression. Nanoplasmonic silencing of EGFP in cells stably expressing EGFP is shown in the fluorescent images in Figure 7b [25]. Nanoplasmonic silencing of endogenous genes using siRNA has also been demonstrated *in vivo* [26]. Thiol-modified siRNA targeting NF- κ B p65 are covalently bound to hollow gold nanoshells, internalized in Hela cells, and transplanted in mice as xenografts. Pulsed illumination is used to photothermally melt gold hollow nanoshells, thereby destabilizing the thiol-gold bond and releasing the siRNA. Nanoplasmonic silencing of NF- κ B p65 is shown in the fluorescent images in Figure 7c [26]. Since key components of RISC have been discovered in both the nucleus and cytosol [37], another model [38] suggests that unbound siRNA transports to the nucleus. In the nucleus, siRNA triggers RISC to unwind the duplex, binds to premature mRNA, and silences gene expression. Other models [38] also propose that RISC actively transports siRNA from the cytosol to the nucleus for degradation.

Future Directions & Conclusions

In order to create widespread acceptance of nanoplasmonic technologies by the biology community, optimal plasmonic properties, colloidal stability, and internalization efficiencies of nanoplasmonic technologies must be demonstrated under physiological conditions. Multilayering cargo-carriers with cationic lipids have been shown improve to colloidal stability and internalization under physiological conditions [25]. Thus, in addition to the importance of cargo conjugation to carriers, additional protective surface coating materials also deserve critical attention [39,40]. It must also be demonstrated to the biology community that nanoplasmonic technologies are indeed biocompatible and nontoxic. Toxicology, biodistribution, and environmental studies on the long term effects of nanoplasmonic materials are necessary to promote these technologies and move the fields of nanoplasmonics-enabled quantitative biology and on-demand gene therapeutics forward.

Finally, both extracellular and intracellular controls are necessary to systematically perturb living systems. This integration of extracellular and intracellular controls can provide unprecedented insight into the dynamic nature of living systems for gene regulation and cell reprogramming. For example, microfluidic technologies can provide precise spatial and temporal control of the local external environment of a cell while nanoplasmonic technologies systematically perturb the intracellular space. Dynamic activities can be controlled and biological questions can be answered that were otherwise previously impossible using conventional methods.

Acknowledgments

The authors acknowledge the Siebel Foundation for graduate support S.E.Lee (Siebel Scholarship, Class of 2010). The authors acknowledge the National Institutes of Health (NIH) Nanomedicine Development Center for the Optical Control of Biological Function (PN2 EY018241) for financial support. The authors thank Prof. Han Lim and Prof. Robert Fischer for insightful discussion on systems biology.

References

1. Ismagilov RF, Maharbiz MM. Can We Build Synthetic, Multicellular Systems by Controlling Developmental Signaling in Space and Time? *Current Opinion in Chemical Biology*. 2007; 11:604–611. [PubMed: 17967432]
2. Hung PJ, Lee PJ, Sabounchi P, Lin R, Lee LP. Continuous Perfusion Microfluidic Cell Culture Array for High Throughput Cell Based Assays. *Biotechnology and Bioengineering*. 2005; 89:1–8. [PubMed: 15580587]
3. Steedman MR, Tao SL, Klassen H, Desai TA. Enhanced differentiation of retinal progenitor cells using microfabricated topographical cues. *Biomedical Microdevices*. 2010
4. LaBarge MA, Nelson CM, Villadsen R, Fridriksdottir A, Ruth JR, Stampfer MR, Peterson OW, Bissell MJ. Human mammary progenitor cell fate decisions are products of interactions with combinatorial microenvironments. *Integrative Biology*. 2009; 1:70–79. [PubMed: 20023793]
5. Little JW. Threshold Effects in Gene Regulation: When Some Is Not Enough. *Proceedings of the National Academy of Sciences*. 2005; 102:5310–5311.
6. Deiters A. Light-activation as a Method of Regulating and Studying Gene Expression. *Current Opinion in Chemical Biology*. 2009; 13:678–686. [PubMed: 19857985] * A review of the most recent advancements in caging technologies for use in cellular systems. Photo-labile groups, otherwise known as caging groups, are initially installed to render biological molecules. UV irradiation is then used to remove the caging and reactivate biological activity. Caging of small molecules, oligonucleotides, and proteins are discussed.
7. Young DD, Lusic H, Lively MO, Yoder JA, Deiters A. Gene Silencing in Mammalian Cells with Light-Activated Antisense Agents. *ChemBioChem*. 2008; 9:2937–2940. [PubMed: 19021142]
8. Mikat V, Heckel A. Light-dependent RNA interference with nucleobase-caged siRNAs. *RNA*. 2007; 13:2341–2347. [PubMed: 17951332]

9. Svoboda K. Biological Applications of Optical Forces. *Annu. Rev. Biophys. Biomol. Struct.* 1994; 23:247–285. [PubMed: 7919782]
10. Cortie M, Xu X, Chowdhury H, Zareie H, Smith G. Plasmonic Heating of Gold Nanoparticles and Its Exploitation. *Proc. SPIE.* 2005; 5649:565–573.
11. Khlebtsov B, Zharov V, Melnikov A, Tuchin V, Khlebtsov N. Optical Amplification of Photothermal Therapy with Gold Nanoparticles and Nanoclusters. *Nanotechnology.* 2006; 17:5167–5179.
12. Link S, El-Sayed M. Shape and Size Dependence of Radiative, Non-Radiative and Photothermal Properties of Gold Nanocrystals. *Int. Reviews in Physical Chemistry.* 2000; 19:409–453.
13. Lee SE, Liu GL, Kim F, Lee LP. Remote Optical Switch for Localized and Selective Control of Gene Interference. *Nanoletters.* 2009; 9:562–570. ** This paper is a demonstration of nanoplasmonic gene silencing using antisense DNA in living cells. Duplex of thiol-modified sense and unmodified antisense DNA is covalently bound to gold nanorods and internalized in human BT474 breast carcinoma cells. Photothermal dehybridization is used to liberate antisense DNA from gold nanorods. Nanoplasmonic gene silencing of ERBB2 expression using antisense DNA is qualitatively demonstrated using immunofluorescence imaging and quantitatively shown using flow cytometry.
14. Skirtach AG, Dejognat C, Braun D, Susha AS, Rogach AL, Parak WJ, Mohwald H, Sukhorukov GB. The Role of Metal Nanoparticles in Remote Release of Encapsulated Materials. *Nanoletters.* 2005; 5:1371–1377.
15. Nikoobakht B, El-Sayed MA. Preparation and Growth Mechanism of Gold Nanorods (NRs) Using Seed-Mediated Growth Method. *Chem. Mater.* 2003; 15:1957–1962.
16. Gou L, Murphy CJ. Fine-Tuning the Shape of Gold Nanorods. *Chem. Mater.* 2005; 17:3668–3672.
17. Barhoumi A, Huschka R, Bardhana R, Knight MW, Halas NJ. Light-Induced Release of DNA from Plasmon-Resonant Nanoparticles: Towards Light-Controlled Gene Therapy. *Chemical Physics Letters.* 2009; 482:171–179. ** This work is excellent motivation for emerging nanoplasmonic antisense gene therapies. Duplex of thiol-modified sense and unmodified antisense DNA is covalently bound to gold nanoshells. The authors demonstrate photothermal dehybridization of antisense DNA from gold nanoshells. The authors demonstrate that antisense DNA dehybridization is an irreversible process, a critical initial step for effective nanoplasmonic gene therapies.
18. Loo C, Lin A, Hirsch L, Lee M, Barton J, Halas N, West J, Drezek R. Nanoshell-enabled Photonic-based Imaging and Therapy of Cancer. *Technology in Cancer Research & Treatment.* 2004; 3:33–40. [PubMed: 14750891]
19. Jones MR, Millstone JE, Giljohann DA, Seferos DS, Young KL, Mirkin CA. Plasmonically Controlled Nucleic Acid Dehybridization with Gold Nanoprisms. *ChemPhysChem.* 2009; 10:1461–1465. [PubMed: 19431161] * In this work, duplexes of thiol-modified DNA oligonucleotides and unmodified complementary DNA oligonucleotides are covalently bound to gold nanoprisms. Photothermal dehybridization is used to release unmodified complementary DNA oligonucleotides from gold nanoprisms. The authors demonstrate that the Au-thiol bond between the thiol-modified DNA oligonucleotides and the gold nanoprisms is indeed stable after light illumination.
20. Zhang D, Neumann O, Wang H, Yuwono VM, Barhoumi A, Perham M, Hartgerink JD, Wittung-Stafshede P, Halas NJ. Gold Nanoparticles Can Induce the Formation of Protein-based Aggregates at Physiological pH. *Nanoletters.* 2009; 9:666–671. * This work demonstrates the significance of surface coatings on nanomaterials. The authors show that bare gold nanoparticles can interact with proteins and induce mis-folding at physiological conditions. Additionally, the authors show that bare gold nanoparticles can induce protein-nanoparticle aggregation at physiological conditions and consequently lose their unique nano-scale optical properties.
21. Demers LM, Mirkin CA, Mucic RC, Reynolds RA, Letsinger RL, Elghanian R, Viswanadham G. A Fluorescence-Based Method for Determining the Surface Coverage and Hybridization Efficiency of Thiol-Capped Oligonucleotides Bound to Gold Thin Films and Nanoparticles. *Anal. Chem.* 2000; 72:5535–5541. [PubMed: 11101228]
22. Brayner R. The Toxicology Impact of Nanoparticles. *Nanotoday.* 2008; 3:48–55.

23. Wijaya A, Schaffer SB, Pallares IG, Hamad-Schifferli K. Selective Release of Multiple DNA Oligonucleotides from Gold Nanorods. *ACS Nano*. 2009; 3:80–86. [PubMed: 19206252] ** This work demonstrates the release of multiple DNA species from gold nanorods of different aspect ratios. Thiol-modified single-stranded DNA is covalently bound to gold nanorods. Photothermal melting of gold nanorods is used to destabilize the gold-thiol covalent bond and release thiol-modified single-stranded DNA. The authors show selective photothermal melting of gold nanorods based on aspect ratio.
24. Chen C, Lin Y, Wang C, Tzeng H, Wu C, Chen Y, Chen C, Chen L, Wu Y. DNA-Gold Nanorod Conjugates for Remote Control of Localized Gene Expression by near Infrared Irradiation. *Journal of the American Chemical Society*. 2006; 128:3709–3715.
25. Braun GB, Pallaoro A, Wu G, Missirlis D, Zasadzinski JA, Tirrell M, Reich NO. Laser-Activated Gene Silencing via Gold Nanoshell-siRNA Conjugates. *ACS Nano*. 2009; 3:2007–2015. [PubMed: 19527019] ** This paper is a demonstration of nanoplasmonic gene silencing using siRNA in living cells. Thiol-modified siRNA recognizing EGFP is covalently bound to gold hollow nanoshells and internalized in mouse endothelial C166 cells stably expressing EGFP. Photothermal melting is used to liberate siRNA from gold hollow nanoshells. Nanoplasmonic gene silencing of EGFP expression is demonstrated using fluorescence microscopy.
26. Lu W, Zhang G, Zhang R, Flores LG, Huang Q, Gelovani JG, Li C. Tumor Site-specific Silencing of NF- κ B p65 by Targeted Hollow Gold Nanosphere-mediated Photothermal Transfection. *Cancer Res*. 2010; 70:3177–3188. [PubMed: 20388791] ** This paper is a demonstration of nanoplasmonic gene silencing using siRNA *in vivo*. Thiol-modified siRNA targeting NF- κ B p65 are covalently bound to hollow gold nanoshells, internalized in HeLa cells, and transplanted in mice as xenografts. Pulsed illumination is used to photothermally melt gold hollow nanoshells, thereby destabilizing the thiol-gold bond and releasing the siRNA. Nanoplasmonic gene silencing of NF- κ B p65 expression is demonstrated using fluorescence microscopy.
27. Horiguchi Y, Niidome T, Yamada S, Nakashima N, Niidome Y. Expression of Plasmid DNA Released from DNA Conjugates of Gold Nanorods. *Chemistry Letters*. 2007; 36:952–953.
28. Takahashi H, Niidome Y, Yamada S. Controlled Release of Plasmid DNA from Gold Nanorods Induced by Pulsed Near-Infrared Light. *Chem. Commun*. 2005:2247–2249.
29. Aguirre CM, Moran CE, Young JF, Halas NJ. Laser-Induced Reshaping of Metallo-dielectric Nanoshells under Femtosecond and 20 Nanosecond Plasmon Resonant Illumination. *J. Phys. Chem. B*. 2004; 108:7040–7045.
30. Rosi NL, Giljohann DA, Thaxton CS, Lytton-Jean AKR, Han MS, Mirkin CA. Oligonucleotide-Modified Gold Nanoparticles for Intracellular Gene Regulation. *Science*. 2006; 312:1027–1030. [PubMed: 16709779]
31. Troutman TS, Leung SJ, Romanowski M. Light-induced Content Release from Plasmon-resonant Liposomes. *Advanced Materials*. 2009; 21:2334–2338.
32. Wu G, Mikhailovsky A, Khant HA, Fu C, Chiu W, Zasadzinski JA. Remotely Triggered Liposome Release by Near-Infrared Light Absorption via Hollow Gold Nanoshells. *Journal of the American Chemical Society*. 2008; 130:8175–8177. [PubMed: 18543914]
33. Skirtach AG, Javier AM, Kreft O, Kohler K, Alberola AP, Mohwald H, Parak WJ, Sukhorukov GB. Laser-Induced Release of Encapsulated Materials inside Living Cells. *Angew. Chem. Int. Ed*. 2006; 45:4612–4617.
34. Febvay S, Marini DM, Belcher AM, Clapham DE. Targeted Cytosolic Delivery of Cell-Impermeable Compounds by Nanoparticle-Mediated, Light-Triggered Endosome Disruption. *Nano Letters*. 2010; 10:2211–2219. [PubMed: 20446663]
35. Dean NM, Bennett CF. Antisense Oligonucleotide-based Therapeutics for Cancer. *Nature Oncogene*. 2003; 22:9087–9096.
36. Marcusson EG, Bhat B, Manoharan M, Bennett CF, Dean NM. Phosphorothioate Oligodeoxyribonucleotides Dissociate from Cationic Lipids Before Entering the Nucleus. *Nucleic Acids Research*. 1998; 26:2016–2023. [PubMed: 9518498]
37. Robb GB, Brown KM, Khurana J, Rana TM. Specific and Potent RNAi in the Nucleus of Human Cells. *Nature Structural and Molecular Biology*. 2005; 12:133–137.

38. Berezhna SY, Supekova L, Supek F, Schultz PG, Deniz AA. siRNA in Human Cells Selectively Localizes to Target RNA sites. *Proceedings of the National Academy of Sciences*. 2006; 103:7682–7687.
39. Eghtedari M, Liopo AV, Copland JA, Oraevsky AA, Motamedi M. Engineering of Hetero-Functional Gold Nanorods for the in vivo Molecular Targeting of Breast Cancer Cells. *Nanoletters*. 2009; 9:287–291.
40. Lee SE, Sasaki DY, Perroud TD, Yoo D, Patel KD, Lee LP. Biologically Functional Cationic Phospholipid-Gold Nanoplasmonic Carriers of RNA. *Journal of the American Chemical Society*. 2009; 131:14066–14074. [PubMed: 19746908]

Fig. 1. Concept of nanoplasmonics-enabled on-demand and systematic intracellular gene regulation for quantitative biology

(a) Extracellular Control. Extracellular input signals, such as chemical stimuli or nutrients, are externally introduced. As this extracellular input signal traverses through each step of the extracellular-to-intracellular cascade, signal distortions are introduced. **(b)** The activity of Y does not satisfy the threshold condition due to signal distortions. **(c)** The output is not produced. This result gives the misleading impression that no relationship exists between the input signal and Y, when in fact they are related. Internal connections are incorrectly mapped. **(d) Intracellular Control.** Nanoplasmonic technologies are used as carriers of oligonucleotide cargo. Using light illumination as a remote trigger to release free oligonucleotides and “activate” their functionality, Y is directly controlled at the intracellular level. Signal distortions are minimized since steps in the extracellular-to-intracellular cascade are essentially bypassed. As a result, the output is reliably produced and internal connections are correctly mapped.

*Nanoplasmonic technology for intracellular control:***a** Gold nanorod carriers **b** Gold nanoshell carriers **Fig. 2. Nanoplasmonic technologies for intracellular control**

(a) Scanning electron microscopy image showing large-scale synthesis with uniform distribution, visual image showing facile tunability of the plasmon resonance wavelength, and UV-VIS absorbance spectra showing large absorption cross-sections of gold nanorod carriers. In the UV-VIS absorbance spectra, the plasmon resonance wavelengths correspond to aspect ratios (AR): (i) 2.3 AR, (ii) 3.5 AR, (iii) 5.0 AR [13] (b) Scanning electron microscopy image showing large-scale synthesis with uniform distribution, visual image showing facile tunability of the plasmon resonance wavelength, and UV-VIS absorbance spectra showing large absorption cross-sections of gold nanoshell carriers. In the UV-VIS absorbance spectra, the plasmon resonance wavelengths correspond to nanoshell thicknesses (60 nm core diameter): (iv) 20 nm, (v) 10 nm, (vi) 5 nm [17, 18]

Photothermal dehybridization:

Fig. 3. Photothermal dehybridization using continuous-wave illumination

(a) Scanning electron microscopy image of a gold nanorod carrier [13]. The duplex of thiol-modified sense and antisense DNA is covalently bound to the gold nanorod carrier. In the presence of continuous-wave illumination, the antisense DNA strands dehybridize while the thiol-modified sense strands remain attached to the carrier. **(b)** Scanning electron microscopy image of a gold nanoshell carrier [17]. The duplex of thiol-modified sense and antisense DNA is covalently bound to the gold nanoshell carrier. In the presence of continuous-wave illumination, the antisense DNA strands dehybridize while the thiol-modified sense strands remain attached to the carrier. **(c)** Scanning electron microscopy image of a gold nanoprism carrier [19]. The duplex of thiol-modified sense and antisense DNA is covalently bound to the gold nanoprism carrier. In the presence of continuous-wave illumination, the antisense DNA strands dehybridize while the thiol-modified sense strands remain attached to the carrier.

Photothermal melting: **Fig. 4. Photothermal melting using pulsed illumination**

(a) Scanning electron microscopy image of gold nanorod carriers [23]. Thiol-modified antisense DNA is covalently bound to the gold nanorod carrier. Pulsed illumination is used to photothermally melt gold nanorods into spheres, thereby destabilizing the thiol-gold bond and releasing the antisense DNA. (b) Scanning electron microscopy image of gold nanorod carriers [24]. Thiol-modified linearized plasmid DNA is covalently bound to the gold nanorod carrier. Pulsed illumination is used to photothermally melt gold nanorods into spheres, thereby destabilizing the thiol-gold bond and releasing the plasmid. (c) Scanning electron microscopy image of gold nanorod carriers [27, 28]. Circular plasmid DNA is

electrostatically attached to the gold nanorod carrier. Pulsed illumination is used to photothermally melt gold nanorods into spheres, thereby destabilizing the electrostatic interaction and releasing the plasmid. **(d)** Scanning electron microscopy image of gold hollow nanoshell carriers [25]. Thiol-modified siRNA is covalently bound to the gold hollow nanoshell carrier. Pulsed illumination is used to photothermally melt gold hollow nanoshells, thereby destabilizing the thiol-gold bond and releasing the siRNA.

a

Nanoplasmonic induction of exogenous foreign genes:

b

Fig. 5. Nanoplasmonic induction of exogenous foreign genes

(a) Concept of nanoplasmonic induction of exogenous EGFP in HeLa cells. In step 1, thiol-modified, linearized EGFP-N1 plasmid DNA is covalently bound to gold nanorods and internalized in HeLa cells. In step 2, pulsed illumination is used to photothermally melt gold nanorods into spheres, thereby destabilizing the thiol-gold bond and releasing the plasmid into the cytosol. In step 3, the plasmid transports to the nucleus and foreign mRNA is then transcribed from the plasmid. In step 4, the foreign mRNA is exported out of the nucleus and translated into the corresponding amino acids. In step 5, this primary sequence of amino

acids is then folded into the final EGFP protein structure. **(b)** Brightfield and fluorescent image after nanoplasmonic induction of EGFP in HeLa cells [24].

a

Nanoplasmonic silencing of endogenous genes using antisense DNA:

b

Fig. 6. Nanoplasmonic silencing of endogenous genes using antisense DNA

(a) Concept of nanoplasmonic silencing of endogenous ERBB2 in BT474 cells using antisense DNA. In step 1, the duplex of thiol-modified sense and unmodified antisense DNA is covalently bound to gold nanorods (GNPs) and internalized in BT474 cells. In step 2, in the presence of continuous-wave illumination, antisense DNA strands dehybridize and release into the cytosol, while the thiol-modified sense strands remain attached to the gold nanorods. In step 3, the unbound antisense DNA then binds to a portion of the mature mRNA corresponding to ERBB2. In step 4, once the mRNA/antisense DNA heteroduplex is formed, it is recognized by RNase H enzymes. In step 5 RNase H enzymes degrade the

heteroduplex, thereby silencing ERBB2. **(b)** Flow cytometric data and immunofluorescent images showing nanoplasmonic silencing of ERBB2 using activated GNPs in BT474 cells. Flow cytometric data and immunofluorescent images of the control showing no silencing using unactivated GNPs [13].

a

Nanoplasmonic silencing of endogenous genes using siRNA:

b

EGFP

c

NF- κ B p65

Fig. 7. Nanoplasmonic silencing of endogenous genes using siRNA

(a) Concept of nanoplasmonic silencing of EGFP using siRNA in C166 cells stably expressing EGFP. In step 1, thiol-modified siRNA targeting EGFP is covalently bound to gold hollow nanoshells (GNPs) and internalized in endothelial cells stably expressing EGFP. In step 2, pulsed illumination is used to photothermally melt gold hollow nanoshells, thereby destabilizing the thiol-gold bond and releasing siRNA into the cytosol. In step 3, the unbound siRNA sequentially triggers cytosolic RISC to unwind the duplex. In step 4, cytosolic RISC binds single-stranded RNA to its complementary mRNA. In step 5, mRNA is degraded and EGFP expression is silenced. (b) Fluorescent image showing

nanoplasmonic silencing of EGFP using activated GNPs in C166 cells stably expressing EGFP. Fluorescent image of the control showing no silencing using unactivated GNPs [25].
(c) Fluorescent image showing nanoplasmonic silencing of NF- κ B p65 using siRNA in HeLa xenografts in mice [26].

Table 1

Carrier and cargo strategies for nanoplasmonic technologies.

Genetic cargo	Target	Cargo length	Cell type	Carrier type	Carrier size	Carrier plasmon resonance	Surface conjugation	NIR light wavelength	Light type	Mechanism	Ref.
dsDNA	ERBB2	15 bp	Human BT474	Gold nanorod	$l = 50$ nm $d = 15$ nm	780 nm	Thiol-gold bond	785 nm	CW	Antisense DNA dehybridization; Gene silencing of ERBB2 in BT474 cells	13
dsDNA	unspecified	30–70 bp	n/a	Gold nanoshell	$d = 120$ nm	800 nm	Thiol-gold bond	800 nm	CW	Antisense DNA dehybridization	17
dsDNA	unspecified	15 bp	n/a	Gold nanoprism	$l = 120$ nm $l = 7.5$ nm	1200 nm	Thiol-gold bond	1064 nm	CW	Antisense DNA dehybridization	19
ssDNA	unspecified	40 bases	n/a	Gold nanorod	$l = 44$ nm $d = 11$ nm	800 nm	Thiol-gold bond	800 nm	Pulse	ssDNA release	23
ssDNA	unspecified	40 bases	n/a	Gold nanorod	$l = 89$ nm $d = 17$ nm	1100 nm	Thiol-gold bond	1100 nm	Pulse	ssDNA release	23
Linearized plasmid DNA	EGFP-N1	4700 bp	Human HeLa	Gold nanorod	$l = 42$ nm $d = 11$ nm	782 nm	Thiol-gold bond	800 nm	Pulse	Linearized plasmid release; Induction of EGFP-N1 expression in HeLa cells	24
Circular plasmid DNA	EGFP-C1	7000 bp	n/a	Gold nanorod	$l = 65$ nm $d = 11$ nm	900 nm	Electrostatic interaction	1064 nm	Pulse	Circular plasmid release	28
Circular plasmid DNA	Luciferase	unspecified	n/a	Gold nanorod	$l = 65$ nm $d = 11$ nm	900 nm	Electrostatic interaction	1064 nm	Pulse	Circular plasmid release; <i>in vitro</i> luciferase expression assay	27
siRNA	EGFP-N1	25 bp	Mouse C166-EGFP	Gold hollow nanoshell	$d = 40$ nm	800 nm	Thiol-gold bond	800 nm	Pulse	siRNA release; Gene silencing of EGFP-N1 in C166-EGFP cells	25
siRNA	NF- κ B p65	21 bp	Human HeLa	Gold hollow nanoshell	$D = 40$ nm	800 nm	Thiol-gold bond	800 nm	Pulse	siRNA release; Gene silencing of NF- κ B p65 in HeLa	26

Genetic cargo	Target	Cargo length	Cell type	Carrier type	Carrier size	Carrier plasmon resonance	Surface conjugation	NIR light wavelength	Light type	Mechanism	Ref.
xenografts of mice											

Abbreviations: continuous-wave (CW), length (l), diameter (d), thickness (t), base pairs (bp)