Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading doi:10.1016/S2221-1691(11)60044-0 © 2011 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Effect of insulin on *Blattella germanica* Linnacus

Abolghasemi E, Moosa Kazem SH*, Abolhasani M, Davoudi M

Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences

ARTICLE INFO

Article history:

Received 19 February 2011

Received in revised form 25 March 2011

Accepted 10 April 2011

Available online 30 April 2011

Keywords:

Insulin

Rearing

Blattella germanica

Insectary

Sensitivity

ABSTRACT

Objective: To determine the sensitivity of *Blattella germanica* L (*B. germanica* L) to different doses of insulin. **Methods:** *B. germanica* were reared in laboratory conditions at $(25 \pm 2)^\circ\text{C}$ and $(50 \pm 5)\%$ relative humidity (RH), and exposure period of 12:12 L/D. Different concentrations, viz. 5, 10, 15, 20 and 25 μ of insulin N, R, (N+R) were prepared and injected to 10 treated cockroaches with another 10 cockroaches which were injected with normal saline as control group. **Results:** Insulin N with a dose of 20 μ caused more than 70% mortality of *B. germanica* in this study. There was a significant difference between 20 μ of insulin N with other doses of 5, 10, 15 and 25 μ , and its comparison with other forms of medication also showed obvious difference ($P < 0.05$). **Conclusions:** It can be concluded that effective drug doses of insulin which can be used as poison bait or gel against German cockroaches could be utilized in the control of *B. germanica* in the future field studies.

1. Introduction

German cockroaches *Blattella germanica* (*B. germanica*) is one of the most potential mechanical vectors and responsible for transmission of many protozoan, bacteria, viruses and intestinal parasites to humans[1].

German cockroaches can also cause irritation and allergies in human hosts[2–6]. Based on mentioned above, control of cockroaches is very important[7,8]. German cockroaches are becoming resistant to pesticides through special mechanisms and survive successfully due to quick adaptation to environment these years[9–16]. The recent research shows that environmental contamination which is regarded as a significant contamination does not affect the cockroaches with pathogenic bacterial[16]. Society urgently needs to find new materials used as a control tool against this pest which is pesticide-resistant arising from high and irregular use of pesticides in the world. Insulin drugs are used to the treatment of diabetic patients. This type of diabetes occurs when the pancreas can not produce enough insulin to help the body target cells. Paul Langerhans, a medical student in 1869 during the study in pancreatic

tissue specialized cells under a microscope found that these cells were mediated by insulin secretion, so pancreatic tissue was called by the name of the first scholar researcher. Minkowski and Joseph Van Mering removed a dog pancreas during the surgery in Germany in 1889. The observation of flies gathering in dog urine showed the presence of sugar in dog urine. The relation between pancreatic function and diabetic diseases was found more specific in 1901. Steiner and Federwisk *et al* cited secret material responsible for regulation of blood glucose (insulin) from the pancreas gland in 1920[17,18].

Currently, common forms of insulin are R (Human regular insulin), NPH (Isophane insulin suspension), N (Insulin isophane, novolin)[18].

So far, several methods have been exploited to the cockroaches control and some control tools such as poisoned bait gel can be seen as an effective tool[19,20].

Poison bait which can be treated in the appropriate places is very effective due to its adsorbent and scentless property. Other pharmaceutical compounds such as tetratoxin found in sex glands, liver, intestines and skin of puffer fish were used as poison bait to the control of cockroaches[23]. It should be mentioned that its combination effect on sodium channels and nerve membranes has been demonstrated previously[21–24]. This combination wipes off neural activity in the species of cockroaches, especially *B. germanica* L and it seems that the killing effects with type 1 pyrethroids,

*Corresponding author: Dr Seyed Hassan Moosa-Kazemi, PhD, Assistant Professor, Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences.

E-mail: moosakazemi@sina.tums.ac.ir

Foundation Project: Supported by Tehran University of Medical Sciences (No: 9855–27–

carbamates, DDT, have less similar characteristics as compared with combination with type 2 pyrethroids, organophosphate, nicotine and muscarine[25,26].

No studies of the effect of insulin on *B. germanica* L have been investigated previously. Therefore, we conduct the study to determine the susceptibility of *B. germanica* L to different doses of insulin.

2. Materials and methods

2.1. Rearing

B. germanica L were reared in the special glass containers containing bread, sugar, water in the laboratory conditions (25 ± 2) °C and (50 ± 5)% relative humidity and exposure period of 12:12 L/D[27]. Then the *B. germanica* strains were kept without any contact with insecticide in Insectary of School of Public Health Tehran University of Medical Sciences and this susceptible strains were used in various studies.

2.2. Test

Effect of insulin was determined by injection of insulin N, R, (N+R) at a dose of 5, 10, 15, 20 and 25 μ into insect coelom. Initially different concentrations of insulin were prepared and injected into 10 divided cockroaches with another 10 as the control which was injected with normal saline. The results were analysed with SPSS software. The standard contact topical application test was used[28]. Different doses of 5, 10, 15, 20 and 25 μ of drugs were determined by the device driver. Then, these cockroaches were anesthetized with CO₂ gas and different doses of insulin, N, R, (N+R), were

injected. The control group was given doses of 5, 10, 15, 20 and 25 μ of 0.9% normal saline.

2.3. Statistical analysis

SPSS ver. 11.5, stata 80, and one way ANOVA test were used to compare the mortality of *B. germanica* L between two groups. A *P*-value less than 0.05 was considered statistically significant.

3. Results

The results of effectiveness of insulin by different doses was shown in Table 1. Significant difference was found between the mortality of German cockroaches exposed to insulin N with a dose of 20 μ in comparison with the control group (*P*<0.05) (Figure 1).

Insulin N with a dose of 20 μ was more effective than other doses. Also this dose was more effective than the same doses of other drugs separately (Figure 1).

Figure 1. The mortality of German cockroaches (*B. germanica* L) exposed to insulin N, R, and N+R with a dose of 20 μ in comparison with the control group.

Table 1

The mortality rate of German cockroaches (*B. germanica* L) exposed to different dosage of insulin (*n*=10).

		Mortality rate (%)	Dead	Alive
N	5 μ	40	4	6
	10 μ	30	3	7
	15 μ	40	4	6
	20 μ	70	7	3
	25 μ	20	2	8
R	5 μ	0	0	10
	10 μ	0	0	10
	15 μ	0	0	10
	20 μ	20	2	8
	25 μ	30	3	7
R+N	5 μ	30	3	7
	10 μ	40	4	6
	15 μ	0	0	10
	20 μ	30	3	7
	25 μ	0	0	10

4. Discussion

This is the first study in relation to effects of insulin on German cockroaches while previous studies stated some

effects of insecticides, chemical pesticides and other drugs on cockroaches[28–31]. Effect of insulin-R reported 30 minutes post injection and the duration was 6–8 hours. The maximum effect reaches 6–8 hours post treatment. Onset of insulin N reported 2 hours post injection with the duration of 18–26 hours. In this report the maximum effect calculated was 4–12 hours post treatment by insulin N[32–41]. It is evident in our research that the common signs of post treatment were the reduction of irritability, decreased activity, less frequencies of drinking water, reduced weakness and eventually death.

Our studies showed that the most mortality rate occurred when *B. germanica* L were exposed to insulin N with a dose of 20 μ . The effectiveness of insulin on *B. germanica* L has not been evaluated yet. Nasirian *et al* reported that imidacloprid gel bait was more effective than the gel of Fipronil[26]. This difference may be due to size, metabolism and physiology of cockroaches.

Many studies reported the resistance of cockroaches to different insecticides, particularly organophosphate and pyrethroides, so the failure of control programs occurred subsequently[28,42–50].

In conclusion, the use of drug instead of common pesticides possesses the advantages of reduced contact or oral toxicity to people, especially to children and the elderly and also less environmental pollution. Our study indicates that insulin N with a dose of 20 μ which can be used as poison bait or gel against German cockroaches was deadly effective for preventing the occurrence of *B. germanica* L resistance. Results of our study show that injection of insulin N at a dose of 20 μ has greater impact than other modes with the common characteristics of decreased stimulation, activity, mobility, desire to drink water, weakness, hunger, and eventually death. The effective doses of insulin which can be used as poison bait or gel against German cockroaches are recommended in the future studies.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors are grateful to Ms Faranak Firoozfar, Fatemeh Erfani, Ms Fatemeh Rafee, Mr Majed Esazadehs, Mr Azim Paksa, MSPH students in School of Public Health, Tehran University of Medical Sciences. This study was financially supported by grant of Tehran University of Medical Sciences.

References

- [1] Kinfu A, Erko B. Cockroaches as carriers of human intestinal parasites in two localities in Ethiopia. *Trans R Soc Trop Med Hyg* 2008; **102**(11): 1143–1147.
- [2] Slater JE, James R, Pongracic JA, Liu AH, Sarpong S, Sampson HA, et al. Biological potency of German cockroach allergen extracts determined in an inner city population. *Clin Exp Allergy* 2007; **37**(7): 1033–1039.
- [3] Tahernejad K. *Sensitivity level of Blattella germanica study collected from several hospitals in Tehran to different insecticides for receive of academic thesis masters degree in the field of medical entomology*. Tehran: Tarbiat Modarres University; 1995, p. 1373–1378.
- [4] Douroudgar A, Asadi M. Sensitivity *Blattella germanica* of Kashan hospitals to insecticides. *J Feiz* 2001; **17**: 70–76.
- [5] Motavaliaghghi F, Sharif M. *Medical importance and methods of insects control in urban areas (Pederos-Blattella)*. Sari: Assistant University of Medical Sciences and Health Services; 2001, p. 51–54.
- [6] Motavaliaghghi IF, Gholami SH, Sedaghat M. Determine the level of sensitivity Sari hospital *Blattella germanica* to insecticides in 1997. *J Mazan Uni Med Sci* 1998; **16**: 42–46.
- [7] Wang C, Bennett GW. Cost and effectiveness of community-wide integrated pest management for German cockroach, cockroach allergen, and insecticide use reduction in low-income housing. *J Econ Entomol* 2009; **102**(4): 1614–1623.
- [8] Nalyanya G, Gore JC, Linker HM, Schal C. German cockroach allergen levels in North Carolina schools: comparison of integrated pest management and conventional cockroach control. *J Med Entomol* 2009; **46**(3): 420–427.
- [9] Lee CY. *Perspective in urban insect pest management in Malaysia*. Penang: Vector Control Research Unit, Universiti of Sains; 2007, p. 104.
- [10] Gore JC, Schal C. Gene expression and tissue distribution of the major human allergen Bla g 1 in the German cockroach, *Blattella germanica* L. (Dictyoptera: Blattellidae). *J Med Entomol* 2004; **41**: 953–960.
- [11] Ladonni H. Susceptibility of *Blattella germanica* to different insecticides in different hospitals in Tehran-Iran. *J Entomol Soc Iran* 1993; **12**: 23–28.
- [12] Ladonni H. Susceptibility of different field strains of *Blattella germanica* to four pyrethroids (Orthoptera: Blattellidae). *Iran J Public Health* 1997; **26**: 35–40.
- [13] Ladonni H, Sadegheyani S. Permethrin toxicity and synergistic effect of piperonil butoxide in the first nymphal stage of *Blattella germanica* (Dictyoptera: Blattellidae). *Iran J Public Health* 1998; **27**: 44–50.
- [14] Nasirian H, Ladonni H, Shayeghi M, Ahmadi MS. Iranian non-responding contact method German cockroach permethrin resistance strains resulting from field pressure pyrethroid spraying. *Pak J Biol Sci* 2009; **12**(8): 643–647.
- [15] Shahi M, Hanafi-Bojd AA, Vatandoost H. Evaluation of five local formulated insecticides against German cockroach (*Blattella germanica* L.) in Southern Iran. *Iran J Arthropod Borne Dis* 2008; **2**(1): 21–27.
- [16] Fu X, Ye L, Ge F. Habitat influences on diversity of bacteria found on German cockroach in Beijing. *J Environ Sci (China)* 2009; **21**(2): 249–254.

- [17] Steiner DF, Oyer PE. The biosynthesis of insulin and a probable precursor of insulin by a human islet cell adenoma. *Proc Natl Acad Sci USA* 1967; **57**(2): 473–480.
- [18] Federwisch M, Dieken ML, De Meyts PE. *Insulin and related proteins—structure to function and pharmacology*. Dordrecht: Kluwer Academic Publishers; 2002, p. 9.
- [19] Wang C, Bennett GW. Efficacy of noviflumuron gel bait for control of the German cockroach, *Blattella germanica* (Dictyoptera: Blattellidae)—laboratory studies. *Pest Manag Sci* 2006; **62**: 434–439.
- [20] Nasirian H. Duration of fipronil and imidacloprid gel baits toxicity against *Blattella germanica* strains of Iran. *Iran J Arthropod Borne Dis* 2007; **1**(2): 40–47.
- [21] Nakamura M, Yasumoto T. Tetrodotoxin derivatives in puffer fish. *Toxicon* 1985; **23**: 271–276.
- [22] Robert H, Coby S. Effects of dietary protein levels on reproduction and food consumption in the German cockroach (Dictyoptera: Blattellidae). *Ann Entomol Soc Am* 1988; **81**(6): 969–976.
- [23] Hamilton R, Coby S. Effects of dextrin and cellulose on feeding and reproduction in female brown-banded cockroach *supella longipalpa*. *Physiol Entomol* 2008b; **18**(1): 57–64.
- [24] Suiter DR, Koehler PG, Patterson RS. Dietary effects of allopurinol and sulfapyrazone on development, survival, and reproduction of German cockroaches (Dictyoptera: Blattellidae). *J Econ Entomol* 1992; **85**(1): 117–122.
- [25] Enayati AA, Motevalli HF. Biochemistry of pyrethroid resistance in German cockroach (Dictyoptera, Blattellidae) from hospitals of Sari. *Iran Biomed J* 2007; **11**(4): 251–258.
- [26] Nasirian H, Ladonni H, Shayeghi M, Vatandoost H, Yaghoobi-Ershadi MR, Rassi Y, et al. Comparison of permethrin and fipronil toxicity against German cockroach (Dictyoptera: Blattellidae) strains. *Iran J Public Health* 2006; **35**: 63–67.
- [27] Derek WG. The action of tetrodotoxin on the cockroach, *Periplaneta americana*: a toxicological and neurophysiological study. *Physiol Entomol* 1987; **3**: 37–42.
- [28] Bracke JW, Cruden DL, Markovetz AJ. Effect of metronidazole on the intestinal microflora of the American cockroach, *Periplaneta americana* L. *Antimicrob Agents Chemother* 1978; **13**(1): 115–120.
- [29] Cochran DG. Mortality and reproductive effects of avermectin B1 fed to German cockroaches. *Entomol Exp Appl* 1985; **37**: 83–88.
- [30] MedicineNet. Health and medical information produced by doctors. [Online] Available from: www.medicinenet.com
- [31] Limoe M, Enayati AA, Ladonni H, Vatandoost H, Baseri H, Oshaghi MA. Various mechanisms responsible for permethrin metabolic resistance in seven field-collected strains of the German cockroach, *Blattella germanica* (L.) (Dictyoptera: Blattellidae) from Iran. *Pestic Biochem Physiol* 2007; **87**: 138–146.
- [32] Cochran DG. Misuse of the tarsal-contact method for detecting insecticide resistance in the German cockroach (Dictyoptera: Blattellidae). *J Econ Entomol* 1997; **90**: 1441–1444.
- [33] Wilma DSC, Kavya N, Kulkarni S. Evaluation of insulin sensitivity status in polycystic ovarian syndrome. *Asian Pac J Trop Dis* 2011; **1**(1): 67–70.
- [34] Firoozfar F, Moosa-Kazemi H, Baniardalani M, Abolhassani M, Khoobdel M, Rafinejd J. Mass rearing of *Lucilia sericata* Meigen (Diptera, Calliphoridae). *Asian Pac J Trop Biomed* 2011; **1**(1): 54–56.
- [35] Radji M, Fauziah S, Aribinuko N. Antibiotic sensitivity pattern of bacterial pathogens in the intensive care unit of Fatmawati Hospital, Indonesia. *Asian Pac J Trop Biomed* 2011; **1**(1): 39–42.
- [36] Muhamad O, Chacharoenkul W, Rungsihirunrat K, Ruengweerayut R, Na-Bangchang K. Assessment of in vitro sensitivity of *Plasmodium vivax* fresh isolates. *Asian Pac J Trop Biomed* 2011; **1**(1): 49–53.
- [37] Erukainure OL, Ajiboye JA, Adejobi RO, Okafor OY, Adenekan SO. Protective effect of pineapple (*Ananas cosmosus*) peel extract on alcohol-induced oxidative stress in brain tissues of male albino rats. *Asian Pac J Trop Dis* 2011; **1**(1): 5–9.
- [38] Thaweboon B, Thaweboon S. Effect of *Phyllanthus emblica* Linn. on candida adhesion to oral epithelium and denture acrylic. *Asian Pac J Trop Med* 2011; **4**(1): 41–45.
- [39] Lu G, Hu XC, Huang C, Lu YJ, Wu LX, Li LH, et al. Effect of artemether, hemin and Fe³⁺ on recombinant lactate dehydrogenase from *Schistosoma japonicum*. *Asian Pac J Trop Med* 2010; **3**(12): 930–933.
- [40] Chakraborty S, Singha S, Chandra G. Mosquito larvicidal effect of orthophosphoric acid and lactic acid individually or their combined form on *Aedes aegypti*. *Asian Pac J Trop Med* 2010; **3**(12): 954–956.
- [41] Damabi NM, Moazedi AA, Seyyednejad SM. The role of α - and β -adrenergic receptors in the spasmolytic effects on rat ileum of *Petroselinum crispum* Latifolium (parsley). *Asian Pac J Trop Med* 2010; **3**(11): 866–870.
- [42] Ladonni H. Evaluation of three methods for detecting permethrin resistance in adult and nymphal *Blattella germanica* (Dictyoptera: Blattellidae). *J Econ Entomol* 2001; **94**: 694–697.
- [43] Cochran DG. Monitoring for insecticide resistance in field-collected strains of the German cockroach (Dictyoptera: Blattellidae). *J Econ Entomol* 1989; **82**: 336–341.
- [44] Phillips Ak, Appel AG, Sims SR. Topical toxicity of essential oils to the German cockroach (Dictyoptera: Blattellidae). *J Econ Entomol* 2010; **103**(2): 448–459.
- [45] Pai HH, Wu SC, Hsu EL. Insecticide resistance in German cockroaches (*Blattella germanica*) from hospitals and households in Taiwan. *Int J Environ Health Res* 2005; **15**(1): 33–40.
- [46] Kristensen M, Hansen KK, Jensen KMV. Cross-resistance between dieldrin and pronil in German cockroach (Dictyoptera: Blattellidae). *J Econ Entomol* 2005; **98**: 1305–1310.
- [47] Guo GZ, Geng YJ, Huang DN, Xue CF, Zhang RL. Level of CYP4G19 expression is associated with pyrethroid resistance in *Blattella germanica*. *J Parasitol Res* 2010; pii: 517534.
- [48] Chang KS, Jung JS, Park C, Lee HI, Lee WG, Lee DK, et al. Insecticide susceptibility and resistance of *Blattella germanica* (Blattaria: Blattellidae) in Seoul, Republic of Korea, 2007. *Entomol Res* 2009; **39**: 243–247.
- [49] Nasirian H. An overview of German cockroach, *Blattella germanica*, studies conducted in Iran. *Pak J Biol Sci* 2010; **13**(22): 1077–1084.
- [50] Chai RY, Lee CY. Insecticide resistance profiles and synergism in field populations of the German cockroach (Dictyoptera: Blattellidae) from Singapore. *J Econ Entomol* 2010; **103**(2): 460–471.