Published in final edited form as:

Metallomics. 2013 January ; 5(1): 29–42. doi:10.1039/c2mt20009k.

Calciomics: integrative studies of Ca^{2+} -binding proteins and their interactomes in biological systems

Yubin Zhou^{a,b}, Shenghui Xue^c, and Jenny J. Yang^b

^aCenter for Translational Cancer Research, Institute of Biosciences and Technology, Texas A&M University System Health Science Center, Houston, TX, 77030 USA

^bDepartment of Chemistry, Georgia State University, Atlanta, GA, 30303 USA

^cDepartment of Biology, Georgia State University, Atlanta, GA, 30303 USA

Abstract

Calcium ion (Ca^{2+}), the fifth most common chemical element in the earth's crust, represents the most abundant mineral in the human body. By binding to a myriad of proteins distributed in different cellular organelles, Ca^{2+} impacts nearly every aspect of cellular life. In prokaryotes, Ca^{2+} plays an important role in bacterial movement, chemotaxis, survival reactions and sporulation. In eukaryotes, Ca^{2+} has been chosen through evolution to function as a universal and versatile intracellular signal. Viruses, as obligate intracellular parasites, also develop smart strategies to manipulate the host Ca^{2+} signaling machinery to benefit their own life cycles. This review focuses on recent advances in applying both bioinformatic and experimental approaches to predict and validate Ca^{2+} -binding proteins and their interactomes in biological systems on genome-wide scale (termed "calciomics"). Calmodulin is used as an example of Ca^{2+} -binding protein (CaBP) to demonstrate the role of CaBPs on the regulation of biological functions. This review is anticipated to rekindle interest in investigating Ca^{2+} -binding proteins and Ca^{2+} -modulated functions at the systems level in the post-genomic era.

Keywords

Ca^{2+} ; interaction; signaling; bioinformatics; proteomics; EF-hand; prediction; binding; metalloproteome

1. Introduction

Ca^{2+} , the fifth most abundant element on earth, constitutes approximately 3.6% of the earth's crust. Ca^{2+} has a flexible coordination number of 4–8, an irregular coordination geometry and rapid binding kinetics, which maximizes its potential to interact with various ligands. Ca^{2+} tends to precipitate both inorganic and organic anions (at ~mM range) so that high Ca^{2+} is incompatible with life and it has to be extruded, sequestered or compartmentalized.¹ These unique physio-chemical properties of Ca^{2+} make it the only available candidate in the environment to perform both micro (as a messenger) and macro (e.g., biomineralization) functions in biological systems. During the evolution from prokaryotes to eukaryotes that began over 2 billion years ago, the $\text{Fe}^{2+}/\text{H}_2\text{S}$ buffering system in the sea was progressively exhausted, along with the increase of oxygen supply and prevalence of oxidative reactions.¹ Ca^{2+} played an important role in facilitating the adaptive changes of prokaryotes to an oxidative environment, and contributed to the evolution of

*Corresponding: jenny@gsu.edu.

more advanced organisms and land colonization.² For example, Ca^{2+} is closely involved in crucial evolutionary events such as the evolution of cytoskeleton, cell motility, the formation of skeletons and hard-shelled eggs (biomineralization), and the ability to supply milk.

By mass, Ca^{2+} represents the most abundant mineral (~1 kg) in the adult human body, with 99% in the form of calcium phosphates in bones. In mammals, specific Ca^{2+} homeostatic system is evolved to set both the extracellular and intracellular Ca^{2+} concentrations at proper levels, and produces a steep Ca^{2+} gradient (>10,000 times) across the plasma membrane.³ The external Ca^{2+} , sensed by the extracellular Ca^{2+} sensing receptor, is maintained at 1.1~1.4 mM through the coordinated actions of three major hormones (i.e., parathyroid hormone, calcitonin and 1,25-dihydroxyvitamin D_3).⁴ Compared to the extracellular Ca^{2+} concentration of about 10^{-3} M, the internal Ca^{2+} concentration is maintained within the narrow range of 10^{-7} to 10^{-8} M in nearly all living cells. In multi-cellular eukaryotes, Ca^{2+} levels in different cellular compartments or organelles are tightly controlled through the molecular choreography of a repertoire of Ca^{2+} signaling molecules, including Ca^{2+} channels, Ca^{2+} transporters, receptors, Ca^{2+} buffers, Ca^{2+} -responsive proteins (e.g., Ca^{2+} -sensitive enzymes and transcription factors) that are distributed in cytoplasm, endoplasmic/sarcoplasmic reticulum (ER/SR), Golgi complex, mitochondria, nucleus or the extracellular matrix (Fig. 1).⁵ Even in the most primitive prokaryotes, the free cytosolic Ca^{2+} is maintained at $\sim 10^{-7}$ M by using a variety of prokaryotic ion transporters (e.g., $\text{Ca}^{2+}/\text{H}^+$ antiporter, $\text{Ca}^{2+}/\text{Na}^+$ antiporter, and Ca^{2+} -ATPase) or channels.⁶ Although bacteria does not possess complex intracellular organelles and Ca^{2+} is not extensively used as an intracellular messenger, accumulating evidence indicates that Ca^{2+} plays an essential role in the regulation of bacterial movements, chemotaxis, survival reactions and sporulation.^{3b, 6c, d, 7}

In eukaryotic cells, Ca^{2+} functions as a universal and versatile signal by interacting with hundreds of proteins over a 10^6 -fold range of affinities (nM to mM) (Fig. 1, left panel). Inside cells, Ca^{2+} -binding proteins, such as calmodulin (CaM) or parvalbumin, have Ca^{2+} -binding affinities in the submicromolar range corresponding to the resting cytosolic Ca^{2+} concentrations at ~100 nM. Ca^{2+} concentrations in organelles such as the ER, Golgi, and endocytic vacuoles are in the hundreds of μM . By modulating the activities of these proteins, Ca^{2+} participates in various biological processes (e.g., hormonal secretion, muscle contraction, neurotransmission and memory formation) over a wide range of timescale, ranging from milliseconds up to days (Fig. 1, right panel). Overall, the Ca^{2+} signaling system exhibits great versatility in speed, amplitude and spatial-temporal patterns. In this review, we focus on how integrative studies can be carried out to probe the sophisticated Ca^{2+} signaling network in biological systems (termed calciomics). First, we describe bioinformatics approaches tailored for predicting Ca^{2+} -binding sites and Ca^{2+} -modulated activities from genomic and structural information. Second, we discuss how experimental approaches can be applied to probe Ca^{2+} -binding properties (stoichiometry, selectivity, affinity, cooperativity etc.). Finally, we present herein, recent progress in identifying the interactomes of CaBPs by using CaM-target interactions as an example.

2. Bioinformatics approaches to predict Ca^{2+} -binding proteins

The last decade we have witnessed an explosion of genomic information that far exceeds the capability of experimental characterization on each gene product. Nearly 30% of all proteins bind various metals. This has created unprecedented need for the Ca^{2+} signaling field to predict Ca^{2+} -binding sites in proteins based on the amino acids sequences translated from nucleotide sequences, or from low-resolution or modeled structures. The prediction of a protein's capability to bind Ca^{2+} may provide important insights into its biological function and guide experimental designs.⁸

2.1 Prediction from primary sequences

Ca²⁺-binding sites in proteins can be divided into two types: continuous and discontinuous ones. Continuous Ca²⁺-binding sites are formed by a stretch of continuous amino acids in a linear sequence with at least some of the binding ligands well conserved. Thus, a pattern (or motif signature) representing the conserved features can be generated, and used to determine whether a protein contains a given Ca²⁺-binding pattern with simple string matching algorithms. A major class of Ca²⁺-binding proteins (CaBPs) contains continuous Ca²⁺-binding sites with an EF-hand structure (Fig. 2).⁹ A typical canonical EF hand (Fig. 2, middle panel) contains a helix-loop-helix topology with the Ca²⁺-binding pocket adopting a pentagonal bipyramidal geometry. Another continuous Ca²⁺-binding motif is called “pseudo” EF-hands that are exclusively found in the N-termini of S100 and S100-like proteins (Fig. 2, lower panel).⁹ The major difference between these two types of EF-hands lies in the Ca²⁺-binding loop (Fig. 2): the 12-residue canonical EF-hand loop binds Ca²⁺ mainly via sidechain carboxylates or carbonyls (loop sequence positions 1, 3, 5, 12), whereas the 14-residue pseudo EF-hand loop chelates Ca²⁺ primarily via backbone carbonyls (positions 1, 4, 6, 9) and the sidechain carboxylates (position 14). The residue at position 9 in a canonical EF-hand, or position 11 in a pseudo EF-hand, coordinates the Ca²⁺ ion through a bridged water molecule. While the majority of the known EF-hand CaBPs contain paired EF-hand motifs, CaBP's with single EF hands have also been discovered in prokaryotes and viruses.¹⁰ In addition, “EF-hand-like motifs” (e.g., Excalibur as shown in Fig. 2, upper panel) have been found in a number of bacterial proteins.^{10a, 11} Although their coordination properties remain similar to the canonical 29-residue helix-loop-helix EF-hand motif, the EF-hand-like motifs differ from EF-hands in that they contain deviations in the secondary structure of the flanking sequences and/or variation in the length of the Ca²⁺ coordinating loop. A series of patterns reflecting these features have been generated with satisfactory predictive accuracy and sensitivity. We have used these patterns to detect putative EF-hand or EF-hand like Ca²⁺-binding motifs in both viral and prokaryotic genomes.¹⁰ A list of all the predicted CaBPs in these primitive genomes can be found in Supplementary Tables 1 and 2. Aside from our contribution, several other groups have developed webserver or libraries using their own patterns or published data to predict and catalog EF-hand CaBPs (see Table 1).

Given that the pattern search method simply looks for Ca²⁺-binding motif signatures in proteins without considering the similarity to known domains, it tends to generate false positives. Another convenient method to predict CaBPs is to search for conserved Ca²⁺-binding domains (CaBDs) by online resources such as the Pfam or InterPro databases.¹² The Pfam database has a comprehensive collection of profile hidden Markov models (HMM),¹³ which contain probability-based consensus information for protein domains. The profile HMMs can be visualized by HMM Logos¹⁴ (Fig. 2) and are compatible with HMMER,¹⁵ a free and effective software package used to assign sequence similarity to a query protein sequence. This prediction strategy is based on sequence homology alignment using a known Ca²⁺-binding protein (such as C2 domain) against a genome. It is extremely useful for identifying non-continuous CaBPs as it allows for the identification of any homologous proteins without the restriction of any gaps in sequences. An excellent introduction to using profile HMMs to predict metalloproteomes has been recently described in a tutorial review in *Metallomics*.¹⁶ As a common limitation of sequence-based predictive computational approaches, one has to be aware that both methods are unlikely to detect potential CaBPs that do not contain either a known Ca²⁺-binding pattern or a well-established CaBD.

All these prediction methods enabled us to make a phylogenetic distribution of typical Ca²⁺-binding motifs or domains found in nature (Table 2). Interestingly, EF-hands or EF-hand like motifs (particularly the Excalibur motif¹¹), are frequently present in prokaryotes in a single unit, which raises the hypothesis that the prokaryotic single EF-hand motif is likely

the precursor for the paired EF-hands found in eukaryotes.^{9–10} In contrast, the pseudo EF-hand is almost exclusively found in vertebrates. Thus, it is reasonable to postulate that pseudo EF-hands are phylogenetically younger and have a shorter history than canonical EF-hands. Likewise, calreticulin and calsequestrin, two signature CaBPs inside ER/SR, are only found in eukaryotes given that endo-membranous structures evolved later and that ER structure is absent in prokaryotes.

2.2 Prediction from 3D structures

Ca²⁺-binding sites do not always follow a set of linear amino acid sequences. Approximately 90% of known Ca²⁺-binding sites in Protein Data Bank (PDB) are non-linear, which means the Ca²⁺-coordinating ligands may come from different loops in a protein, or different subunits, or even several different proteins. Prediction of those Ca²⁺-binding sites requires three-dimensional structural information from either determined structures deposited in PDB or from homology modeling. Fortunately, key structural factors governing Ca²⁺-binding share some common features with defined ligand types, charge distribution, Ca²⁺-ligand distances, bond angles and geometric configurations.¹⁷ This enables the development of a number of computational methods to predict non-continuous Ca²⁺-binding sites based on 3D structures (summarized in Table 1). Those prediction algorithms employ structural parameters as distinguishing factors between Ca²⁺-binding and non-binding sites. Frequently, multiple algorithms are combined, typically using some types of scoring function with a grid system or other methods of spatial analysis (e.g., graph theory) to evaluate physicochemical features systematically and identify either binding ligands or binding sites based on comparative scoring. It is worthy of note that these functions are dependent on *a priori* statistical analyses to define or constrain function parameters based on structural, chemical or thermodynamic data. Selected examples are briefly discussed below.

An earlier example of using combinatory methods to predict Ca²⁺-binding sites is the Fold-X algorithm.¹⁸ This method combines geometric pattern matching and energy calculations based on Fold-X empirical force field, where Ca²⁺-binding sites are constrained to four to six ligand oxygen atoms.¹⁸ The Fold-X force field is used for energy calculation and for the optimization of the position of the predicted Ca²⁺. This algorithm can also coarsely predict binding affinity based on energy-optimized placement of Ca²⁺ to distinguish between low and high affinity sites, as well as high affinity Mg²⁺-binding sites. Although these results indicate higher prediction rates than previous efforts,¹⁹ this method was less successful at identifying either multiple binding sites or sites with lower coordination numbers (CN<4).

Another commonly used predicting strategy is based on the machine learning method (e.g., SVMProt,²⁰ SitePredict²¹ and FINDSITE-METAL²²). Machine learning is particularly effective at classifying metal-binding ligands and determining whether the target atom is close enough for chelating the ion. In this method, a window including neighboring residues slides through the sequence so that each successive residue is viewed as a target residue in turn, to identify binding residues. The training procedures include applying the same learning machine to a training dataset consisting of sequences with known metal-binding residues. Classification features include evolutionary information, location in the primary sequence, secondary structure, metal-binding propensity, and solvent accessibility. Additional information on machine learning methods for predicting CaBPs can be found through the webservers listed in Table 1.

Toward the goal of predicting Ca²⁺-binding sites with improved accuracy and sensitivity, we have developed several prediction algorithms during the recent five years. The first-generation program GG (Graph theory and Geometry) presented a new approach to prediction with the combination of graph theory and geometric parameters.²³ Based on the *a*

priori statistical parameters that govern Ca^{2+} -binding, the GG algorithm identifies oxygen clusters as the basis for predicting Ca^{2+} -binding sites (Fig. 4). In this method, a graph $G(V, E)$ was first constructed with oxygen atoms as its vertices and edges between two vertices if their spatial distance was no more than 6.0 Å. Oxygen clusters corresponding to cliques in $G(V, E)$ consisting of exactly four vertices were identified, and the Ca^{2+} center was determined at an equidistant center within each cluster if the distance ranged from 1.8 to 3.0 Å. This algorithm did not include a procedure to merge overlapping, predicted Ca^{2+} locations, resulting in multiple, adjacent predicted sites near the documented Ca^{2+} -binding site. Nonetheless, this algorithm produces rapid results, mainly due to the absence of a grid algorithm, and achieves approximately 90% site sensitivity and 80% site selectivity on testing datasets. To fully utilize the Ca^{2+} -binding geometric properties, we have also developed the next-generation MUG (Multiple Geometries) program, which is able to predict Ca^{2+} -binding sites with different coordination numbers in proteins at atomic resolution.²⁴ MUG requires that potential Ca^{2+} -binding sites consisting of at least four oxygen atoms although MUG further allows oxygen atoms from either water or cofactors in addition to amino acid residues. Spatial analysis is achieved based on graph theory modeling as well as imposition of a grid system. MUG also has the capability of predicting both low coordination sites and multiple Ca^{2+} -binding sites that share a common ligand. Very recently, we further developed the third-generation MUG^C algorithm²⁵ to predict Ca^{2+} -binding sites in both X-ray and NMR structures with flexibility in the binding region or low occupancy in a Ca^{2+} -binding site. Using second shell carbon atoms, and without explicit reference to side-chain oxygen ligand coordinates, MUG^C is able to achieve 94% sensitivity with 76% selectivity on a dataset of X-ray structures comprised of 43 Ca^{2+} -binding proteins. Additionally, prediction of Ca^{2+} -binding sites in NMR structures was obtained by MUG^C using a different set of parameters determined by analysis of both Ca^{2+} -constrained and unconstrained Ca^{2+} -loaded structures derived from NMR data. The geometric arrangement of the second-shell carbon cluster is sufficient for both accurate identification of Ca^{2+} -binding sites in NMR and X-ray structures, and for selective differentiation between Ca^{2+} and other relevant divalent cations.

While the majority of the computational algorithms are tailored for predicting CaBPs in their metal-bound (holo) forms, predicting Ca^{2+} -binding sites from apo-structures is equally important but much more challenging due to two facts: *first*, a large number of protein structures are determined in the absence of metal ions; *second*, the binding of Ca^{2+} may trigger conformational changes and cause the rearrangement of the binding pocket. Tremendous efforts have been directed to this end with steady progress. A FEATURE program, with machine learning that utilizes a Bayesian approach to classify potential Ca^{2+} -binding sites, has been recently reported.²⁶ Following the identification of distributions of predefined Ca^{2+} -binding characteristics, the query protein structure was embedded in a 3D-grid of 2 Å grid cubes, and a score was computed for each grid (e.g., query site) based on a scoring function that uses Bayes' Rule to combine prior known distributions and observed frequencies (e.g., derived from a training dataset and the query site) to indicate the likelihood of Ca^{2+} -binding in the queried site. Besides being applied to static X-ray crystal structures, FEATURE has recently been applied to multiple conformations of the protein parvalbumin β generated from molecular dynamics (MD) simulation.²⁷ For MD generated conformations of both holo and apo forms, FEATURE correctly identifies the Ca^{2+} -binding sites and non-sites with some interesting fluctuations in score. The combination of machine learning techniques such as FEATURE with MD simulations may provide new insights into Ca^{2+} -binding affinity, persistence and the stability of Ca^{2+} -binding sites. More recently, we have improved the MUG algorithm (MUG^{SR}) to accurately predict CaBPs with multiple-binding sites that undergo local conformational change or side chain rotations upon Ca^{2+} -binding.²⁸ Given an apo (unbound) protein structure, this new approach conducts a side chain rotation procedure according to a rotamer library to find binding sites with possible

side-chain movements (local conformational changes) caused by Ca^{2+} -binding. Given an apo protein structure, MUG^{SR} is able to capture Ca^{2+} -binding sites that may undergo local conformational changes, which is validated by comparison with the corresponding holo protein structure sharing more than 98% sequence similarity with the apo protein.

2.3 Toward the prediction of the CaBP-target interactomes

Upon the stimulation of signals or alteration in the membrane potential, the cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_c$) could be elevated by 100 folds from 10^{-8} M to 10^{-6} M.⁵ The elevation of $[\text{Ca}^{2+}]_c$ can be effectively sensed by Ca^{2+} sensors inside cells. Ca^{2+} signaling is a fast process and takes place in milliseconds. To convert this transient Ca^{2+} signals into more sustained physiological processes, CaBPs, such as calmodulin (CaM), undergo conformational changes and subsequently interact with a variety of protein kinases (e.g., CaM-dependent kinase I) and phosphatase (e.g., calcineurin), both of which are capable of covalently modifying downstream effectors by adding or removing a phosphate group.²⁹ The time scale of this process usually lasts for seconds or even minutes. In addition, to more efficiently modulate the signaling cascades, Ca^{2+} -CaM (sometimes apo-CaM) targets to enzymes or proteins (e.g., adenylate cyclases, phosphodiesterase, ion channels, and GPCRs) involved in the conversion of other second messengers, such as cAMP and cGMP. These enzymes or proteins themselves do not respond to changes in Ca^{2+} concentration. However, CaBPs, such as CaM, can sense the intracellular Ca^{2+} concentration changes. Binding or releasing of Ca^{2+} induces conformational changes of these intracellular “ Ca^{2+} sensors proteins” (e.g., CaM), which further alter the binding affinity of “ Ca^{2+} sensor proteins” to these enzyme and proteins. The reciprocal interactions between CaBP and signaling proteins will regulate the function of these proteins. Thus, CaBPs like CaM relay the Ca^{2+} signaling to a bewildering number of target enzymes or proteins to carry out corresponding cellular functions. More importantly, the Ca^{2+} signal itself can even be modulated by CaM through CaM's interaction with membrane receptors, ion channels and pumps. CaM resides in almost every subcellular compartment and targets a wide range of proteins (Fig. 5). The complex CaBP-target interactions lead to the divergent cellular responses to transient Ca^{2+} signals in different cell types. Thus, the prediction of CaBP-target interactomes would not only provide a testable hypothesis and functional information in Ca^{2+} -regulated biological processes, but would also lay the cornerstone for the ultimate computational modeling of the sophisticated Ca^{2+} signaling network.

The most successful example of predicting the CaBP-target interactome is the CaM target database.³⁰ Although low sequence identity has been observed among different sequences, CaM target peptides share some key features: the hydrophobic residues with bulky sidechains that anchor the target sequence to the hydrophobic cleft of CaM are arranged in particular spacing patterns with 2–6 positively charged residues interspersed in between. According to the distribution of anchoring residues in the continuous polypeptide sequence, the binding motifs can be divided into five classes: 1–10 class, 1–14 class, 1–16 class, IQ class and others.^{30–31} Among them, the first three classes have been found to interact with CaM almost exclusively in a Ca^{2+} -dependent way. Additional binding mode of 1–7 and 1–17 for CaM interaction of ryanodine receptor (RyR) and NMDA receptor were identified respectively and were shown to be also Ca^{2+} -dependent.³² The IQ class motifs, however, are activated or inhibited by CaM in a Ca^{2+} -independent fashion and CaM seems to be a constitutive part of its functional unit. Recent works by Shea and Chazin's groups have shown that the C-domain of apo-CaM interacts with several voltage dependent sodium channels via binding to IQ motifs.³³ In addition, there are several unclassified CaM binding motifs that interact with CaM when the four Ca^{2+} -binding sites are only half occupied. All of these common features enable one to gain more confidence when confronted by the following three questions: 1) are the proteins under investigation capable of interacting with

CaM? If so, 2) is it Ca^{2+} -dependent or independent? And 3) How the CaM-mediated function is affected by intracellular Ca^{2+} concentration change? An online server dedicated to predicting CaM binding sites has been generated by Ikura's group.³⁰ We have successful experience using this prediction algorithm to predict putative CaM target sequences in a viral protease domain,³⁴ the gap junction protein connexins³⁵ and the intracellular domain of the Ca^{2+} -sensing receptor,³⁴ and to further demonstrate the functional importance of such CaM-target interactions. CaM has been shown to use either N or C-domain to sense local or global Ca^{2+} concentration change to regulate a number of Ca^{2+} permeable channels.³⁶ If only using CaM variants with removed/decreased Ca^{2+} -binding affinities, it is difficult to unambiguously distinguish between the effects of Ca^{2+} directly on the channel and the indirect effect of Ca^{2+} on its target mediated by CaM. Hence, we engineered a series of mutant CaMs to individually tune the Ca^{2+} affinity of each of CaM's EF-hands by increasing the number of acidic residues in Ca^{2+} chelating positions.³⁷ Domain-specific Ca^{2+} affinities of each CaM variant were determined by equilibrium fluorescence titration. Mutations in sites I (T26>D) or II (N60>D) in CaM's N-terminal domain had little effect on CaM Ca^{2+} affinity and regulation of RyR1. However, the site III mutation (N97>D) increased the Ca^{2+} -binding affinity of CaM's C-terminal domain and caused CaM to inhibit RyR1 at a lower Ca^{2+} concentration than wild type CaM. Conversely, the site IV mutation (Q135>D) decreased the Ca^{2+} -binding affinity of CaM's C-terminal domain and caused CaM to inhibit RyR1 at higher Ca^{2+} concentrations. These results support the hypothesis that Ca^{2+} -binding to CaM's C-terminal acts as the switch converting CaM from a RyR1 activator to a channel inhibitor. These results further indicate that targeting CaM's Ca^{2+} affinity may be a valid strategy to tune the activation profile of CaM-regulated ion channels.³⁷

Overall, the rapid progress in bioinformatics has led to the constant improvement and expanding of prediction algorithms, databases and tools that are rapidly becoming indispensable to the metallomics field. The availability of genomic information (both 1D sequences and 3D structures), prediction algorithms and software for modeling structural and energetic characteristics has facilitated an important new phase in calciomic research, allowing for pre-experimental evaluation of hypotheses and the rational design of experimental plans (Fig. 3).

3. Experimental approaches to detect Ca^{2+} -binding proteins

To understand the molecular mechanism of Ca^{2+} modulated biological processes, we have to answer at least three important questions with experimental evidence: *i)* What is the entirety of Ca^{2+} -binding proteins? *ii)* What is the exact Ca^{2+} /protein stoichiometry? *iii)* What are the site-specific Ca^{2+} -binding affinities for Ca^{2+} -binding proteins, particularly those that utilize multiple coupled Ca^{2+} -binding sites to respond to sharp changes in cellular Ca^{2+} concentration? However, finding answers to these questions is often hampered by challenges listed below. *First*, while X-ray analysis has been the "gold standard" for revealing the metal-binding sites, Ca^{2+} has often been missing in the determined structure, particularly for those proteins with weak Ca^{2+} -binding affinities. This is possibly due to a rapid off rate owing to the low affinity of the Ca^{2+} -binding and the existence of multiple conformations that are in equilibrium with one another. *Second*, Ca^{2+} is spectroscopically silent, making it impossible to trace by fluorescence or other spectroscopic techniques that are amenable to study other metal ions. Fortunately, this could be partially overcome by using radioactive ^{45}Ca , [^{103}Ru azido ruthenium] (Section 3.1), or Ca^{2+} analogs that are spectroscopically visible (e.g., Eu^{3+} and Tb^{3+}) (Section 3.3). *Third*, most experimental methods such as dialysis coupled with mass spectrometry (Section 3.2) are only sensitive to the total Ca^{2+} content, instead of the protein-bound fraction of Ca^{2+} . *Fourth*, overcoming the persistent background contamination of Ca^{2+} during the preparation of Ca^{2+} -free sample for

proteins with strong Ca^{2+} -binding affinities is not a trivial task. *Last*, obtaining information for site-specific Ca^{2+} -binding is difficult due to complexities encountered in cooperative, multi-site systems, and the use of Ca^{2+} -binding energy to bring about conformational changes. Unfortunately, the use of peptide models has been limited by lack of well-defined conformation of peptides in solution. To circumvent these limitations, we have developed a grafting approach coupled with luminescence resonance energy transfer (LRET) to probe intrinsic metal binding properties of continuous Ca^{2+} -binding sites (Section 3.3).

3.1 Proteomic approaches based on gel electrophoresis

The first question regarding the entirety of CaBPs can be partially answered by taking proteomic approaches (two dimensional gel electrophoresis (2D-GE) coupled with mass spectrometry) and the additional use of radioactive metal ions, such as ^{45}Ca or ^{103}Ru thorium.

The ^{45}Ca overlay method, first described by Maruyama *et al.*,³⁸ has been widely used for detecting CaBPs from crude mixtures for almost three decades. Protein from various sources are separated by means of either one-(1D) or two-dimensional (2D) gel electrophoresis, transferred to a membrane (nitrocellulose or PVDF), and overlaid with radioactive ^{45}Ca . The “Stains-all” method described by Campbell *et al.* provides a simple and efficient way to probe CaBPs.³⁹ CaBPs, such as troponin C, S-100 and calsequestrin, resolved in SDS-PAGE gel can be stained as blue and purple color. The dye has a maximum absorbance at 615 nm, indicating that this dye is in the anionic sites when it associates with Ca^{2+} -binding proteins and then forms a J complex. Interestingly, “stains-all” also interacts with native proteins with a maximal absorbance around 600 nm. Further studies showed that the “stains-all” dye could cause changes in both the UV/vis absorbance and the circular dichroism signals when it interacts with Ca^{2+} -binding motifs.⁴⁰

The advent of powerful mass spectrometry techniques further extends the applicability of this methodology to efficiently identify unknown CaBPs. This method has been recently adapted to investigate the repertoire of CaBPs associated with the plasma membrane of *Arabidopsis thaliana*⁴¹ or human sperm.⁴² These techniques have also been used to characterize CaBPs such as p95, Calreticulin, synaptophysin and γ -Crystallin.⁴³ When coupled with equilibrium dialysis, the ^{45}Ca overlay method can be further used to determine the stoichiometry and binding affinity of a Ca^{2+} -binding protein.

The recent use of azido ruthenium (AzRu or PhotoCal) represents a new avenue of studying Ca^{2+} -dependent processes.^{42,44} AzRu was shown to specifically interact with Ca^{2+} -binding proteins and remain covalently bound after ultraviolet irradiation. By taking advantage of this unique feature, Israelson *et al.* generated an AzRu-cellulose column to purify CaBPs with affinity chromatography, and developed AzRu-based biosensor chips for identifying previously unknown CaBPs.⁴² Moreover, radioactive [^{103}Ru] AzRu can be used to detect CaBPs in biological samples after SDS-PAGE and autoradiography. It is anticipated that the wide use of this reagent will pave the way for generating a comprehensive Ca^{2+} -targeted protein database and detecting defective CaBPs implicated in various disorders and diseases associated with dysregulated Ca^{2+} signaling. Nonetheless, this method has two potential caveats: it has not been fully established that AzRu mimics the coordination chemistry of all known CaBPs and the metal selectivity (except for Mg^{2+}) has not been rigorously tested.

In summary, the gel electrophoresis and autoradiography-based method is well-known for its high sensitivity and accuracy. However, its disadvantages are also obvious: *First*, it requires the handling of radioactive products and involves relatively labor-intensive and time-consuming procedures. *Second*, the amount of proteins encoded by rare genes is far from sufficient to be detected by SDS-PAGE. *Third*, the demanding gel electrophoresis step

is likely to disrupt the native Ca^{2+} -binding sites in protein samples. *Last but critically*, CaBPs with weak affinities for Ca^{2+} may not be readily detected with this method.

3.2 Mass spectrometry approaches

The determination of Ca^{2+} /protein stoichiometry and metal selectivity in protein samples can be routinely performed using electrospray ionization mass spectrometry (ESI-MS) or inductively coupled plasma mass spectrometry (ICP-MS). ESI-MS is widely used to capture non-covalent interactions as seen in Ca^{2+} : protein complexes. Metal-bound biological samples can be mildly ionized and transferred to the gas-phase while maintaining their native states. This technique is very sensitive to CaBPs with a strong affinity to Ca^{2+} . The comparison of the mass/charge peaks corresponding to the same sample under native or denatured conditions gives directly the mass of bound metal. Nonetheless, ESI-MS has a poor tolerance toward nonvolatile salt buffers and solubilizing agents, which are often necessary to maintain the structural integrity and functionality of CaBPs. Furthermore, ESI-MS often cannot provide quantitative information on the amount of bound metals. Complementary to ESI-MS, ICP-MS can quantitatively detect the total metal contents in a biological sample with high sensitivity regardless of its molecular environment, and therefore, is often used in speciation analysis coupled with liquid chromatographic techniques.⁴⁵ A detailed description of the application of these mass spectrometry techniques in bioinorganic analytical chemistry can be found in a comprehensive review by Lobinski *et al.*^{45a}

3.3 A grafting approach to probe isolated continuous Ca^{2+} -binding sites

To answer the third question listed above, we have developed a grafting approach to dissect the key structural factors that control Ca^{2+} -binding affinity, conformational change and cooperativity.⁴⁶ In principle, the key determinants for Ca^{2+} affinity can be systematically introduced into a stable host protein frame and evaluated by eliminating or minimizing the contribution of conformational change. We have shown that the domain 1 of CD2 (CD2.D1) is an excellent scaffold protein.⁴⁷ It retains its native structure following insertion of the EF-hand motif with proper flanking linkers. This provides the foundation for measuring the intrinsic metal-binding affinity with minimized contribution of protein conformational change. More importantly, the aromatic residues (particularly Trp32) in CD2, which is juxta-positioned to the grafted Ca^{2+} -binding loop, enable us to obtain Tb^{3+} binding affinity with LRET using fluorescence spectroscopy. Tb^{3+} is commonly used to probe Ca^{2+} -binding because of their similar ionic radii (Ca^{2+} , 1.00 Å; Tb^{3+} , 0.92 Å) and metal coordination chemistry.⁴⁸ A Ca^{2+} competition assay can be further performed to obtain Ca^{2+} -binding affinity. This approach has been successfully applied to determine the site-specific metal-binding affinities of calmodulin, a ubiquitous Ca^{2+} -sensing protein with four EF-hands. Based on the energetics relationship, the cooperativity of coupled Ca^{2+} -binding sites can also be estimated after obtaining the intrinsic Ca^{2+} -binding affinity for each site.^{46a} Another advantage of this approach lies in the fact that it circumvents the expression and purification of the original proteins that harbor the predicted Ca^{2+} -binding sites. The same method has also been used to probe the site-specific metal-binding properties of putative EF-hand motifs in viral and bacterial proteins,^{10, 49} as well as predicted continuous Ca^{2+} -binding sites in the family C of G-protein coupled receptors (GPCRs)^{35a, 50} and the ER-resident stromal interaction molecule 1.⁵¹ By taking advantage of the sensitive LRET signal and the dropping cost of mini-gene synthesis, we foresee ample opportunities to further extend this approach to systematically validate putative linear Ca^{2+} -binding sites in a high-throughput manner with fluorescence microplate readers. For example, GST-tagged engineered proteins containing various putative Ca^{2+} -binding sites can be enriched from *E. coli* cell lysates with agarose beads conjugated with glutathione and subjected to LRET measurement in 96-well format.

3.4. Approaches to map the CaBP-target interactome

We use the most extensively studied CaM-target interactions as an example here. Several robust methods have been developed to screen CaM-binding proteins (CaMBPs) on the proteome-wide scale. A CaM binding overlay (CAMBOT) technique that requires labeling of CaM with ^{35}S and the use of SDS-PAGE was established by O'Day *et al.*⁵² In this technique, the bacteria-expressed ^{35}S -CaM was used as the probe to detect CaMBPs and compare the CaMBPs expression level under different conditions. Under Ca^{2+} -saturated and Ca^{2+} -free conditions, it is possible, in theory, to detect both Ca^{2+} -dependent and Ca^{2+} -independent CaMBPs. This method can be potentially applied to verify and profile the CaMBPs, to assess the effects of drugs on CaM-CaMBP networks, and to evaluate the effects of heavy metal ion in the environment on the CaM-CaMBP network.

An alternative way to screen CaMBP is explored by Shen *et al.*⁵³ This large-scale screening method utilizes the mRNA display technique. mRNA-displayed enriched proteome library can be selected by biotinylated CaM, which will be further captured by streptavidin-agarose beads. Another round of selection involving six steps (PCR, in vitro transcription, crosslinking, cell-free translation and fusion, reverse transcription and affinity chromatography) can be repeated to diminish unspecific interactions. Screening of CaMBPs in the human and *C. elegans* proteome with this method reveals a large body of previously undetected proteins or uncharacterized proteins, such as ribosomal proteins, proteasome 26S subunits, deubiquitinating enzymes, and leucine zipper proteins.

In another method, combining affinity chromatography and tandem mass spectrometry, Berggard *et al.* detected 140 putative CaM binding proteins in the mouse brain.⁵⁴ Recently, a protein microarray based method was used to map plant proteins that interact with Ca^{2+} -CaM or CaM-like (CaML) protein in *Arabidopsis thaliana*.⁵⁵ The protein array strategy overcomes at least two limitations of the above-mentioned methods. *First*, all the target proteins are presented in distinct locations and identification of secondary targets is minimized. *Second*, the protein array platform is compatible with transmembrane proteins, including GPCRs and ion channels which are often closely implicated in Ca^{2+} signaling. A total of 1,133 high-quality *Arabidopsis* proteins were screened for their capability to interact with CaM that is conjugated with Alexa Fluor 647 or Alexa Fluor 594. This medium-content array identified 173 novel CaM/CaML targets in plant. A high-content protein array has been applied to profile CaM-target interactions in the human proteome.⁵⁶ This protein array contains 37,200 redundant proteins and over 10,000 unique human proteins expressed from a human brain cDNA library. One of the novel findings from this study is the identification of a high-affinity interaction between CaM and the ER-resident single-pass transmembrane proteins, stromal interaction molecule (STIM), which plays instrumental roles in the store operated Ca^{2+} entry process by translocating from ER toward plasma membrane to open the ORAI1 store operated Ca^{2+} channel.⁵⁷

4. Conclusions

The Ca^{2+} signaling field has been constantly reinvigorated with the continuous discovery of new components and expansion of the Ca^{2+} -signaling toolkits. The combination of powerful bioinformatics tools and innovative methodologies is expected to enable us to explore a growing list of Ca^{2+} -binding proteins and Ca^{2+} -modulated interactomes that are currently underrepresented. Our ability to investigate the structure-function correlations of CaBPs has advanced in leaps with the evolution of algorithms capable of recognizing or predicting Ca^{2+} -binding sites. It is optimistically anticipated that the development of tools to predict Ca^{2+} -binding sites in apo-structures may become routine in the near future. Moving toward rational protein design based on computational prediction, it is highly expected that the engineering of metal binding proteins with precise affinity and metal selectivity will

significantly improve our ability to provide proteins with tailored functions for diagnostic or therapeutic purposes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We appreciate the critical review by Natalie White and Kathryn Meenach. This work is supported in part by research from NIH GM081749 to JJY and a Special Fellow Award (3013-12) from the Leukemia & Lymphoma Society to YZ.

References

1. Williams RJ. The evolution of calcium biochemistry. *Biochim Biophys Acta*. 2006; 1763:1139–46.10.1016/j.bbamcr.2006.08.042 [PubMed: 17023069]
2. Ross MD. The influence of gravity on structure and function of animals. *Adv Spce Res*. 1984; 4:305–14.
3. (a) Stini WA. Calcium homeostasis and human evolution. *Coll Antropol*. 1998; 22:411–25. [PubMed: 9887597] (b) Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhratsky A. Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system. *Cell Calcium*. 2007; 42:345–50.10.1016/j.ceca.2007.05.001 [PubMed: 17574670]
4. Brown EM. Clinical lessons from the calcium-sensing receptor. *Nat Clin Pract Endocrinol Metab*. 2007; 3:122–33.10.1038/ncpendmet0388 [PubMed: 17237839]
5. Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol*. 2003; 4:517–29.10.1038/nrm1155 [PubMed: 12838335]
6. (a) Gangola P, Rosen BP. Maintenance of intracellular calcium in *Escherichia coli*. *J Biol Chem*. 1987; 262:12570–4. [PubMed: 2442165] (b) Rosen BP. Bacterial calcium transport. *Biochim Biophys Acta*. 1987; 906:101–10. [PubMed: 2436666] (c) Vyas NK, Vyas MN, Quiocho FA. A novel calcium binding site in the galactose-binding protein of bacterial transport and chemotaxis. *Nature*. 1987; 327:635–8.10.1038/327635a0 [PubMed: 3600760] (d) Watkins NJ, Knight MR, Trewavas AJ, Campbell AK. Free calcium transients in chemotactic and non-chemotactic strains of *Escherichia coli* determined by using recombinant aequorin. *Biochem J*. 1995; 306(Pt 3):865–9. [PubMed: 7702585] (e) Reusch RN, Huang R, Bramble LL. Poly-3-hydroxybutyrate/polyphosphate complexes form voltage-activated Ca²⁺ channels in the plasma membranes of *Escherichia coli*. *Biophys J*. 1995; 69:754–66.10.1016/S0006-3495(95)79958-1 [PubMed: 8519976] (f) Durell SR, Guy HR. A putative prokaryote voltage-gated Ca(2+) channel with only one 6TM motif per subunit. *Biochem Biophys Res Commun*. 2001; 281:741–6.10.1006/bbrc.2001.4408 [PubMed: 11237720]
7. Kippert F. Endocytobiotic coordination, intracellular calcium signaling, and the origin of endogenous rhythms. *Ann N Y Acad Sci*. 1987; 503:476–95. [PubMed: 3304083]
8. Kirberger M, Wang X, Zhao K, Tang S, Chen G, Yang JJ. Integration of Diverse Research Methods to Analyze and Engineer Ca-Binding Proteins: From Prediction to Production. *Curr Bioinform*. 2010; 5:68–80.10.2174/157489310790596358 [PubMed: 20802832]
9. Kawasaki H, Nakayama S, Kretsinger RH. Classification and evolution of EF-hand proteins. *Biometals*. 1998; 11:277–95. [PubMed: 10191494]
10. (a) Zhou Y, Yang W, Kirberger M, Lee HW, Ayalasomayajula G, Yang JJ. Prediction of EF-hand calcium-binding proteins and analysis of bacterial EF-hand proteins. *Proteins*. 2006; 65:643–55.10.1002/prot.21139 [PubMed: 16981205] (b) Zhou Y, Frey TK, Yang JJ. Viral calciomics: interplays between Ca²⁺ and virus. *Cell Calcium*. 2009; 46:1–17.10.1016/j.ceca.2009.05.005 [PubMed: 19535138]
11. (a) Rigden DJ, Woodhead DD, Wong PW, Galperin MY. New structural and functional contexts of the Dx[DN]xDG linear motif: insights into evolution of calcium-binding proteins. *PLoS One*. 2011; 6:e21507.10.1371/journal.pone.0021507 [PubMed: 21720552] (b) Rigden DJ, Jedrzejewski MJ, Galperin MY. An extracellular calcium-binding domain in bacteria with a distant relationship to

- EF-hands. *FEMS Microbiol Lett.* 2003; 221:103–10. [PubMed: 12694917] (c) Michiels J, Xi C, Verhaert J, Vanderleyden J. The functions of Ca(2+) in bacteria: a role for EF-hand proteins? *Trends Microbiol.* 2002; 10:87–93. [PubMed: 11827810]
12. (a) Hunter S, Jones P, Mitchell A, Apweiler R, Attwood TK, Bateman A, Bernard T, Binns D, Bork P, Burge S, de Castro E, Coggill P, Corbett M, Das U, Daugherty L, Duquenne L, Finn RD, Fraser M, Gough J, Haft D, Hulo N, Kahn D, Kelly E, Letunic I, Lonsdale D, Lopez R, Madera M, Maslen J, McAnulla C, McDowall J, McMenamin C, Mi H, Mutowo-Muellenet P, Mulder N, Natale D, Orengo C, Pesseat S, Punta M, Quinn AF, Rivoire C, Sangrador-Vegas A, Selengut JD, Sigrist CJ, Scheremetjew M, Tate J, Thimmajananathan M, Thomas PD, Wu CH, Yeats C, Yong SY. InterPro in 2011: new developments in the family and domain prediction database. *Nucleic Acids Res.* 2012; 40:D306–D312.10.1093/nar/gkr948 [PubMed: 22096229] (b) Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer EL, Eddy SR, Bateman A, Finn RD. The Pfam protein families database. *Nucleic Acids Res.* 2012; 40:D290–D301.10.1093/nar/gkr1065 [PubMed: 22127870]
 13. Krogh A, Brown M, Mian IS, Sjolander K, Haussler D. Hidden Markov models in computational biology. Applications to protein modeling. *J Mol Biol.* 1994; 235:1501–31.10.1006/jmbi.1994.1104 [PubMed: 8107089]
 14. Schuster-Bockler B, Schultz J, Rahmann S. HMM Logos for visualization of protein families. *BMC Bioinformatics.* 2004; 5:7.10.1186/1471-2105-5-7 [PubMed: 14736340]
 15. Finn RD, Clements J, Eddy SR. HMMER web server: interactive sequence similarity searching. *Nucleic Acids Res.* 2011; 39:W29–37.10.1093/nar/gkr367 [PubMed: 21593126]
 16. Bertini I, Cavallaro G. Bioinformatics in bioinorganic chemistry. *Metallomics.* 2010; 2:39–51.10.1039/b912156k [PubMed: 21072373]
 17. (a) Kirberger M, Wang X, Deng H, Yang W, Chen G, Yang JJ. Statistical analysis of structural characteristics of protein Ca²⁺-binding sites. *J Biol Inorg Chem.* 2008; 13:1169–81.10.1007/s00775-008-0402-7 [PubMed: 18594878] (b) Yang W, Lee HW, Hellinga H, Yang JJ. Structural analysis, identification, and design of calcium-binding sites in proteins. *Proteins.* 2002; 47:344–56. [PubMed: 11948788] (c) Pidcock E, Moore GR. Structural characteristics of protein binding sites for calcium and lanthanide ions. *J Biol Inorg Chem.* 2001; 6:479–89. [PubMed: 11472012]
 18. Schymkowitz JW, Rousseau F, Martins IC, Ferkinghoff-Borg J, Stricher F, Serrano L. Prediction of water and metal binding sites and their affinities by using the Fold-X force field. *Proc Natl Acad Sci U S A.* 2005; 102:10147–52.10.1073/pnas.0501980102 [PubMed: 16006526]
 19. (a) Nayal M, Di Cera E. Predicting Ca(2+)-binding sites in proteins. *Proc Natl Acad Sci U S A.* 1994; 91:817–21. [PubMed: 8290605] (b) Yamashita MM, Wesson L, Eisenman G, Eisenberg D. Where metal ions bind in proteins. *Proc Natl Acad Sci U S A.* 1990; 87:5648–52. [PubMed: 2377604]
 20. Lin HH, Han LY, Zhang HL, Zheng CJ, Xie B, Cao ZW, Chen YZ. Prediction of the functional class of metal-binding proteins from sequence derived physicochemical properties by support vector machine approach. *BMC Bioinformatics.* 2006; 7(Suppl 5):S13.10.1186/1471-2105-7-S5-S13 [PubMed: 17254297]
 21. Bordner AJ. Predicting small ligand binding sites in proteins using backbone structure. *Bioinformatics.* 2008; 24:2865–71.10.1093/bioinformatics/btn543 [PubMed: 18940825]
 22. Brylinski M, Skolnick J. FINDSITE-metal: integrating evolutionary information and machine learning for structure-based metal-binding site prediction at the proteome level. *Proteins.* 2011; 79:735–51.10.1002/prot.22913 [PubMed: 21287609]
 23. Deng H, Chen G, Yang W, Yang JJ. Predicting calcium-binding sites in proteins - a graph theory and geometry approach. *Proteins.* 2006; 64:34–42.10.1002/prot.20973 [PubMed: 16617426]
 24. Wang X, Kirberger M, Qiu F, Chen G, Yang JJ. Towards predicting Ca²⁺-binding sites with different coordination numbers in proteins with atomic resolution. *Proteins.* 2009; 75:787–98.10.1002/prot.22285 [PubMed: 19003991]
 25. Zhao K, Wang X, Wong HC, Wohlhueter R, Kirberger MP, Chen G, Yang JJ. Predicting Ca(2+)-binding sites using refined carbon clusters. *Proteins.* 2012; 80:2666–79.10.1002/prot.24149 [PubMed: 22821762]

26. Liang MP, Banatao DR, Klein TE, Brutlag DL, Altman RB. WebFEATURE: An interactive web tool for identifying and visualizing functional sites on macromolecular structures. *Nucleic Acids Res.* 2003; 31:3324–7. [PubMed: 12824318]
27. Glazer DS, Radmer RJ, Altman RB. Combining molecular dynamics and machine learning to improve protein function recognition. *Pac Symp Biocomput.* 2008:332–43. [PubMed: 18229697]
28. Wang X, Zhao K, Kirberger M, Wong H, Chen G, Yang JJ. Analysis and prediction of calcium-binding pockets from apo-protein structures exhibiting calcium-induced localized conformational changes. *Protein Sci.* 2010; 19:1180–90.10.1002/pro.394 [PubMed: 20512971]
29. Hoeflich KP, Ikura M. Calmodulin in action: diversity in target recognition and activation mechanisms. *Cell.* 2002; 108:739–42. [PubMed: 11955428]
30. Yap KL, Kim J, Truong K, Sherman M, Yuan T, Ikura M. Calmodulin target database. *J Struct Funct Genomics.* 2000; 1:8–14. [PubMed: 12836676]
31. Radivojac P, Vucetic S, O'Connor TR, Uversky VN, Obradovic Z, Dunker AK. Calmodulin signaling: analysis and prediction of a disorder-dependent molecular recognition. *Proteins.* 2006; 63:398–410.10.1002/prot.20873 [PubMed: 16493654]
32. (a) Ataman ZA, Gakhbar L, Sorensen BR, Hell JW, Shea MA. The NMDA receptor NR1 C1 region bound to calmodulin: structural insights into functional differences between homologous domains. *Structure.* 2007; 15:1603–17. S0969-2126(07)00410-8 [pii]. 10.1016/j.str.2007.10.012 [PubMed: 18073110] (b) Maximciuc AA, Putkey JA, Shamoo Y, Mackenzie KR. Complex of calmodulin with a ryanodine receptor target reveals a novel, flexible binding mode. *Structure.* 2006; 14:1547–56. S0969-2126(06)00361-3 [pii]. 10.1016/j.str.2006.08.011 [PubMed: 17027503]
33. (a) Feldkamp MD, Yu L, Shea MA. Structural and energetic determinants of apo calmodulin binding to the IQ motif of the Na(V)1.2 voltage-dependent sodium channel. *Structure.* 2011; 19:733–47. S0969-2126(11)00096-7 [pii]. 10.1016/j.str.2011.02.009 [PubMed: 21439835] (b) Chagot B, Chazin WJ. Solution NMR structure of Apo-calmodulin in complex with the IQ motif of human cardiac sodium channel NaV1.5. *J Mol Biol.* 2011; 406:106–19. S0022-2836(10)01276-3 [pii]. 10.1016/j.jmb.2010.11.046 [PubMed: 21167176]
34. Zhou Y, Tzeng WP, Wong HC, Ye Y, Jiang J, Chen Y, Huang Y, Suppiah S, Frey TK, Yang JJ. Calcium-dependent association of calmodulin with the rubella virus nonstructural protease domain. *J Biol Chem.* 2010; 285:8855–68.10.1074/jbc.M109.097063 [PubMed: 20086014]
35. (a) Zhou Y, Yang W, Lurtz MM, Ye Y, Huang Y, Lee HW, Chen Y, Louis CF, Yang JJ. Identification of the calmodulin binding domain of connexin 43. *J Biol Chem.* 2007; 282:35005–17.10.1074/jbc.M707728200 [PubMed: 17901047] (b) Zhou Y, Yang W, Lurtz MM, Chen Y, Jiang J, Huang Y, Louis CF, Yang JJ. Calmodulin mediates the Ca²⁺-dependent regulation of Cx44 gap junctions. *Biophys J.* 2009; 96:2832–48.10.1016/j.bpj.2008.12.3941 [PubMed: 19348766] (c) Chen Y, Zhou Y, Lin X, Wong HC, Xu Q, Jiang J, Wang S, Lurtz MM, Louis CF, Veenstra RD, Yang JJ. Molecular interaction and functional regulation of connexin50 gap junctions by calmodulin. *Biochem J.* 2011; 435:711–22.10.1042/BJ20101726 [PubMed: 21320072]
36. (a) Tadross MR, Dick IE, Yue DT. Mechanism of local and global Ca²⁺ sensing by calmodulin in complex with a Ca²⁺ channel. *Cell.* 2008; 133:1228–40. DOI: S0092-8674(08)00684-3 [pii] 10.1016/j.cell.2008.05.025. [PubMed: 18585356] (b) Dick IE, Tadross MR, Liang H, Tay LH, Yang W, Yue DT. A modular switch for spatial Ca²⁺ selectivity in the calmodulin regulation of CaV channels. *Nature.* 2008; 451:830–4. DOI: nature06529 [pii] 10.1038/nature06529. [PubMed: 18235447]
37. Jiang J, Zhou Y, Zou J, Chen Y, Patel P, Yang JJ, Balog EM. Site-specific modification of calmodulin Ca(2)(+) affinity tunes the skeletal muscle ryanodine receptor activation profile. *Biochem J.* 2010; 432:89–99. DOI: BJ20100505 [pii] 10.1042/BJ20100505. [PubMed: 20815817]
38. Maruyama K, Mikawa T, Ebashi S. Detection of calcium binding proteins by ⁴⁵Ca autoradiography on nitrocellulose membrane after sodium dodecyl sulfate gel electrophoresis. *J Biochem.* 1984; 95:511–9. [PubMed: 6715311]
39. Campbell KP, MacLennan DH, Jorgensen AO. Staining of the Ca²⁺-binding proteins, calsequestrin, calmodulin, troponin C, and S-100, with the cationic carbocyanine dye “Stains-all”. *J Biol Chem.* 1983; 258:11267–73. [PubMed: 6193121]

40. Sharma Y, Gopalakrishna A, Balasubramanian D, Fairwell T, Krishna G. Studies on the interaction of the dye, stains-all, with individual calcium-binding domains of calmodulin. *FEBS Lett.* 1993; 326:59–64. DOI: 0014-5793(93)81761-N [pii]. [PubMed: 8325390]
41. Ide Y, Nagasaki N, Tomioka R, Suito M, Kamiya T, Maeshima M. Molecular properties of a novel, hydrophilic cation-binding protein associated with the plasma membrane. *J Exp Bot.* 2007; 58:1173–83.10.1093/jxb/erl284 [PubMed: 17264065]
42. Naaby-Hansen S, Diekmann A, Shetty J, Flickinger CJ, Westbrook A, Herr JC. Identification of calcium-binding proteins associated with the human sperm plasma membrane. *Reprod Biol Endocrinol.* 2010; 8:6.10.1186/1477-7827-8-6 [PubMed: 20078857]
43. (a) Gilchrist JS, Pierce GN. Identification and purification of a calcium-binding protein in hepatic nuclear membranes. *J Biol Chem.* 1993; 268:4291–9. [PubMed: 8440713] (b) Milner RE, Baksh S, Shemanko C, Carpenter MR, Smillie L, Vance JE, Opas M, Michalak M. Calreticulin, and not calsequestrin, is the major calcium binding protein of smooth muscle sarcoplasmic reticulum and liver endoplasmic reticulum. *J Biol Chem.* 1991; 266:7155–65. [PubMed: 2016321] (c) Rajini B, Shridas P, Sundari CS, Muralidhar D, Chandani S, Thomas F, Sharma Y. Calcium binding properties of gamma-crystallin: calcium ion binds at the Greek key beta gamma-crystallin fold. *J Biol Chem.* 2001; 276:38464–71.10.1074/jbc.M102164200 M102164200 [pii] [PubMed: 11502736] (d) Rehm H, Wiedenmann B, Betz H. Molecular characterization of synaptophysin, a major calcium-binding protein of the synaptic vesicle membrane. *EMBO J.* 1986; 5:535–41. [PubMed: 3086086]
44. Israelson A, Arzoine L, Abu-hamad S, Khodorkovsky V, Shoshan-Barmatz V. A photoactivable probe for calcium binding proteins. *Chem Biol.* 2005; 12:1169–78.10.1016/j.chembiol.2005.08.006 [PubMed: 16298296]
45. (a) Lobinski R, Schaumloffel D, Szpunar J. Mass spectrometry in bioinorganic analytical chemistry. *Mass Spectrom Rev.* 2006; 25:255–89.10.1002/mas.20069 [PubMed: 16273552] (b) Montes-Bayon M, DeNicola K, Caruso JA. Liquid chromatography-inductively coupled plasma mass spectrometry. *J Chromatogr A.* 2003; 1000:457–76. [PubMed: 12877184]
46. (a) Ye Y, Lee HW, Yang W, Shealy S, Yang JJ. Probing site-specific calmodulin calcium and lanthanide affinity by grafting. *J Am Chem Soc.* 2005; 127:3743–50.10.1021/ja042786x [PubMed: 15771508] (b) Yang JJ, Gawthrop A, Ye Y. Obtaining site-specific calcium-binding affinities of calmodulin. *Protein Pept Lett.* 2003; 10:331–45. [PubMed: 14529487] (c) Ye Y, Shealy S, Lee HW, Torshin I, Harrison R, Yang JJ. A grafting approach to obtain site-specific metal-binding properties of EF-hand proteins. *Protein Eng.* 2003; 16:429–34. [PubMed: 12874376]
47. Ye Y, Lee HW, Yang W, Shealy SJ, Wilkins AL, Liu ZR, Torshin I, Harrison R, Wohlhueter R, Yang JJ. Metal binding affinity and structural properties of an isolated EF-loop in a scaffold protein. *Protein Eng.* 2001; 14:1001–13. [PubMed: 11809931]
48. Horrocks WD Jr. Luminescence spectroscopy. *Methods Enzymol.* 1993; 226:495–538. [PubMed: 8277879]
49. (a) Zhou Y, Tzeng WP, Yang W, Ye Y, Lee HW, Frey TK, Yang J. Identification of a Ca²⁺-binding domain in the rubella virus nonstructural protease. *J Virol.* 2007; 81:7517–28.10.1128/JVI.00605-07 [PubMed: 17475644] (b) Yanyi C, Shenghui X, Yubin Z, Jie YJ. Calciomics: prediction and analysis of EF-hand calcium binding proteins by protein engineering. *Sci China Chem.* 2010; 53:52–60. [PubMed: 20802784]
50. Jiang Y, Huang Y, Wong HC, Zhou Y, Wang X, Yang J, Hall RA, Brown EM, Yang JJ. Elucidation of a novel extracellular calcium-binding site on metabotropic glutamate receptor 1{alpha} (mGluR1{alpha}) that controls receptor activation. *J Biol Chem.* 2010; 285:33463–74.10.1074/jbc.M110.147033 [PubMed: 20705606]
51. Huang Y, Zhou Y, Wong HC, Chen Y, Wang S, Castiblanco A, Liu A, Yang JJ. A single EF-hand isolated from STIM1 forms dimer in the absence and presence of Ca²⁺ *FEBS J.* 2009; 276:5589–97.10.1111/j.1742-4658.2009.07240.x [PubMed: 19694801]
52. O'Day DH. CaMBOT: profiling and characterizing calmodulin-binding proteins. *Cell Signal.* 2003; 15:347–54. [PubMed: 12618209]
53. (a) Shen X, Valencia CA, Gao W, Cotten SW, Dong B, Huang BC, Liu R. Ca(2+)/Calmodulin-binding proteins from the *C. elegans* proteome. *Cell Calcium.* 2008; 43:444–56.10.1016/j.ceca.2007.07.008 [PubMed: 17854888] (b) Shen X, Valencia CA, Szostak JW, Dong B, Liu R.

- Scanning the human proteome for calmodulin-binding proteins. *Proc Natl Acad Sci U S A*. 2005; 102:5969–74.10.1073/pnas.0407928102 [PubMed: 15840729]
54. Berggard T, Arrigoni G, Olsson O, Fex M, Linse S, James P. 140 mouse brain proteins identified by Ca²⁺-calmodulin affinity chromatography and tandem mass spectrometry. *J Proteome Res*. 2006; 5:669–87.10.1021/pr050421l [PubMed: 16512683]
 55. Popescu SC, Popescu GV, Bachan S, Zhang Z, Seay M, Gerstein M, Snyder M, Dinesh-Kumar SP. Differential binding of calmodulin-related proteins to their targets revealed through high-density Arabidopsis protein microarrays. *Proc Natl Acad Sci U S A*. 2007; 104:4730–5.10.1073/pnas.0611615104 [PubMed: 17360592]
 56. O'Connell DJ, Bauer M, Linse S, Cahill DJ. Probing calmodulin protein-protein interactions using high-content protein arrays. *Methods Mol Biol*. 2011; 785:289–303.10.1007/978-1-61779-286-1_20 [PubMed: 21901608]
 57. (a) Hogan PG, Lewis RS, Rao A. Molecular basis of calcium signaling in lymphocytes: STIM and ORAI. *Annu Rev Immunol*. 2010; 28:491–533.10.1146/annurev.immunol.021908.132550 [PubMed: 20307213] (b) Zhou Y, Meraner P, Kwon HT, Machnes D, Oh-hora M, Zimmer J, Huang Y, Stura A, Rao A, Hogan PG. STIM1 gates the store-operated calcium channel ORAI1 in vitro. *Nat Struct Mol Biol*. 2010; 17:112–6.10.1038/nsmb.1724 [PubMed: 20037597]
 58. Haiech J, Moulhaye SB, Kilhoffer MC. The EF-Handome: combining comparative genomic study using FamDBtool, a new bioinformatics tool, and the network of expertise of the European Calcium Society. *Biochim Biophys Acta*. 2004; 1742:179–83.10.1016/j.bbamcr.2004.10.001 [PubMed: 15590068]
 59. Horst JA, Samudrala R. A protein sequence meta-functional signature for calcium binding residue prediction. *Pattern Recognit Lett*. 2010; 31:2103–2112.10.1016/j.patrec.2010.04.012 [PubMed: 20824111]
 60. Pradeep KN, Piyush R, Pooja K. MetalloPred: A tool for hierarchical prediction of metal ion binding proteins using cluster of neural networks and sequence derived features. *Journal of Biophysical Chemistry*. 2011; 2:112–123.10.4236/jbpc.2011.22014
 61. Roche DB, Tetchner SJ, McGuffin LJ. FunFOLD: an improved automated method for the prediction of ligand binding residues using 3D models of proteins. *BMC Bioinformatics*. 2011; 12:160.10.1186/1471-2105-12-160 [PubMed: 21575183]
 62. Huang B. MetaPocket: a meta approach to improve protein ligand binding site prediction. *Omics*. 2009; 13:325–30.10.1089/omi.2009.0045 [PubMed: 19645590]
 63. Sodhi JS, Bryson K, McGuffin LJ, Ward JJ, Wernisch L, Jones DT. Predicting metal-binding site residues in low-resolution structural models. *J Mol Biol*. 2004; 342:307–20.10.1016/j.jmb.2004.07.019 [PubMed: 15313626]
 64. Bhardwaj N, Stahelin RV, Langlois RE, Cho W, Lu H. Structural bioinformatics prediction of membrane-binding proteins. *J Mol Biol*. 2006; 359:486–95.10.1016/j.jmb.2006.03.039 [PubMed: 16626739]
 65. Dahiyat BI, Mayo SL. De novo protein design: fully automated sequence selection. *Science*. 1997; 278:82–7. [PubMed: 9311930]
 66. Hellinga HW, Richards FM. Construction of new ligand binding sites in proteins of known structure. I. Computer-aided modeling of sites with pre-defined geometry. *J Mol Biol*. 1991; 222:763–85. [PubMed: 1749000]

Figure 1. Schematics of the Ca^{2+} signaling machinery, the range of Ca^{2+} -binding affinities and the timescale of Ca^{2+} modulated activities

The extracellular Ca^{2+} homeostasis is maintained by the coordinated actions of hormones, bone cells and balanced uptake and excretion of Ca^{2+} in intestine and kidney. The internal Ca^{2+} homeostasis is achieved through the exquisite choreography of the Ca^{2+} signaling toolkits. Under resting conditions, cytosolic Ca^{2+} is maintained at submicromolar range by extruding Ca^{2+} outside of the plasma membrane via plasma membrane Ca^{2+} -ATPase (PMCA) and $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX), or by pumping Ca^{2+} back into internal stores through sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase (SERCA) or secretory pathway Ca^{2+} -ATPase (SPCA). Upon extracellular stimulation, the free cytosolic Ca^{2+} rapidly increases by the entry of extracellular Ca^{2+} across the plasma membrane via Ca^{2+} channels, including voltage-operated channels (VOC), receptor-operated channels (ROC), transient receptor potential ion-channel (TRP) and store-operated channels (SOC), or by the release of Ca^{2+} from internal stores (e.g., endoplasmic reticulum (ER) and Golgi complex) through inositol-1,4,5-triphosphate receptors (IP₃R) and ryanodine receptors (RyR) due to activation of membrane receptors (G protein coupled receptors (GPCRs) and receptor tyrosine kinase [RTK]) and the subsequent synthesis of IP₃. In mitochondria, Ca^{2+} can easily pass through outer mitochondrial membrane pores and cross the inner mitochondrial membrane through the membrane-embedded Ca^{2+} uniporter (MCU). Ca^{2+} exits mitochondria through the opening of a nonselective high-conductance channel permeability transition pore (PTP) in the inner mitochondrial membrane and the $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX). The Ca^{2+} signals are delivered by affecting the activity of Ca^{2+} buffers, Ca^{2+} effectors and Ca^{2+} -regulated enzymes. Ca^{2+} -binding proteins have Ca^{2+} affinities that vary by 10^6 -fold or more depending upon their locations and functions (*left panel*). Ca^{2+} can exert short-term effects

by triggering neurotransmitter release within microseconds. The signals can also elicit “long-term” effects by modulating gene expression (*right panel*).

Figure 2. Profile HMMs for three representative continuous Ca^{2+} -binding motifs

The numbers on top of the amino acids indicate the positions of ligands in the Ca^{2+} -binding loop. The Pfam entries for EF-hand like (Excalibur), canonical EF-hand and pseudo EF-hand motifs are PF05901, PF00036 and PF01023, respectively. The letter size is positively correlated to the distribution probability of amino acids at each position. See text for detailed discussions (“Section 2.1 Prediction from primary sequences”).

Figure 3. Schematic diagram of approaches used to study calciomics

Starting from protein amino acid sequences, continuous Ca^{2+} -binding sites can be simply detected by pattern search, whereas some non-continuous Ca^{2+} -binding domains (CaBDs) can be predicted by domain assignment using Pfam. Complementary to these methods, prediction can be made from 3D structural information using several online web servers (e.g., MUG, as listed in Table 1). The proteome for Ca^{2+} -binding proteins (CaBPs) can be obtained by standard proteomic approaches coupled with the use of radioactive metals (^{45}Ca or ^{103}Ru). Quantitative mass spectrometry techniques (e.g., ESI-MS and ICP-MS) can be further applied to characterize the metal/protein stoichiometry of CaBPs. To determine the Ca^{2+} -binding affinities, one can use equilibrium dialysis coupled with ICP-MS (or ^{45}Ca) under circumstances that the predicted CaBP can be readily produced to high purity while retaining its function. For continuous Ca^{2+} -binding sites, a grafting approach can be used to probe metal binding properties without the need to purify the predicted protein. An important aspect of calciomics is to define the CaBP-target interactome by combining both computational and high-throughput experimental approaches. Next, the cellular localization of putative CaBPs can be traced by tagging the CaBP of interest with fluorescent proteins. Lastly, to establish possible functional correlations of Ca^{2+} -modulated activities, loss-of-function experiments should be carried out by removing the predicted Ca^{2+} -binding ligands.

Figure 4. Prediction of Ca^{2+} -binding sites by the geometry and graph (GG) algorithm program

To predict Ca^{2+} -binding sites in proteins, oxygen ions (black) from protein 3D-structure are extracted, while other types of ions are excluded. Next, the distance between any of two oxygen ions is calculated, and an edge will be assigned if the calculated oxygen distance is below a cutoff value. A potential Ca^{2+} -binding site contains oxygen clusters in which every oxygen ion is linked to each other with an O-O distance less than the cutoff and every oxygen ions is linked to all other oxygen atoms by the assigned edges in this oxygen cluster. The Ca^{2+} center is determined at an equidistant center within each oxygen cluster if the distance ranges from 1.8 to 3.0 Å. Adapted from Deng et al.,²³ with permission from John Wiley and Sons.

Figure 5. The interaction of Ca^{2+} -CaM with target proteins

Temporal and spatial changes of the Ca^{2+} concentration in different compartments of cells affect the regulation of cellular signaling, by modulating the activity of a large body of CaBPs. Upon binding to Ca^{2+} , CaBP such as calmodulin (CaM) undergoes substantial conformational changes and subsequently activate or inactivate over 100 functional enzymes, cellular receptors and ion channels through direct protein-protein interactions.

Table 1List of the bioinformatics resources for probing Ca²⁺ signaling.

Resource (Reference)	Type of resource	Web address
<i>Predicting Ca²⁺-binding sites based on protein primary sequences</i>		
CaPS ^{10a}	Webserver for prediction of EF-hand or EF-hand like Ca ²⁺ -binding motifs	http://chemistry.gsu.edu/faculty/Yang/Calciomics.htm
EF-hand CaBPs data library	Database of sequence and structural information on EF-hand CaBPs	http://structbio.vanderbilt.edu/cabp_database/cabp.html
EF-Handome ⁵⁸	A collection of gene, mRNA and protein information of EF-hand proteins in human, rat and mouse	Available from the authors
MFSCa ⁵⁹	A meta-functional signature algorithm for Ca ²⁺ -binding residue prediction	Available from the authors
MetalloPred ⁶⁰	Webserver for prediction of metal-binding sites using cascade of neural networks from sequence derived features	http://www.juit.ac.in/assets/MetalloPred/
CalPred	Web tools for EF-hand CaBP prediction and calcium binding region identification	http://www.bioinformatics.org/calpred/index.html
<i>Predicting Ca²⁺-binding sites based on protein structures (PDB entries or modeled structures)</i>		
GG ²³ , MUG ²⁴ , MUG ^{SR(28)}	Tools for predicting Ca ²⁺ -binding sites based on graph theory and geometric analyses	http://chemistry.gsu.edu/faculty/Yang/Calciomics.htm
Fold-X ¹⁸	A computations algorithm based on empirical force field to predict the position of metal ions in protein	http://foldx.crg.es/
SVMProt ²⁰	Webserver for assigning protein functions (including Ca ²⁺ -binding) with support vector machine learning	http://bidd.cz3.nus.edu.sg/cgi-bin/svmprot.cgi
WebFEATURE ²⁶	Webserver for automated function prediction (including Ca ²⁺ -binding sites) in protein structures with machine learning	http://feature.stanford.edu/webfeature/
SitePredict ²¹	Webserver for predicting metal ion binding sites with the Random Forest machine learning method	http://sitepredict.org/index.php
FunFOLD ⁶¹	An integrated web resource for ligand binding site prediction	http://www.reading.ac.uk/bioinf/FunFOLD/FunFOLD_form.html
FINDSITE-metal ²²	A threading-based method to detect metal-binding site in modeled structures by integrating evolutionary information and machine learning	http://cssb.biology.gatech.edu/findsite-metal
MetaPocket ⁶²	A consensus method to predict ligand binding sites by integrating LIGSITE, PASS, Q-SiteFinder and SURFNET, Fpocket, GHECOM, ConCavity and POCASA	http://projects.biotec.tu-dresden.de/metapocket/index.php
MetSite ⁶³	An automatic approach for detecting metal-binding residues in low-resolution 3D models by with neural network classifiers	http://bioinf.cs.ucl.ac.uk/metsite
<i>Predicting Ca²⁺-modulated functions</i>		
Calmodulin Target Database ³⁰	Webserver for predicting calmodulin binding sites from protein sequences	http://calcium.uhnres.utoronto.ca/ctdb/ctdb/home.html
MeTaDor ⁶⁴	Webserver for predicting membrane targeting domains (e.g., Ca ²⁺ -dependent C2 domain)	http://proteomics.bioengr.uic.edu/metador/predict.html

Resource (Reference)	Type of resource	Web address
ORBIT ⁶⁵	De novo design of protein sequence based on a desired backbone fold	Available from the authors
DEZYMER ⁶⁶	Design of metal binding sties by selecting suitable ligands	Available from the authors

Table 2

Distribution of predicted major Ca^{2+} -binding motifs/domains in different organisms *

Motifs/CaBPs (Pfam entry)	Locations & Functions	Organisms						
		Viruses	Archaea	Bacteria	Fungi	Yeast	Plants	Metazoa
EF-hand like (e.g.Excalibur (PF05901))	External: enzymes	93**	-	662	-	-	-	-
EF-hand (PF13499)	Internal: triggers, buffers, enzymes, messengers	-	6	509	2303	192	3540	6471
S100 (PF01023)	Internal or external: triggers, buffers, messengers	-	-	-	-	-	-	393
C2 domains (PF00168)	Internal: membrane-linked enzymes	-	1	8	1435	257	2180	6700
Annexins (PF00168)	Internal: binds to membrane phospholipids; triggers	-	-	3	336	-	823	2765
Calreticulin (PF00262)	Calcium stores in reticula; chaperone	-	-	-	106	20	158	352
Calsequestrin (PF01216)	Calcium stores in reticula	-	-	-	-	-	-	105

* Note: the values indicate the numbers of predicted Ca^{2+} -binding motifs in protein sequences annotated by Pfam (<http://pfam.sanger.ac.uk/>) as of the year of 2011. One protein sequence may contain multiple predicted motifs.

** The number includes both EF-hand and EF-hand like motifs according to Zhou *et al.* ^{10b}.