Published in final edited form as:

J Allergy Clin Immunol. 2013 April ; 131(4): 1058–1063. doi:10.1016/j.jaci.2012.10.023.

Abuse during Childhood and Adolescence and Risk of Adult-Onset Asthma in African American Women

Patricia F. Coogan, Sc.D.^a, Lauren A. Wise, Sc.D.^a, George T. O'Connor, M.D.^b, Timothy A. Brown, Psy.D.^c, Julie R. Palmer, Sc.D.^a, and Lynn Rosenberg, Sc.D.^a

^aSlone Epidemiology Center at Boston University

^bBoston University School of Medicine

^cCenter for Anxiety and Related Disorders, Boston University

Abstract

Background—Experiences of violence contribute to the occurrence of childhood asthma but there is little information on the effect of early life abuse on adult-onset asthma.

Objective—We prospectively assessed the relation between physical and sexual abuse during childhood and adolescence and the incidence of adult-onset asthma in the Black Women's Health Study.

Methods—We followed 28,456 women from 1995 through 2011 with biennial mailed questionnaires. Experiences of physical and sexual abuse that occurred during childhood and adolescence were obtained in 2005. Cox regression models were used to estimate incidence rate ratios (IRR) and 95% confidence intervals (CI).

Results—During 417,931 person-years of follow-up, 1160 participants reported physician diagnosed asthma and concurrent use of asthma medication. Compared with women who experienced no abuse during childhood or adolescence, the multivariable IRR for any childhood abuse was 1.24 (95% CI 1.06–1.45) and for any adolescent abuse it was 1.10 (95% CI 0.88–1.36). The IRR was higher for childhood physical abuse (IRR=1.29, 95% CI 1.07–1.49) than for childhood sexual abuse (IRR=1.15, 95% CI 0.88–1.49). IRRs for physical and sexual abuse during adolescence were compatible with 1.0. The association between childhood abuse and asthma incidence was stronger in older compared with younger women.

Conclusion—In this large cohort of African American women, there was a positive association between adult-onset asthma and childhood physical abuse, and weaker associations for childhood sexual abuse and any abuse during adolescence. Given the high prevalence of asthma and of childhood abuse, the association is of public health importance.

Keywords

Asthma; child abuse; adolescent abuse; incidence; longitudinal studies; cohort studies; African Americans; women

© 2012 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

Corresponding author: Patricia F. Coogan, Slone Epidemiology Center at Boston University, 1010 Commonwealth Ave., Boston, MA 02215, (tel) 617 734 6006, (fax) 617 738 5119, pcoogan@bu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

The high and increasing prevalence of asthma in the United States (8% of the population in 2009¹) poses enormous burdens to the affected individuals and to society.² African American women suffer disproportionately from asthma. In 2001–2003 National Health Interview Survey data, the average annual prevalence was 9.5% among black women compared with 8.4% among white women.³ The average annual rate of death from asthma among adult asthmatics was 4.6/10,000 among black women and 2.7/10,000 among white women.³

The prevalence of childhood and adolescent abuse has been reported to be twice as high among black children as white children.⁴ Cross-sectional studies show that the prevalence and severity of childhood asthma are associated with experiences of violence.^{5–7} In addition, two prospective studies suggest that exposure to violence may increase asthma incidence in children.^{8, 9} However, there is little information on the effect of childhood abuse on adult-onset asthma. A meta-analysis indicated that respiratory problems are more common among adults who experienced abuse as children,¹⁰ but most studies have been small and cross-sectional, and have grouped respiratory problems together. In two studies that assessed asthma specifically, both prevalence¹¹ and incidence¹² were twice as high in subjects who reported childhood abuse as those who did not. The hypothesized mechanism linking abuse to asthma incidence is stress and its physiological consequences, particularly effects on the immune system.^{6, 13, 14}

To determine whether physical or sexual abuse during childhood or adolescence is associated with increased risk of adult onset asthma, we conducted a prospective analysis of abuse during childhood and adolescence in relation to the incidence of adult-onset asthma in the Black Women's Health Study (BWHS), a large cohort study of African American women who have been followed for more than 15 years.

METHODS

Study population

The BWHS is a prospective cohort study established in 1995, when 59,000 African-American women aged 21 through 69 years were recruited mainly from subscribers to Essence magazine, a general readership magazine targeted to black women.¹⁵ The baseline questionnaire elicited information on demographic and lifestyle factors, reproductive history, and medical conditions. The cohort is followed biennially by mailed and web health questionnaires. Follow-up has averaged over 80 percent of the original cohort through seven questionnaire cycles. The study protocol was approved by the Institutional Review Board of Boston University.

This analysis uses data from the baseline questionnaire and eight subsequent follow-up cycles (1995 through 2011). The 2005 questionnaire elicited information on abuse experienced during childhood, adolescence, and adulthood. We excluded women who did not complete the 2005 questionnaire (n=15,826) or answer any of the questions about abuse (n=8412), reported prevalent asthma (n=3347) or lung cancer (n=18) at baseline, or reported asthma during childhood (n=749).

Diagnosis of asthma

On all questionnaires from 1997 through 2011 participants were asked whether they had been diagnosed with asthma in the previous two year interval, the year in which they had first been diagnosed, and whether they used "inhalers or pills" for asthma at least 3 days per week. Cases were defined as women who reported an initial physician diagnosis of asthma

after 1995 together with concurrent use of asthma medication. We excluded women who reported incident asthma but did not report concurrent asthma medication use (n=1208) and women who reported asthma medication use but no diagnosis of asthma (n=985). After exclusions, 28,456 women remained.

Ascertainment of Abuse

On the 2005 questionnaire, participants were asked multiple questions about experiences of abuse that occurred “as a child (up to age 11)”, “as a teenager (age 12–18)”, and as an adult (“age 19 to present”). The 9-item abuse assessment instrument was adapted from the Conflict Tactics Scales¹⁶ and the Pregnancy Abuse Assessment Screen.¹⁷ Response categories were never, 1–3 times, and 4 times. Methods for creating the summary abuse variables have been described elsewhere.¹⁸ In brief, a physical abuse summary score was created to indicate frequency of occurrence of non-sexual abuse (e.g., someone “pushed, grabbed, or shoved me”; “kicked, bit, or punched me”); the resulting score ranged from 0 to 9 and was categorized as low (score=1), intermediate (score=2), or high (score=3). A sexual abuse summary score was created from two questions that ascertained sexual abuse (i.e., someone has “been sexual with me against my will”, or “exposed genitals against my will”) with categories of never, 1–3 incidents and 4+ incidents. Participants were also asked how often they felt in danger in the home, “physically or sexually”, separately as a child and as a teenager (never, 1–3 times, and 4+ times).

Ascertainment of Covariates

Data on weight and height were obtained in 1995 and weight was updated on each subsequent follow-up questionnaire. Data on smoking history were obtained in 1995 and smoking status and amount smoked were updated on each follow-up questionnaire. Alcohol consumption, hours/week spent in vigorous physical activity, parity, menopausal status and use of menopausal female hormones were ascertained in 1995 and updated every questionnaire cycle. In 1997 respondents were asked about exposure to secondhand tobacco smoke at home at ages 0–10, 11–20, 21–30, 31–40, and currently; they were asked for exposure at the office for ages 21 and older. Exposure was defined as being “in the same room with a smoker for at least an hour a day for 12 consecutive months or more”. Information was also obtained on history of asthma in either parent, whether the participant had health insurance and a regular doctor (obtained in 1997); household income and number of people supported by the income (obtained in 2003); marital status (updated in 2005); and whether the participant’s mother smoked when pregnant with her (obtained in 2009).

Statistical Analysis

We used Cox proportional hazards regression models to estimate incidence rate ratios (IRR, synonymous with the relative risk) and 95% confidence intervals (CI) for the association between abuse victimization and incident asthma. The reference category for all analyses was women who reported no abuse during childhood or adolescence. The basic model was adjusted for age and calendar time. Separate variables for physical and sexual abuse during childhood and adolescence were included simultaneously in the basic model. We then added potential confounders: parental history of asthma, body mass index (BMI, weight in kg divided by height in m squared), hours/week of vigorous physical activity, menopausal female hormone use, household income, number of people supported by the income, education, exposure to secondhand smoke, current smoking, pack years of smoking, and alcohol consumption. Indicator variables were used to model missing covariate data. The following variables did not materially change the results and therefore were not included in the final model: having access to health insurance, having a regular doctor, marital status, height, and whether the participant’s mother smoked when pregnant with her.

In a separate model we assessed the effect of four covariates that may mediate the relation between childhood or adolescent abuse and asthma incidence because they may be consequences of early abuse and in the pathway linking abuse and asthma: adult abuse and scores on the depression, stress and coping scales. The 2005 questionnaire included scales that assessed symptoms of depression (The Center for Epidemiological Studies-Depression (CES-D) scale¹⁹), symptoms of stress (a short form of the Perceived Stress Scale²⁰), and coping skills (a short form of the Brief COPE scale²¹). For each scale, a simple composite score was created by summing the items.

We conducted analyses stratified by age, BMI, smoking status, adult abuse, and scores on the CES-D, stress and coping scales. We tested for statistical interaction with the likelihood ratio test comparing models with and without interaction terms. Analyses were conducted with SAS statistical software version 9.3 (SAS Institute, Inc., Cary, NC).

RESULTS

During 16 years of follow-up comprising 417,931 person-years, 1160 women reported incident adult-onset asthma together with asthma medication use. The mean age at diagnosis was 47 years (SD=11) (range, 23–82).

Compared with women who were not abused during childhood or adolescence, abused women were younger, more likely to be current drinkers and to have been exposed to secondhand smoke, and to have reported more pack-years of smoking, a higher prevalence of adult abuse, and more symptoms of depression and stress (table 1).

IRRs for asthma were increased among women who reported childhood abuse (Table 2), with a multivariable IRR for any childhood abuse of 1.24 (95% CI 1.06–1.45). The multivariable IRR for childhood physical abuse only was 1.26 (95% CI 1.07–1.49) and it was 1.15 (95% CI 0.88–1.49) for childhood sexual abuse only. All IRRs for adolescent abuse were consistent with 1.0.

Adding the potential mediators to the multivariable model did not materially change the IRRs for childhood physical abuse (data not shown). For example, the IRR for the highest category of frequency of childhood physical abuse decreased from 1.35 (95% CI 1.08–1.69) to 1.28 (95% CI 1.02–1.61) when terms for adult abuse and scores on the stress, coping, and CES-D scales were added.

IRRs for women who felt in danger in the home in addition to being abused as children (38% of those reporting child abuse) were higher than for those who were abused but did not feel in danger (table 3). For example, the IRR was 1.53 (95% CI 1.15–2.03) for the highest frequency of physical abuse among women who felt in danger and 1.26 (95% CI 0.97–1.64) among women who did not feel danger in the home.

The IRRs for child abuse were higher in women who were age 40 and older at baseline compared with those among women younger than age 40 (table 4). The IRR for any child abuse was 1.43 (95% CI 1.14–1.78) among older women compared to 1.07 (95% CI 0.85–1.33) among younger women (p for interaction = 0.02); the IRR in the highest category of frequency of physical abuse was 1.54 (95% CI 1.11–2.14) in the older women and it was 1.18 (95% CI 0.86–1.62) in the younger women (p for interaction > 0.05). Regardless of age at asthma diagnosis, the difference in IRRs by baseline age persisted.

To assess whether recall bias influenced the effect estimates, we calculated IRRs for childhood physical abuse confining follow-up to the years after abuse was ascertained in 2005 (table 4). IRRs from the later follow-up period were higher than those from the entire

follow-up period. The variation in IRRs by age was also apparent in the late follow-up period (data not shown).

The association of childhood physical abuse with asthma incidence was present within strata of smoking (ever/never), CES-D score (<16, ≥16), COPE and stress scales (< median, ≥ median), baseline BMI (<25, 25–29, 30+), and presence of adult abuse (ever/never) (all p for interaction >0.05) (data not shown).

When we confined the cases to those who reported use of an asthma preventative (e.g., fluticasone) and thus may have had more severe asthma (769 cases), the IRRs were similar to the main results (e.g., IRR for any child abuse = 1.23, 95% CI 1.02–1.49). When we used a less stringent case definition and included as cases all women who reported physician-diagnosed asthma regardless of the use of asthma medications (2268 cases), the IRRs for childhood abuse were also similar to the main results (e.g., IRR for any child abuse = 1.22, 95% CI 1.09–1.37). The results were also unchanged in analyses that included cases who reported childhood asthma (1267 total cases).

DISCUSSION

In this large cohort of African American women, exposure to physical abuse during childhood was associated with an increased incidence of adult-onset asthma of more than 20%. The risk was higher among women who also felt in danger in the home as a child. IRRs were weaker for sexual abuse during childhood, and there was little evidence that abuse during adolescence increased asthma incidence. The adverse effect of childhood physical abuse was confined to older women.

Two previous studies have assessed the relation of childhood abuse with adult asthma. In a small cross-sectional study of 354 women sampled from a community in New Zealand,¹¹ the prevalence of asthma ($n=33$) was twice as high in women who reported any sexual abuse before age 16 compared with those who did not; asthma was not associated with childhood physical abuse. Results from the much larger cross-sectional 10-country World Mental Health study¹² are consistent with those of the present study. Based on 649 cases of adult-onset asthma reported by 18,303 participants who were interviewed about experiences of childhood abuse, the hazard ratio for adult asthma was 1.92 (95% CI 1.32–2.81) among those who reported physical abuse and 1.26 (95% CI 0.84–1.82) among those who reported childhood sexual abuse.¹²

A study of war-related stressors and asthma incidence in Kuwait following the 1990 Iraqi invasion supports the notion that stress can contribute to adult-onset asthma.²² In that study, adults reporting the highest stress exposure were more than twice as likely to report incident asthma over 13 years of follow-up as those reporting no stressors. In addition, several cross-sectional studies have found that trauma and/or abuse during adulthood was positively associated with adult asthma prevalence.^{23–26} We did not assess asthma incidence in relation to adult abuse because we did not know which came first until 2005, when abuse was ascertained; in analyses confined to follow-up after 2005 there was no relation between adult abuse and asthma incidence. When we included adult abuse as a potential mediator, the IRRs for childhood physical abuse were not materially changed.

The observed association of childhood physical abuse with adult-onset asthma could result from the stress caused by the abuse persisting into adulthood^{13, 14} and adversely affecting the immune system, the airways, and/or the autonomic nervous system. Stress has been shown to be associated with more severe airway inflammation and airflow obstruction in an allergen inhalation challenge among subjects with established asthma.²⁷ In a mouse model of allergic airway inflammation, stress was observed to worsen the histopathologic changes

of inflammation in the airways.²⁸ Alternatively, stress in childhood could lead to long-lasting or even permanent alterations in the immune system, the airways, or the autonomic nervous system that predispose to the development of asthma decades after the abuse occurred. During infancy and early childhood, adaptive immunity is undergoing maturation and the airways are undergoing growth and epithelial changes.²⁹ Animal data also suggest that the innervation of the airways continues to develop in early post-natal life and may be altered by environmental exposures,³⁰ thereby influencing the risk of airway hyperresponsiveness. Environmental factors and their physiologic consequences, including stress, that affect these developmental processes in childhood – especially early childhood – could have sequelae many years later that influence asthma risk. Exposures during this key developmental period could have a greater impact, even on outcomes occurring in later adulthood, than exposures encountered in adolescence or young adulthood.

There is no ready explanation for our observation that the adverse effect of child abuse was stronger in women age 40 and older at baseline. Lung function declines with age,³¹ and perhaps the effect of early life stress manifests most strongly in the aging lung.

Asthma was self-reported in the present study. There is no gold standard for defining asthma,^{32, 33} and in a large national study like BHWS it is infeasible to clinically examine all cases to verify asthma. It is also impossible to examine all noncases to identify undiagnosed asthma. Thus self-reported physician-diagnosed asthma has been recommended as the preferred outcome for use in large population-based studies³⁴ and has indeed been used in many prospective epidemiologic studies.^{35–38} To increase specificity, we included use of an asthma medication in the case definition. In a validation study, among 43 women who met our case definition and who gave us permission to contact their physician, 39 (91%) were confirmed by their physician as asthma cases. In the present analysis, the association between childhood abuse and asthma incidence was robust to both more stringent and less stringent case definitions.

Detection bias might have occurred if women who suffered childhood abuse were more likely to go to the doctor and to complain of asthma symptoms than women not abused. Evidence against detection bias is as follows: almost all BWHS participants have access to care and health insurance regardless of abuse, adjustment for these factors did not change the results, and results were similar when we restricted cases to those who may have had more severe asthma.

We relied on self-reported abuse and were unable to validate abuse occurrence, type, or severity. However, the abuse questions were based on instruments that have been widely used and have demonstrated high reproducibility both within our cohort¹⁸ and other studies.^{16, 17} Recall bias could have occurred if women who developed asthma over-reported abuse or were more likely to remember abuse compared with women who did not develop asthma. The fact that the IRRs from the late follow-up period, based on cases that developed after the report of child abuse in 2005, were higher than the IRRs from the main analysis argues against recall bias.

Our analyses included many comparisons, which increases the chance that some significant associations may be spurious. Nevertheless, our finding of an association of childhood physical abuse with adult-onset asthma is credible in light of similar findings from the World Mental Health Study.¹² Confirmation from other prospective studies is desirable.

In conclusion, ours is the first longitudinal study to provide evidence of a positive association between childhood physical abuse and adult onset asthma. This relation has not previously been assessed in African American women. According to the Department of Health and Human Service's National Child Abuse and Neglect Data System⁴, in 2010

approximately 695,000 children aged 0–17 were determined by state Child Protective Service agencies to be neglected or abused, of which 18% was physical abuse. The true prevalence of child abuse may be much higher given underreporting.^{39,40} In light of the high prevalence of abuse and of asthma in the United States, the association is of public health importance. Clinicians should be vigilant to screen for and to intervene to stop child abuse not only to prevent acute physical injuries and psychological effects but also to prevent long-term medical sequelae such as asthma.

Acknowledgments

This work was funded by a grant from the National Heart, Lung, and Blood Institute of the National Institutes of Health, HL107314.

Abbreviations

BMI	Body Mass Index
BWHS	Black Women's Health Study
CES-D	Center for Epidemiological Studies - Depression
CI	Confidence Interval
COPD	Chronic Obstructive Pulmonary Disease
IRR	incidence rate ratio

References

- Centers for Disease Control and Prevention. [Accessed 6/21/2012] Vital Signs. May. 2011 <http://www.cdc.gov/vitalS>
- Cisternas MG, Blanc PD, Yen IH, Katz PP, Earnest G, Eisner MD, et al. A comprehensive study of the direct and indirect costs of adult asthma. *J Allergy Clin Immunol*. 2003; 111:1212–8. [PubMed: 12789219]
- Moorman JE, Rudd RA, Johnson CA, King M, Minor P, Bailey C, et al. National surveillance for asthma--United States, 1980–2004. *MMWR Surveill Summ*. 2007; 56:1–54. [PubMed: 17947969]
- U.S. Department of Health and Human Services. Child Maltreatment 2010. Children's Bureau; 2011. Administration on Children, Youth, and Families. Available from <http://www.acf.hhs.gov/program/cb/stats-research/index.htm#can>
- Wright RJ, Mitchell H, Visness CM, Cohen S, Stout J, Evans R, et al. Community violence and asthma morbidity: the Inner-City Asthma Study. *Am J Public Health*. 2004; 94:625–32. [PubMed: 15054016]
- Wright RJ, Steinbach SF. Violence: an unrecognized environmental exposure that may contribute to greater asthma morbidity in high risk inner-city populations. *Environ Health Perspect*. 2001; 109:1085–9. [PubMed: 11675274]
- Cohen RT, Canino GJ, Bird HR, Celedon JC. Violence, abuse, and asthma in Puerto Rican children. *Am J Respir Crit Care Med*. 2008; 178:453–9. [PubMed: 18556622]
- Suglia SF, Enlow MB, Kullowatz A, Wright RJ. Maternal intimate partner violence and increased asthma incidence in children: buffering effects of supportive caregiving. *Arch Pediatr Adolesc Med*. 2009; 163:244–50. [PubMed: 19255392]
- Sternthal MJ, Jun HJ, Earls F, Wright RJ. Community violence and urban childhood asthma: a multilevel analysis. *Eur Respir J*. 2010; 36:1400–9. [PubMed: 20413538]
- Wegman HL, Stetler C. A meta-analytic review of the effects of childhood abuse on medical outcomes in adulthood. *Psychosom Med*. 2009; 71:805–12. [PubMed: 19779142]

11. Romans S, Belaise C, Martin J, Morris E, Raffi A. Childhood abuse and later medical disorders in women. An epidemiological study. *Psychother Psychosom*. 2002; 71:141–50. [PubMed: 12021556]
12. Scott KM, Von Korff M, Alonso J, Angermeyer MC, Benjet C, Bruffaerts R, et al. Childhood adversity, early-onset depressive/anxiety disorders, and adult-onset asthma. *Psychosom Med*. 2008; 70:1035–43. [PubMed: 18941133]
13. McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med*. 1998; 338:171–9. [PubMed: 9428819]
14. Heim C, Newport DJ, Heit S, Graham YP, Wilcox M, Bonsall R, et al. Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *JAMA*. 2000; 284:592–7. [PubMed: 10918705]
15. Rosenberg L, Adams-Campbell L, Palmer JR. The Black Women's Health Study: a follow-up study for causes and preventions of illness. *J Am Med Womens Assoc*. 1995; 50:56–8. [PubMed: 7722208]
16. Straus MA. Measuring intrafamily conflict and violence: the Conflict Tactics (CT) scales. *J Marriage Fam*. 1979; 41:75–88.
17. McFarlane J, Parker B, Soeken K, Bullock L. Assessing for abuse during pregnancy. Severity and frequency of injuries and associated entry into prenatal care. *Jama*. 1992; 267:3176–8. [PubMed: 1593739]
18. Wise LA, Palmer JR, Rothman EF, Rosenberg L. Childhood abuse and early menarche: findings from the black women's health study. *Am J Public Health*. 2009; 99 (Suppl 2):S460–6. [PubMed: 19443822]
19. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psych Measurement*. 1977; 1:385–401.
20. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983; 24:385–96. [PubMed: 6668417]
21. Carver CS. You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med*. 1997; 4:92–100. [PubMed: 16250744]
22. Wright RJ, Fay ME, Suglia SF, Clark CJ, Evans JS, Dockery DW, et al. War-related stressors are associated with asthma risk among older Kuwaitis following the 1990 Iraqi invasion and occupation. *J Epidemiol Community Health*. 2010; 64:630–5. [PubMed: 20231738]
23. Coker AL, Smith PH, Fadden MK. Intimate partner violence and disabilities among women attending family practice clinics. *J Womens Health (Larchmt)*. 2005; 14:829–38. [PubMed: 16313210]
24. Loxton D, Schofield M, Hussain R, Mishra G. History of domestic violence and physical health in midlife. *Violence Against Women*. 2006; 12:715–31. [PubMed: 16861329]
25. Spitzer C, Barnow S, Volzke H, John U, Freyberger HJ, Grabe HJ. Trauma, posttraumatic stress disorder, and physical illness: findings from the general population. *Psychosom Med*. 2009; 71:1012–7. [PubMed: 19834051]
26. Subramanian SV, Ackerson LK, Subramanyam MA, Wright RJ. Domestic violence is associated with adult and childhood asthma prevalence in India. *Int J Epidemiol*. 2007; 36:569–79. [PubMed: 17329314]
27. Liu LY, Coe CL, Swenson CA, Kelly EA, Kita H, Busse WW. School examinations enhance airway inflammation to antigen challenge. *Am J Respir Crit Care Med*. 2002; 165:1062–7. [PubMed: 11956045]
28. Deshmukh A, Kim BJ, Gonzales X, Caffrey J, Vishwanatha J, Jones HP. A murine model of stress controllability attenuates Th2-dominant airway inflammatory responses. *J Neuroimmunol*. 2010; 225:13–21. [PubMed: 20462642]
29. Holt PG, Upham JW, Sly PD. Contemporaneous maturation of immunologic and respiratory functions during early childhood: implications for development of asthma prevention strategies. *J Allergy Clin Immunol*. 2005; 116:16–24. [PubMed: 15990766]
30. Wu ZX, Hunter DD, Kish VL, Benders KM, Batchelor TP, Dey RD. Prenatal and early, but not late, postnatal exposure of mice to sidestream tobacco smoke increases airway

- hyperresponsiveness later in life. *Environ Health Perspect.* 2009; 117:1434–40. [PubMed: 19750110]
31. Degens H, Maden-Wilkinson TM, Ireland A, Korhonen MT, Suominen H, Heinonen A, et al. Relationship between ventilatory function and age in master athletes and a sedentary reference population. *Age (Dordr)*. 2012 Apr 28. [Epub ahead of print].
 32. Peat JK, Toelle BG, Marks GB, Mellis CM. Continuing the debate about measuring asthma in population studies. *Thorax*. 2001; 56:406–11. [PubMed: 11312411]
 33. Pekkanen J, Pearce N. Defining asthma in epidemiological studies. *Eur Respir J*. 1999; 14:951–7. [PubMed: 10573248]
 34. Burr ML. Diagnosing asthma by questionnaire in epidemiological surveys. *Clin Exp Allergy*. 1992; 22:509–10. [PubMed: 1628247]
 35. Beckett WS, Jacobs DR Jr, Yu X, Iribarren C, Williams OD. Asthma is associated with weight gain in females but not males, independent of physical activity. *Am J Respir Crit Care Med*. 2001; 164:2045–50. [PubMed: 11739133]
 36. Camargo CA Jr, Weiss ST, Zhang S, Willett WC, Speizer FE. Prospective study of body mass index, weight change, and risk of adult-onset asthma in women. *Arch Intern Med*. 1999; 159:2582–8. [PubMed: 10573048]
 37. Nystad W, Meyer HE, Nafstad P, Tverdal A, Engeland A. Body mass index in relation to adult asthma among 135,000 Norwegian men and women. *Am J Epidemiol*. 2004; 160:969–76. [PubMed: 15522853]
 38. Romieu I, Avenel V, Leynaert B, Kauffmann F, Clavel-Chapelon F. Body mass index, change in body silhouette, and risk of asthma in the E3N cohort study. *Am J Epidemiol*. 2003; 158:165–74. [PubMed: 12851230]
 39. Flaherty EG, Sege RD, Griffith J, Price LL, Wasserman R, Slora E, et al. From suspicion of physical child abuse to reporting: primary care clinician decision-making. *Pediatrics*. 2008; 122:611–9. [PubMed: 18676507]
 40. Raphael KG, Widom CS, Lange G. Childhood victimization and pain in adulthood: a prospective investigation. *Pain*. 2001; 92:283–93. [PubMed: 11323150]

Key Messages

- In this large cohort of African American women, there was a positive association between childhood abuse and adult-onset asthma.
- The association was stronger for childhood physical than sexual abuse, and was stronger in older compared with younger women.
- The hypothesized mechanism linking abuse to asthma incidence is stress and its physiological consequences, particularly effects on the immune system.

Table I

Selected baseline characteristics of 28,456 BWHS participants according to abuse victimization reported during childhood and adolescence ^{*,†}

Characteristic (% unless noted)	No abuse as child or teen (n=11,980)	Any child abuse (n=13,885)	Any teen abuse (n=10,497)	Abuse as both child and teen (n=7906)
Age at baseline, years (mean ± SD)	40.2 ± 11.0	37.7 ± 10.0	37.6 ± 9.7	37.4 ± 9.6
College graduate	51	50	46	47
Income of \$50,000 or more (2003)	54	54	51	51
Has health insurance (1997)	88	88	87	87
Had a regular MD (1997)	83	83	83	83
Parental history of asthma (1999)	6	7	7	7
Presence of sleep apnea (2001)	2	2	2	3
Female hormone use, ever	14	15	15	16
BMI, kg/m ² (mean ± SD)	27.3 ± 6.2	27.8 ± 6.7	28.0 ± 6.7	28.1 ± 6.8
5+hours/week of vigorous exercise	12	14	14	14
Past drinker	12	15	16	17
Current drinker	22	26	27	26
Current smoker	13	15	16	17
Pack-years of smoking among current/past smokers (mean ± SD)	10.7 ± 11.2	12.6 ± 12.6	12.7 ± 12.5	13.1 ± 12.8
Exposure to secondhand smoke (1997)				
Ages 0–10	44	51	51	53
Ages 11–20	42	50	52	53
Ages 21–30	44	51	53	54
Ages 31–40	30	33	35	35
Currently	17	17	19	18
Any abuse as an adult (2005)	25	51	59	62
Scored 16+ on CES-D scale (2005) (high level depressive symptoms)	20	29	32	33
Highest quartile Perceived Stress Scale (2005) (most stressed)	21	30	32	34
Lowest quartile COPE score (2005)(fewest coping skills)	25	24	23	23

* Age-adjusted in 5 year intervals

† Ascertained at baseline in 1995 unless a different year is noted in parenthesis.

Table II

Abuse during childhood and adolescence and risk of adult-onset asthma, BWHS, 1995–2011

	Cases/person-years	Basic IRR* (95% CI)	Multivariable IRR [†] (95% CI)
STAGE OF ABUSE			
No abuse as child or teen	411/176185	1.0	1.0
Any child abuse	646/203626	1.28 (1.09–1.49)	1.24 (1.06–1.45)
Any teen abuse	492/153657	1.18 (0.95–1.47)	1.10 (0.88–1.36)
Child abuse only	257/88088	1.27 (1.09–1.49)	1.24 (1.06–1.45)
Teen abuse only	103/38120	1.18 (0.95–1.46)	1.10 (0.88–1.36)
Both child and teen abuse	389/115537	1.48 (1.29–1.70)	1.29 (1.12–1.48)
CHILDHOOD ABUSE			
Type of childhood abuse			
Physical abuse only	406/128517	1.28 (1.09–1.51)	1.26 (1.07–1.49)
Sexual abuse only	73/25451	1.20 (0.92–1.56)	1.15 (0.88–1.49)
Physical and sexual abuse	167/49657	1.34 (1.08–1.67)	1.24 (1.00–1.24)
Frequency of childhood physical abuse			
Low	236/79922	1.23 (1.02–1.47)	1.22 (1.01–1.46)
Intermediate	134/39229	1.40 (1.11–1.76)	1.38 (1.10–1.73)
High	203/59024	1.41 (1.12–1.76)	1.35 (1.08–1.69)
Frequency of childhood sexual abuse			
1–3 incidents	151/51708	1.13 (0.85–1.51)	1.09 (0.82–1.46)
4+ incidents	89/23400	1.43 (1.00–2.04)	1.28 (0.90–1.83)
ADOLESCENT ABUSE			
Type of teen abuse			
Physical abuse only	228/72763	1.19 (0.93–1.52)	1.12 (0.87–1.43)
Sexual abuse only	121/42657	1.12 (0.87–1.43)	1.05 (0.82–1.34)
Physical and sexual abuse	143/38238	1.39 (1.06–1.84)	1.23 (0.93–1.62)
Frequency of teen physical abuse			
Low	182/60686	1.10 (0.82–1.48)	1.06 (0.79–1.43)
Intermediate	83/23424	1.27 (0.88–1.82)	1.16 (0.81–1.66)
High	106/26891	1.38 (0.96–1.98)	1.22 (0.85–1.76)
Frequency of teen sexual abuse			
1–3 incidents	211/66359	1.10 (0.84–1.45)	1.04 (0.79–1.37)
4+ incidents	53/14536	1.16 (0.77–1.73)	1.04 (0.70–1.57)

* Adjusted for age and calendar time. Child abuse variables are adjusted for teen abuse and teen abuse variables are adjusted for child abuse. Sexual abuse variables are adjusted for physical abuse and physical abuse variables are adjusted for sexual abuse.

[†] Adds to the basic model parental history of asthma (yes, no, unknown), income (<=\$25,000, \$25,001–\$50,000, \$50,001–\$100,000, >\$100,000, missing), number in household supported by the income (1, 2, 3–4, 5+, missing), years of education (<=12, 13–15, 16, 17+), years of hormone therapy use (never, <5 years, >=5 years, missing), BMI (<25, 25–29, 30–34, 35–39, 40+, missing), hours/week vigorous exercise (none, <5, 5+, missing), 2nd hand smoke exposure at ages 0–10, 11–20, 21–30, 31–40 and currently (yes, no, missing), current smoking (yes, no), pack-years of smoking (never smoke, <5, 5–14, 15–24, 25+, missing), and alcohol consumption (never, past, current 1–3/week, 4–6/week, 7–13/week, 14+/week, missing).

Table IIIIRR_s for asthma according to child abuse and whether danger was felt in the home, BWHS, 1995–2011

	Cases/person-years	Multivariable IRR* (95% CI)
No danger felt in home and no child or teen abuse	389/167031	1.0
NEVER FELT IN DANGER IN HOME		
Any child abuse	400/134666	1.18 (1.03–1.35)
Any physical abuse as child	359/119200	1.24 (1.04–1.47)
Frequency of child physical abuse		
Low	185/63320	1.24 (1.02–1.50)
Intermediate	84/26232	1.32 (1.02–1.71)
High	90/29648	1.26 (0.97–1.64)
Any sexual abuse as child	96/33093	1.12 (0.83–1.51)
YES, FELT IN DANGER IN HOME		
Any child abuse	237/66100	1.31 (1.11–1.56)
Any physical abuse as child	206/56694	1.40 (1.12–1.74)
Frequency of child physical abuse		
Low	46/15347	1.25 (0.88–1.76)
Intermediate	49/12601	1.63 (1.16–2.30)
High	111/28746	1.53 (1.15–2.03)
Any sexual abuse as child	140/40819	1.17 (0.83–1.64)

* Adjusted for age, calendar time, parental history of asthma (yes, no, unknown), income ($\leq$ \$25,000, \$25,001–\$50,000, \$50,001–\$100,000, >\$100,000, missing), number in household supported by the income (1, 2, 3–4, 5+, missing), years of education ($\leq$ 12, 13–15, 16, 17+), years of hormone therapy use (never, <5 years, $\geq$ 5 years, missing), BMI (<25, 25–29, 30–34, 35–39, 40+, missing) hours/week vigorous exercise (none, <5, 5+, missing), 2nd hand smoke exposure at ages 0–10, 11–20, 21–30, 31–40 and currently (yes, no, missing), current smoking (yes, no), pack-years of smoking (never smoke, <5, 5–14, 15–24, 25+, missing), alcohol consumption (never, past, current 1–3/week, 4–6/week, 7–13/week, 14+/week, missing), and adolescent abuse (ever, never). Physical abuse variables are adjusted for sexual abuse and any sexual abuse is adjusted for physical abuse.

Table IV

Abuse during childhood and risk of adult-onset asthma by age, BWHS, 1995–2011

	Cases/person-years	Multivariable IRR* (95% CI)
Age less than 40 at baseline		
No abuse as child or teen	199/89083	1.0
Any child abuse	320/120223	1.07 (0.85–1.33)
Type of child abuse		
Physical abuse only	190/74071	1.06 (0.84–1.35)
Sexual abuse only	37/15016	1.01 (0.70–1.46)
Physical and sexual abuse	93/31136	1.13 (0.84–1.53)
Frequency of child physical abuse		
Low	103/45200	0.96 (0.73–1.26)
Intermediate	69/22846	1.27 (0.92–1.75)
High	111/37162	1.18 (0.86–1.62)
Age 40 and older at baseline		
No abuse as child or teen	212/87102	1.0
Any child abuse	326/83402	1.43 (1.14–4.78)
Type of child abuse		
Physical abuse only	216/54446	1.48 (1.17–1.86)
Sexual abuse only	36/10435	1.27 (0.87–1.84)
Physical and sexual abuse	74/18521	1.35 (0.99–1.86)
Frequency of child physical abuse		
Low	133/34723	1.49 (1.16–1.91)
Intermediate	65/16383	1.46 (1.06–2.02)
High	92/21862	1.54 (1.11–2.14)

* Adjusted for age, calendar time, parental history of asthma (yes, no, unknown), income ($\leq$ \$25,000, \$25,001–\$50,000, \$50,001–\$100,000, >\$100,000, missing), number in household supported by the income (1, 2, 3–4, 5+, missing), years of education ($\leq$ 12, 13–15, 16, 17+), years of hormone therapy use (never, <5 years, $\geq$ 5 years, missing), BMI (<25, 25–29, 30–34, 35–39, 40+, missing) hours/week vigorous exercise (none, <5, 5+, missing), 2nd hand smoke exposure at ages 0–10, 11–20, 21–30, 31–40 and currently (yes, no, missing), current smoking (yes, no), pack-years of smoking (never smoke, <5, 5–14, 15–24, 25+, missing), alcohol consumption (never, past, current 1–3/week, 4–6/week, 7–13/week, 14+/week, missing) and adolescent abuse (ever, never). Physical abuse variables are adjusted for sexual abuse and sexual abuse is adjusted for physical abuse.

Table V

Abuse during childhood and risk of adult-onset asthma during later years of follow-up, BWHS, 2005–2011

	Cases/person-years	Multivariable IRR* (95% CI)
No abuse as child or teen	83/58155	1.0
Any child abuse	138/67542	1.34 (0.95–1.88)
Type of childhood abuse		
Physical abuse only	92/42606	1.46 (1.03–2.09)
Sexual abuse only	11/8462	0.90 (0.47–1.72)
Physical and sexual abuse	35/16474	1.32 (0.83–2.13)
Frequency of childhood physical abuse		
Low	56/26626	1.42 (0.96–2.10)
Intermediate	27/12973	1.39 (0.84–2.29)
High	44/19481	1.47 (0.90–2.38)
Frequency of childhood sexual abuse		
1–3 incidents	29/17263	0.99 (0.50–1.95)
4+ incidents	17/7673	1.22 (0.54–2.79)

* Adjusted for age, calendar time, parental history of asthma (yes, no, unknown), income ($\leq$ \$25,000, \$25,001–\$50,000, \$50,001–\$100,000, >\$100,000, missing), number in household supported by the income (1, 2, 3–4, 5+, missing), years of education ($\leq$ 12, 13–15, 16, 17+), years of hormone therapy use (never, <5 years, $\geq$ 5 years, missing), BMI (<25, 25–29, 30–34, 35–39, 40+, missing) hours/week vigorous exercise (none, <5, 5+, missing), 2nd hand smoke exposure at ages 0–10, 11–20, 21–30, 31–40 and currently (yes, no, missing), current smoking (yes, no), pack-years of smoking (never smoke, <5, 5–14, 15–24, 25+, missing), alcohol consumption (never, past, current 1–3/week, 4–6/week, 7–13/week, 14+/week, missing) and adolescent abuse (ever, never). Physical abuse variables are adjusted for sexual abuse and sexual abuse is adjusted for physical abuse.