Published in final edited form as:

Biochem Biophys Res Commun. 2013 March 29; 433(1): 73–78. doi:10.1016/j.bbrc.2013.02.063.

Translational Efficiency of *rpoS* mRNA from *Borrelia burgdorferi*: Effects of the Length and Sequence of the mRNA Leader Region

Linda Archambault, Joshua Linscott, Nicholas Swerdlow, Kathleen Boyland, Eammon Riley, and Paula Schlax*

Program in Biological Chemistry, Bates College, 5 Andrews Road, Lewiston Maine 04240

Abstract

Regulation of the enzootic cycle in *Borrelia burgdorferi* requires a shift to the RNA polymerase alternative sigma factor, RpoS. We used *in vitro* and *in vivo* assays to assess the relative importance of the putative Shine-Dalgarno sequence and its sequestration for the translational efficiency of *rpoS*. We created mutant leader regions in which we either removed the Shine-Dalgarno sequence, disrupted the secondary structure or both. Binding assays and toeprint assays demonstrated that both the presence and the availability of the Shine-Dalgarno sequence are important to the efficiency and specificity of ribosome binding. Adding a DsrA_{BB} mimic in the form of a single-stranded DNA oligonucleotide increased the level and specificity of binding of RL, presumably by making the Shine-Dalgarno sequence available for binding. In *in vivo* assays we confirmed that the Shine-Dalgarno sequence must be both present and un-sequestered in order for translation to proceed efficiently. The RL transcript was significantly better translated in *B. burgdorferi* at 37°C than at 26°C, lending support to the hypothesis that DsrA_{BB} acts as a temperature-dependent stimulator of translation. These studies demonstrate that translational regulation of gene expression in *B. burgdorferi* may be an important mechanism for responding to environmental signals important in the enzootic cycle.

Keywords

Borrelia burgdorferi; translational regulation; *rpoS*

Introduction

The spirochete, *Borrelia burgdorferi*, is the causative agent of Lyme disease [1-4]. In nature, the spirochete undergoes significant shifts in gene expression patterns as it cycles between an *Ixodes* tick vector and a vertebrate host [5-8]. Temperature [9,10], pH [4,11-13] and mammalian host factors [14,15] play a role in the shifts in gene expression patterns (reviewed by Samuels (2011)[7]).

The alternative sigma factor RpoS is induced during tick feeding and is responsible for global changes in gene expression associated with transmission of the spirochete to a

© 2013 Elsevier Inc. All rights reserved.

*Corresponding Author pschlax@bates.edu, (207)786-6290 .

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

vertebrate host [7,8,16,17]. Two *rpoS* transcripts have been identified in *B. burgdorferi* [18]. At high cell density ($\sim 1 \times 10^8$ cells/ml) a transcript, RS, with a 5' end located 50 nucleotides upstream of the start codon is found [18]. At lower cell density ($1-3 \times 10^7$ cells/ml), a longer transcript, RL, with a 5' end 171 nucleotides upstream of the start codon, is found [18].

The two transcripts are predicted to have different translational efficiencies. (Model presented in Fig. 1.) In *Escherichia coli*, the interaction of the 3' end of the 16S rRNA in the 30S ribosomal subunit with the Shine-Dalgarno sequence (SD) [19] of the mRNA may be rate determining in translational initiation [20]. A strong inverse correlation between translational efficiency and the thermodynamic stability of secondary structure that sequesters the SD sequence has been observed [20-25]. In the model presented by Lybecker and Samuels, the RS transcript is predicted to have an unstructured leader sequence allowing the SD sequence to interact with the 3' end of the 16S rRNA in the 30S ribosomal subunit for efficient translational initiation (Fig. 1a) [18]. The RL transcript is predicted to have extensive secondary structure that involves the base-pairing of the ribosome binding site to an upstream region in the transcript's 5' untranslated region (UTR) [18] to form a stable helix that inhibits translational initiation (Fig. 1b).

The RL transcript contains a region that is complementary to a small non-coding RNA, DsrA_{Bb} [18]. DsrA_{Bb}, in the presence of a *B. burgdorferi*-specific Hfq_{Bb} chaperone [26], up-regulates *rpoS* expression post-transcriptionally with temperature upshift by an as yet unknown mechanism [7,18,26]. Base-pairing of DsrA_{Bb} to the 5' UTR of the mRNA is predicted to change the conformation of the mRNA and increase translational efficiency (Fig. 1c) [18]. In *B. burgdorferi*, two alternative pairings of the 3' end of the 16S rRNA to the leader region of the *rpoS* mRNA are possible (Fig. 2a). Although both are further from the start codon than what is typically observed (<13 nucleotides) [27] for a SD sequence, these pairings share 5 nucleotides in common which will be considered the putative SD sequence for the purpose of these studies.

The role of the length, sequence and structure of the *B. burgdorferi rpoS* 5' UTR on translational efficiency is the subject of these studies. We have examined the translational efficiency of *E. coli* 30S ribosomal subunits (30S_{Ec}) with *rpoS*_{Bb} variants (Fig. 2b) using toe-printing assays and compared the levels of luciferase expression controlled by *rpoS*_{Bb} leader variants in *E. coli* and in *B. burgdorferi*. Our results indicate that the presence of the putative SD sequence is essential for strong ribosome affinity and that sequestration of the sequence reduces translational efficiency. Our results support the models of Lybecker and Samuels [7,18,26] that the RL transcript is poorly translated and that translational efficiency is modulated by a conformational change in the mRNA facilitated by base-pairing a nucleic acid to the 5' UTR of *rpoS* mRNA. .

Materials and Methods

Bacterial culture conditions and construction of variants used in these studies are described in the Supplemental Materials and Methods. All reagents were Molecular Biology grade or better.

rpoS and variant mRNA transcripts

RNA transcripts were prepared using T7 RNA polymerase (Promega, Madison, WI) from PCR fragments and purified as described previously [25]. 1 μ M RNA was denatured in standard renaturation buffer (30 mM Tris-Cl (pH 7.3) and 100 mM potassium acetate) at 65°C for 5 minutes and slow cooled to temperatures below 37°C. In assays involving primer extension, the appropriate 5' γ -³²P labeled primer was added to the RNA before denaturation. In assays involving the DsrA_{Bb}-like DNA oligonucleotide, the DsrA_{Bb}-like

DNA oligonucleotide was added to the RNA to a final concentration of 1 μ M prior to denaturation.

30S subunits

E. coli 30S subunits were purified as previously described [25]. 30S_{Ec} were renatured in binding buffer (20 mM Tris-Cl pH 7.5 at 4 °C, 100 mM NH₄Cl, 10.5 mM magnesium acetate, 0.5 mM EDTA, 3 mM 2-mercaptoethanol) at 37°C for 20 minutes before use.

Toeprint Assays

In toeprint assays [28], varying concentrations of 30S_{Ec} (from 100-1000nM), 1 μ M tRNA_f^{Met} and renatured mRNA transcripts (typically 2 picomoles) were incubated together in toeprint buffer (100 mM potassium acetate, 30 mM Tris-Cl, 1 mM Spermidine, 10 mM MgCl₂, 1 mM DTT) for 15 minutes at 37°C. 0.5-1 units of AMV Reverse Transcriptase (Promega, Madison, WI) were added and reactions were incubated another 20 minutes. Sequencing reactions consisted of ~ 2 picomoles of renatured mRNA transcript annealed to the appropriate radioactive primer, 1 unit of AMV Reverse Transcriptase, ddNTP mix (48 μ M each NTP, 4.8 μ M each ddNTP) (USB/Affymetrix, Santa Clara, CA), 50 mM Tris-HCl, pH 8.3, 10 mM MgCl₂, 0.5 mM spermidine, and 10 mM DTT). Reactions were incubated for 20 min. at 42°C. All reactions were quenched on ice. All reactions were placed in loading buffer (95% Formamide, 18 mM EDTA, 0.025% SDS, and Bromophenol Blue), heated to 90°C for 3 minutes and run on 10% (0.5× TBE) denaturing acrylamide gels. Gels were dried and exposed overnight to phosphorimager plates. The plates were scanned with a Fuji FLA-5000 imaging system.

Luciferase Assay

Luciferase assays were performed in a Glomax Multi Jr. luminometer using a Luciferase Assay System kit (Promega, Madison, WI). Transformed DH5 α *E. coli* were grown to an OD₅₉₀ of 0.4 to 0.6 at 37°C with shaking. *B. burgdorferi* were grown to a density of 7.5×10^6 to 8.5×10^7 cells/mL at 35°C or 26°C and 5% CO₂. Cells were harvested by centrifugation in aliquots of 4×10^7 cells, and treated according to the manufacturer's protocol. Luciferase activity was measured in relative luminescence units (RLUs) for 10s after a 2s delay.

Results and Discussion

Induction of RpoS expression is required for transmission of *B. burgdorferi* from a tick to a vertebrate host [7,8,15,17]. It has been proposed that the folded RL transcript, found at low cell density and temperature mimicking the midgut of an unfed tick host, has poor ribosome binding site accessibility and that interactions with DsrA_{Bb} increase its translational efficiency [18]. Additionally, the RS transcript, which does not form the same inhibitory structure and is found at higher cell density and temperature, should have a relatively high translational efficiency [18] (Fig. 1).

To examine the effects of the putative SD sequence and the mRNA secondary structure on translational efficiency, a series of sequence variants were constructed. The sequence variants were designed to replace the SD sequence with its complement (designated by -SD) or to replace the 5' UTR sequence that pairs to the putative SD sequence with its complement. The double variant (restoring the base-pairing) was also generated (Fig. 2b). The variants were named by the position of their 5' end (-171 or -50), the expectation that the sequence would be single stranded (SS) or double stranded (DS) and the presence (+SD) or absence of the SD sequence (-SD).

To determine that ribosome binding is occurring at the initiation codon and to determine the level of translational efficiency *in vitro*, we performed toeprint assays. The relative abundance of the signal from the toeprint fragment to the signal from the full length fragment is a measure of the translational efficiency in single time point kinetic assays. We observed that primer extension of RS or RL in the absence of either tRNA^{met} or 30S_{Ec} resulted in a full length primer extension product (Fig. 3). The pattern and intensity of toeprint signals depends upon the secondary structure of the mRNA. For the single stranded constructs containing a SD sequence, RS and -171 SS+SD, intense toeprint signals are observed at the expected position, +16 (numbered with +1 corresponding to the translation start site AUG) (e.g Figs. 3 and S3). The RL transcript results in two low intensity toeprint signals at residues +16 and +31 relative to the initiation codon, corresponding to occupancy of the downstream AUG alternative ribosome binding site (Figs 3 and S3). Variants missing the SD sequence (-171 SS-SD, -50 SS-SD, -171 DS-SD) all showed toeprint signals at +16 corresponding to occupancy of the initiation codon of the *rpoS* gene, although at reduced intensity, and at position +31 (Fig. 3). These data suggest that under our conditions, the SD sequence is not essential, but is helpful for formation of a stable *in vitro* translation initiation complex.

The toeprint signals at positions +16 and +31 require an AUG codon 16 or 15 nucleotides respectively upstream (Fig. S3). Other toeprint signals, further downstream of additional AUG sequences in the coding region of the *rpoS* gene were also observed (data not shown). Although alternative ribosome binding sites complicate quantitative analysis of translational efficiency, the intensity of the toeprint signal arising from occupancy at the *rpoS* initiation codon is always higher for constructs with limited secondary structure and functional SD sequences (RS and SS+SD) at all concentrations of 30S_{Ec} tested. Our results indicate that sequestering the SD sequence reduces translational efficiency. They also support a hypothesis that the ribosome occupies a stand-by site on the mRNA of a highly structured message to facilitate rapid occupancy of the appropriate ribosome binding site on the time scale associated with the kinetics of secondary structure shifts in the mRNA [29].

A DNA oligonucleotide that mimicked the action of the *E. coli* DsrA increased ribosome accessibility on *E. coli rpoS* leader sequences [25]. We observe a similar effect here. Annealing the DsrA_{Bb}-like DNA oligonucleotide (DsrA_{Bb} DNA) to RL before primer extension results in inhibition of transcription of the full length product and in formation of truncated primer extension products corresponding to the annealing site of the *rpoS*: DsrA_{Bb} DNA duplex (Fig. 3 and S3). This result suggests that the DNA:RNA hybrid is resistant to dissociation by AMV Reverse Transcriptase. In the presence of 30S_{Ec} and initiator tRNA^{met}, toeprint signals appear downstream of the initiation codon. The intensity of the toeprint signal arising from occupancy of the initiation codon is increased and the signal arising from occupancy of the downstream site (the alternative toeprint signal) is reduced in the presence of DsrA_{Bb} DNA (Fig. 6 and S3), suggesting that melting out mRNA secondary structure by interacting with the DsrA_{Bb} DNA increases 30S_{Ec} binding to the initiation codon and reduces the binding at the alternative ribosome binding site. These data strongly suggest that the ribosome accessibility for RS and the RL:DsrA_{Bb} DNA duplex are higher than for RL. Experiments (supplementary data) performed with 30S_{Ec} indicate a larger fraction of the mRNA participates in ribosome binding for the RS fragment and for the RL fragment annealed to DsrA_{Bb} DNA than to RL. Affinities are similar to those previously reported [30]. Structure mapping data further supports the hypothesis that DsrA_{Bb} DNA alters the mRNA secondary structure to increase the single-stranded nature of the SD sequence. (Fig. S2).

In order to assess the importance of sequences and the potential for forming secondary structures that sequester the SD sequence *in vivo*, translational fusions were constructed

using the leader sequences of the *rpoS* variants described above. The *rpoS* leader regions were subcloned into the NdeI site of pJSB541 downstream of a *flaB* promoter and upstream of a luciferase gene codon optimized for expression in *Borrelia burgdorferi* [31] as described in Supplemental Materials and Methods. The levels of luciferase expression were measured for the RL, RS, -171 SS+SD, -171 SS-SD, -171 DS-SD and -50 SS-SD translational fusion constructs in both *E. coli* DH5 α and in *B. burgdorferi* B31 cfarm (Fig. 4). As expected, the SD sequence and the predicted secondary structure of the mRNA played a significant role in determining the amount of luciferase expressed. In *E. coli* and *B. burgdorferi*, high levels (nearly identical) of luciferase were made from constructs containing SD sequences and that were predicted to be single stranded (RS and -171 SS+SD). In *E. coli*, very low levels of luciferase were expressed from constructs lacking a SD sequence (-171 DS-SD, -171 SS-SD, -50 SS-SD) or predicted to have significant secondary structure (RL, -171 DS-SD). Results in *B. burgdorferi* were similar to the results in *E. coli* with the exception of expression from the construct containing the RL sequence. In *B. burgdorferi*, but not *E. coli*, RL levels were significantly higher than constructs lacking a SD sequence. *B. burgdorferi* contains the trans-acting regulators DsrA_{Bb} and HfqBb [7,18,26] necessary for melting out the secondary structure which sequesters the SD sequence in this construct. This effect is specific to the RL construct and not observed in constructs lacking a SD sequence or significant secondary structure (RS, -171 DS-SD, -50 SS-SD, -171 SS-SD, -171 SS+SD).

Lybecker and Samuels have postulated that the effects of DsrA_{Bb} on *rpoS* expression are temperature dependent [18]. They proposed that at elevated temperatures (35°C) DsrA_{Bb} has a positive effect on expression that is not present at lower temperatures. Within the uncertainty of our experiments, only the constructs which are predicted to be double stranded (RL and -171 DS-SD) show a difference (>4-fold and >2-fold respectively) in the expression levels at 35°C and 26°C (Fig. 4) indicating that the temperature-dependent *rpoS* induction requires the presence of an extended 5' leader region containing the binding site for DsrA_{Bb} and involves base pairing interactions that sequester the region upstream of the *rpoS* start codon.

Samuels and coworkers have reported that in an RpoN mutant (where RS is not transcribed) RpoS levels are low suggesting poor translational efficiency of RL, even in the presence of DsrA_{Bb} and HfqBb [7]. Our *in vitro* results indicate that RL hybridized to a DsrA_{Bb}-like oligonucleotide has a high translational efficiency with 30S_{Ec}: the toeprint signal arising from occupancy of the initiation codon of RL hybridized to DsrA_{Bb}DNA is nearly the same as that observed in the RS construct. Although our *in vitro* experiments use 30S_{Ec}, the 3' ends of the 16S rRNA in *E. coli* (5'...tggatcacctcctta 3' NCBI Reference Sequence: NC_000913.2) and in *B. burgdorferi* (5'...tggatcacctccttt 3' NCBI Reference Sequence: NC_001318.1) are identical in all but the terminal nucleotide. Thus we expect that the first steps in translational initiation, 30S:mRNA interactions and binding of the initiator tRNA [20] are likely to be similar in *E. coli* and *B. burgdorferi*. It is more likely that our high levels of translational efficiency observed *in vitro* result from the use of DsrA_{Bb}DNA. DsrA_{Bb}DNA does have a larger effect on translational efficiency than a DsrA_{Bb} transcript (data not shown), suggesting that the hybridization *in vitro* with this synthetic oligonucleotide is more efficient than the natural regulator. *In vivo*, expression of RL translational fusions is lower than RS-fusions, but still moderately high. Our translational fusions may be slightly more efficient than the natural transcript because they are truncated at the initiation codon and may lack sequences that are essential for translational regulation *in vivo* or that influence rates of mRNA degradation or processing. Regardless, the data presented indicate that the secondary structure of RL sequesters a sequence that is important for translational efficiency, and that altering the structure of the RNA through annealing a

nucleic acid to the upstream region of the mRNA where DsrA_{Bb} binds increases translational efficiency.

Very few mechanisms for transcriptional regulation in *B. burgdorferi* have been identified despite extensive efforts [7]. Our studies demonstrate that translational regulation of gene expression in *B. burgdorferi* may be an important mechanism for responding to environmental signals important in the enzootic cycle.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This project was supported by grants from the National Center for Research Resources (5P20RR016463) and the National Institute of General Medical Sciences (8 P20 GM103423) from the National Institutes of Health. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

Megan Lybecker and D. Scott Samuels provided RL, RS and DsrA PCR products and frequent advice. Jon Blevins provided plasmid pJSB541. Melissa Caimano and Justin Radolf provided *Borrelia burgdorferi* strain CE16 B31 cFarm. Michael Gilbert provided frequent advice and assistance with protocols. Transformations were performed in Peter Brooks' Laboratory at Maine Medical Center Research Institute, Scarborough Maine.

Abbreviations

DsrA_{Bb} DNA	DsrA _{Bb} DNA oligonucleotide
30S_{Ec}	E. coli 30S ribosomal subunits
SD	Shine-Dalgarno
UTR	Untranslated region

Reference List

- [1]. Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. Lyme disease - A tick-borne spirochetosis? Science. 1982; 216:1317–1319. [PubMed: 7043737]
- [2]. Benach JL, Bosler EM, Hanrahan JP, Coleman JL, Habicht GS, Bast TF, Cameron DJ, Ziegler JL, Barbour AG, Burgdorfer W, Edelman R, Kaslow RA. Spirochetes isolated from the blood of two patients with Lyme disease. N. Engl. J. Med. 1983; 308:740–742. [PubMed: 6828119]
- [3]. Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W, Schmid GP, Johnson E, Malawista SE. The spirochetal etiology of Lyme disease. N. Engl. J. Med. 1983; 308:733–740. [PubMed: 6828118]
- [4]. Radolf JD, Salazar JC, Dattwyler RJ. *Borrelia*: Molecular Biology, Host Interaction and Pathogenesis. Lyme disease in humans. 2010:487–533.
- [5]. Singh SK, Girschick HJ. Molecular survival strategies of the Lyme disease spirochete *Borrelia burgdorferi*. Lancet Infectious Diseases. 2004; 4:575–583. [PubMed: 15336225]
- [6]. Radolf JD, Caimano MJ, Stevenson B, Hu LT. Of ticks, mice and men: Understanding the dual-host lifestyle of Lyme disease spirochaetes. Nat. Rev. Microbiol. 2012; 10:87–99. [PubMed: 22230951]
- [7]. Samuels DS. Gene regulation in *borrelia burgdorferi*. Annu. Rev. Microbiol. 2011; 65:479–499. [PubMed: 21801026]
- [8]. Skare JT, Carroll JA, Yang XF, Samuels DS, Akins DR. *Borrelia*: Molecular Biology, Host Interaction and Pathogenesis. Gene regulation, transcriptomics and proteomics. 2010:67–101.
- [9]. Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. Induction of an outer surface protein on *Borrelia burgdorferi* during tick feeding. Proc. Natl. Acad. Sci. U. S. A. 1995; 92:2909–2913. [PubMed: 7708747]

- [10]. Ojaimi C, Brooks C, Casjens S, Rosa P, Elias A, Barbour A, Jasinskas A, Benach J, Katona L, Radolf J, Caimano M, Skare J, Swingle K, Akins D, Schwartz I. Profiling of temperature-induced changes in *Borrelia burgdorferi* gene expression by using whole genome arrays. *Infect. Immun.* 2003; 71:1689–1705. [PubMed: 12654782]
- [11]. Carroll JA, Cordova RM, Garon CF. Identification of 11 pH-regulated Genes in *Borrelia burgdorferi* localizing to linear plasmids. *Infect. Immun.* 2000; 68:6677–6684. [PubMed: 11083781]
- [12]. Carroll JA, El-Hage N, Miller JC, Babb K, Stevenson B. *Borrelia burgdorferi* RevA antigen is a surface-exposed outer membrane protein whose expression is regulated in response to environmental temperature and pH. *Infect. Immun.* 2001; 69:5286–5293. [PubMed: 11500397]
- [13]. Ramamoorthy R, Scholl-Meeker D. *Borrelia burgdorferi* proteins whose expression is similarly affected by culture temperature and pH. *Infect. Immun.* 2001; 69:2739–2742. [PubMed: 11254645]
- [14]. Akins DR, Bourell KW, Caimano MJ, Norgard MV, Radolf JD. A new animal model for studying Lyme disease spirochetes in a mammalian host-adapted state. *J. Clin. Invest.* 1998; 101:2240–2250. [PubMed: 9593780]
- [15]. Caimano MJ, Iyer R, Eggers CH, Gonzalez C, Morton EA, Gilbert MA, Schwartz I, Radolf JD. Analysis of the RpoS regulon in *Borrelia burgdorferi* in response to mammalian host signals provides insight into RpoS function during the enzootic cycle. *Mol. Microbiol.* 2007; 65:1193–1217. [PubMed: 17645733]
- [16]. Caimano MJ, Eggers CH, Hazlett KRO, Radolf JD. RpoS is not central to the general stress response in *Borrelia burgdorferi* but does control expression of one or more essential virulence determinants. *Infect. Immun.* 2004; 72:6433–6445. [PubMed: 15501774]
- [17]. Dunham-Ems SM, Caimano MJ, Eggers CH, Radolf JD. *Borrelia burgdorferi* requires the alternative sigma factor RpoS for dissemination within the vector during tick-to-mammal transmission. *PLoS Pathog.* 2012; 8
- [18]. Lybecker MC, Samuels DS. Temperature-induced regulation of RpoS by a small RNA in *Borrelia burgdorferi*. *Mol. Microbiol.* 2007; 64:1075–1089. [PubMed: 17501929]
- [19]. Shine J, Dalgarno L. The 3' terminal sequence of *Escherichia coli* 16S ribosomal RNA: complementarity to nonsense triplets and ribosome binding sites. *Proc. Natl. Acad. Sci. U. S. A.* 1974; 71:1342–1346. [PubMed: 4598299]
- [20]. Schlax PJ, Worhunsky DJ. Translational repression mechanisms in prokaryotes. *Mol. Microbiol.* 2003; 48:1157–1169. [PubMed: 12787346]
- [21]. De Smit MH, Van Duin J. Secondary structure of the ribosome binding site determines translational efficiency: A quantitative analysis. *Proc. Natl. Acad. Sci. U. S. A.* 1990; 87:7668–7672. [PubMed: 2217199]
- [22]. De Smit MH, Van Duin J. Translational initiation on structured messengers. Another role for the Shine-Dalgarno interaction. *J. Mol. Biol.* 1994; 235:173–184. [PubMed: 8289239]
- [23]. De Smit MH. Translational control by mRNA structure in eubacteria: Molecular biology and physical chemistry. *RNA Structure and Function.* 1998:495–540.
- [24]. Draper DE, Gluick TC, Schlax PJ. Pseudoknots, RNA folding, and translational regulation. *RNA Structure and Function.* 1998:415–436.
- [25]. Worhunsky DJ, Godek K, Litsch S, Schlax PJ. Interactions of the non-coding RNA DsrA and RpoS mRNA with the 30 S ribosomal subunit. *J. Biol. Chem.* 2003; 278:15815–15824. [PubMed: 12600997]
- [26]. Lybecker MC, Abel CA, Feig AL, Samuels DS. Identification and function of the RNA chaperone Hfq in the Lyme disease spirochete *Borrelia burgdorferi*. *Mol. Microbiol.* 2010; 78:622–635. [PubMed: 20815822]
- [27]. Chen H, Bjerknes M, Kumar R, Jay E. Determination of the optimal aligned spacing between the Shine-Dalgarno sequence and the translation initiation codon of *Escherichia coli* mRNAs. *Nucleic Acids Res.* 1994; 22:4953–4957. [PubMed: 7528374]
- [28]. Spedding G, Draper DE. Allosteric mechanism for translational repression in the *Escherichia coli* α operon. *Proc. Natl. Acad. Sci. U. S. A.* 1993; 90:4399–4403. [PubMed: 7685102]

- [29]. De Smit MH, Van Duin J. Translational standby sites: How ribosomes may deal with the rapid folding kinetics of mRNA. *J. Mol. Biol.* 2003; 331:737–743. [PubMed: 12909006]
- [30]. Schlax PJ, Xavier KA, Gluick TC, Draper DE. Translational repression of the *Escherichia coli* α operon mRNA. Importance of an mRNA conformational switch and a ternary entrapment complex. *J. Biol. Chem.* 2001; 276:38494–38501. [PubMed: 11504736]
- [31]. Blevins JS, Revel AT, Smith AH, Bachlani GN, Norgard MV. Adaptation of a luciferase gene reporter and lac expression system to *Borrelia burgdorferi*. *Appl. Environ. Microbiol.* 2007; 73:1501–1513. [PubMed: 17220265]
- [32]. Zuker M. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 2003; 31:3406–3415. [PubMed: 12824337]

Highlights

- B. burgdorferi*'s *rpoS* mRNA leader sequence's secondary structure influences gene expression
- A putative Shine-Dalgarno region is required for strong gene expression
- Modifying RNA sequence and structure can regulate gene expression in *B. burgdorferi*

Figure 1. Model: Translational Regulation of the *Borrelia burgdorferi* *rpoS* mRNA transcript

In this model, the 30S ribosomal subunit is represented by a grey oval. The 3' end of the 16S rRNA is represented as a line within the “ribosome”. Trace drawings [32], of the *rpoS* mRNA leader region are represented as a line drawing.

- The –50 RS mRNA, expressed at high cell density, lacks an extended upstream sequence and is predicted to have an accessible Shine-Dalgarno sequence that pairs to the 3' end of the 16S rRNA.
- The –171 RL mRNA, expressed at low cell density, is predicted to have an extended structure that sequesters the Shine-Dalgarno sequence in an intramolecular base-pairing interaction, preventing the intermolecular interaction with the 16S rRNA.
- At low cell density and elevated temperature, the –171 RL mRNA transcript is predicted to interact with the non-coding RNA DsrA_{Bb} (represented by a dark bar along the mRNA). This interaction is expected to increase the extent of the interaction with the 30S ribosomal subunit, and therefore, the translational efficiency of the mRNA.

Figure 2. Predicted Shine-Dalgarno Interactions in *rpoS* mRNA Sequence Variants

a) The 5' leader region of *rpoS* mRNA [18] has two potential Shine-Dalgarno (SD) sequences that would allow pairing to the 3' end of the 16S rRNA in the 30S ribosomal subunit (grey oval) of *B. burgdorferi*.

b) The putative SD sequence is predicted to be base-paired to an upstream sequence in the long *rpoS* transcript RL. Shown is the sequence and potential base-pairing of this helix for RL and for a series of sequence variants that alter the sequence of the putative SD region and/or the trans-acting upstream sequence which sequesters it. Sequences predicted to lack secondary structure upstream of the initiation codon are RS, -50 SS-SD, -171 S-SD and -171 SS+SD. RL and -171 DS-SD are predicted to have a paired helix upstream of the initiation codon. Constructs RS, RL, and -171 SS+SD are expected to have functional SD sequences.

Figure 3. Interactions of *E. coli* 30S subunits with Sequence and Structure Variants of *rpoS* mRNA

Sequence variants altering the secondary structure and/or eliminating the presence of the Shine-Dalgarno sequence influence the intensities of the primary and alternative toeprint signals. Annealing the DsrA_{Bb}-like DNA oligonucleotide increases the signal of the primary toeprint. The location of the full length transcripts for the -171 constructs (RL, FC and SC) and the -50 construct (RS) are indicated as well as the location of the primary toeprint located at +16 and a secondary, alternative toeprint located at position +31.

Figure 4. *In vivo* Expression Data for Translational Fusions Controlling Luciferase Expression
Translational efficiency of *rpoS* leader region variants as measured by luciferase activity is shown for *B. burgdorferi* grown at 35°C (n=4) and 26°C (-50RS: n=5, -171RL: n=4, others: n=3) and for *E. coli* at 37°C (-171RL: n=3, others: n=4). Luciferase activity was measured in relative luminescence units (RLUs). Constructs in which the Shine-Dalgarno sequence was expected to be available for ribosome binding (-50 RS, -171 SS+SD) showed high levels of expression. In constructs in which the Shine-Dalgarno sequence was either missing (-50 SS-SD, -171SS-SD, -171 DS-SD) or expected to be sequestered (-171 RL), expression levels were greatly reduced. The exception to this trend is seen in the -171 RL construct in *B. burgdorferi* grown at 35°C, where expression levels are somewhat enhanced relative to the same construct in *E. coli* and in *B. burgdorferi* at 26°C.