Published in final edited form as:

Org Lett. 2013 February 15; 15(4): 736–739. doi:10.1021/ol303272w.

Towards a Unified Approach for the Lycopodines: Synthesis of 10-Hydroxylycopodine, Deacetylpaniculine and Paniculine

Mrinmoy Saha and Rich G. Carter

Department of Chemistry, 153 Gilbert Hall, Oregon State University, Corvallis, OR 97331

Rich G. Carter: rich.carter@oregonstate.edu

Abstract

The enantioselective syntheses of 10-hydroxylycopodine, deacetylpaniculine and paniculine have been accomplished through use of a common intermediate. Key steps in the synthetic sequence towards these *lycopodium* alkaloids include formation of the tricyclic core via a conformationally-accelerated, intramolecular Mannich cyclization and an organocatalyzed, intramolecular Michael addition to form the C₇–C₁₂ linkage.

The *lycopodium* alkaloids represent a sizable collection of natural products with wide-ranging and significant biological activity – consisting of over two hundred known compounds from four sub-families (the lycopodines, lycodines, fawcettimines and phlegmarines).¹ The lycopodine and lycodine sub-families have attracted considerable attention from numerous laboratories² including our own.³ Interestingly, despite these efforts, there are wide sections of the *lycopodium* alkaloids that remain unexplored (e.g. compounds **2–5** in Figure 1). Herein, we disclose the first total syntheses of three members of the lycopodine sub-family: 10-hydroxylycopodine (**2**),⁴ deacetylpaniculine (**3**)⁵ and paniculine (**4**).⁵ These natural products **2–4** were isolated from a Chilean club moss *lycopodium confertum*.⁴ In addition, compounds **3** and **4** were also isolated from *lycopodium paniculatum*.^{4,5}

Our unified approach to the synthesis of natural products **2–4** is shown in Scheme 1. We planned to employ a common intermediate strategy using tricycle **7** which should be accessible from the sulfone **6**. Sulfone **6** in turn could be constructed via a tandem sulfone migration followed by intramolecular Mannich cyclization of imine **8** – a process first developed for our total synthesis of lycopodine (**1**).^{3b–c} The imine **8** would be derived from keto sulfone **9**. The key C₇–C₁₂ linkage could be accessed via a diastereoselective intramolecular Michael addition. The proposed strategy builds on our prior work in the total synthesis of **1**,^{3b–c} however, the presence of the additional stereochemistry and functionality

Correspondence to: Rich G. Carter, rich.carter@oregonstate.edu.

Supporting Information Available Complete experimental procedures are provided, including ¹H and ¹³C spectra, of all new compounds. x-ray crystallographic data (CIF) for compounds **9**, **17** and **19** are also provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

at C₁₀ complicates the approach. While natural products **2–4** represent the only known *lycopodium* alkaloids containing oxygenation at the C₁₀ position, this functionality could provide an important handle for accessing compounds possessing an additional C-C bond at this position (e.g. compound **5**).

We commenced our synthesis of the key cyclohexanone **9** with known diol **13**⁶ (Scheme 2). Compound **13** is available in enantioenriched form in 3 steps from 1-bromo-3-butene via sulfone incorporation, epoxidation and Jacobsen hydrolytic kinetic resolution.⁷ Acetonide formation under standard conditions followed by sulfone ester coupling with ester **11**^{3b-c} cleanly provided the coupled material **14**. Next, cross metathesis with 3-penten-2-one and Hoveyda-Grubbs 2nd generation catalyst (**15**) produced the enone **10**. Intramolecular Michael addition of the keto sulfone **10** was smoothly facilitated by the acylsulfonamide catalyst **16** developed in our lab^{3c,8} to provide cyclohexanone **9** in 85% yield and modest 3:1 diastereoselectivity (51% isolated yield of the pure **9** after crystallization). The stereochemistry of **9** was conclusively established by x-ray crystallographic analysis. Addition of 1% EtOH was key to this transformation as its presence led to a noticeable acceleration in reaction rate. It should also be noted that in the absence of this catalyst or the use of an alternate base, the conversion and diastereoselectivity were significantly reduced. In fact, use of our *i*-Pr₂NH, IPA/CH₂Cl₂ conditions (which worked smoothly in lycopodine synthesis^{3b-c}) resulted in <50% conversion and low diastereoselectivity (1.5:1 dr). This divergence in reactivity is related to the presence of the additional functionality at C₁₀ which likely counteracts the directing influence of the C₁₅ methyl moiety.

With the functionalized cyclohexanone in hand, we turned our attention to the synthesis of the key common intermediate (Scheme 3). Acetonide deprotection was facilitated using aqueous HCl in dioxane to give the hemiketal **17**. Sulfonate activation at C₉ followed by azide displacement provided compound **20** along with the ketal **19**. We attribute the formation of **19** to the close conformational proximity of the C₁₃ alcohol and the C₉ mesitylate. In fact, both mesitylate **18** and the azide **20** were prone to forming the ketal **19** upon prolonged storage. Compound **19** could easily be recycled by treatment with aqueous acid to reveal the hemiketal **17**. The hemiketal **20** was sequentially silylated with TBSOTf followed by TIPSOTf to reveal the ketone **21**. Aza-Wittig reaction provided the cyclization precursor **8**. Treatment with Zn(OTf)₂ provided the unexpected tricycle **22** in 57% yield. While compound **22** was a productive intermediate in the synthetic sequence, we had anticipated that the rearranged sulfone **6** would be the major product.^{3b-c} We attribute this divergence in the expected and observed reactivity to the presence of the C₁₀ stereocenter. This additional stereochemistry likely increases the conformational preference of conformer **8** over conformer **8a** (as compared to the C₁₀ deoxy series) for placement of both the C₁₂ sulfone and the C₇ sidearm in the pseudoaxial positions. This conformationally-accelerated intramolecular Mannich cyclization proceeded significantly faster than the C₁₀ deoxy series - leading to preferential Mannich cyclization prior to sulfone rearrangement.^{3b-c} The location of the sulfone moiety is inconsequential as desulfurization using sodium/mercury amalgam provided the key common intermediate **7**.

The synthesis of the tetracyclic core for the *lycopodium* alkaloids is shown in Scheme 4. *N*-Alkylation using 3-iodopropanol provided the primary alcohol **23**. Reaction time for this alkylation appeared to be critical as less than 10 h provided more recovered starting material and longer reaction times led to overalkylation. The tandem Oppenauer oxidation / intramolecular aldol condensation produced the desired enone **24** in 71% yield. The use of freshly prepared potassium *t*-butoxide and freshly distilled benzene were necessary to achieve a reproducibly high yield in this transformation. A small amount of the retro-Michael product **7** (17%) was also isolated in this reaction. Hydrogenation using Adams' catalyst in methanol cleanly yielded the desired product **25**.^{2s,2u}

With the tetracycle in hand, we shifted our attention to the completion of the total syntheses of 10-hydroxylycopodine (**2**), deacetylpaniculine (**3**) and paniculine (**4**) (Scheme 5). TAS-F removal of the TIPS protecting group cleanly provided 10-hydroxylycopodine (**2**). DIBAL-H reduction of ketone **25** cleanly proceeded to provide the axial alcohol as a single stereoisomer.⁹ Acylation using Ac₂O and pyridine in the presence of DMAP yielded the C₅ acetate. Cleavage of the C₁₀ TIPS ether was best accomplished using aqueous HCl to produce the des-silyl product as its HCl salt. Treatment with pH 10 buffer provided paniculine (**4**) and treatment with aqueous sodium hydroxide yielded deacetylpaniculine (**3**).

One important aspect of this work is the clarification of the structural assignments and spectral data for this sub-family of natural products. For example, optical rotation data for these natural products was not provided in the isolation papers.^{4,5} Our synthetic material provides a reference for future isolation chemists to confirm the absolute configuration by comparison to our reported synthetic data. We hypothesize that the absolute stereochemistry should be assigned based on analogy to other *lycopodium* alkaloids (e.g. **1**) for which we have confirmed synthetically.^{3b-c} For compound **2**, the isolation paper only provided limited spectroscopic data for comparison.⁴ From what data was provided, our synthesized material was in good agreement with the natural product. To garner additional confirmation of the original structural assignment of 10-hydroxylycopodine (**2**), we sought to link **2** with other *lycopodium* alkaloids isolated from the same plant. Synthesis of related members of this family (compounds **3** and **4**) showed excellent agreement between the synthesized material and the literature data.⁵ For compound **3**, the isolation paper^{5e} provided a tabular listing of the spectral data which shows both H_{1eq} and H_{14eq} as doublets of doublets at 2.61 ppm; however, careful analysis of actual ¹H NMR spectra provided by the isolation chemists revealed that this data was tabulated incorrectly in the isolation paper.

The first total syntheses of three related members of the lycopodine-subfamily have been accomplished. Key aspects to this work include the development of acylsulfonamide-mediated intramolecular Michael reaction to incorporate the C₇ and C₁₂ stereocenters, a conformationally-accelerated intramolecular Mannich cyclization to construct the tricyclic core and development of a common intermediate **7** that should provide access to additional members of this family of alkaloids.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the National Institutes of Health (NIH) (GM63723) and Oregon State University). Professor Orlando Muñoz (University of Chile) is acknowledged for providing a copy of spectra data for **3** and **4**. The authors are grateful to Professor Claudia Maier & Jeff Morré (OSU) for mass spectra data and Dr. Lev Zakharov (OSU) for x-ray crystallographic analysis of **9**, **17** and **19**. Finally, the authors thank Professor James D. White (OSU), Mr. Nathan Collett (OSU) and Dr. Roger Hanselmann (Rib-X Pharmaceuticals) for their helpful discussions.

References

1. For recent reviews, see: Ma X, Gang DR. Nat. Prod. Rep. 2004; 21:752–772. [PubMed: 15565253] Hirasawa Y, Kobayashi J, Morita H. Heterocycles. 2009; 77:679–729. Kitajima M, Takayama H. Top. Curr. Chem. 2012; 309:1–32. [PubMed: 21452079]
2. (a) Stork G, Kretchmer RA, Schlessinger RH. J. Am. Chem. Soc. 1968; 90:1647–1648. [PubMed: 5636803] (b) Ayer WA, Bowman WR, Joseph TC, Smith PJ. Am. Chem. Soc. 1968; 90:1648–1650. (c) Colvin EW, Martin J, Parker W, Raphael RA, Shroot B, Doyle M. J. Chem. Soc. Perkin Trans. 1. 1972:860–870. (d) Kim S, Bando Y, Horii Z. Tetrahedron Lett. 1978:2293–2294. (e) Heathcock

- CH, Kleinman EF, Binkly ES. *J. Am. Chem. Soc.* 1982; 104:1054–1068.(f) Schumann D, Mueller HJ, Naumann A. *Lebigs Ann. Chem.* 1982:1700–1705.(g) Wenkert E, Broka CA. *J. Chem. Soc. Chem. Commun.* 1984:714–715.(h) Kraus GA, Hon YS. *Heterocycles.* 1987; 25:377–386.(i) Qian L. *Tetrahedron Lett.* 1989; 30:2089–2090.(g) Padwa A, Brodney MA, Marino JP Jr, Sheehan SM. *J. Org. Chem.* 1997; 62:78–87. [PubMed: 11671366] (j) Wu B, Bai D. *J. Org. Chem.* 1997; 62:5978–5981.(k) Grieco PA, Dai Y. *J. Am. Chem. Soc.* 1998; 120:5128–5129.(k) Mori M, Hori K, Akashi M, Hori M, Sato Y, Nishida M. *Angew. Chem Int. Ed.* 1998; 37:637–638.(l) Evans DA, Scheerer JR. *Angew. Chem Int. Ed.* 2005; 44:6038–6042.(m) Bisai SP, West R, Sarpong. *J. Am. Chem. Soc.* 2008; 130:7222–7223. [PubMed: 18479095] (n) West SP, Bisai A, Lim AD, Narayan RR, Sarpong R. *J. Am. Chem. Soc.* 2009; 131:11187–11194. [PubMed: 19591469] (o) Fisher DF, Sarpong R. *J. Am. Chem. Soc.* 2010; 132:5926–5927. [PubMed: 20387895] (p) Laemmerhold KM, Breit B. *Angew. Chem. Int. Ed.* 2010; 49:2367–2370.(q) Tsukano C, Zhao L, Takemoto Y, Hirama M. *Eur. J. Org. Chem.* 2010:4198–4200.(r) Liau BB, Shair MD. *J. Am. Chem. Soc.* 2010; 132:9594–9595. [PubMed: 20568766] (s) Lin H-Y, Snider BB. *Org. Lett.* 2011; 13:1234–1237. [PubMed: 21271726] (t) Nakahara K, Hirano K, Maehata R, Kita Y, Fujioka H. *Org. Lett.* 2011; 13:2015–2017. [PubMed: 21388171] (u) Lin H-Y, Causey R, Garcia GE, Snider BB. *J. Org. Chem.* 2012; 77:7143–7156. [PubMed: 22443298]
3. (a) Carlson EC, Rathbone LK, Yang H, Collett ND, Carter RG. *J. Org. Chem.* 2008; 73:5155–5158. [PubMed: 18529081] (b) Yang H, Carter RG, Zakharov LN. *J. Am. Chem. Soc.* 2008; 130:9238–9239. [PubMed: 18582046] (c) Yang H, Carter RG. *J. Org. Chem.* 2010; 75:4929–4938. [PubMed: 20586477] (d) Collett ND, Carter RG. *Org. Lett.* 2011; 13:4144–4147. [PubMed: 21749086] (e) Veerasamy N, Carlson EC, Carter RG. *Org. Lett.* 2012; 14:1596–1599. [PubMed: 22372610]
4. Muñoz O, Castillo M. *Heterocycles.* 1982; 19:2287–2290.
5. (a) Castillo M, Morales G, Loyola LA, Singh I, Calvo C, Holland HL, MacLean DB. *Can. J. Chem.* 1976; 54:2900–2908.(b) Morales G, Loyola LA, Castillo M. *Phytochemistry.* 1979; 18:1719–1720.(c) Garland MT, Muñoz O. *J. Appl. Cryst.* 1982; 15:112–114.(d) Manríquez V, Quintana R, Muñoz O, Castillo M, von Schnering HG, Peters K. *Acta Cryst.* 1988; C44:165–166.(e) Muñoz OM, Castillo M, San Feliciano A. *J. Nat. Prod.* 1990; 53:200–203.(f) Muñoz, OM.; M, S. Thesis. Universidad de Chile; 1981. (g) Muñoz OM. private communication.
6. Jin C, Ramirez RD, Gopalan AS. *Tetrahedron Lett.* 2001; 42:4747–4750.
7. (a) Schaus SE, Brandes BD, Larrow JF, Tokunaga M, Hansen KB, Gould AE, Furrow ME, Jacobsen EN. *J. Am. Chem. Soc.* 2002; 124:1307–1315. [PubMed: 11841300] (b) Tokunaga M, Larrow JF, Kakiuchi F, Jacobsen EN. *Science.* 1997; 277:936–938. [PubMed: 9252321]
8. Yang H, Carter RG. *Org. Lett.* 2008; 10:4649–4652. [PubMed: 18808137]
9. It was imperative that this reaction be quenched with saturated aqueous Rochelle's salt for an extended period in order to fully break-up the aluminium complex. We attribute this reactivity to the presence of the neighbouring basic nitrogen. In support of this hypothesis, this product **26** readily forms the DCl salt upon standing in CDCl₃.

Figure 1.
Lycopodine-type *Lycopodium* Alkaloids

Scheme 1.
Retrosynthetic Analysis.

Scheme 2.
Organocatalyzed Intramolecular Michael Reaction.

Scheme 3.
Synthesis of the Common Intermediate

Scheme 4.
Synthesis of the Tetracycline

Scheme 5.

Total Syntheses of 10-Hydroxylycopodine, Deacetylpaniculine & Paniculine