Published in final edited form as:

J Int Neuropsychol Soc. 2011 January ; 17(1): 14–23. doi:10.1017/S1355617710001177.

Evaluation of Specific Executive Functioning Skills and the Processes Underlying Executive Control in Schizophrenia

Gauri N. Savla¹, Elizabeth W. Twamley^{1,2}, Wesley K. Thompson^{1,3}, Dean C. Delis¹, Dilip V. Jeste^{1,3}, and Barton W. Palmer¹

¹Department of Psychiatry, University of California, San Diego, La Jolla, California

²Center of Excellence for Stress and Mental Health, Psychiatry Service, VA San Diego Healthcare System, San Diego, California

³The Sam and Rose Stein Center for Research on Aging, University of California School of Medicine, La Jolla, California

Abstract

Schizophrenia is associated with executive dysfunction. Yet, the degree to which executive functions are impaired differentially, or above and beyond underlying basic cognitive processes is less clear. Participants included 145 matched pairs of individuals with schizophrenia (SCs) and normal comparison subjects (NCs). Executive functions were assessed with 10 tasks of the Delis-Kaplan Executive Function System (D-KEFS), in terms of “achievement scores” reflecting overall performance on the task. Five of these tasks (all measuring executive control) were further examined in terms of their basic component (e.g., processing speed) scores and contrast scores (reflecting residual higher order skills adjusted for basic component skills). Group differences were examined via multivariate analysis of variance. SCs had worse performance than NCs on all achievement scores, but the greatest SC-NC difference was that for the Trails Switching task. SCs also had worse performance than NCs on all basic component skills. Of the executive control tasks, only Trails Switching continued to be impaired after accounting for impairments in underlying basic component skills. Much of the impairment in executive functions in schizophrenia may reflect the underlying component skills rather than higher-order functions. However, the results from one task suggest that there might be additional impairment in some aspects of executive control.

Keywords

Cognition; Executive function; Schizoaffective disorder; Psychotic disorders; Trail Making Test; D-KEFS

INTRODUCTION

Executive functions are often a key focus in contemporary neurocognitive models of schizophrenia, due to both the early recognition of executive functioning deficits in patients with the illness (Kraepelin, 1913/1971) and their clear relevance to independent and social functioning (Green, 1996; Palmer & Savla, 2007; Velligan & Bow-Thomas, 1999). Executive functions, historically referred to as “higher mental functions,” are considered to

be “uniquely human” (Luria, 1966), and have been extensively studied in schizophrenia. Early documentations of the striking “concreteness” of people with schizophrenia (see Zec, 1995, for review) were followed by numerous empirical studies exploring tasks measuring abstract thinking and logical reasoning among people with the illness (e.g., Cameron, 1939; Goldstein, 1959). However, a second sense of the construct of executive functions is that of executive control, defined as “the ability to flexibly and dynamically adjust one’s performance to changing environmental demands and internal goal states” (Barch, Braver, Carter, Poldrack, & Robbins, 2009).

Some studies indicate that executive functions may be more impaired than other cognitive functions in schizophrenia (e.g., Kremen, Seidman, Faraone, & Tsuang, 2001), but others suggest that impairment in executive functions may be less than that associated with attention and verbal learning (Atbasoglu, Ozguven, Saka, & Olmez, 2005). This inconsistency may partially reflect the substantial heterogeneity among schizophrenia patients (cf., Braff et al., 1991), but key conceptual and methodological issues also make consensus difficult. Specifically, the construct *executive functions* comprises several skills, which are each measured by multiple neuropsychological tests; neither the set of skills nor the ways in which they are measured has been consistent across studies. Whereas some executive skills such as abstraction (vs. concreteness) have no immediate connotation of self-regulation, those such as executive control emphasize the cybernetic (i.e., pilot or *governor*) aspects of executive function, and their interaction with more basic (non-executive) processes (Royall et al., 2002). Furthermore, the skills that typically make up the executive functions domain are hierarchical in nature, building upon more basic cognitive skills in other domains, such as attention or processing speed, which may themselves be impaired among people with schizophrenia (cf., Dickinson, Ramsey, & Gold, 2007). Because each specific executive function and other (non-executive) cognitive ability likely reflects at least partially distinct brain systems or circuits (Lichter & Cummings, 2001), it is important to disentangle the effects of schizophrenia on various types of executive functions, as well as to distinguish deficits in these functions from the more basic cognitive processes upon which they draw. The lack of consensus in the definition of the construct, compounded by the lack of psychometric equivalence of commonly used tests makes it difficult to draw firm conclusions about the degree of differential deficits and interrelationships among specific executive functions, as well as those between executive and other cognitive abilities (Chapman & Chapman, 1973).

A partial solution to the above confounds is provided by the Delis-Kaplan Executive Function System (D-KEFS; Delis, Kaplan, & Kramer, 2001). The D-KEFS is a battery of nine psychometrically comparable tests (and several subtests) designed to measure multiple types of executive functions. The tests were co-normed on a large national standardization sample, which allows for a head-to-head comparison of the tests. In addition to the achievement scores, which indicate overall performance on each of the D-KEFS tasks, a subset (specifically, those measuring executive control) also include contrast scores, which reflect the residual higher-order cognitive skills after adjusting for performance on lower-level or basic component skills. This allows for a more direct discernment of the extent to which impairment on a task intended as a measure of an executive function may reflect impairment on the more basic (non-executive) cognitive skill upon which the executive function also draws. For example, the achievement score of the Number-Letter Switching task of the D-KEFS Trail Making Test is the total time taken to complete the task. Successful completion of the task, however, requires both basic component skills such as visual scanning, processing speed, and motor speed as well as higher-order skills such as executive control or set-switching. The contrast scores of this task would reflect the residual of the achievement score after controlling for each of these basic component skills.

To our knowledge, there have been no prior published studies employing the full D-KEFS, or any other co-normed battery of executive tests to examine a range of specific executive functions in schizophrenia with the goal of determining differential deficits (if any) among these functions. Thus, in the present study we evaluated the types and levels of executive functioning impairments among persons with schizophrenia or schizoaffective disorder (SC) relative to normal comparison (NC) participants, individually matched on demographic variables and crystallized verbal abilities. Our most general hypothesis was that, relative to NCs, the SCs would have worse mean performance on the selected total achievement scores on D-KEFS. We also examined whether the SC-NC differences on the executive function scores differed among the types of executive function scores, that is, whether there was a differential degree of schizophrenia-related impairment among the specific executive functions. Given the lack of prior studies comparing types of executive functions in schizophrenia, these were intended as exploratory analyses. In addition to the above, we also conducted a comparison of the NCs and SCs on the contrast scores of the five executive control tasks, so that we could test the hypothesis that schizophrenia-related deficits in executive control are non-specific, that is, deficits in higher-order skills can be fully accounted for by the degree of deficit in the underlying basic skills. Finally, given the expected heterogeneity among people with schizophrenia, we also conducted *post hoc* correlational analyses to explore the relationships between clinical variables and the executive functioning tasks among our schizophrenia sample.

METHOD

Participants

Participants included 145 community-dwelling individuals with schizophrenia or schizoaffective disorder and 145 normal comparison participants with no neurological or psychiatric diagnosis. Data for the schizophrenia sample were partly obtained from two ongoing studies at the University of California, San Diego (UCSD; $n=124$), both led by one of the co-authors (E.W.T.), and partly collected prospectively ($n=21$). Patient participants were diagnosed by their treating psychiatrists clinically or with a structured clinical interview. Specifically, diagnoses for the 21 patients whose data were collected prospectively for this study were established with the MINI-International Neuropsychiatric Interview (or MINI; Sheehan, Lecrubier, & Sheehan, 1998). Diagnoses for the other 124 patient subjects were established by their treating clinicians, and then confirmed via chart reviews by formally trained research associates and postdoctoral fellows. Inclusion criteria were (a) Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV) diagnosis of schizophrenia or schizoaffective disorder; (b) age 18 or older at the time of enrollment; and (c) the ability to give written, informed consent to participate in the studies. Exclusion criteria were (a) a concurrent DSM-IV diagnosis of dementia, delirium, or mental retardation; (b) history of head injury with loss of consciousness > 30 min; and (c) substance abuse or dependence per DSM-IV criteria within 1 month of enrollment.

Since the parent studies included patients with either schizophrenia or schizoaffective disorder, and prior research indicates these two groups to be neurocognitively equivalent (reviewed in Palmer & Savla, 2009), we included individuals with either diagnosis in the patient group. *T* tests and chi-square tests confirmed that there were no demographic (age, education, ethnicity, and gender) or clinical (American National Adult Reading Test [ANART] estimated Verbal IQ, Positive and Negative Syndrome Scale [PANSS], or Hamilton Depression Rating Scale [HAM-D] scores) differences between the schizophrenia and schizoaffective disorder sub-samples. The Wilks' λ omnibus test statistic indicated no significant differences between the two groups, $F(10, 134) = 1.767$, $p = .072$, across the 10 D-KEFS executive functioning variables.

Data for the normal comparison sample ($n = 145$) were obtained from the D-KEFS national standardization sample (total $N = 18$ years of age = 916), selected based on propensity score matching to maximize demographic similarity to the SC group. (The method by which we calculated propensity scores and used them to match the samples is described in detail in the data analyses section.)

The parent studies as well as the current study were approved by the UCSD Human Research Protections Program and all participants gave written consent for use of their data.

Each participant's age, highest level of education, gender, ethnicity, and living situation were recorded. For the patient sample, age of onset of illness, duration of illness, and medication information (current medication type, i.e., conventional, atypical, or both) were recorded.

Assessments

Psychopathology—Severity of positive symptoms (e.g., hallucinations, delusions) and negative symptoms (e.g., blunted affect, alogia, apathy) were assessed using the PANSS (Kay, Opler, & Fiszbein, 1987). Severity of depressive symptoms was measured with the 17-item HAM-D (Hamilton, 1967).

Crystallized verbal functioning—As a measure of crystallized verbal knowledge, reading scores are widely used to estimate premorbid Verbal IQ (e.g., Baade & Schoenberg, 2004); thus, using a previously validated and published formula (Grober & Sliwinski, 1991), we estimated each patient's premorbid Verbal IQ based on his or her ANART performance. We did not have ANART data for the NCs; however, we used their Verbal IQ estimates from the Wechsler Abbreviated Scale of Intelligence (computed from WASI verbal subtests, i.e., Vocabulary and Similarities; Wechsler, 1999) as a comparable verbal ability measure.

Executive functioning—The D-KEFS (Delis et al., 2001) was designed to incorporate principles of the *process approach* to neuropsychological assessment whereby basic processes required for successful performance of the higher order tasks can be assessed separately and then built upon over successive conditions. For example, to ensure that any observed deficits on a mental flexibility task such as the Number-Letter Switching condition of the Trail Making Test (analogous to the widely used Trail Making Test, Part B; Reitan & Wolfson, 1993) are not attributable to more basic skills, those basic skills (e.g., visual scanning, number sequencing, letter sequencing, and motor speed) are also assessed separately.

Participants in the present study completed the full D-KEFS, but our analyses were focused on 10 higher-order conditions across the nine tests. In particular, we used the following tasks:

1. Trial Making (Switching, total correct): Described above.
2. Verbal Fluency (Category Switching, total correct): Participants are asked to rapidly name words from two categories, alternating between the two categories with each successive response.
3. Design Fluency (Switching, total correct): Participants are given 60 s to connect pre-printed filled or unfilled dots with a set number of lines to generate as many different designs as they can think of. In the switching condition, they must alternate the connections between filled and unfilled dots.
4. Color-Word Interference (Inhibition and Inhibition/Switching, total correct): The former is analogous to the Color-Word Interference trial of the Stroop task (Golden

& Freshwater, 2002). In the latter, some of the color words are printed inside small boxes, others are not. Participants are required to read the word for boxed words, and to name the color of ink for unboxed words.

5. Sorting (Free Sorting, confirmed correct). Participants are given six cards that vary across several dimensions (such as shape, number of sides, color, content, and so forth), which they must sort into different piles multiple times, using as many different sorting strategies as they can think of.
6. Twenty Questions (total achievement score): Participants are shown a large card with a variety of objects on it, and they must use a series of yes or no questions to determine the target object in as few questions as possible.
7. Word Context (total consecutively correct): Participants are shown a sequence of five sentences with one non-word (but with a pronounceable, plausible combination of consonants and vowels) embedded within each sentence. The participant must use the context of the word within each sentence to determine the word's meaning.
8. Tower (total achievement score): Analogous to the widely used Tower of London/Hanoi tasks (Sullivan, Riccio, & Castillo, 2009).
9. Proverb (Free Inquiry, total achievement score): Analogous to the proverb questions on the Comprehension subtest of the Wechsler intelligence scales (c.f., Wechsler, 2008).

For each of these tasks, we examined achievement scores reflecting overall performance. We further examined Trails Switching, Category Switching, Design Fluency Switching, Color-Word Inhibition, and Color-Word Inhibition/Switching (all tasks measuring executive control) in terms of their underlying processes, that is, basic component skills such as processing speed, visual scanning, and motor speed) and contrast scores (residual scores after adjusting for "basic" component skills). Basic component skills included Number Sequencing, Category Fluency, Combined Filled and Empty Dots, Color Naming, and Color-Word Inhibition, respectively, for each of the five executive control tasks, and contrast scores were calculated by subtracting achievement scores from scores on the basic component skills for each task. Age-corrected scaled scores for the achievement scores, basic component task scores, and contrast scores based on the D-KEFS normative sample were used for analyses.

Analyses

Propensity score calculation and matching—We computed propensity scores (probability of being in the SC group; Rosenbaum & Rubin, 1983) based on variables deemed *a priori* likely to confound executive functioning performance: age, education, ethnicity, and estimated premorbid Verbal IQ. Propensity score matching is a technique that attempts to reduce bias in estimates of group effects due to unequal distribution of confounding variables; subjects with the same propensity score also have the same probability of belonging to either the SC or comparison group; matching pairs of subjects with similar propensity scores can therefore substantially reduce covariate imbalances across groups. We used logistic regression with SC group status as the dependent variable and the *a priori* chosen confounders as independent variables, along with all pairwise interactions of these to calculate these scores. The scores were then used to create 145 matched pairs of SC and NC subjects using a propensity score matching program (Painter, 2010) that uses syntax and macros in PASW (Predictive Analytics SoftWare, version 18.0). We tested the resulting SC and NC samples for equivalence on the potential confounding variables after pair matching.

Analyses of target variables—The distributions of all variables were checked for normality; none of the variables required transformation to meet assumptions for parametric analyses. We examined the differences between the NC and SC participants on the 10 executive functioning tasks via a between-subjects (1 factor [diagnostic status], 2-level [NC vs. SC]) multivariate analysis of variance (MANOVA) to account for correlation among the variables. Following a significant omnibus test yielded by the MANOVA, we conducted *post hoc* analyses of variance (ANOVAs) between the NC and SC groups to determine group differences between the individual executive functioning tasks. We used .005 as the Bonferroni-adjusted level for significant differences. We used Cohen's benchmarks (Cohen, 1988) to categorize the magnitude of the effect sizes (partial η^2 s; .01 = *small*, .06 = *medium*, and .14 = *large*).

Next, we compared the relative strength of differences between the NC and SC groups on the 10 executive functioning tasks. A difference score was computed on each of the 10 tasks, for each of the 145 NC-SC pairs. The resulting difference scores were bootstrapped using 10,000 resamples with replacement. Mean values and 95% confidence intervals were extracted from the resulting bootstrap distributions. These were Bonferroni-adjusted using $\alpha = .05/10 = .005$ as the cutoff level of significance testing difference scores against zero. Finally, we computed all 45 pairwise comparisons of the 10 executive functioning tasks, computing means and 95% confidence intervals from the bootstrap distributions. These 95% confidence intervals were Bonferroni-adjusted using $\alpha = .05/45 = .001$.

We conducted separate MANOVAs to examine differences between the NCs and SCs on basic component skills and contrast scores (higher-order/multilevel processes isolated from these basic processes). The MANOVAs were followed by *post hoc* univariate ANOVAs after significant omnibus tests.

Relationships with clinical variables—For the SC sample only, we conducted bivariate correlational analyses to examine relationships, if any between clinical variables (positive and negative symptoms [PANSS], and depressive symptoms [HAM-D]) and the D-KEFS tasks (both total achievement scores and contrast scores).

RESULTS

As shown in Table 1, the NC and SC paired samples did not differ on mean age, education, and estimated premorbid Verbal IQ. Relative to the SC group, the NC group had significantly more Caucasians ($p = .002$) and women ($p = .003$). As a group, the SC participants were clinically stable, with mild psychopathology.

Using Wilks' λ as the omnibus test statistic, the combined D-KEFS achievement scores resulted in a significant main (and *large*) effect of diagnostic group (NC vs. SC), $F(10, 279) = 16.38$, $p < .001$, partial $\eta^2 = .370$. Follow-up univariate ANOVAs revealed significant differences between the NC and SC groups on all 10 D-KEFS achievement scores, with SCs consistently performing worse than NC subjects (Table 2). Effect sizes for the individual achievement scores ranged from medium (partial $\eta^2 = .098$ and .099 for Verbal Fluency Switching and Sorting, respectively) to large (partial $\eta^2 = .268$ and .193 for Trails Switching and Color-Word Inhibition, respectively).

Results of the ranking and bootstrapping method to statistically examine the relative magnitude of differences between the NC and SC groups on the executive functioning tasks are presented in Table 3. The SC-NC difference for Trails Switching was significantly greater than differences for Verbal Fluency, Design Fluency, Color-Word Inhibition/Switching, Twenty Questions, Tower, and Proverb tests. Color-Word Inhibition was

differentially more impaired than Color-Word Inhibition/Switching and Word Context tests. The magnitudes of SC-NC differences among the other D-KEFS tasks were not statistically significant.

Separate follow-up MANOVAs for the basic component skills and contrast scores were conducted to evaluate the levels of impairment underlying the achievement scores on five D-KEFS executive functioning tasks. As shown in Table 4, using Wilks' λ as the omnibus test statistic, the combined basic component skills resulted in a significant and large main effect for diagnostic group (SC vs. NC), $F(5, 284) = 26.02$, $p < .001$, partial $\eta^2 = .314$, indicating a large effect size of diagnostic group on performance. Follow-up ANOVAs indicated that SCs did consistently worse than did the NCs on all basic component skills (all p s $< .001$), with large effect sizes, except for Trails Number Sequencing (medium effect size).

As shown in Table 5, using Wilks' λ as the omnibus test statistic, the combined contrast scores resulted in a significant large main effect of diagnostic group on performance (SC vs. NC), $F(5, 284) = 9.43$, $p < .001$, partial $\eta^2 = .142$. Follow-up univariate ANOVAs indicated SCs performed significantly worse than NCs on the Trails Switching contrast score (partial $\eta^2 = .101$, $p < .001$), with diagnostic group having a large effect on switching performance. However, SCs performed marginally (but at statistically significant level) *better* than the NCs on Color-Word Inhibition/Switching contrast score (partial $\eta^2 = .031$, $p = .003$). These differences continued to be significant after Bonferroni adjustments for multiple comparisons, that is, $.05/5 = .01$. There were no differences between the SCs and NCs on the other tasks.

Bivariate correlations revealed small effect size but statistically significant associations between severity of positive symptoms and the total achievement scores on the Trails Number-Letter Switching, Verbal Fluency, and Color-Word Inhibition tasks ($r = -.230$, $-.261$, and $.188$, respectively; all p s $> .026$), between severity of negative symptoms and the Word-Context task achievement score ($r = .171$, $p = .042$), and between severity of depressive symptoms and the achievement scores on Trails Number-Letter Switching and Color-Word Inhibition tasks ($r = -.234$ and $-.239$, respectively, both p s $= .005$). There were no other statistically significant correlations between severity of symptoms and D-KEFS achievement scores, nor between severity of symptoms and any of the D-KEFS contrast scores.

DISCUSSION

In this study, we examined executive impairment in schizophrenia in terms of three categories of scores: (1) the primary achievement scores on executive functioning tasks, (2) scores on basic component skills, and (3) contrast scores, that is, residual executive control skills isolated from basic component skills. Consistent with our *a priori* hypotheses, SC patients had worse functioning than did NCs on the achievement scores on all executive function tasks and all basic skills. They had significantly greater impairments on the Trail Making Switching task than on any other D-KEFS executive functioning task. The SC group continued to do poorly on Trails Switching even after accounting for processing speed (on which they were also significantly impaired). However, there were no group differences on the other executive control tasks (i.e., Category Fluency Switching, Design Fluency Switching, and on Color-Word Inhibition, after accounting for basic skills (which were all impaired among the SCs). The findings from the contrast scores suggest that the difficulties on switching/cognitive flexibility commonly observed among people with schizophrenia or schizoaffective disorder at least partially reflect impairment in more basic or component skills also measured by these tasks, such as processing or psychomotor speed (Dickinson et al., 2007).

Contrary to our expectations, the SCs in fact did slightly *better* than did NCs on the isolated Color-Word Inhibition/Switching component, that is, once the score was adjusted for performance on the more basic Color Word Inhibition task. This finding is counter-intuitive as we can think of no model of the cognitive deficits associated with schizophrenia that would lead one to predict a difference in this direction. It may best serve as a reminder of the complex challenges involved in deconstructing higher-order cognitive processes, and the need for replication before drawing definitive conclusions about any particular cognitive deficit or strength among the entire population of individuals with schizophrenia.

Symptoms of psychopathology among the SCs, (particularly severity of positive and depressive symptoms) were significantly related with some achievement scores of some of the executive functioning tasks, but with not with any of the contrast scores. This suggests that psychopathology was likely related to the more basic component processes involved in those tasks, but not the switching processes. In contrast, there were no significant relationships among negative symptoms and any executive functions (with the exception of Word Context, wherein the relationship was small). Our sample, on average, had minimal psychopathology with little variance in scores, and the small magnitude of relationships (when present) was expected, but these findings may not fully generalize to SC patients in acutely psychotic states and/or to those in long-term institutional settings.

With regard to the more basic cognitive processes, it is hardly surprising to find that schizophrenia diagnosis had large effects on fluency deficits; both verbal and spatial fluency were significantly impaired among the SCs and this is consistent with previous findings (e.g., Moore et al., 2006). It has been previously suggested that people with schizophrenia tend to show greater-than-normal speed when naming the ink color in the Color Naming task on the Stroop test (reviewed by Perlstein, Carter, Barch, & Baird, 1998). This was not found to be the case with our current sample of SCs, whose performance on color naming was significantly impaired. Surprisingly, their performance on the response inhibition component of the Color-Word task was equivalent to that among the NCs, indicating that they seemed to have the most difficulty with processing speed, even on relatively simple tasks such as color naming. We did not, however, have data on whether SCs made more or equivalent numbers of errors on the Color-Word tasks—it would have been interesting to determine whether SCs made a speed/accuracy tradeoff on these tasks, in a qualitatively different way than did NCs.

Notably, the only executive control task that SCs did worse on than NCs, after accounting for processing speed, was Trails Switching. This task may be a particularly good measure of overall cerebral integrity in schizophrenia; it was designed to measure switching ability primarily, but also measures inhibition (by incorporating “capture stimuli” or close foils) and other processes measured by the standard Trail Making Test, Part B, including attention, motor and processing speed, sequencing, and working memory. Working memory is especially overloaded in this task, involving not only switching but also consecutively maintaining the sequence of numbers and letters and inhibiting automatic responses (i.e., avoiding capture stimuli). This finding is supported by the extensive literature on Trail Making Test Part B versus Part A, which suggests that patients consistently do worse on Trails B (Braff et al., 1991; Saykin et al., 1991).

The results of our study are particularly interesting because (a) people with schizophrenia, even those with relatively higher levels of educational achievement and premorbid verbal functioning, had difficulties with executive functioning, (b) an examination of underlying processes indicates that the primary difficulty may be with processing speed, although complex, multi-level skills which demand the simultaneous processing of a set of rules may also be difficult for people with schizophrenia, and (c) the Number-Letter Switching

component of the Trail Making Test, of all executive control tasks, may be differentially impaired among people with schizophrenia.

To our knowledge, this is the first study to comprehensively characterize executive functioning abilities in schizophrenia using a large battery of co-normed measures including the standard multilevel tasks and measures of the basic abilities underlying each task. Executive functions have never been previously examined in this manner among people with schizophrenia, despite suggestions in the literature that they may be particularly important predictors of everyday functioning (Green, Kern, Braff, & Mintz, 2000). Another strength of this study is that we were able to individually match the schizophrenia and normal comparison samples on age, education, ethnicity, and crystallized verbal knowledge, which allowed us to make more plausible conclusions about schizophrenia-related cognitive impairments.

There were some limitations in the present study. This study was largely based on secondary analyses of existing data, resulting in several constraints on the data; the SC and NC data were not collected in parallel, so the possibility of examiner or cohort effects cannot be completely ruled out. However, the D-KEFS was administered according to standardized procedures; also, use of data from the D-KEFS standardization sample permitted one-to-one matching of the SCs and NCs, resulting in a level of comparability not typically achieved in prospectively collected case control samples. Also, the measures estimating crystallized verbal knowledge differed in the SC and NC samples, the former being assessed with the ANART and the latter, with the WASI. There are no published data, to our knowledge, that directly compare the ANART and WASI scores in schizophrenia patients, but using the scores from these two measures permitted a gross matching of participants in terms of overall premorbid verbal functions as part of the propensity score calculations used in matching NC to SC patients.

Our SC sample comprised outpatients with chronic illness (with a mean duration of illness of 25 years), who were relatively stable on their medication, had low levels of psychopathology, and had higher-than-expected estimated crystallized verbal knowledge. It is possible that the present pattern of results may not readily generalize to individuals with markedly different demographic or clinical characteristics. On the other hand such characteristics have not generally shown strong associations with level or pattern of cognitive deficits among people with schizophrenia (reviewed in Palmer, Dawes, & Heaton, 2009).

Our SC sample also consisted of a large proportion of patients with schizoaffective disorder, which could be considered by some to be a separate comparison group with schizophrenia, given its mood component. However, a review of the literature [including a prior study from our group (Evans et al., 1999)] suggests that the two subgroups are neurocognitively equivalent (Palmer & Savla, 2009). Thus, it appears unlikely that the present pattern of results is affected by the proportion of patients with schizoaffective disorder.

Although the D-KEFS has many strengths, it is also limited in the scope of executive functions and their underlying processes it is able to measure. The broader construct of executive functions embodies such socially-mediated processes as environmental monitoring, goal-setting, foresight and planning, and self-regulation based on internal and external states (Barkley, 2001). Also, some of the D-KEFS tests (i.e., those measuring abstraction, logical reasoning, or planning aspects of executive functions) did not allow for the deconstruction of component processes. Unlike the executive control tasks, wherein higher-order tasks naturally draw on lower-level cognitive processes, there is no such

analogue to these tasks. An examination of the processes underlying impairments in these functions would be a useful complement to those evaluated in the present study.

The above limitations noted, however, the current findings nonetheless have some clear theoretical and pragmatic implications. Executive functions are universally acknowledged as vital to optimal daily functioning, yet have so far remained a nebulous and poorly defined construct. On a theoretical level, the identification of specific types of executive functioning skills contributes to overall efforts to identify meaningful neurocognitive subtypes, and to link cognitive deficits to the underlying pathology in neurological systems or processes. The present findings suggest value in distinguishing between achievement scores and underlying process scores, especially in regard to executive functioning tasks. On a clinical or pragmatic level, this study sheds light on the importance of including a wide array of executive functioning measures in a comprehensive neuropsychological assessment in schizophrenia, and considering the relative strengths and weaknesses within the executive functioning domain to identify targets for rehabilitation/vocational planning for patients with schizophrenia.

Despite more than a century of research, little is known about schizophrenia's effect on the brain, partly because of the heterogeneity among individuals with illness, and partly because of lack of adequate methodology to study its complex presentation. Future directions include studies to examine the relationships between the specific executive functions and domains of social cognition, as well as real life functioning among patients with schizophrenia, in areas such as medication adherence, decision making capacity, driving ability, academic/vocational functioning, and other instrumental activities of daily living to better determine the types of strategies needed to rehabilitate patients with the illness. This study also suggests the potential role of both structural and functional neuroimaging studies in shedding light on frontal-subcortical neural pathways that mediate executive (as well as other cognitive) functions, and differential deficits and brain abnormalities associated with schizophrenia.

Acknowledgments

Gauri N. Savla, Department of Psychiatry, University of California, San Diego; Elizabeth W. Twamley, Department of Psychiatry, University of California, San Diego, Center of Excellence for Stress and Mental Health, Psychiatry Service, VA San Diego Healthcare System; Wesley K. Thompson, Department of Psychiatry, University of California, San Diego, The Sam and Rose Stein Center for Research on Aging, University of California, San Diego School of Medicine; Dean C. Delis, Department of Psychiatry, University of California, San Diego; Dilip V. Jeste, Department of Psychiatry, University of California, San Diego, The Sam and Rose Stein Center for Research on Aging, University of California, San Diego School of Medicine; Barton W. Palmer, Department of Psychiatry, University of California, San Diego.

This study was conducted as part of Dr. Savla's dissertation at the San Diego State University/University of California, San Diego Joint Doctoral Program in Clinical Psychology.

This work was supported by the National Institutes of Health/National Institute of Mental Health (grant numbers MH19934, MH064722, and MH080002) and a grant from the National Alliance for Research on Schizophrenia and Depression.

REFERENCES

- Atbasoglu EC, Ozguven HD, Saka MC, Olmez S. Relative sparing of executive functions in the early phase of schizophrenia. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 2005; 17:510–516. [PubMed: 16387991]
- Baade LE, Schoenberg MR. A proposed method to estimate premorbid intelligence utilizing group achievement measures from school records. *Archives of Clinical Neuropsychology*. 2004; 19:227–243. [PubMed: 15010088]

- Barch DM, Braver TS, Carter CS, Poldrack RA, Robbins TW. CNTRICS final task selection: Executive control. *Schizophrenia Bulletin*. 2009; 35:115–135. [PubMed: 19011235]
- Barkley RA. The executive functions and self-regulation: An evolutionary neuropsychological perspective. *Neuropsychology Review*. 2001; 11:1–28. [PubMed: 11392560]
- Braff DL, Heaton R, Kuck J, Cullum M, Moranville J, Grant I, Zisook S. The generalized pattern of neuropsychological deficits in outpatients with chronic schizophrenia with heterogeneous Wisconsin Card Sorting Test Results. *Archives of General Psychiatry*. 1991; 48:891–898. [PubMed: 1929758]
- Cameron N. Schizophrenic thinking in a problem-solving situation. *Journal of Mental Science*. 1939; 85:1012–1035.
- Chapman LJ, Chapman JP. Problems in the measurement of cognitive deficits. *Psychological Bulletin*. 1973; 79:380–385. [PubMed: 4707457]
- Cohen, J. Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc; 1988.
- Delis, DC.; Kaplan, E.; Kramer, JH. Delis-Kaplan Executive Function Scale (D-KEFS): Examiner's manual. San Antonio, TX: The Psychological Corporation; 2001.
- Dickinson D, Ramsey ME, Gold JM. Overlooking the obvious: A meta-analytic comparison of Digit Symbol Coding tasks and other cognitive measures in schizophrenia. *Archives of General Psychiatry*. 2007; 64:532–542. [PubMed: 17485605]
- Evans JD, Heaton RK, Paulsen JS, McAdams LA, Heaton SC, Jeste DV. Schizoaffective disorder: A form of schizophrenia or affective disorder? *Journal of Clinical Psychiatry*. 1999; 60:874–882. [PubMed: 10665641]
- Golden, CJ.; Freshwater, SM. Stroop color and word test: A manual for clinical and experimental uses. 2nd ed.. IL: Stoelting Co; 2002.
- Goldstein K. Concerning the concreteness in schizophrenia. *Journal of Abnormal and Social Psychology*. 1959; 59:146–148.
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *American Journal of Psychiatry*. 1996; 153:321–330. [PubMed: 8610818]
- Green MF, Kern RS, Braff DL, Mintz J. Neurocognitive deficits and functional outcome in schizophrenia: Are we measuring the “right stuff”? *Schizophrenia Bulletin*. 2000; 26:119–136. [PubMed: 10755673]
- Grober E, Sliwinski M. Development and validation of a model for estimating premorbid verbal intelligence in the elderly. *Journal of Clinical and Experimental Neuropsychology*. 1991; 13:933–949. [PubMed: 1779032]
- Hamilton M. Development of a rating scale for primary depressive illness. *The British Journal of Social and Clinical Psychology*. 1967; 6:278–296. [PubMed: 6080235]
- Kay, SR.; Opler, LA.; Fiszbein, A. Positive and Negative Syndrome Scale (PANSS) rating manual. San Rafael, CA: Social and Behavioral Sciences Documents; 1987.
- Kraepelin, E. Dementia praecox and paraphrenia. Huntington, NY: Krieger; [Translated in 1919 from the eighth German edition of the ‘Test-book of Psychiatry,’ vol. iii, part ii, section on Endogenous Dementias published in 1913]
- Kremen WS, Seidman LJ, Faraone SV, Tsuang MT. Intelligence quotient and neuropsychological profiles in patients with schizophrenia and in normal volunteers. *Biological Psychiatry*. 2001; 50:453–462. [PubMed: 11566163]
- Lichter, DG.; Cummings, JL. Introduction and overview. Frontal-subcortical circuits in psychiatric and neurological disorders. Lichter, DG.; Cummings, JL., editors. New York: Guilford Press; 2001. p. 1-43.
- Luria, AR. Higher cortical functions in man. New York: Basic Books; 1966. [B. Haigh Trans]
- Moore DJ, Savla GN, Woods SP, Jeste DV, Palmer BW. Verbal fluency impairments among middle-aged and older outpatients with schizophrenia are characterized by deficient switching. *Schizophrenia Research*. 2006; 87:254–260. [PubMed: 16854567]
- Painter JS. SPSS propensity matching program description. 2010 Retrieved from www.unc.edu/~painter.

- Palmer BW, Dawes SE, Heaton RK. What do we know about the neuropsychological aspects of schizophrenia? *Neuropsychology Review*. 2009; 19:365–384. [PubMed: 19639412]
- Palmer BW, Savla GN. The association of specific neuropsychological deficits on capacity to consent to research or treatment. *Journal of the International Neuropsychological Society*. 2007; 13:1047–1059. [PubMed: 17942022]
- Palmer, BW.; Savla, GN. Neuropsychological deficits in schizoaffective disorder. In: Yip, KS., editor. *Schizoaffective disorders: International perspectives on understanding, intervention and rehabilitation*. New York: Nova Science; 2009.
- Perlstein WM, Carter CS, Barch DM, Baird JW. The Stroop task and attention deficits in schizophrenia: A critical evaluation of card and single-trial Stroop methodologies. *Neuropsychology*. 1998; 12:414–425. [PubMed: 9673997]
- Reitan, RM.; Wolfson, D. *The Halstead-Reitan neuropsychological test battery: Theory and clinical interpretation*. 2nd ed. Tucson, AZ: Neuropsychology Press; 1993.
- Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika*. 1983; 70:41–55.
- Royall DR, Lauterback EC, Cummings JL, Reeve A, Rummans TA, Kaufer DI, Coffey CE. Executive control function: A review of its promise and challenges for clinical research: A report from the Committee on Research of the American Neuropsychiatric Association. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 2002; 14:377–405. [PubMed: 12426407]
- Saykin AJ, Gur RC, Gur RE, Mozley PD, Mozley LH, Resnick SM, Stafiniak P. Neuropsychological function in schizophrenia: Selective impairment in memory and learning. *Archives of General Psychiatry*. 1991; 48:618–624. [PubMed: 2069492]
- Sheehan DV, Lecrubier Y, Sheehan KH. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*. 1998; 59:22–33. [PubMed: 9881538]
- Sullivan JR, Riccio CA, Castillo CL. Concurrent validity of the tower tasks as measures of executive function in adults: A meta-analysis. *Applied Neuropsychology*. 2009; 16:62–75. [PubMed: 19205950]
- Velligan DI, Bow-Thomas CC. Executive function in schizophrenia. *Seminars in Clinical Neuropsychiatry*. 1999; 4:24–33. [PubMed: 10229790]
- Wechsler, D. *Wechsler Abbreviated Scale of Intelligence (WASI)*. San Antonio, TX: Psychological Corporation; 1999.
- Wechsler, D. *Wechsler Adult Intelligence Scale-Fourth Edition (WAIS-IV)*. San Antonio, TX: Pearson, Inc; 2008.
- Zec RF. Neuropsychology of schizophrenia according to Kraepelin: Disorders of volition and executive functioning. *European Archives of Psychiatry and Clinical Neuroscience*. 1995; 245:216–223. [PubMed: 7578284]

Table 1

Description of the samples

	Normal comparison participants (NCs) Mean (SD) or %	Schizophrenia participants (SCs) Mean (SD) or %	<i>t</i> -Test or chi-square differences
Age at testing (years)	48.7 (11.7)	48.5 (8.6)	$p = .891$
Education level, % 12 years	84.1%	77.2%	$p = .544$
Gender, % women	53.8%	36.6%	$p = .003$
Ethnicity, % Caucasian	75.9%	54.5%	$p = .002$
Living situation	Not available		Not applicable
% Independently		76.4	
% Board and care facility		16.0	
Verbal IQ estimate (WASI VIQ for NCs; ANART estimated IQ for SCs)	105.9 (9.6)	105.2 (14.4)	$p = .600$
Diagnosis	Not applicable		Not applicable
% Schizophrenia		48.3%	
% Schizoaffective disorder		51.7%	
Age of onset of illness (years), $N = 141$	Not applicable	23.6 (10.03)	Not applicable
Duration of illness (years), $N = 141$	Not applicable	25.0 (12.0)	Not applicable
PANSS positive symptom score, $N = 142$	Not available	15.8 (5.8)	Not applicable
PANSS negative symptom score, $N = 142$	Not available	14.8 (5.0)	Not applicable
HAM-D (17-item total), $N = 135$	Not available	11.9 (6.6)	Not applicable
Type of antipsychotic medication	Not applicable		Not applicable
% Conventional		4.8%	
% Atypical		84.1%	
% Both		4.1%	
% None		6.9%	

Note. Where not indicated, $N = 145$;

ANART = American National Adult Reading Test; WASI = Wechsler Abbreviated Scale of Intelligence; PANSS = Positive and Negative Syndrome Scale; HAM-D = Hamilton Depression Rating Scale; VIQ = Verbal IQ.

Table 2

Group differences on D-KEFS executive functioning tasks

Target variables	Estimated marginal means			Between-subjects effects		
	Mean	SE		F (1, 288)	p	Partial η^2
Trails Number-Letter Switching	NC	10.45	.266	105.55	<.001	.268
	SC	6.58	.266			
Verbal Fluency Switching	NC	9.98	.268	31.16	<.001	.098
	SC	7.72	.268			
Design Fluency Switching	NC	10.670	.255	58.02	<.001	.168
	SC	8.24	.255			
Color-Word Inhibition	NC	10.03	.289	68.72	<.001	.193
	SC	6.64	.289			
Color-Word Inhibition/Switching	NC	10.28	.260	42.93	<.001	.130
	SC	7.87	.260			
Sorting	NC	10.48	.283	31.73	<.001	.099
	SC	8.22	.283			
Twenty Questions	NC	10.90	.280	35.28	<.001	.109
	SC	8.55	.280			
Word Context	NC	10.66	.276	65.16	<.001	.185
	SC	7.50	.276			
Tower	NC	10.59	.263	40.85	<.001	.124
	SC	8.21	.263			
Proverb	NC	10.79	.237	64.60	<.001	.183
	SC	8.09	.237			

Note. Cohen's benchmarks for η^2 : .01 = *small*, .06 = *medium*, .14 = *large*. Variables indicate achievement scores; all are age-scaled scores.

D-KEFS = Delis-Kaplan Executive Function System; NC = normal comparison participants; SC = schizophrenia participants.

Table 3
Mean Schizophrenia-Normal comparison pair differences on D-KEFS executive functioning tasks

Differences in mean values, SC minus NC (Bonferroni-adjusted <i>p</i> values)							
Trails	VF	DF	CW-I	CW-I/S	Sorting	TQ	Tower
Trails							
VF	-2.30 (<.001)						
DF	-1.58 (.018)	-.29 (1)					
CW-I	-0.31 (1)	-1.11 (.117)					
CW-I/S	-1.41 (<.001)	-0.90 (1)	-26 (1)	-1.08 (<.001)			
Sorting	-1.37 (.072)	-0.72 (1)	-.04 (1)	.40 (1)			
TQ	-1.22 (<.001)	-0.32 (1)	-1.24 (1)	-0.02 (1)	.50 (1)		
WC	-0.90 (.207)	-1.24 (.099)	-.4 (1)	.64 (<.001)	-.51 (1)	-0.72 (.783)	
Tower	-1.76 (.045)	.27 (1)	-.09 (1)	-1.45 (.099)	.30 (1)	-.51 (1)	-0.61 (.558)
Proverb	-1.36 (.027)	-.51 (1)	.09 (1)	-.60 (1)	-0.43 (1)	-0.30 (1)	-.19 (1)
							-.90 (.225)
							-.60 (1)

Note. Bolded values denote significant differences between tasks.

Trails = Trails Switching Total Time; VF = Verbal Fluency Category Switching Total Correct; DF = Design Fluency Switching Total Correct; CW-I = Color-Word Inhibition Total Correct; CW-I/S = Color-Word Inhibition/Switching Total Correct; Sorting = Sorting – Confirmed Correct Sorts; TQ = Twenty Questions – Total Achievement; WC = Word Context – Total Consecutively Correct; Tower = Tower – Total Achievement; Proverb = Proverb – Total Achievement.

Table 4

Group differences on basic component skills underlying D-KEFS executive control skills

Cognitive skills	Estimated marginal means		Between-subjects effects		
	Mean	SE	F(1, 288)	p	Partial η^2
Trails Number Sequencing	NC	10.39 .249	23.26	<.001 *	.075
	SC	8.69 .249			
Verbal Fluency Category Fluency	NC	10.32 .271	54.63	<.001 *	.159
	SC	7.49 .271			
Design Fluency Combined Filled and Empty Dots	NC	10.74 .191	69.76	<.001 *	.195
	SC	8.48 .191			
Color-Word—Color Naming	NC	10.22 .264	79.93	<.001 *	.211
	SC	6.95 .264			
Color-Word—Inhibition	NC	10.03 .289	68.72	<.001 *	.193
	SC	6.64 .289			

Note. Age Scaled Scores were used in these analyses. Cohen's benchmarks for partial η^2 : .01 = *small*/but not trivial, .06 = *medium*, .14 = *large*.

* Significant differences after Bonferroni correction for multiple comparisons.

NC = normal comparison participants; SC = schizophrenia participants.

Table 5
Group differences on contrast scores, i.e., executive control skills adjusted for the basic component skills

(Residual) executive control skills	Estimated marginal means			Between-subjects effects		
	Mean	SE		F(1, 288)	p	Partial η^2
Trails Switching – Number Sequencing	NC	9.48	.242	18.71	<.001*	.101
	SC	8.00	.242			
Verbal Fluency Category Switching – Category Fluency	NC	9.57	.279	2.88	.091	.010
	SC	10.23	.279			
Design Fluency Switching – Combined Filled and Empty Dots	NC	9.98	.228	.41	.522	.001
	SC	9.77	.228			
Color-Word Inhibition – Color Naming	NC	9.79	.229	.07	.782	.000
	SC	9.70	.229			
Color-Word Inhibition/Switching – Inhibition	NC	10.26	.247	8.87	.003*	.031
	SC	11.30	.247			

Note. Age-corrected scaled scores were used in these analyses. Cohen's benchmarks for η^2 : .01 = small, .06 = medium, .14 = large.

* Significant differences after Bonferroni correction for multiple comparisons.

NC = normal comparison participants; SC = schizophrenia participants.