Published in final edited form as:

J Addict Med. 2013 ; 7(2): 116–121. doi:10.1097/ADM.0b013e31828230ca.

Pubertal Status, Sensation-Seeking, Impulsivity, and Substance Use in High-School-Aged Boys and Girls

Grace Kong, PhD^a, Anne E. Smith, PhD, Thomas J. McMahon, PhD^{a,b}, Dana A. Cavallo, PhD^a, Ty S. Schepis, PhD^c, Rani A. Desai, PhD, MPH^d, Marc N. Potenza, MD, PhD^{a,b,e}, and Suchitra Krishnan-Sarin, PhD^a

^aDepartment of Psychiatry, Yale University School of Medicine, New Haven, CT 06519

^bChild Study Center, Yale University School of Medicine, New Haven, CT 06519

^cDepartment of Psychology, Texas State University, San Marcos, TX 78666

^dDepartment of Psychiatry, Yale University School of Medicine, VA Connecticut Healthcare System, West Haven, CT 06546

^eDepartment of Neurobiology, Yale University School of Medicine, New Haven, CT 06519

Abstract

Objective—To examine how factors such as pubertal status, sensation-seeking, and impulsivity are related to substance use (cigarettes, alcohol and marijuana) in high-school students and examine these associations by gender.

Methods—Ten public high schools in Connecticut participated in a survey of high-risk behaviors. Adolescents from grades 9 to 12 ($N = 3068$) completed measures of physical development (Pubertal Development Scale), perceived pubertal timing, impulsivity and sensation-seeking (Zuckerman-Kuhlman Personality Questionnaire-Form III), and cigarette, marijuana, and alcohol use in the past 30 days.

Results—Logistic regression analyses modeling each substance use (cigarettes, marijuana, alcohol) and gender separately showed that (1) early perceived pubertal timing was associated with cigarette use but Pubertal Development Scale was not associated with any substance use, (2) sensation-seeking was associated with use of all three substances, (3) gender differences were

---

Corresponding author: Grace Kong, PhD. Department of Psychiatry, Yale University School of Medicine 34 Park Street, Room S-201, New Haven CT 06519. Tel: (203) 974-7601. Fax: (203) 974-7606. grace.kong@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflicts of Interest/Dislosures: The study sponsors did not have any role in the study design, the collection, analysis and interpretation of data, the writing of the manuscript and the decision to submit the manuscript for publication. Dr. Potenza consults for and is an advisor to Boehringer Ingelheim; has consulted for and has financial interests in Somaxon; has received research support from the National Institutes of Health, Veteran's Administration, Mohegan Sun Casino, the National Center for Responsible Gaming and its affiliated Institute for Research on Gambling Disorders, and Forest Laboratories, Ortho-McNeil, Oy-Control/Biotie, Psyadon and Glaxo-SmithKline pharmaceuticals; has participated in surveys, mailings or telephone consultations related to drug addiction, impulse control disorders or other health topics; has consulted for law offices and the federal public defender's office in issues related to impulse control disorders; provides clinical care in the Connecticut Department of Mental Health and Addiction Services Problem Gambling Services Program; has performed grant reviews for the National Institutes of Health and other agencies; has guest-edited journal sections; has given academic lectures in grand rounds, CME events and other clinical or scientific venues; and has generated books or book chapters for publishers of mental health texts. The rest of the authors have no financial conflicts of interest with respect to the content of this manuscript.

detected: impulsivity was associated with use of all three substances among girls, but this was not found among boys.

Conclusion—Future studies should examine the relationship among pubertal status, sensation-seeking, and impulsivity from pre-adolescence through late adolescence to better understand how these relationships to substance use may change with time and growth.

Keywords

puberty; impulsivity; sensation-seeking; substance use; gender

Current epidemiological survey indicates that substance use is problematic among high-school-aged youth; among 12th graders, 22.6% reported using marijuana, 40.0% alcohol and 18.7% cigarettes in the past 30 days (Johnston et al., 2012). Adolescence is a critical developmental period when drug experimentation occurs, with use increasing with age through adolescence to young adulthood and then declining thereafter (Chen & Jacobson, 2012).

One of major correlates of substance use is the onset of puberty when significant neuroendocrinological changes occur, such as increases in adrenal and gonadal hormones, which are associated with the development of secondary sexual characteristics and brain functioning (Spear, 2000). Anatomical neuroimaging studies have shown that prefrontal areas are the last brain regions to mature (Giedd, 2004; Luna & Sweeney, 2004). As such, greater impulsivity, sensation-seeking, and substance use are evident during adolescence when prefrontal cortex may still be undergoing changes. Delayed prefrontal functioning has been linked with substance use (See review by Goldstein & Volkow, 2002).

Epidemiological evidence also shows that the pattern of substance use differs by gender during adolescence (Chen & Jacobson, 2012; Johnston, et al., 2012). For instance, girls have higher levels of substance use in early adolescence and boys show greater changes over time with higher levels of use in mid-adolescence and early adulthood (Chen & Jacobson, 2012). The different pattern of drug use may be linked to pubertal development, as well as elevated levels of impulsivity and sensation-seeking. Females start and complete physical development earlier than males (Sun et al., 2002), and males have higher impulsivity and sensation-seeking than females (Cross et al., 2011; Rosenblitt et al., 2001; Zuckerman, 1994).

Pubertal Status and Substance Use

Increases in substance use occur with biological changes that accompany puberty. As such, physical pubertal development is associated with alcohol, cigarette, and marijuana use among adolescents, with most data indicating that early-maturing adolescents have greater likelihood of initiation and faster progression of substance use (Biehl et al., 2007; Chung et al., 2005; Dick et al., 2000; Downing & Bellis, 2009; Ge et al., 2006; Hayatbakhsh et al., 2009). This pattern is consistent with the maturation-deviance hypothesis, which states that adolescents whose development deviates from the norm experience heightened emotional distress, which in turn, influences substance use behaviors (Tschann et al., 1994). Findings also point to gender differences in these associations; early-maturing girls report greater substance use (Ge, et al., 2006; Graber et al., 1997), but less consistent findings exist for boys. Some studies find that late development for boys is related to greater risk for increased substance use (Wichstrøm, 2001), while others observe that both boys' and girls' early development is related to substance use (Westling et al., 2008).

The link between pubertal development based on physical and associated objective pubertal status variables and substance use is inherently limited by reduced variations in overt signs of development among older adolescents, such as high-school-aged youth (Graber, et al., 1997). Among adults, early puberty is not associated with increased substance use (Viner et al., 2006); therefore, the association between early puberty with substance use may not be significant in a high-school sample whose pubertal development is complete or near complete. Given this possibility, we examined both pubertal timing based on self-reported physical development and subjective appraisal of one's pubertal development relative to peers to assess both aspects of pubertal status.

Pubertal timing based on perceptions of one's relative development compared to one's peers may more strongly influence the course of substance use behaviors than would actual pubertal development in high-school (Graber, et al., 1997; Storvoll et al., 2008) and young adult samples (Graber et al., 2004). Although also likely to correlate with physical pubertal development based on physician ratings of physical characteristics, perceptions of pubertal timing relative to peers may encompass aspects that are not captured by the ratings of physical characteristics (Dubas et al., 1991). For instance, perceptions may include more subtle psychosocial traits or milestones that adolescents associate with development, including being teased or rejected, which may generate stress and low self-esteem and ultimately influence behaviors like engagement in sexual activity and substance use.

Impulsivity/Sensation-Seeking and Substance Use

Impulsivity and sensation-seeking dramatically increases in adolescence and they can enhance our understanding of the link between pubertal status and substance use, especially among a high-school-aged sample when overt physical development may be less salient. Sensation-seeking is defined as willingness to take risks for novel experience or excitement (Zuckerman, 1994). Sensation-seeking is positively associated with the presence of sex hormones that is found at the onset of pubertal development (Zuckerman et al., 1980) and the association between pubertal status and substance use can be attributed to sensation-seeking in early-developing adolescents (Martin et al., 2002). Sensation-seeking may attract adolescents to engage in activities that are novel, impulsive, and potentially risky, including alcohol, cigarette, and marijuana use (Arnett, 1992; Crawford et al., 2003; Malmberg et al., 2010).

Impulsivity, defined as acting on impulse without thinking or planning (Zuckerman, 1994), is another trait that is increased in adolescence and has been linked to substance use (Verdejo-García et al., 2008). Sensation-seeking and impulsivity are conflated yet dissociable constructs; high sensation-seeking does not necessarily result from impulsive actions and not all impulsive behaviors are aimed at eliciting novel, exciting experiences (Steinberg et al., 2008). Sensation-seeking and impulsivity may have different associations with pubertal timing and substance use in a high school-aged sample because they follow different developmental trajectories; sensation-seeking sharply increases between ages 10 and 15 and slowly decreases thereafter, and impulsivity steadily increases from age 10 and into young adulthood (Steinberg, et al., 2008). However, studies examining impulsivity and substance use that take pubertal status into account are sparse. Given that impulsivity increases over the course of adolescence (Steinberg, et al., 2008), it is possible that along with sensation-seeking, impulsivity may also be correlated with substance use.

Hypotheses

We hypothesized that early perceived pubertal timing, high impulsivity, and high sensation-seeking would be associated with substance use (alcohol, cigarettes, marijuana), but pubertal timing based on self-report of physical development would not be associated with substance

use among high-school-aged adolescents. To explore gender differences in these relationships, we examined these associations separately by gender.

Methods

Procedures

This study analyzed a cross-sectional survey of high-school students conducted in 2006. Study procedures were described previously (Cavallo et al., 2010; Potenza et al., 2011). Briefly, all public 4-year high schools in the state of Connecticut were invited to participate. Ten schools representing a wide geographical region of the state and the three tiers of the state's district reference groups (DRGs) participated. DRGs represent the socioeconomic status of the families in the school district, and sampling of schools across DRGs helped to enhance the generalizability of the data.

Following approval from the school principal, boards and/or superintendents, a passive consent procedure was used to obtain permission from a parent or legal guardian. Research assistants administered the survey to the entire student body in assemblies, health or English classes and informed students of the confidentiality and the voluntary nature of their participation. The survey took about 30 minutes to complete and all students were provided a pen for their participation. This study was reviewed and approved by the Yale Institutional Review Board.

Measures

The Pubertal Development Scale—(PDS; Petersen et al., 1988) is a self-report scale that assesses children's stages of physical growth on height, skin changes, and pubic hair. Boys complete additional items on facial hair and voice changes; girls complete additional items on breast growth and menarche. Item responses fall on a scale from "1 = not started" to "4 = completed" with the exception of menarche, which is dichotomous ("yes/no"). The PDS is widely used in longitudinal and school-based studies and has been shown to reliably capture actual physical development in adolescents (Patton et al., 2004; Petersen, et al., 1988). PDS scores have shown moderate reliability (alpha coefficients ranging from 0.68 to 0.83; Petersen et al., 1988) and correlations between 0.61 and 0.67 with physical exams (Brooks-Gunn et al., 1987). Responses were averaged across the items to retain metric, resulting in a score indicative of stage of development. The Cronbach's alpha of this scale in this sample was 0.79.

Perceived pubertal timing—(PPT) relative to peers was assessed by averaging two questions ($\alpha = 0.85$): "Do you believe your development is (was) any earlier or later than most other boys/girls your age?" (Dubas, et al., 1991) and "Compared with other kids your age, when did you start puberty?" Participants indicated their development on a 5-point Likert scale ranging from "much earlier/a lot before" to "much later/a lot later." The group who reported not knowing how their development compared with that of their peers ($n = 514$, 11.1%) was excluded from comparisons.

Cigarettes, marijuana, and alcohol use—was measured by asking about each substance use in the past 30 days. As there was limited variability in frequencies of substance use, items were dichotomized to indicate use (once or more) or no use of substances in the past 30 days.

Impulsivity and sensation-seeking—was assessed using the 19 items from the Zuckerman-Kuhlman Personality Questionnaire (Zuckerman et al., 1993). The impulsivity score was computed by adding the "true" responses of the 8 items of this subscale (e.g., "I

often do things on impulse.”) and sensation-seeking score was computed by adding 11 items (e.g., “I like to have new and exciting experiences and sensations even if they are a little frightening.”). Factor analysis of this measure supported the 5-factor model of personality, in which impulsivity and sensation-seeking were two unique factors, and these subscales have shown good internal consistency (alphas ranging from 0.64 – 0.77; Zuckerman, 2002). In our sample, we detected Cronbach’s alpha of 0.77 for the sensation-seeking scale and 0.71 for the impulsivity scale. We used mean substitution to compute missing scores for the participants missing less than 50% of the items that comprised each subscale. Both list-wise deletion and mean substitution detected the same significant results; therefore, we presented the results from mean substitution.

Participants

The sample consisted of 4637 students in grades 9 through 12. Of the total sample, 3068 participants (66.2%) with complete data on age, gender, PDS, PPT, cigarettes, marijuana and alcohol use, sensation-seeking, and impulsivity were included in the analyses.

Those with incomplete and complete data did not differ on age ($t(4510) = 0.49, p = 0.62$), sensation-seeking ($t(3788) = -0.73, p = 0.47$), and PPT ($t(3423) = 1.81, p = 0.07$), but those with incomplete data were more likely to be boys ($\chi^2(1, 4468) = 231.67, p < .001$), use cigarettes ($\chi^2(1, 4637) = 135.88, p < 0.001$), alcohol ($\chi^2(1, 4637) = 4.61, p = .03$) and marijuana ($\chi^2(1, 4637) = 148.23, p < 0.001$), and be more impulsive ($t(3826) = 3.11, p = 0.002$) and develop later based on PDS scores ($\chi^2(1, 3989) = -3.87, p < .001$).

Statistical Analyses

Chi-squares and t-tests were used to assess for differences between study variables and gender, as well as data completeness and all study variables. Those with complete data were selected for logistic regression to assess the use of cigarettes, marijuana, and alcohol in the previous 30 days separately for boys and girls. Each substance use was modeled separately and all models contained sensation-seeking, impulsivity, PDS, and PPT as independent variables while controlling for age. To correct for multiple comparisons, Bonferroni-determined p -values equal to or less than 0.008 were considered statistically significant.

Results

See Table 1 for the demographic and study variables for study participants. Boys and girls did not differ on alcohol use, but more girls reported cigarette use and more boys reported marijuana use. Compared to girls, boys had greater impulsivity and higher PDS scores. The scores on PPT and sensation-seeking did not differ by gender.

PDS and PPT correlated weakly ($r(3028) = 0.14, p < 0.001$) and sensation-seeking and impulsivity correlated moderately ($r(3028) = 0.57, p < 0.001$).

Logistic regression analysis modeling the use of three substances (alcohol, cigarettes, marijuana) with PDS, PPT, sensation-seeking, and impulsivity as independent variables separately for boys and girls showed that sensation-seeking is associated with substance use for both boys and girls (Table 2). Different relationship between impulsivity and substance use was detected for boys and girls: among girls, impulsivity was associated with use of all three substances, but among boys, it was not associated with any substance use. Pubertal status differentially associated with use of substances; early PPT was associated with cigarette smoking for both boys and girls, but PDS was not associated with any substance use for both genders.

Discussion

This study examined the role of pubertal timing (based on self-reported physical development score [PDS] and on perceptions of pubertal timing relative to one's peers [PPT]), sensation-seeking, and impulsivity in relation to substance use (alcohol, cigarette, marijuana) for high-school-aged boys and girls.

Consistent with our hypothesis, physical development determined by the PDS was not associated with past 30-day use of alcohol, cigarettes and marijuana. This finding suggests that the relationship between early pubertal development and substance use may be more salient earlier in adolescent development, at initial stages of puberty. This interpretation is consistent with recent findings that the negative effects of early physical maturation attenuated as adolescents reached young adulthood (Copeland et al., 2010). This sample of adolescents may have already progressed in pubertal stages, and therefore, other predictors specific to adolescence such as perceived pubertal timing, sensation-seeking and impulsivity may be more strongly related to substance use.

Interestingly, we found substance-specific findings: among boys and girls, early perceived pubertal timing was associated with use of cigarettes but not use of alcohol or marijuana. The perception of early maturation compared to one's peers may affect the social interactions and self-image more so than would actual physical maturation. Smoking cigarettes may represent one way that an adolescent expresses an image of the mature self; adolescent girls who smoke consider themselves sophisticated and worldly and that smoking promotes this image (Lloyd et al., 1997). As such, meanings that adolescents construct about smoking may become integrated into their social identity and may be important in making decisions to smoke. Alcohol and marijuana use may be less contingent upon one's subjective appraisal of pubertal development; alcohol use may be more normative and acceptable in high-school-aged youth (close to 50% reported drinking in the past 30 days) and marijuana use may be seen as more deviant and less linked to one's identity.

We found that for both boys and girls, sensation-seeking was independently related to substance use (alcohol, cigarette, and marijuana). This is consistent with other findings that indicated that sensation-seeking motivates adolescents toward behaviors that are novel and exciting such as substance use (Arnett, 1992; Crawford, et al., 2003; Malmberg, et al., 2010). Because nicotine, alcohol, and marijuana elicit rewarding effects (Chambers et al., 2003), those with strong sensation-seeking drives may continue to use substances to elicit such responses.

Even though boys had greater impulsivity than girls, impulsivity was significantly associated with use of all three substances among girls only. This finding corresponds to the adult literature that have also reported on gender differences in the association between impulsivity and substance use (Waldeck & Miller, 1997). For example, impulsivity prospectively predicted the increase in smoking in women and alcohol consumption in men and women (Granö et al., 2004). It is unclear why impulsivity in girls but not in boys is related to greater substance use. It is possible that the association between impulsivity and substance use among boys in this sample may not truly represent high-school-aged boys because the excluded because of missing data were more likely to be boys and impulsive. The potential bias that may exist in our data precludes us from making conclusive statements about the role of impulsivity, and its role in substance use among boys should not be ruled out without further research. Despite this limitation, we were still able to detect an association between early perceived pubertal timing and cigarette use, and sensation-seeking and past 30-day use of cigarettes, alcohol, marijuana in boys and girls.

Several other limitations should be taken into account when interpreting the findings of this study. This is a cross-sectional self-reported survey, and thus, accuracy of adolescents' actual substance use cannot be verified and directionality of the associations cannot be established. In addition, biological markers or physician ratings of pubertal development using tools such as Tanner staging were not used to determine pubertal development. Using the PDS to estimate actual physical developmental status is inherently subjective and adolescents' ratings may not accurately represent their actual physical development. However, in contexts such as school settings where a physical exam is not feasible and the use of illustrations of body developmental stages may be concerning to school districts and/or parents, it represents an appropriate and valid substitution (Dorn et al., 2006), especially considering the PDS correlations with Tanner staging (Patton, et al., 2004).

This study has several strengths: (1) examination of constructs that have been linked to prefrontal cortex and specific in adolescence such as puberty, sensation-seeking, impulsivity, and substance use, (2) a simultaneous comparison of two measures that encompass pubertal development and perceptions of pubertal timing, (3) a focus on past-30-day use of three of the most commonly used substances in adolescence: alcohol, cigarettes, and marijuana, and (4) exploration of these factors separated by gender in a non-clinical sample of high-school-aged adolescents.

In conclusion, sensation-seeking is associated with substance use for all adolescents; therefore, drug prevention efforts that use more effective anti-drug messages targeting high sensation-seekers may be useful. Because sensation-seeking is biologically-based, better understanding of the neural mechanisms underlying this trait may inform targeted drug interventions. Our findings also suggest that among high-school-aged adolescents, perceived pubertal timing compared to peers rather than physical pubertal development was associated with cigarette smoking. Perhaps, parents, educators, and clinicians who work with adolescents can focus on promoting a positive self-image that does not include cigarette smoking. Finally, future studies using longitudinal design should investigate how changes in physical development and subjective experience of pubertal timing influence sensation-seeking and impulsivity from early adolescence through late adolescence to better understand how these relationships to substance use may change with time and growth.

Acknowledgments

Funding: This work was supported in part by National Institutes P50 AA15632, P50 DA09421, T32 DA07238, the Connecticut Department of Mental Health and Addiction Services.

References

- Arnett J. Reckless behavior in adolescence: A developmental perspective. *Developmental Review*. 1992; 12:339–373.
- Biehl M, Natsuaki M, Ge X. The Influence of Pubertal Timing on Alcohol Use and Heavy Drinking Trajectories. *Journal of Youth and Adolescence*. 2007; 36(2):153–167.
- Brooks-Gunn J, Warren MP, Rosso J, Gargiulo J. Validity of self-report measures of girls' pubertal status. *Child Development*. 1987; 58(3):829–841. [PubMed: 3608653]
- Cavallo D, Smith AE, Schepis TS, Desai R, Potenza MN, Krishnan-Sarin S. Smoking expectancies, weight concerns, and dietary behaviors in adolescence. *Pediatrics*. 2010; 126(1):e66–e72. [PubMed: 20547649]
- Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: A critical period of addiction vulnerability. *The American Journal of Psychiatry*. 2003; 160:1041–1052. [PubMed: 12777258]

- Chen P, Jacobson KC. Developmental trajectories of substance use from early adolescence to young adulthood: Gender and racial/ethnic differences. *Journal of Adolescent Health*. 2012; 50(2):154–163. [PubMed: 22265111]
- Chung H, Park Y, Lanza ST. Latent transition analysis with covariates: Pubertal timing and substance use behaviours in adolescent females. *Statistics in Medicine*. 2005; 24(18):2895–2910. [PubMed: 16134129]
- Copeland W, Shanahan L, Miller S, Costello J, Angold A, Maughan B. Do the negative effects of early pubertal timing on adolescent girls continue into young adulthood? *American Journal of Psychiatry*. 2010; 167:1218–1225. [PubMed: 20478880]
- Crawford AM, Pentz MA, Chou C, Li C, Dwyer JH. Parallel developmental trajectories of sensation seeking and regular substance use in adolescents. *Psychology of Addictive Behaviors*. 2003; 17(3):179–192. [PubMed: 14498812]
- Cross CP, Copping LT, Campbell A. Sex differences in impulsivity: A meta-analysis. *Psychological Bulletin*. 2011; 137:97–130. [PubMed: 21219058]
- Dick DM, Rose RJ, Viken RJ, Kaprio J. Pubertal timing and substance use: Associations between and within families across late adolescence. *Developmental Psychology*. 2000; 36(2):180–189. [PubMed: 10749075]
- Dorn LD, Dahl RE, Woodward HR, Biro F. Defining the boundaries of early adolescence: A user's guide to assessing pubertal status and pubertal timing in research with adolescents. *Applied Developmental Science*. 2006; 10(1):30–56.
- Downing J, Bellis MA. Early pubertal onset and its relationship with sexual risk taking, substance use and anti-social behaviour: A preliminary cross-sectional study. *BMC Public Health*. 2009; 9:446. [PubMed: 19958543]
- Dubas JS, Graber JA, Petersen AC. A longitudinal investigation of adolescents' changing perceptions of pubertal timing. *Developmental Psychology*. 1991; 27(4):580–586.
- Ge X, Jin R, Natsuaki MN, Gibbons FX, Brody GH, Cutrona CE, et al. Pubertal maturation and early substance use risks among African American children. *Psychology of Addictive Behaviors*. 2006; 20(4):404–414. [PubMed: 17176175]
- Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Annals of the New York Academy of Sciences*. 2004; 1021:77–85. [PubMed: 15251877]
- Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: Neuroimaging evidence for the involvement of the frontal cortex. *American Journal of Psychiatry*. 2002; 159:1642–1652. [PubMed: 12359667]
- Graber JA, Lewinsohn PM, Seeley JR, Brooks-Gunn J. Is psychopathology associated with the timing of pubertal development? *Journal of the American Academy of Child & Adolescent Psychiatry*. 1997; 36(12):1768–1776. [PubMed: 9401339]
- Graber JA, Seeley JR, Brooks-Gunn J, Lewinsohn PM. Is pubertal timing associated with psychopathology in young adulthood? *Journal of the American Academy of Child & Adolescent Psychiatry*. 2004; 43(6):718–726. [PubMed: 15167088]
- Granö N, Virtanen M, Vahtera J, Elovainio M, Kivimäki M. Impulsivity as a predictor of smoking and alcohol consumption. *Personality and Individual Differences*. 2004; 37(8):1693–1700.
- Hayatbakhsh MR, Najman JM, McGee TR, Bor W, O'Callaghan MJ. Early pubertal maturation in the prediction of early adult substance use: A prospective study. *Addiction*. 2009; 104(1):59–66. [PubMed: 18983626]
- Johnston, LD.; O'Malley, PM.; Bachman, JD.; Schulenberg, JE. Monitoring the Future national survey results on drug use. 1975–2011. Volume II: College students and adults ages 19–50. Ann Arbor: Institute for Social Research, The University of Michigan; 2012.
- Lloyd B, Lucas K, Fernbach M. Adolescent girls' constructions of smoking identities: Implications for health promotion. *Journal of Adolescence*. 1997; 20(1):43–56. [PubMed: 9063774]
- Luna B, Sweeney JA. The emergence of collaborative brain function: FMRI studies of the development of response inhibition. *Annals of the New York Academy of Sciences*. 2004; 1021:296–309. [PubMed: 15251900]

- Malmberg M, Overbeek G, Monshouwer K, Lammers J, Vollebergh WA, Engels RC. Substance use risk profiles and associations with early substance use in adolescence. *Journal of Behavioral Medicine*. 2010; 33(6):474–485. [PubMed: 20625809]
- Martin CA, Kelly TH, Rayens MK, Brogli BR, Brenzel A, Smith WJ, et al. Sensation seeking, puberty, and nicotine, alcohol, and marijuana use in adolescence. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2002; 41(12):1495–1502. [PubMed: 12447037]
- Martin CA, Logan TK, Leukefeld C, Milich R, Omar H, Clayton R. Adolescent and young adult substance use: Association with sensation seeking, self esteem and retrospective report of early pubertal onset. A preliminary examination. *International Journal of Adolescent Medicine and Health*. 2001; 13:211–219.
- Patton GC, McMorris BJ, Toumbourou JW, Hemphill SA, Donath S, Catalano RF. Puberty and the onset of substance use and abuse. *Pediatrics*. 2004; 114(3):300–306.
- Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*. 1988; 17(2):117–133.
- Potenza MN, Wareham JD, Steinberg MA, Rugle L, Cavallo DA, Krishnan-Sarin S, et al. Correlates of at-risk/problem internet gambling in adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2011; 50(2):150–159. [PubMed: 21241952]
- Rosenblitt JC, Soler H, Johnson SE, Quadagno DM. Sensation seeking and hormones in men and women: Exploring the link. *Hormones and Behavior*. 2001; 40(3):396–402. [PubMed: 11673912]
- Spear LP. The adolescent brain and age-related behavioral manifestations. *Neuroscience and Biobehavioral Reviews*. 2000; 24:417–463. [PubMed: 10817843]
- Steinberg L, Albert D, Cauffman E, Banich M, Graham S, Woolard J. Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: Evidence for a dual systems model. *Developmental Psychology*. 2008; 44(6):1764–1778. [PubMed: 18999337]
- Storvoll EE, Pape H, Rossow I. Use of commercial and social sources of alcohol by underage drinkers: The role of pubertal timing. *Addictive Behaviors*. 2008; 33(1):161–166. [PubMed: 17553624]
- Sun SS, Schubert CM, Chumlea WC, Roche AF, Kulin HE, Lee PA, et al. National estimates of the timing of sexual maturation and racial differences among US children. *Pediatrics*. 2002; 110(5):911–919. [PubMed: 12415029]
- Tschann JM, Adler NE, Irwin CE, Millstein SG, Turner RA, Kegeles SM. The roles of pubertal timing and emotional distress. *Health Psychology*. 1994; 13(4):326–333. [PubMed: 7957011]
- Verdejo-García A, Lawrence AJ, Clark L. Impulsivity as a vulnerability marker for substance-use disorders: Review of findings from high-risk research, problem gamblers and genetic association studies. *Neuroscience & Biobehavioral Reviews*. 2008; 32(4):777–810. [PubMed: 18295884]
- Viner RM, Cole TJ, Taylor B. What are the implications of early pubertal development? Findings in adulthood from a national birth cohort. *Archives of Disease in Childhood*. 2006; 91(Supplement 1):A86.
- Waldeck TL, Miller LS. Gender and impulsivity differences in licit substance use. *Journal of Substance Abuse*. 1997; 9(0):269–275. [PubMed: 9494954]
- Westling E, Andrews JA, Hampson SE, Peterson M. Pubertal timing and substance use: The effects of gender, parental monitoring and deviant peers. *Journal of Adolescent Health*. 2008; 42(6):555–563. [PubMed: 18486864]
- Wichstrøm L. The impact of pubertal timing on adolescents' alcohol use. *Journal of Research on Adolescence*. 2001; 11(2):131–150.
- Zuckerman, M. *Behavioral Expressions and Biosocial Bases of Sensation Seeking*. New York: Cambridge University Press; 1994.
- Zuckerman, M. Zuckerman-Kuhlman Personality Questionnaire (ZKPQ): An alternative five-factor model. In: De Raad, B.; Perugini, M., editors. *Big Five Assessment*. Seattle, WA: Hogrefe & Huber; 2002. p. 377–396.
- Zuckerman M, Buchsbaum MS, Murphy DL. Sensation seeking and its biological correlates. *Psychological Bulletin*. 1980; 88:187–214.
- Zuckerman M, Kuhlman MD, Joireman J, Teta P, Kraft M. A comparison of three structural models for personality: The Big Three, the Big Five, and the Alternative Five. *Journal of Personality and Social Psychology*. 1993; 65:757–768.

Table 1

Sample characteristics on study variables.

	Boys (<i>n</i> = 1202)	Girls (<i>n</i> = 1826)	
	%	%	χ^2/t
Age			
13 – 14	11.6	14.3	13.71 *
15–16	54.1	55.3	
17 – 18 and older	34.3	30.4	
Grade			
9 th	28.2	28.0	0.01
10 th	27.4	27.9	
11 th	27.7	28.2	
12 th	16.8	16.0	
Race/Ethnicity			
Caucasian	74.0	75.5	0.05
Multi-race	2.3	2.6	
Hispanic	11.5	11.8	
African American	5.8	5.7	
Asian	3.4	1.9	
Other	3.0	2.5	
Cigarette Use in last 30 days	19.2	22.2	3.97 *
Alcohol Use in last 30 days	47.8	49.2	0.64
Marijuana Use in last 30 days	28.5	25.0	4.52 *
PDS (M, SD)	3.19, 0.64	2.97, 0.45	10.85 **
PPT (M, SD)	3.16, 0.87	3.12, 0.87	1.09
Sensation-seeking (M, SD)	6.60, 2.87	6.64, 2.83	–0.44
Impulsivity (M, SD)	3.76, 2.18	3.22, 2.18	6.75 **

Note: PDS = Pubertal Development Scale; PPT = Perceived Pubertal Timing; M = mean; SD = standard deviation.

*
p 0.05;**
p 0.01.

Logistic Regressions Associating the Risk of Substance Use and Impulsivity, Sensation Seeking, and Pubertal Timing Stratified by Gender.

Table 2

	No Use vs. Use					
	Cigarette		Alcohol		Marijuana	
	OR	95% CI	OR	95% CI	OR	95% CI
Boys (n = 1202)						
Pubertal Developmental Scale	0.73	0.58, 0.93	1.05	0.86, 1.28	0.88	0.71, 1.09
Perceived Pubertal Timing	1.28*	1.08, 1.51	0.99	0.86, 1.14	1.17	1.01, 1.36
Sensation-seeking	1.16**	1.09, 1.25	1.16**	1.10, 1.22	1.13**	1.01, 1.17
Impulsivity	1.10	1.01, 1.20	1.07	1.00, 1.15	1.09	1.01, 1.17
Girls (n = 1826)						
Pubertal Developmental Scale	1.21	0.91,1.61	1.22	0.96, 1.54	0.94	0.73, 1.22
Perceived Pubertal timing	1.22*	1.06, 1.40	1.12	1.00, 1.26	1.14	1.00, 1.29
Sensation-seeking	1.22**	1.16, 1.29	1.25**	1.19, 1.30	1.17**	1.11, 1.23
Impulsivity	1.16**	1.09, 1.24	1.11**	1.05, 1.17	1.16**	1.08, 1.22

Note: All analysis controlled for age and Bonferroni correction was applied for significance testing. PDS = Pubertal Development Scale; OR = odds ratio; SE = standard error;

* p 0.008;
** p 0.001.