CASE REPORT

Disseminated *Mycobacterium haemophilum* infection in a 72-year-old patient with rheumatoid arthritis on infliximab

C Scott Collins,¹ Christine Terrell,² Paul Mueller¹

¹Department of Internal Medicine, Mayo Clinic, Rochester, Minnesota, USA
²Department of Infectious Disease, Mayo Clinic, Rochester, Minnesota, USA

Correspondence to
 Dr Charles Scott Collins,
 Collins.Charles@mayo.edu

SUMMARY

Mycobacterium haemophilum is a slow growing, aerobic, fastidious mycobacterium requiring hemin and a temperature of 30–32° C for optimal growth that is ubiquitous in nature. Disease in immunocompromised adults typically manifests as skin lesions such as papules, pustules and ulcerations. This organism also causes lymphadenitis in immunocompetent children. Antitumour necrosis factor- α (anti-TNF- α) therapy with agents such as infliximab, etanercept and adalimumab is increasingly being used for immunosuppression in patients with various autoimmune conditions. These agents are known to place patients at increased risk for tuberculosis and other granulomatous diseases. However, little is known about illness caused by *M haemophilum* in patients on immunosuppression with anti-TNF- α therapy. We describe a case of disseminated *M haemophilum* manifesting as skin lesions in a 72-year-old man with rheumatoid arthritis on infliximab and methotrexate.

BACKGROUND

This paper demonstrates the clinical and laboratory processes involved in diagnosing *Mycobacterium haemophilum* infection. Our case involves an elderly man with multiple medical comorbidities with unexplained skin lesions who is immunocompromised with an antitumour necrosis factor- α agent. Antimicrobial management in this patient is prolonged, quite complex and has limited evidence in the literature. As such, this paper should be of interest to a broad readership including those interested in infections in immunocompromised hosts, mycobacteriology and clinical microbiology.

CASE PRESENTATION

A 72-year-old man with seronegative rheumatoid arthritis of 10 years duration that had previously been well controlled on infliximab and methotrexate for the past 5 years and intermittently with glucocorticoids came to our institution for another opinion regarding an 8-month history of red nodules and plaques on the upper and lower extremities and face. He was from the South Central USA. He used to work in oil fields, but more recently he supervised oil well cleanup sites. Other pertinent medical history included dilated cardiomyopathy with mild coronary artery disease, atrial fibrillation (and resultant long-term warfarin anticoagulation), status-post placement of a dual chamber pacemaker following repeated episodes of syncope and a prolonged QT interval.

Twenty months prior to presentation to our institution, he developed a nodular lesion on the right medial forearm. Biopsy at that time showed panniculitis consistent with erythema nodosum, and this lesion resolved on its own. He did relatively well until 8 months prior to presentation, at which time he developed nodular and vesicular lesions on the dorsum of the foot with extension to the left calf and posterior medial thigh. Eventually, lesions developed on the right arm near the elbow and then the right cheek. The lesions waxed and waned in intensity and disappeared and reappeared in the same area. The overlying skin was generally erythematous and the nodules mildly tender. They were treated on two or three occasions with bursts of steroids which seemed to help temporarily.

Despite the above skin lesions, he had no fevers, chills or night sweats. His appetite was excellent. He had no involuntary weight loss. Though fatigued, he did not feel ill. He had no significant cough or sputum production. He had had slowly progressive shortness of breath that was stable and had been attributed to his cardiomyopathy. He had lower extremity oedema treated with diuretics and compression. He had no vomiting, abdominal pain or diarrhoea. He did feel like his rheumatoid arthritis was poorly controlled with increased arthralgias. He did not use hot tubs or whirlpools. The only swimming he had done in recent years was in an above-ground swimming pool. He had had several purified protein derivative (PPDs) testing, and they had all been negative. He had no recent foreign travel. He was in Hawaii a few years prior. He was neither a fisherman, nor had he been in contact with fish. He had not had any pedicures.

INVESTIGATIONS

Multiple biopsies of the above skin lesions were performed locally on two additional occasions within 3 months of each other (8 and 5 months prior to presentation to our institution). Most of these biopsies were interpreted as showing erythema nodosum just as the initial one from 20 months prior. This finding and his long term anti-TNF- α therapy prompted placement of another PPD which was negative. He had no other symptoms to suggest systemic processes such as sarcoidosis or inflammatory bowel disease or infection. However, CT of the chest, abdomen and pelvis was performed locally at that time and was unremarkable. In addition to the finding of erythema nodosum, a single leg lesion was interpreted as showing CD30 lymphoproliferative disease and

To cite: Collins CS, Terrell C, Mueller P. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2012-008034

possibly lymphomatoid papulosis. He had no other clinical evidence suggestive of systemic lymphoma, and his negative imaging and lack of symptoms were noted.

In the mean time, he had been placed on prednisone 10 mg daily and non-steroidal anti-inflammatory drugs for the working diagnosis of erythema nodosum with indeterminate cause, poorly controlled rheumatoid arthritis and possible lymphomatoid papulosis. Since his inflammatory arthritis remained poorly controlled, his infliximab was discontinued 1.5 months prior to coming to our institution (with no effect on his lesions). He was then started on abatacept and received one dose of abatacept 2 weeks prior to our meeting him.

On our examination, he was non-toxic but appeared chronically ill and deconditioned. There was no lymphadenopathy. He did have swelling primarily in the right wrist and the second, third and fourth metacarpophalangeal on the right with positive squeeze test. There was also bilateral knee swelling and soft tissue swelling present in both ankles. He had negative metatarsophalangeal squeeze testing. He had no acute inflammation of his joints. Skin examination revealed tender, erythematous nodules and plaques on the right elbow region and the bilateral inner thighs and posterior calves. He had postinflammatory hyperpigmentation and scars from prior lesions and mild purple papules on the dorsal feet as well.

Diagnostic testing at our institution revealed unremarkable complete blood count with differential, electrolytes, sedimentation rate and C reactive protein. Aerobic and anaerobic bacterial blood cultures were negative. Blood cultures for mycobacteria, however, became positive for *M haemophilum* at 38 days. Serologies were negative for fungal antigens and antibodies, acute tick borne diseases and viral hepatitis. Additionally, he was found to be seronegative for rheumatoid arthritis (negative rheumatoid factor, citrullinated peptide antibodies and antinuclear antibodies), had negative monoclonal protein and cryoglobulin studies, and peripheral blood phenotyping was within normal limits. Plain film x-rays of the hands showed evidence of degenerative changes in the MCP joints.

Skin biopsies at our institution were obtained from a lesion on his right forearm and another from the left upper thigh. Immunohistochemical stains (CD3, CD4, CD8 and CD30) demonstrated a CD3 T-cell population with a mixture of CD4 and CD8 positive cells. CD30 stain was negative. Gram stains were negative for microorganisms. Given his long-term exposure to anti-TNF- α therapy and risk for mycobacterial infection, acid-fast staining was performed and showed dermal and subcutaneous granulomatous inflammation with several acid-fast bacilli seen (figure 1). These slides were compared to skin biopsy specimens obtained at the outside institution 8 and 5 months previously. Review of those biopsy specimens revealed superficial and deep dermal perivascular and periadnexal granulomatous inflammation and several acid-fast bacilli. *Mycobacterium tuberculosis* complexes' PCR was negative. Cultures of the biopsy specimens were done, and *M haemophilum* was identified after 23 days of growth. Susceptibilities were performed at an outside reference lab. They were pending at the time a management decision had to be made as this organism usually takes about 4–5 weeks to grow enough to allow for susceptibility testing. The organism ultimately showed susceptibility to clarithromycin (MIC 15 μ g/ml), trimethoprim/sulfamethoxazole (MIC 0.5/9.5 μ g/ml, linezolid (MIC 6 μ g/ml), ciprofloxacin (2 μ g/ml) and doxycycline (MIC 6 μ g/ml). The organism was resistant in vitro to rifampin, amikacin and minocycline.

TREATMENT

Our preferred empiric treatment of this infection was a prolonged course (at least 6–12 months) of azithromycin or

clarithromycin, rifampin and ciprofloxacin.¹ However, the patient's prolonged QT interval and anticoagulation with warfarin posed increased risk for arrhythmia and sudden death and difficulty controlling his international normalised ratio (INR) if this combination were used. Given his clinical stability, we had initially planned to observe him and await susceptibilities before planning a definitive treatment course. However, blood cultures for mycobacteria also were positive for *M haemophilum* at 38 days, which added urgency to commence treatment.

Given the above, we recommended that he would be treated empirically with azithromycin 250 mg daily, rifampin 600 mg daily and intravenous amikacin at a dose of 8–10 mg/kg daily. Azithromycin, like ciprofloxacin, also has the potential to prolong QT interval, but since macrolides are the cornerstone of treatment for this particular organism and many other non-tuberculous mycobacterial organisms we elected to use it. We recommended baseline audiometry since the amikacin can cause tinnitus and hearing loss, particularly given the patient's history of hearing loss. We also recommended monitoring of blood chemistries and close observation of his INR as azithromycin can increase the INR and rifampin can decrease the INR in the setting of warfarin anticoagulation. The duration of the treatment course was recommended to be a minimum of 6–12 months, as he was immunocompromised. The first treatment endpoint was determined to be the resolution of his lesions.

OUTCOME AND FOLLOW-UP

He completed 5 weeks of the empiric regimen prior to susceptibilities becoming available. At that time, his lesions had completely resolved. He had not had any new skin lesions, fevers, chills or sweats. His appetite remained good, and his weight was stable. Amikacin was discontinued at that time to avoid side effects and because the patient had responded so well clinically. It is also notable that he had completed a prolonged prednisone taper, and was no longer on any disease modifying antirheumatic drugs at that time. His arthritis was essentially clinically inactive. He remained on azithromycin and rifampin.

Given his response, we suspect that amikacin and/or rifampin were helpful despite the in vitro susceptibility pattern, and we are sceptical that his excellent response was because of azithromycin alone. We also added a third drug in case his organism was truly resistant to rifampin. We decided against fluoroquinolones because of his prolonged QTc. We did not want to use linezolid for several months because of its propensity to cause bone marrow suppression and other significant side effects. We considered trimethoprim-sulfamethoxazole, but decided against that because the risk of hyperkalemia would be high as he was already on quinapril, spironolactone and supplemental potassium for his cardiovascular issues. For these reasons we elected to add doxycycline 100 mg twice daily to his regimen.

In summary, the final recommendations offered were for azithromycin, rifampin and doxycycline for a total treatment course of at least 6 and more likely 12 months. We will consider antimycobacterial prophylaxis following therapy should he be started on a TNF inhibitor for his inflammatory arthritis in the future.

DISCUSSION

M. haemophilum was first described as a human pathogen in Israel in 1978 by Sompolinsky² from the cutaneous lesions of a woman with Hodgkin's disease and may have possibly caused lesions in a series of renal transplant patients reported prior to that in 1976. This low-virulence organism does not cause disease in immunocompetent adults, but does cause disease as

Figure 1 (A) Photo at $\times 20$ of H&E section from lesion on right elbow showing areas of dermal granulomatous inflammation. (B) Closer up photos ($\times 40$) of same area (H&E section from lesion on right elbow) showing areas of dermal granulomatous inflammation. (C) Corresponding area at $\times 40$ with acid-fast bacilli (AFB) stain. Note red filamentous organisms. (D) AFB stain at $\times 60$. Dozens of acid-fast organisms within area of granulomatous inflammation are visible. Courtesy of Carilyn N Wieland, MD, Department of Dermatology, Mayo Clinic, Rochester, Minnesota, USA.

an opportunistic pathogen in immunocompromised adults such as those with HIV infection, immunosuppression for organ transplants, those on chemotherapy for malignancy and those on disease modifying antirheumatic drugs such as methotrexate or anti-TNF- α therapy.^{3–4} These types of immunocompromised patients are thought to be at risk because of impaired cell-mediated immunity and difficulty with granuloma formation and maintenance. This organism is also known to cause lymphadenitis in immunocompetent children.^{5–6}

Infections caused by *M haemophilum* typically present with skin lesions though other locations such the joints, bone, lymphatics, the blood and the lungs are known to be affected as well. The cutaneous lesions are often initially nodules or erythematous papules which are tender and progress to painful draining ulcers. Other less common cutaneous lesions caused by this organism are cysts, scales or plaques. These lesions are most commonly found on the extremities and usually in close proximity to the joints. It is thought that the lesions manifest in the extremities and in the skin versus more central locations because of low-temperature requirements of the organism for maximal growth.⁷

Our reported case follows the above described typical pattern of an immunosuppressed individual with an autoimmune disease (rheumatoid arthritis) who develops skin lesions on the extremities while on anti-TNF- α therapy. The correct diagnosis in our case was delayed because the initial biopsy tissue was not stained for acid-fast bacilli. Consequently, the patient remained on methotrexate and anti-TNF- α therapy and was additionally placed on glucocorticoids, potentially exacerbating the underlying infection. This demonstrates the importance of having a

high index of suspicion for *M haemophilum* in this type of patient with cutaneous lesions on the extremities.

There is no specific antigenic test for *M haemophilum*, and PPD testing is not a reliable indicator of infection with *M haemophilum*. Clinical microbiology laboratories make the diagnosis of *M haemophilum* by detecting it in biopsy specimens through acid-fast staining, mycobacterial culturing at two temperatures using media with and without iron, and through molecular methods.⁸ Full description of these methods is beyond the scope of this paper.

Once the diagnosis of *M haemophilum* infection is made, there are no specific guidelines for empiric treatment.^{8–9} Thus, case specific susceptibilities are essential for this organism to guide therapy. No optimal duration of treatment or therapeutic regimen has been established. Treatment with multiple antibiotics is recommended by experts. Regimens usually include some combination of clarithromycin (or azithromycin), ciprofloxacin, a tetracycline, rifampin, trimethoprim-sulfamethoxazole and an aminoglycoside (particularly amikacin). The patient in our case initially was treated with azithromycin, amikacin and rifampin as described above. Methotrexate and the anti-TNF- α inhibitor were discontinued and prednisone was tapered off. After 5 weeks of treatment, he had had an excellent response with resolution of his lesions. Amikacin was discontinued, and the lesions did not recur. Several weeks later, the susceptibilities (listed in the case report above) became available. Azithromycin and rifampin were continued and doxycycline was added as a third agent. It is important to emphasise that treating the underlying cause of immunosuppression or halting/decreasing immunosuppression is essential in addition to antibiotic therapy.

There is debate as to whether it is the lifting of immunosuppression or the direct effect of antibiotics that leads to resolution of infection. With proper treatment, the prognosis for these patients is quite good, particularly in patients with only cutaneous manifestations. Most patients similar to ours reviewed in the literature had similar outcomes.

M haemophilum appears to be an increasingly common infection in those who are immunosuppressed, and it should be considered in the differential diagnosis of any such patient with chronic, recurrent or unexplained skin lesions on the extremities. Awareness of this organism by clinicians and laboratory workers and a high index of suspicion are essential as to not miss this diagnosis.

Learning points

- ▶ *Mycobacterium haemophilum* appears to be an increasingly common infection in those who are immunosuppressed.
- ▶ *M haemophilum* should be considered in the differential diagnosis of any immunosuppressed patient with chronic, recurrent or unexplained skin lesions on the extremities.
- ▶ Awareness of *M haemophilum* by clinicians and laboratory workers and a high index of suspicion are essential as to not miss this diagnosis.
- ▶ *M haemophilum* treatment requires an extended course of a multidrug regimen with no optimal duration recommended that may need to be adjusted using the judgment of the clinician based on their particular patient's medical comorbidities.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kelley CF, Armstrong WS, Eaton ME. Disseminated *Mycobacterium haemophilum* infection. *Lancet Infect Dis* 2011;11:571–8.
- 2 LaBombardi VJ, Nord JA. Clinical and laboratory aspects of *Mycobacterium haemophilum* infections. *Rev Med Microbiol* 1998;9:49–54.
- 3 Saubolle MA, Kiehn TE, White MH, et al. *Mycobacterium haemophilum*: microbiology and expanding clinical and geographic spectra of disease in humans. *Clin Microbiol Rev* 1996;9:435–47.
- 4 Swart RM, Van Ingen J, Van Soolingen D, et al. Nontuberculous mycobacteria infection and tumor necrosis factor- α antagonists. *Emerg Infect Dis* [Case Reports Letter] 2009;15:1700–1.
- 5 Sompolsky D, Lagziel A, Naveh D, et al. *Mycobacterium haemophilum* Sp-Nov, a new pathogen of humans. *Int J Syst Bacteriol* 1978;28:67–75.
- 6 Winthrop KL, Chang E, Yamashita S, et al. Nontuberculous mycobacteria infections and anti-tumor necrosis factor- α therapy. *Emerg Infect Dis* [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S. Review]. 2009;15:1556–61.
- 7 Shah MK, Sebt A, Kiehn TE, et al. *Mycobacterium haemophilum* in immunocompromised patients. *Clin Infect Dis* 2001;33:330–7.
- 8 Lindeboom JA, Van Coppenraet L, Van Soolingen D, et al. Clinical manifestations, diagnosis, and treatment of *Mycobacterium haemophilum* infections. *Clin Microbiol Rev* 2011;24:701.
- 9 Griffith DE, Aksamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med* 2007;175:367–416.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow