Published in final edited form as:

J Am Soc Mass Spectrom. 2012 August ; 23(8): 1351–1363. doi:10.1007/s13361-012-0409-8.

Cascade Dissociations of Peptide Cation-Radicals. Part2. Infrared Multiphoton Dissociation and Mechanistic Studies of z- Ions from Pentapeptides

Aaron R. Ledvina^a, Thomas W. Chung^c, Renjie Hui^d, Joshua J. Coon^{a,b}, and Frantisek Turecek^c

^aDepartment of Chemistry, University of Wisconsin, Madison, WI, USA

^bDepartment of Biomolecular Chemistry, University of Wisconsin, Madison, WI, USA

^cDepartment of Chemistry, Bagley Hall, Box 351700, University of Washington, Seattle, WA 98195-1700, USA

^dDCMR, Ecole Polytechnique, Palaiseau, Cedex, France

Abstract

Dissociations of z_4 ions from pentapeptides AAXAR, where X = H, Y, F, W, and V, produce dominant z_2 ions that account for >50% of the fragment ion intensity. The dissociation has been studied in detail by experiment and theory and found to involve several isomerization and bond-breaking steps. Isomerizations in z_4 ions proceed by amide transcis rotations followed by radical-induced transfer of a β -hydrogen atom from the side chain, forming stable C_β radical intermediates. These undergo rate-determining cleavage of the C_α —CO bond at the X residue followed by loss of the neutral AX fragment, forming x_2 intermediates. The latter were detected by energy-resolved resonant excitation collision-activated dissociation (CAD) and infrared multiphoton dissociation (IRMPD) experiments. The x_2 intermediates undergo facile loss of HNCO to form z_2 fragment ions, as also confirmed by energy-resolved CAD and IRMPD MS⁴ experiments. The loss of HNCO from the x_2 ion from AAHWR is kinetically hampered by the Trp residue that traps the OCNH radical group in a cyclic intermediate.

Introduction

Dissociations of fragment ions from electron transfer dissociation (ETD) of peptide cations continue to attract considerable attention because of their potential for providing peptide primary sequence information.^{a1} In the preceding report^{a2} we analyzed resonant excitation collision-activated dissociation (CAD) mass spectra of z-type ions produced by ETD of a set of over 40 penta-, nona- and decapeptides of the tryptic type that were C-terminated with arginine and contained different amino acid residues in the peptide chain. Upon collisional activation, the z ions underwent radical-induced dissociations that depended on the nature of the side chains in the amino acid residues. In particular, z_n ions with amino acid residues that contained benzylic (His, Phe, Tyr, Trp) or tertiary (Val) hydrogens in the C_β positions showed enhanced backbone dissociations in positions shifted two or three amino acid residues toward the C-terminus, forming abundant z_{n-2} or z_{n-3} ions in the MS³ mass spectra.^{a2} This type of dissociation was thought to be analogous to cascade dissociations of peptide ions upon electron capture^{a3} that can be driven by the substantial energy associated with electronion recombination. In keeping with previous studies^{a4–a6} we adopted the

proposed mechanism of backbone dissociations, involving radical-induced hydrogen migration from the side chain forming intermediate radical **I** (Scheme 1). According to Scheme 1, a transfer of a C β hydrogen atom activates the adjacent C α —CO bond in **I** for dissociation forming an **x** intermediate which eliminates HNCO forming the truncated **z_{n-2}** or **z_{n-3}** fragment ion.

Although this multistep mechanism is chemically plausible, it raises several questions not addressed by previous survey-like studies. First, the Scheme 1 mechanism involves a hydrogen atom migration forming intermediate **I**, raising the question of the pertinent transition state (TS) energies and relative stabilities of the reactants and the (**I**) intermediate. A second question is the relative stability of the **x** ions and the TS for the HNCO loss. Finally, to assess the overall kinetics of these cascade dissociations, it is necessary to compare the reaction flux through the transition states and obtain the pertinent rate constants that allow one to pinpoint the rate determining step.

In this Part 2 report we address these questions by experiment and theory. **z**-Type ions from AAHWR were studied using energy-resolved collision-activated and infrared multiphoton dissociation to assess the energetics and kinetics of the consecutive dissociations. Electronic structure theory calculations using two different density functionals and a perturbational Moller-Plesset theory were used to locate transition states and obtain activation energies for the isomerizations and dissociations of **z₄** ions from AAHAR, AAHWR, and AAYAR. The energy data were used for Rice-Ramsperger-Kassel-Marcus calculations of unimolecular rate constants to assess the branching ratios for the dissociations.

Experimental

Materials and Methods

The peptides were custom-synthesized by NEOPeptide Laboratories (Cambridge, MA) or CHI Scientific (Maynard, MA) and used as received. ETD mass spectra were measured on a dual-cell quadrupole linear ion trap (QLT) outfitted with a chemical ionization source for the generation of fluoranthene anion radicals for use as an ETD reagent (LTQ XL, Thermo Fisher Scientific, San Jose, CA, USA). Selected measurements were carried out using a hybrid mass spectrometer comprising both ion trap (QLT) and orbitrap mass analyzers and likewise equipped with a chemical ionization source for the generation of fluoranthene radicals for use as an ETD reagent, (LTQ Orbitrap Velos, Thermo Fisher Scientific, San Jose, CA, USA). Peptide solutions (10 μ M) in 50/50/1 water/methanol/acetic acid were electrosprayed to form precursor cations for ETD. Electrospray ionization of the pentapeptides produced mainly doubly protonated ($M + 2H$)²⁺ ions and minor singly protonated ($M + H$)⁺ ions, typically in a 10:1 ratio.

IRMPD and energy-resolved CAD experiments were carried out on a modified dual cell QLT outfitted with a CI source for the generation of fluoranthene anion radicals as described previously.^{a7} The mass spectrometer was modified to allow for the introduction of IR photons to the ion trapping region of the QLT. We inserted a ZnSe window on the rear flange of the instrument to transmit IR photons through a stainless steel aperture (diameter = 0.065") concentric with the trapping volume of the QLT. To allow the passage of IR photons to the dual cell QLT, we the ion volume of the CI source with the addition of a hole (diameter = 0.070") in the section of the ion volume farthest from the ion trap and the enlargement of the hole in the section of the ion volume closest to the QLT, both also concentric with the QLT trapping volume. IR photons were generated using a Firestar T-100 Synrad 120-W CO₂ continuous wave laser (Mukilteo, WA). Doubly protonated peptide precursor ions were produced by electrospray and subjected to ETD (~70 ms reaction with 5×10^5 fluoranthene radical anions) to form **z₄** ions. The **z₄** ions were re-isolated and activated

using resonant-excitation collisional activation (CID) within the high-pressure QLT (bath gas is He). For the energy-resolved experiments, CAD normalized collision energies (NCE) of 0 to 50 correspond roughly to supplemental AC excitation voltages of 0 to ~2.5 V. Infrared multiphoton dissociation was carried out by operating the laser in a mode whereby the laser was externally triggered *via* a TTL pulse from pin 14 of the J1 connector. Laser power was set as indicated within the text.

Calculations

All calculations were performed using the Gaussain 09 suite of programs.^{c1} Initial guesses for the *z* ion structures were taken from the optimized peptide cation-radicals under the assumption that the fragments preserve the local hydrogen bonding patterns in the charge-reduced ions.^{cx} Geometries were optimized with density functional theory calculations using the hybrid B3LYP functional^{c2-c4} in the spin-unrestricted format and the 6-31+G(d, p) basis set. The optimized structures were confirmed as local energy minima or first-order saddle points by harmonic frequency calculations. Three sets of single point energies were calculated using the larger 6-311++G(2d, p) basis set and B3LYP and M06-2X functionals.^{c5} The latter functional has been claimed to give improved transition state energies for kinetic calculations. In addition, Moller-Pleset theory,^{c6} UMP2(frozen core), with the 6-311++G(2d, p) basis set was used to obtain single point energies, which were corrected for contribution of higher spin state by the standard spin annihilation procedure.^{c7, c8} The UB3LYP and PMP2 single-point energies were averaged (B3-PMP2) to provide improved relative energies for cation-radicals, as reported previously.^{c9-c11} Unimolecular rate constants were calculated for reactions occurring on the calculated B3-PMP2 and M06-2X potential energy surface, including zero-point vibrational energy corrections, within the formalism of the Rice-Ramsperger-Kassel-Marcus (RRKM) theory.^{c12} These RRKM calculations used Hase's program^{c13} that was modified to deal with larger molecular systems with up to 1000 atoms, and recompiled for Windows XP^{c14} and Windows 7. The RRKM rate constants were obtained by direct count of quantum states at internal energies that were increased in 2 kJ mol⁻¹ steps from the transition state up to 400 kJ mol⁻¹ above the reactant. Rotations were treated adiabatically, and the calculated microscopic rate constants $k(E, J, K)$ were then Boltzmann-averaged over the thermal distribution of rotational states at 298 K.

Results

Consecutive Dissociations of *z*₄ Ions

We first present and discuss the features of *z* ion dissociations as observed in the MS³ mass spectra. The *z*₄ and *z*₅ fragment ions produced by ETD of (AAXAR + 2H)²⁺ (X = F, Y, H, W, and V) (Figure 1A) undergo collision-activated dissociations producing dominant *z*₂ ions that amounted to 15–65% of the total fragment intensity in the MS³ mass spectra (Figure 1B).^{a2} A peculiar feature of these dissociations, when starting from *z*₄ ions, was the presence of minor *x*₂ peaks (at 8–10% relative to the *z*₂ ion intensity) whose *m/z* value was reduced by 0.2 units from the expected *m/z* 273.1 nominal value. This phenomenon was observed for MS³ dissociations of all these *z*₄ ions, as illustrated for AAHAR (Figure 1B), AAYAR, AAFAR, and AAVAR (Figure S1, Supplementary Data) and reported previously for AAWAR.^{a2} We speculated that the *x*₂ ion was metastable and dissociated during ejection from the linear ion trap, so that its detection was slightly offset during the scan, resulting in a lower apparent *m/z* value. This was consistent with the CAD mass spectra obtained at longer detection times on an Orbitrap instrument which showed the *x*₂ peak at the correct *m/z* 273.1427 but at a substantially reduced relative intensity (0.9% relative to the *z*₂ ion at *m/z* 230.1370, Figure S2, Supplementary Data). The difference in the exact masses, $\Delta m = 43.0057$ Da, fits the theoretical mass of HNCO (43.0058 Da).

In contrast to CAD of z_4 ions from ETD of the F, Y, H, W, and V-containing AAXAR peptides, the z_4 ions from AAHWR (m/z 553) showed an MS^3 spectrum that contained both the x_2 and z_2 ions at correct m/z values (388.1844 and 345.1786, respectively) and with substantial relative intensities (Figure 2). This enabled the re-isolation of the x_2 ion at m/z 388 and investigation of its dissociations in MS^4 experiments and also to study the consecutive dissociations of the z_4 ions as a function of the excitation energy.

The energy-resolved MS^3 mass spectra of the z_4 ion were measured at nominal laboratory collision energies (NCE) ranging from 0 to 30 eV. The energy-dependent relative intensities of the z_4 , x_2 , and z_2 ions are shown in Figure 3. The graph shows different onsets for the formation of the x_2 ion at NCE = 11 eV and for the z_2 ion at NCE ~ 14 eV. At NCE = 15 eV, the dissociation mainly produces the x_2 ions at an intensity ratio of $[x_2]/[z_2] = 8.8$. This ratio drops to 1.7 at NCE = 30 eV and further decreases at higher collision energies. These results are consistent with the consecutive nature of the z_2 ion formation, whereby the internal energy of the x_2 intermediate increases at higher excitations of the z_4 parent ion to drive the loss of HNCO.

The results were even more dramatic for IRMPD which non-selectively deposits internal energy to all absorbing ions in the ion trap rather than just the mass-selected z_4 precursor ion. The relative intensities of the z_4 , x_2 , and z_2 ions were monitored at three laser power levels (36, 48, and 60 W) and at variable irradiation times ranging from 0 to 50 ms. Figure 4A–C illustrates the time-dependent relative intensities at the 36, 48, and 60 W power setting. The x_2 ion shows an early onset which is followed by that of the z_2 ion. The ion intensities reach maxima at different irradiation times, which were ca. 15 ms for the x_2 ion and ca. 25 ms for the z_2 ion at 48 W (Figure 4B). Control experiments were performed to assess the photo-dissociative depletion of the x_2 and z_2 ions. In this measurement, the ions were independently generated by MS^3 -CAD and irradiated with the CO_2 laser at 60 W. Figure S3 (Supplement) shows very similar depletion curves which indicate that the measured relative intensities are not strongly affected by secondary dissociations. This result is consistent with the calculated IR spectra of the x_2 and z_2 ions which show only weak absorption bands in the region of CO_2 laser wavelengths (900–1100 cm^{-1} , Figure S4).

The CAD and IRMPD MS^4 mass spectra of mass-selected x_2 ions at m/z 388 showed a dominant dissociation by loss of HNCO forming the z_2 ion at m/z 345 (Figure S5). The other fragments in the CAD and IRMPD spectra in Figure S5, appearing at m/z 301, 300, 257, 230, 200, and 158, were mainly produced by consecutive dissociations of the z_2 ion. This was corroborated by the IRMPD spectrum of mass-selected m/z 345 ions which gave the same fragments (Figure S6) as those in the MS^3 spectra of the x_2 ions.

Mechanistic Considerations for the Consecutive $z_4 \rightarrow z_2$ Dissociations

The experimental data showed different dissociations of z_4 ions from the AAXAR (X = H, F, Y, W, V) and AAHWR peptides in that the former proceeded rapidly to z_2 fragments, whereas the latter formed an abundant x_2 intermediate which needed further activation to eliminate HNCO. To rationalize these observations we performed detailed calculations exploring the potential energy surfaces along several reaction paths in z_4 ions from AAHAR and AAYAR and the x_2 ions from AAHWR. The z_4 ion from AAHAR is used here as a case in point. The relative and dissociation energies were obtained at several levels of theory including two density functionals and a perturbational Møller-Plesset (MP2, frozen core) treatment. The energies calculated at all levels of theory are shown in Table 1. The data used in text refer to the averaged B3LYP and spin-projected MP2 (B3-PMP2) values which provided relative and transition state energies at the chemical accuracy limit (± 4 kJ mol $^{-1}$) for a number of radical systems.^{c10,c11} The B3-PMP2 and M06-2X relative energies are also

plotted in the comprehensive potential energy diagram, collating several pathways we investigated (Figure 5). The energy data and ion structures for the AAYAR system are given in Table S1 and Scheme S1 of the Supplement.

Pathways to C $_{\alpha}$ —CO Bond Dissociation in AAHAR

The trans-Pathway—To form an x_2 intermediate, the z_4 ion from AAHAR must break the C $_{\alpha}$ —CO bond at the His residue. Activation of this bond can involve the formation of a C $_{\beta}$ radical at the His residue by β -hydrogen atom transfer to the original C $_{\alpha}$ radical site in the z_4 reactant as postulated previously^{a4,a5} and analyzed in detail for a z_4 ion from AHDAL.^{a6} Inspection of the z_4 ion structure shows that the His β -hydrogen atom transfer is sterically impossible in the amide all-*trans* conformer (**1**) which must first undergo a *trans* $\rightarrow$ *cis* rotation of the terminal Ala amide group (Scheme 2).^{a6} The relevant transition state for the rotation (**TS1**) was calculated to be 36 kJ mol⁻¹ above **1** and led to *cis*-amide isomer **2**. Note that the **TS1** energy is substantially lower than typical activation energies for amide group rotations in peptides (80–100 kJ mol⁻¹)^{r1-r3}. This lowering of the amide rotational barrier in z ions is caused by the conjugation of the C $_{\alpha}$ radical with the adjacent amide group that results in the weakening of the CO—NH double bond character, as discussed previously.^{a6} The *cis* isomer **2** is comparably stable as **1** (Table 1). The His H $_{\beta}$ migration in **2** requires 69 kJ mol⁻¹ in **TS2**, forming a His C $_{\beta}$ radical **3**. Radical **3** is more stable than **1** and **2** at all levels of theory. It can gain further stabilization by *cis* $\rightarrow$ *trans* rotation of the terminal Ala amide group, forming a *trans* isomer (**4**) at -38 kJ mol⁻¹ relative to **1**. The amide rotation in **3** forming **4** proceeds through **TS3** which is 90 kJ mol⁻¹ above **4**. Note that the CO—NH bond of the rotating amide group in **3** is not weakened by non-conjugated the C $_{\beta}$ radical, and therefore the **TS3** energy is close to values typical for peptides.^{r1-r3}

In contrast, the C $_{\beta}$ radical in **4** weakens the adjacent C $_{\alpha}$ —CO bond which can dissociate through **TS4** which is 124 kJ mol⁻¹ above **4** and 86 kJ mol⁻¹ above **1** (Scheme 3). Dissociation of the C $_{\alpha}$ —CO bond forms a complex (**5**) of the incipient x_2 ion (**6**) and a truncated neutral Ala-His fragment (**7**). Complex **5** was slightly less stable than **1**. However, since we did not study more conformers of **5**, we cannot exclude the existence of other, lower energy structures. The important feature is that the dissociation of **5** to **6** and **7**, which requires the incipient fragment separation but no covalent bond cleavage, is substantially endothermic. This makes the thermochemical threshold for these fragments the highest point on the potential energy surface of the above-described *trans* pathway.

The cis-Pathway—We considered several alternatives to the *trans*-pathway. One obvious alternative is for the *cis*- β -radical **3** to undergo a C $_{\alpha}$ —CO bond cleavage to form an ion molecule complex (**8**) of the incipient *cis*-Ala-His neutral fragment and the x_2 ion (**6**) (Scheme 3). The relevant **TS5** (108 kJ mol⁻¹ relative to **3**) requires a substantially higher energy than does **TS3** for the competing amide rotation in **3** (63 kJ mol⁻¹ relative to **3**), see Table 1. Likewise, the *cis*-complex **8** is 11 kJ mol⁻¹ above the *trans*-complex **5**. However, the dissociation threshold for the formation of the *cis*-Ala-His neutral fragment (**9**) and **6** is somewhat lower (124 kJ mol⁻¹ relative to **1**) than for the *trans* fragment **7** (128 kJ mol⁻¹ relative to **1**). Thus, if the fragment separation in complexes **5** and **8** is the rate determining step for the formation of the x_2 ion **6**, the *cis*-pathway is kinetically preferred.

The *cis*-pathway, which was also studied for the isomerizations and dissociations of the z_4 ion from AAYAR (**y1**, Scheme S1), showed some noteworthy similarities and differences from the z_4 ion from AAHAR. The migration of the tyrosine benzylic hydrogen ion in the *cis*- z_4 ion (**y2**) to form the benzylic radical **y3** required 82 kJ mol⁻¹ in **TS1y** (Table S1) which was only slightly higher than the **TS2** energy (Table 1). However, the relative energy of the benzylic radical **y3** was higher than that of the histidine-C $_{\beta}$ radical **3**. The C $_{\alpha}$ —CO

bond cleavage in **y3** can proceed through **TS2y** which is at 111 kJ mol⁻¹ relative to **y1** and thus higher than the analogous **TS5** from AAHAR. The largest difference between the two systems was in the relative energies of the complexes, **y4** and **8**, where the former was substantially less stable than **y1**. This can be attributed to the different type of non-covalent bonding in these complexes. Complexes **5** and **8** gain stabilization by hydrogen bonding between the histidine imidazole ring and the amide hydrogen of the dangling carbamoyl group (Scheme 3). No such interaction occurs between the tyrosine phenol group and the incipient **x2** fragment in complex **t4** (Scheme S1).

The Enol Pathway—Another alternative pathway we addressed differs from the previous two in that the first step, the His β -H transfer, proceeded to the oxygen of the terminal amide group rather than to the C α position (Scheme S2). The amide group in **1** is conjugated with the C α radical and displays non-negligible spin density at the oxygen atom (11%) which presumably can make it an H-atom receptor in a radical driven migration. Note that the His β -H transfer to the neighboring amide oxygen can proceed in **1** without a *trans* $\rightarrow$ *cis* amide rotation. However, the **TS6** energy (127 kJ mol⁻¹) is higher than that of **TS2**, and the enol intermediate **10** is substantially less stable than **1**. This indicates that the enol pathway would have a reversible first step favoring a reverse **10** $\rightarrow$ **1** hydrogen migration, which would adversely affect the kinetics of the overall dissociation. Perhaps the most serious factor disfavoring the enol pathway is the high threshold energy for the formation of an enol form of the Ala-His neutral fragment (**11**) at 187 kJ mol⁻¹ relative to **1** (Table 1).

A common feature of the above-mentioned pathways is the formation of the **x2** intermediate (**6**) which is experimentally observed as a low-intensity metastable *m/z* 273 fragment ion. Its further dissociation by loss of O=C=NH is calculated to be slightly exothermic (Table 1), but requires additional 70 kJ mol⁻¹ in **TS7** and involves a stable ion-molecule complex (**12**, Scheme 3).

The Cyclization Cascade Pathway

The previous study of cascade dissociations in ECD by O'Connor and coworkers^{a3} proposed radical-driven cyclization reactions to create new radical centers in the peptide fragments that would activate adjacent bonds for further dissociations. We analyzed similar mechanisms for the dissociation of the **z4** ion from AAHAR, as shown in Scheme 4. A cyclization reaction between the Ala C α radical and the His amide group must involve amide *trans* $\rightarrow$ *cis* rotation through **TS1** which is a low energy process (Scheme 2). The *cis*-amide **2** can attack the His amide oxygen atom in **TS8** forming a cyclic aminoketyl radical (**13**) which is substantially less stable (+97 kJ mol⁻¹) than **2** (Table 1). The cyclic radical **13** can undergo competitive ring opening back to **2** or proceed by N—C α bond cleavage through **TS9**, yielding directly the **z2** fragment ion and the cyclic neutral fragment **14**. We also investigated the possibility of a C α radical attack at the amide nitrogen atom followed by N—C α bond cleavage (Scheme 4). However, the pertinent **TS10** for the radical attack at the amide nitrogen required a very high energy (Table 1).

Dissociation Kinetics—The efficiency of the cascade dissociation pathways depends on the forward and reverse reaction flux through the pertinent transition states as quantified by the respective rate constants. The rate constants were calculated using RRKM and the B3-PMP2 TS energies and used to express the molar fractions of **z2** fragment ions produced on the time scale of the experiment which was chosen as 300 ms. The calculated rate constants pertinent to the formation of the **x2** intermediate in the *cis* pathway (log *k*, s⁻¹) are shown in Figure 6A. As expected, the amide *cis*-*trans* rotations are fast (*k*₁, *k*₋₁), leading to rapid equilibration of **1** and **2**. The rate constants for the reversible H β migration through **TS2** (*k*₂, *k*₋₂) favor the formation of the C β radical **3**. The rate determining steps are the C α —CO

bond cleavage through **TS5** (k_3) and the subsequent separation of the incipient products in complex **8** forming the x_2 fragment ion. The consecutive loss of HNCO from the x_2 ion is fast and shows a rapid rise of the pertinent rate constant (k_4) with the x_2 internal energy.

The cyclization pathway shows different kinetics (Figure 6B). The rate determining step (k_5) is the cyclization through **TS8** forming intermediate **13**. The latter can undergo competitive C_α —CO bond cleavage through **TS9** to directly produce the z_2 fragment (k_6) or undergo reverse ring opening (k_{-5}). The k_{-5} and k_6 rate constants indicate that the reaction branching in **13** should favor the ring opening to cis-**z4**, not fragmentation to z_2 .

The overall reaction flux on the 300 ms time scale is illustrated by the calculated molar fractions of **6**, and the z_2 fragments produced by the Scheme 2 (cis) and Scheme 4 (cyclization) pathways (Figure 6C). Note that while the formation of **8** was treated rigorously by solving the complex kinetics of reversible, competitive, and consecutive reactions, the rate constant for the subsequent fragment separation, $8 \rightarrow 6 + 9$, was not calculated, because this dissociation is likely to be continuously endothermic without a well-defined saddle point on the potential energy surface. We estimated the pertinent rate constant as being equal to k_3 (Figure 6A), because both the $3 \rightarrow 8$ and $8 \rightarrow 6 + 9$ reactions are simple dissociations requiring similar energies (Table 1). A reverse fragment recombination in the complex, $8 \rightarrow 3$, was considered slow because of entropy loss in the transition state. Under these assumptions, the overall formation of the x_2 intermediate showed a substantial kinetic shift (Figure 6C, *blue curve*), reaching 50% conversion to **6** at 223 kJ mol⁻¹ internal energy, which was 109 kJ mol⁻¹ above the thermochemical threshold. The dissociation was nearly complete (>99%) at >242 kJ mol⁻¹ internal energy. We presumed that fragments **6** and **9** formed from **8** equi-partitioned the available non-fixed internal energy according to the number of their vibrational degrees of freedom (108/177 or 61% for **6**). For example, **1** having 223 kJ mol⁻¹ of internal energy and dissociating at 50% efficiency is estimated to form the intermediate **6** with $(223 - 114) \times 0.61 = 66$ kJ mol⁻¹ internal energy and likewise for dissociations of **1** having higher internal energies. Vibrational excitation in **6** exceeding the 70 kJ mol⁻¹, which was required to reach **TS7**, was calculated to lead to a rapid dissociation by loss of HNCO which showed a steep increase of the rate constant with internal energy (k_4 , Figure 6A). Hence, intermediate **6** formed by dissociation of **1** having >260 kJ mol⁻¹ internal energy was calculated to undergo nearly 100% loss of HNCO on the 300 ms time scale forming the z_2 fragment ion (Figure 6C, *red curve*). The narrow interval of kinetic metastability for **6** (ca. 240–261 kJ mol⁻¹) was consistent with the low relative intensity of the x_2 (m/z 273) ions in the MS³ mass spectrum. In other words, most **z4** ions that acquired excessive internal energy to dissociate on the time-scale of the experiment produced x_2 intermediates with internal energies that were sufficient to drive their fast subsequent dissociation by loss of HNCO.

The formation of the z_2 fragment ion through the cyclization pathway displays a very large kinetic shift, reaching 50% conversion at internal energies >400 kJ mol⁻¹ (Figure 6C *purple curve*). This indicates that the cyclization pathway may not be kinetically competitive. We note that although the calculated rate constants strongly depend on the quality of the potential energy surface, the conclusion regarding the preference of the *cis* over the cyclization pathway also makes chemical sense. Schemes 1 and 2 show that the dissociation is facilitated by the C β -H transfer which explains the activating role of the His, Phe, Tyr, Trp and Val residues. In contrast, the role of the His side-chain group in the Scheme 4 mechanism appears to be one of a spectator with no obvious involvement in the rate-determining ring cyclization.

The Tryptophan Effect on Dissociations of z_4 Ions from AAHWR—Dissociation of this z_4 ion produced an unusually stable x_2 intermediate at m/z 388 (Figure 2). We

reasoned that this effect could be due to the radical-trapping properties of the indole ring^{r4-r6} and investigated the potential energy surface in the pertinent x_2 cation-radical (**w1**). The optimized structures are shown in Scheme 5, the relative energies are given in Table 2. The dissociation by loss of HNCO from **w1** was found to have a **TSw1** energy (75 kJ mol⁻¹, Table 2) that was nearly identical to that for the HNCO loss of from the x_2 ion from AAHAR (**TS7**, Table 1). Thus, the effect of the Trp residue on **TSw1** was weak. However, the optimized structure of **w1** showed that the O=C-NH radical group can reach to the C-2', C-3' and C-10' positions of the indole ring for radical-driven cyclizations. The TS energy for cyclization to C-2' was calculated as 54 kJ mol⁻¹ (**TSw2**) to give the most stable radical **w2** which was -24 kJ mol⁻¹ relative to **w1** (Table 2, Scheme 4). Cyclizations at C-3' and C-10' analogously gave radicals **w3** and **w4**, respectively, which, however, were comparably stable as **w1** (Table 2). This order of tryptophan radical stabilities is quite analogous to those reported for H-atom adducts which also preferred a C-2' adduct as the lowest energy isomer.^{r6}

The existence of the low energy isomer **w2** and **TSw2** for cyclization have a pronounced effect on the dissociation kinetics of the x_2 ion. RRKM calculations on the B3-PMP2 and M06-2X potential energy surfaces (Figure 7, panels A and C, respectively) show that the cyclization in **w1** (k_7) forming **w2** was faster than both the loss of HNCO (k_8) and the ring opening in **w2** (k_{-7}) to reform **w1**. Solving the kinetics for the competitive and consecutive reactions gave the breakdown diagrams for m/z 388 ions which are plotted as combined molar fractions of **w1** and **w2** as a function of the internal energy of **w1** at reaction times ranging from 50 μ s to 300 ms. (Figure 7, panels B and D). The breakdown diagrams show substantial kinetic shifts for the depletion of **w1** and **w2** by loss of HNCO even at the 300 ms reaction time. The B3-PMP2 data (Figure 7B) indicate a kinetic shift of $\Delta E_{\text{kin}} = 85$ kJ mol⁻¹ for 50% depletion of the m/z 388 ion (**w1** + **w2**) after 300 ms. The M06-2X data (Figure 7D), which gave higher TS energies for the HNCO loss and ring opening in **w2** (Table 2), indicate a 150 kJ mol⁻¹ kinetic shift under the same conditions. This is to be compared with the HNCO loss from **6** which shows 50% dissociation over 300 ms at 86 kJ mol⁻¹ internal energy, giving a much smaller kinetic shift of 16 kJ mol⁻¹.

In summary, the reason for the lack of dissociation of **w1** is the competing radical cyclization to isomer **w2**. Ions **w1** and **w2** appear at the same m/z 388 and therefore are not distinguished in the mass spectrum. The calculated energy-dependent ratio of **w1/w2** molar fractions shows that the majority of the non-dissociating x_2 ions should be the **w2** isomer (Figure S7, Supplement).

Conclusions

Experimental and computational analysis of cascade dissociations of z_4 ions from pentapeptides AAHAR, AAYAR, and AAHWR indicate that these proceed with involvement of β -hydrogen atoms in the benzylic positions of the side chains. Cyclization mechanisms, analogous to those postulated previously, are not compatible with the dissociation kinetics for z_4 ions. Intermediates of the x_2 type were detected by IRMPD and energy-resolved CAD experiments in complete agreement with the kinetic analysis. The rate determining step in the dissociation is the C_α —CO bond dissociation at the activating residue (H, Y, F, V, W) in the z_4 ion. Loss of HNCO from the x_2 intermediate is fast, leaving a narrow energy window of metastability for the x_2 ions. The loss of HNCO is hampered by the Trp residue that functions as a kinetic trap for the OCNH radical group.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Research at University of Washington has been supported by the NSF (Grant CHE-01055132) and the work at Wisconsin was supported by National Institutes of Health (NIH) grant R01 GM080148.

References

- a1. Li W, Song C, Bailey DJ, Tseng GC, Coon JJ, Wysocki VH. Statistical Analysis of Electron Transfer Dissociation Pairwise Fragmentation Patterns. *Anal Chem*. 2011; 83:9540–9545. [PubMed: 22022956]
- a2. Chung TW, Renjie Hui R, Ledvina AR, Coon JJ, Turecek F. Cascade Dissociations of Peptide Cation-Radicals. Part 1. Scope and Effects of Amino Acid Residues in Penta-, Nona- and Decapeptides. *J Am Soc Mass Spectrom*. Apr.2012 submitted.
- a3. Leymarie N, Costello CE, O'Connor PB. Electron Capture Dissociation Initiates a Free Radical Reaction Cascade. *J Am Chem Soc*. 2003; 125:8949–8958. [PubMed: 12862492]
- a4. Liu J, Liang X, McLuckey SA. On the Value of Knowing a z. *Ion for What It Is*. *J Proteome Res*. 2008; 7:130–137. [PubMed: 17997514]
- a5. Sun Q, Nelson H, Stolz BM, Julian RR. Side Chain Chemistry Mediates Backbone Fragmentation in Hydrogen Deficient Peptide Radicals. *J Proteome Res*. 2009; 8:958–966. [PubMed: 19113886]
- a6. Chung TW, Turecek F. Backbone and Side-Chain Specific Dissociations of z Ions from Non-Tryptic Peptides. *J Am Soc Mass Spectrom*. 2010; 21:1279–1295. [PubMed: 20299240]
- a7. Ledvina AR, Beauchene NA, McAlister GC, Syka JEP, Schwartz JC, Griep-Raming J, Westphall MS, Coon JJ. Activated-Ion Electron Transfer Dissociation Improves the Ability of Electron Transfer Dissociation to Identify peptides in a Complex Mixture. *Anal Chem*. 2010; 82:10068–10074. [PubMed: 21062032]
- c1. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, GA.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, HP.; Izmaylov, AF.; Bloino, J.; Zheng, G.; Sonnenberg, JL.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, JA., Jr; Peralta, JE.; Ogliaro, F.; Bearpark, M.; Heyd, JJ.; Brothers, E.; Kudin, KN.; Staroverov, VN.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, JC.; Iyengar, SS.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, JM.; Klene, M.; Knox, JE.; Cross, JB.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, RE.; Yazyev, O.; Austin, AJ.; Cammi, R.; Pomelli, C.; Ochterski, JW.; Martin, RL.; Morokuma, K.; Zakrzewski, VG.; Voth, GA.; Salvador, P.; Dannenberg, JJ.; Dapprich, S.; Daniels, AD.; Farkas, O.; Foresman, JB.; Ortiz, JV.; Cioslowski, J.; Fox, DJ. Gaussian 09, Revision A.02. Gaussian, Inc; Wallingford CT: 2009.
- cx. Moss CL, Chamot-Rooke J, Brown J, Campuzano I, Richardson K, Williams J, Bush M, Bythell B, Paizs B, Turecek F. Assigning Structures to Gas-Phase Peptide Cations and Cation-Radicals. An Infrared Multiphoton Dissociation, Ion Mobility, Electron Transfer and Computational Study of a Histidine Peptide Ion. *J Phys Chem B*. 2012; 106:3445–3456. [PubMed: 22364440]
- c2. Becke AD. New Mixing of Hartree-Fock and Local Density-Functional Theories. *J Chem Phys*. 1993; 98:1372–1377.
- c3. Becke AD. Density Functional Thermochemistry. III. The Role of Exact Exchange. *J Chem Phys*. 1993; 98:5648–5652.
- c4. Stephens PJ, Devlin FJ, Chabalowski CF, Frisch MJ. Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields. *J Phys Chem*. 1994; 98:11623–11627.
- c5. Zhao Y, Truhlar DG. The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals. *Theor Chem Acc*. 2008; 120:215–241.
- c6. Møller C, Plesset MS. A Note on an Approximation Treatment for Many-Electron Systems. *Phys Rev*. 1934; 46:618–622.

- c7. Schlegel HB. Potential Energy Curves Using Unrestricted Moller-Plesset Perturbation Theory with Spin Annihilation. *J Chem Phys.* 1986; 84:4530.
- c8. Mayer I. Spin-Projected UHF Method. IV. Comparison of Potential Curves Given by Different One-Electron Methods. *Adv Quantum Chem.* 1980; 12:189.
- c9. Turek F. Proton Affinity of Dimethyl Sulfoxide and Relative Stabilities of C₂H₆OS Molecules and C₂H₇OS⁺ Ions. A Comparative G2(MP2) ab Initio and Density Functional Theory Study. *J Phys Chem A.* 1998; 102:4703.
- c10. Polášek M, Turek F. Hydrogen Atom Adducts to Nitrobenzene. Formation of the Phenylnitronic Radical in the Gas Phase and Energetics of Wheland Intermediates. *J Am Chem Soc.* 2000; 122:9511–9524.
- c11. Wolken JK, Yao C, Turek F, Polce MJ, Wesdemiotis C. Cytosine neutral molecules and cation-radicals in the gas-phase. structures, energetics, ion chemistry, and neutralization-reionization mass spectrometry. *Int J Mass Spectrom.* 2007; 267:30–42.
- c12. Gilbert, RG.; Smith, SC. *Theory of Unimolecular and Recombination Reactions.* Blackwell Scientific Publications; Oxford: 1990. p. 52-132.
- c13. Zhu, L.; Hase, WL. *Quantum Chemistry Program Exchange.* Indiana University; Bloomington: 1994. Program No. QCPE 644
- c14. Frank AJ, Sadílek M, Ferrier JG, Turek F. Sulfur Oxyacids and Radicals in the Gas Phase. A Variable-Time Neutralization-Photoexcitation-Reionization Mass Spectrometric and Ab Initio/RRKM Study. *J Am Chem Soc.* 1997; 119:12343.
- r1. Jhon JS, Kang YK. Imide Cis-Trans Isomerization of N-Acetyl-N'-methylprolineamide and Solvent Effects. *J Phys Chem A.* 1999; 103:5436–5439.
- r2. Shi T, Spain SM, Rabenstein DL. Unexpectedly Fast Cis-Trans Isomerization of Xaa-Pro Peptide Bonds in Disulfide-Constrained Cyclic Peptides. *J Am Chem Soc.* 2004; 126:790–796. [PubMed: 14733553]
- r3. Fischer G. Chemical aspects of peptide bond isomerization. *Chem Soc Rev.* 2000; 29:119–127.
- r4. Byers JH, Campbell JE, Knapp FH, Thissell JG. Radical aromatic substitution via atom-transfer addition. *Tetrahedron Lett.* 1999; 40:2677–2680.
- r5. Walden SE, Wheeler RA. Distinguishing features of indolyl radical and radical cation: implications for tryptophan radical studies in proteins. *J Phys Chem.* 1996; 100:1530–1535.
- r6. Gregersen JA, Turek F. Mass-Spectrometric and Computational Study of Tryptophan Radicals (Trp + H)[•] Produced by Collisional Electron Transfer to Protonated Tryptophan in the Gas Phase. *Phys Chem Chem Phys.* 2010; 12:13434–13447. [PubMed: 20830385]

Figure 1. (A) ETD mass spectrum of (AAHAR + 2H)²⁺ and (B) CID-MS³ mass spectrum of the *z*₄ ion at *m/z* 438.

Figure 2.
CID-MS³ mass spectrum of the m/z 553 z_4 ion from AAHWR.

Figure 3. Energy-resolved CID-MS³ mass spectrum of the m/z 553 z_4 ion from AAHWR.

Figure 4. IRMPD-MS³ mass spectra of the m/z 553 z_4 ion from AAHWR at the indicated laser power levels.

Figure 5.

Potential energy surface (kJ mol^{-1}) for isomerizations and dissociations of z_4 ions from AAHAR. The energies are from B3-PMP2/6-311++G(2d, p) single-point calculations and include zero-point corrections. The energies in parentheses are from single-point M06-2X/6-311++G(2d, p) calculations including zero-point corrections. The color-coding is as follows: Black lines: cis pathway; blue lines: trans pathway; Pink lines: enol pathway.

Figure 6.

RRKM kinetics of z_4 and x_2 ion isomerizations and dissociations for the AAHAR peptide. Panel A: RRKM rate constants ($\log k$, s^{-1}) for reactions: k_1 : $1 \rightarrow \text{TS1}$; k_{-1} : $2 \rightarrow \text{TS1}$; k_2 : $2 \rightarrow \text{TS2}$; k_{-2} : $3 \rightarrow \text{TS2}$; k_3 : $3 \rightarrow \text{TS5}$; k_4 : $6 \rightarrow \text{TS7}$. Panel B: RRKM rate constants for reactions: k_5 : $2 \rightarrow \text{TS8}$; k_{-5} : $13 \rightarrow \text{TS8}$; k_6 : $13 \rightarrow \text{TS9}$. Panel C: Energy-dependent molar fractions of x_2 intermediate (blue curve) and z_2 fragment ions (red curve) at 300 ms reaction time. The purple curve shows the molar fraction of z_2 ions formed at 300 ms by the cyclization cascade pathway.

Figure 7.

RRKM kinetics x_2 ion isomerizations and dissociations for the AAHWR peptide. Panels A and C: Rate constants for reactions: k_7 : $\mathbf{w1} \rightarrow \mathbf{TSw2}$; k_{-7} : $\mathbf{w2} \rightarrow \mathbf{TSw2}$; k_8 : $\mathbf{w1} \rightarrow \mathbf{TSw1}$, on the B3-PMMP2/6-311++G(2d, p) (A) and M06-2X/6-311++G(2d, p) (C) potential energy surfaces. Panels B (B3-PMP2) and D (M06-2X): Calculated breakdown curves for m/z 388 ($\mathbf{w1} + \mathbf{w2}$) ions at 50 μs , 1 ms, 10 ms, 100 ms, and 300 ms reaction times.

Scheme 1.

Previously proposed mechanism for β -hydrogen transfer triggering C_α —CO bond dissociation.

Scheme 2.

B3LYP/6-31+G(d, p) optimized structures of z_4 ions from AAHAR. The atoms are color-coded as follows: Green = C, blue = N, red = O, gray = H. Hydrogen bonds are denoted by double-ended blue arrows and distances are given in Ångströms.

Scheme 3.

B3LYP/6–31+G(d, p) optimized structures of reaction intermediates from AAHAR. The atoms are color-coded as in Scheme 2.

Scheme 4.

Structures of x_2 ion intermediates and transition states from AAHWR.

Table 1

Relative and dissociation energies of z ions from AAHAR.

Species/reaction	Relative Energy ^{a,b}				
	B3LYP	B3LYP ^c	PMP2 ^c	B3-PMP2 ^d	M06-2X ^c
	6-31+G(d, p)			6-311++G(2d, p)	
1 → TS1	45	44	27	36	28
1 → 2	12	12	-1	5	-0.4
2 → TS2	65	67	71	69	81
1 → 3	-10	-10	-9	-10	-11
3 → TS3	66	64	61	63	64
1 → 4	-47	-47	-28	-38	-31
4 → TS4	123	118	129	124	136
1 → 5	-4	-6	17	5	22
1 → 6 + 7	109	102	155	128	146
3 → TS5	104	101	115	108	120
1 → 8	6	2	31	16	36
1 → 6 + 9	97	89	140	114	132
1 → TS6	120	121	132	127	135
1 → 10	89	87	90	89	81
1 → 6 + 11	198	187	245	187	227
6 → z ₂ + O=C=NH	-18	-31	4	-13	4
6 → TS7	63	59	81	70	89
6 → 12	-44	-53	-23	-38	-22
1 → TS8	135	138	127	132	
1 → 13	105	109	86	97	
1 → TS9	141	143	133	138	
1 → TS10	206	210	179	194	

^aIn units of kJ mol⁻¹.

^bIncluding B3LYP/6-31+G(d, p) zero-point energies scaled by 0.963 and referring to 0 K.

^cFrom single point energy calculations on B3LYP/6-31+G(d, p) optimized geometries.

d From averaged B3LYP and spin-projected PMP2 single-point energies.

Table 2

Relative and dissociation energies of z ions from AAHWR.

Species/reaction	Relative Energy ^{a, b}					
	B3LYP			M06-2X ^c		
	6-31+G(d, p)	6-31+G(d, p)	6-31+G(d, p)	6-31+G(d, p)	6-31+G(d, p)	6-31+G(d, p)
w1 → TS1	69	65	85	75	93	93
w1 → TS2	46	48	61	54	54	54
w1 → w2	-39	-36	-12	-24	-38	-38
w1 → w3	7	10	11	10	-5	-5
w1 → w4	2	6	9	7	-4	-4

^aIn units of kJ mol⁻¹.

^bIncluding B3LYP/6-31+G(d, p) zero-point energies scaled by 0.963 and referring to 0 K.

^cFrom single point energy calculations on B3LYP/6-31+G(d, p) optimized geometries.

^dFrom averaged B3LYP and spin-projected PMP2 single-point energies.