Research

Cite this article: Refardt D, Bergmiller T, Kümmerli R. 2013 Altruism can evolve when relatedness is low: evidence from bacteria committing suicide upon phage infection. *Proc R Soc B* 280: 20123035.
<http://dx.doi.org/10.1098/rspb.2012.3035>

Received: 20 December 2012

Accepted: 21 February 2013

Subject Areas:

behaviour, evolution, microbiology

Keywords:

altruistic suicide, host defence, parasite transmission, structured population, Hamilton's rule, kin selection

Author for correspondence:

Dominik Refardt
e-mail: dominik.refardt@gmail.com

[†]Present address: Zürich University of Applied Sciences, Institute of Natural Resource Sciences, Grüental, 8820 Wädenswil, Switzerland.

Altruism can evolve when relatedness is low: evidence from bacteria committing suicide upon phage infection

Dominik Refardt^{1,†}, Tobias Bergmiller^{1,2} and Rolf Kümmerli^{3,4}

¹Department of Environmental Systems Science, Institute of Integrative Biology, Swiss Federal Institute of Technology, Universitätsstrasse 16, 8092 Zürich, Switzerland

²Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

³Environmental Microbiology, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133, 8600 Dübendorf, Switzerland

⁴Institute of Plant Biology, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

High relatedness among interacting individuals has generally been considered a precondition for the evolution of altruism. However, kin-selection theory also predicts the evolution of altruism when relatedness is low, as long as the cost of the altruistic act is minor compared with its benefit. Here, we demonstrate evidence for a low-cost altruistic act in bacteria. We investigated *Escherichia coli* responding to the attack of an obligately lytic phage by committing suicide in order to prevent parasite transmission to nearby relatives. We found that bacterial suicide provides large benefits to survivors at marginal costs to committers. The cost of suicide was low, because infected cells are moribund, rapidly dying upon phage infection, such that no more opportunity for reproduction remains. As a consequence of its marginal cost, host suicide was selectively favoured even when relatedness between committers and survivors approached zero. Altogether, our findings demonstrate that low-cost suicide can evolve with ease, represents an effective host-defence strategy, and seems to be widespread among microbes. Moreover, low-cost suicide might also occur in higher organisms as exemplified by infected social insect workers leaving the colony to die in isolation.

1. Introduction

Kin-selection theory [1] predicts that an altruistic behaviour is favoured if its fitness cost (c) to the actor is outweighed by its fitness benefit (b) to the recipient times the genetic relatedness (r at the cooperative trait) between the actor and the recipient ($rb > c$, Hamilton's rule). Consistent with theory, the importance of relatedness for the evolution of altruism has been demonstrated in numerous studies [2]. It most clearly emerged in comparative work showing that high relatedness was key for the evolution of reproductive altruism in social insects [3,4], birds [5] and mammals [6]. However, Hamilton's rule can also predict the evolution of altruism when relatedness is low, for example, in situations where the cost of altruism is minor compared with its benefit. In the extreme case where cost approaches zero, Hamilton's rule reduces to $rb > 0$, such that altruism can be favoured even when r tends towards zero.

Here, we demonstrate evidence of such a low-cost altruistic act in *Escherichia coli* bacteria, which express an abortive infection system (Abi) that responds to lethal phage attacks by causing infected cells to die together with the infecting phage. From an evolutionary perspective, the expression of an Abi system could reflect an altruistic act that allows infected bacteria to commit suicide in order to prevent parasite transmission to nearby relatives. Bacteria possess a number of Abi mechanisms, which usually involve joint actions by a prophage (a phage integrated into the bacterial genome) or a plasmid that encodes the genes for the abortive infection, and the bacterium, which transcribes and synthesizes the abortive machinery [7–11]. Important to note is that in many cases

the interests of the bacterium and an already resident prophage become completely aligned upon obligately lytic phage infection (i.e. infection by phages that do not insert into the bacterial genome, but kill the bacterium instantly), such that they can be regarded as one genetic entity [12].

Although Abi systems have long been considered by some to reflect altruistic acts [8], a rigorous test involving the measurement of parameters of Hamilton's rule is lacking. We accomplished this by measuring the benefit and cost of suicide, as well as the effect of relatedness on suicide in *E. coli* carrying the prophage λ (henceforth *E. coli* λ), which encodes the Rex abortive infection system [13]. The Rex system codes for an intracellular sensor (RexA) that is activated upon infection by the obligately lytic phage T4rII [14]. RexA then activates a membrane-anchored ion channel (RexB), which immediately results in the drop of the membrane potential and the cellular ATP levels, finally leading to the death of the bacterium, its prophage and the infecting phage [8]. To assess the benefit of the Rex system, we infected altruistic *E. coli* λ cells with phage T4rII in liquid-shaken medium and compared their fitness with an infected, non-altruistic *E. coli* strain (carrying the control prophage HK97, henceforth *E. coli* HK97). To measure intrinsic (maintenance) and extrinsic (inefficiency) costs of the Rex system, we competed *E. coli* λ against *E. coli* HK97 in the absence and presence of phages. In both types of competition assays, costs of suicide should manifest in *E. coli* HK97 outcompeting *E. coli* λ . Finally, we manipulated relatedness at the altruistic trait by competing *E. coli* λ against *E. coli* HK97 in medium containing different amounts of agarose. Bacterial dispersal is limited in more solid medium [15,16], which increases relatedness among interacting individuals (i.e. Rex-carriers interact more often among themselves than expected based on their relative frequency in the population).

2. Material and methods

(a) Strains

We used *E. coli* str. K-12 substr. MG1655 in all experiments. Two strains carrying different prophages were constructed by infection with either temperate phage λ PaPa or HK97 (henceforth referred to as *E. coli* λ and *E. coli* HK97). *Escherichia coli* λ expresses the suicide trait, whereas *E. coli* HK97 does not. Because both prophages belong to the same immunity group, they do not transmit in a co-culture, and hence the phenotypic difference between the two *E. coli* strains is reduced to the suicide trait. Both strains were marked with plasmids pGFP and pDsRed carrying genes for fluorescent proteins under an inducible promoter. Specifically, the GFPmut3 variant from plasmid pMA3 and a fast-maturing DsRed1 from pCLR7 were PCR-amplified with primers containing *Kpn*I and *Hind*III restriction sites (GFPmut3) or *Bam*HI *Hind*III restriction sites (DsRed), and cloned into pGDR11,2 a pQE31-based vector with an isopropyl β -D-1-thiogalactopyranoside (IPTG)-inducible promoter. Colonies that appeared bright green or red with 100 μ M IPTG were picked, and the insert was verified by digestion and PCR. Primer sequences (5'–3') for PCR verification were AGGTACCT AGTAAAGGAGAAGAACTTTT (5'-GFP), GATAAGCTTTTATT TGTAGAGCTCATCCA (3'-GFP), CGCGGATCCTGCATCCACC GAGGACGTCAT (5'-DsRed), GGGAAGCTTCTACAGGAACAG GTGGTGGC (3'-DsRed).

We used phage T4rII, a mutant strain of phage T4 that is susceptible to the Rex system of λ lysogens of *E. coli* [14]. The strain was plated on a lawn of *E. coli*, a single plaque was cored and

used to obtain a high-titer stock via plate lysis. The stock was stored in sodium, magnesium, gelatin buffer (0.1 M NaCl, 8 mM MgSO₄, 50 mM Tris–HCl pH 7.5, 0.01% gelatin) at 4°C. A drop of chloroform was added to prevent contamination by other micro-organisms.

(b) Experimental conditions

We carried out all experiments in lysogeny broth (10 g l^{−1} tryptone, 5 g l^{−1} yeast extract, 10 g l^{−1} NaCl; Sigma, Switzerland) supplemented with 100 μ g ml^{−1} ampicillin (Sigma) to prevent plasmid loss and 200 μ M IPTG (Sigma) to induce fluorescent proteins. Where indicated, agarose (Invitrogen) was added to impose population structure. All experiments were initiated using overnight cultures of *E. coli* λ and *E. coli* HK97.

To assess the benefit of host suicide, monocultures of *E. coli* λ and *E. coli* HK97 (5×10^6 cells) were infected with 0, 5, 50, 5×10^2 , 5×10^3 , 5×10^4 , 5×10^5 , 5×10^6 , 5×10^7 or 5×10^8 pfu of phage T4rII in 100 μ l medium in 96-well plates. All treatments were replicated four times; controls without T4rII were replicated eight times. Cultures were incubated in a plate reader (Molecular Devices, Sunnyvale, CA, USA) at 37°C and shaken at regular intervals (2 min), at which optical densities (600 nm) were measured. Here, we used strains carrying pGFP.

To estimate intrinsic (maintenance) costs of encoding the suicide system, we mixed *E. coli* λ and *E. coli* HK97 in a 50:50 ratio (5×10^5 cells in total) in 500 μ l medium in a 96-deep well plate in the absence of T4rII. Cultures were replicated 24 times for each marker combination, whereby each culture was based on an independent overnight culture. Cultures were incubated overnight at 37°C in an orbital shaker (400 r.p.m., a glass bead was added to each well to ensure proper mixing). Frequencies of both strains were measured before and after overnight growth using flow cytometry and used to calculate fitness of *E. coli* λ relative to *E. coli* HK97 for every replicate.

The Rex system might also incur extrinsic costs, for example because of mistakes (e.g. mortality rate upon T4rII infection might be lower than 100%, such that suicide is sometimes in vain resulting in the loss of direct reproduction). We estimated putative extrinsic costs of suicide in well-shaken medium. Here, the relatedness among interacting individuals is expected to be low, and any costs associated with suicide should result in the relative increase in *E. coli* HK97, which does not pay the cost, but still benefits from suicide (see [17,18] for theory). Specifically, we mixed *E. coli* λ and *E. coli* HK97 in various frequencies (5×10^5 cells in total) in 100 μ l medium in the presence (10^4 pfu, multiplicities of infection (MOI) = 0.02) or absence of T4rII. The volumetric mixing frequencies of *E. coli* λ were 10, 50, 90, 95, 99 and 99.9 per cent, which corresponded to the actual mixing frequencies of 18.6, 66.3, 95.7, 98.0, 99.1 and 99.8 per cent (carrying pGFP), and 11.5, 51.0, 90.5, 95.5, 98.4 and 99.5 per cent (carrying pDsRed) assessed by flow cytometry. All treatments were replicated four times for both marker combinations. Cultures were incubated for 8 h in a plate reader at 37°C and shaken in 2 min intervals, after which optical densities (600 nm) were measured. Final frequencies of both strains were measured using flow cytometry.

To measure the fitness consequences of host suicide in structured environments, we mixed strains (10^5 cells) in volumetric frequencies of 0.1 per cent and 50 per cent *E. coli* λ (actual frequencies assessed by flow cytometry were 0.38 per cent and 46.6 per cent, carrying pGFP; and 0.56 per cent and 43.0 per cent, carrying pDsRed) in 20 μ l medium. The mix was supplemented with 0, 10^3 , 10^4 , 10^5 , 10^6 and 10^7 pfu of phage T4rII, and then added on top of 1 ml medium containing 0.4 per cent agarose in 24-well plates. All treatments were replicated four times for both marker combinations. Cultures were incubated at 37°C overnight, followed by strain frequency estimation using flow cytometry.

To measure the fitness consequences of host suicide in environments with reduced population structure, the same 50 : 50 mix as above was used. Cells (10^5) in 20 μ l medium were supplemented with 0 or 10^4 pfu of phage T4rII and added into 1 ml medium containing 0, 0.025, 0.05, 0.1, 0.2 and 0.4 per cent agarose in 24-well plates. All treatments were replicated four times for both marker combinations. Cultures were incubated at 37 °C overnight in an orbital shaker (250 r.p.m.), followed by strain frequency estimation using flow cytometry. Because cultures grown with T4rII in 0 per cent agarose went extinct, no frequencies could be measured at this agarose concentration.

(c) Flow cytometry

Following competition, bacterial co-cultures were diluted in phosphate-buffered saline (Sigma), and 20 000 cells per sample were analysed (FACSCalibur, BD Biosciences, Franklin Lakes, NJ, USA). Samples were excited at a wavelength of 488 nm, whereas emission was detected at 510 (GFP) and 560 nm (DsRed). The use of GFP and DsRed is ideal here, because both markers can be excited with the same wavelength. Data were filtered by gating based on forward and side scatter, after which cells were counted based on fluorescence intensity using WEASEL v. 3.0.1 (Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia).

(d) Statistical analysis

All statistical analyses were carried out with R v. 2.15.0. [19]. Maximum doubling times were estimated by fitting a Gompertz growth model [20] to the monoculture growth data obtained during the first 180 min following incubation. Doubling times were compared across treatments with a one-way analysis of variance (ANOVA), including the dose of phage T4rII as a discrete explanatory variable. Post hoc pairwise comparisons were carried out with Tukey's honestly significant difference (HSD) test (95% confidence interval, CI).

We first calculated the Malthusian fitness (m) of *E. coli* λ as $m = \log([f_{E. coli \lambda \text{ end}} \cdot f_{E. coli HK97 \text{ start}}] / [f_{E. coli \lambda \text{ start}} \cdot f_{E. coli HK97 \text{ end}}])$ [21,22], where f with the respective subscript is the frequency of *E. coli* λ or *E. coli* HK97 at the beginning or the end of the competition in the same replicate. Fitness is thus based on the shift in strain frequencies that occurred during the time of competition, with fitness values of $m > 0$ or $m < 0$ signifying competitive advantages of *E. coli* λ or *E. coli* HK97, respectively.

Analysis of this first measure of fitness revealed significant effects of the marker plasmids in the absence of the phage. Strains with pGFP generally performed better than their competitors with pDsRed (average m of *E. coli* λ pGFP $\pm$ 95% CI: 1.06 ± 0.82 ; *E. coli* λ pDsRed: -0.95 ± 0.78 ; Student's t -test, $t_{46} = 17.2$, $p < 0.0001$), suggesting that pDsRed bears higher costs. However, this effect was consistent and independent of the strain itself: m of *E. coli* λ pGFP did not significantly differ from the absolute $|-m|$ of *E. coli* λ pDsRed (Student's t -test, $t_{46} = 0.95$, $p = 0.35$). To control for this marker effect in the presence of the phage, we used a modified formula for m , which compared the final frequencies of *E. coli* λ (carrying a given plasmid marker) between competitions with ($f_{E. coli \lambda \text{ treat}}$) and without ($f_{E. coli \lambda \text{ ctrl}}$) phage T4rII: $m = \log([f_{E. coli \lambda \text{ treat}} \cdot f_{E. coli HK97 \text{ ctrl}}] / [f_{E. coli \lambda \text{ ctrl}} \cdot f_{E. coli HK97 \text{ treat}}])$. Because replicates in treatment and control competitions were not paired, the final frequency in each treatment replicate was compared with the average final frequency across all control replicates.

We used Student's t -tests to compare fitness estimates or to test whether they were significantly different from 0. Furthermore, we fitted full factorial ANOVAs to fitness values (response variable), including the factors 'dose of T4rII' (continuous), 'initial frequency of *E. coli* λ ' (discrete), 'agarose concentration' (continuous) and 'marker colour' (discrete). Our data is deposited at Dryad: doi:10.5061/dryad.b1q2n.

Figure 1. (a) *Escherichia coli* HK97 populations without a suicidal abortive system are completely eradicated within a short period of time even when infected with very low numbers (indicated above lines) of phage T4rII. (b) *Escherichia coli* λ populations with the Rex abortive system show unhindered growth when infected with a MOI ≤ 0.1 , demonstrating the benefit of suicide in clonal populations. At a MOI ≈ 1 , growth was delayed, and at higher MOIs, no growth occurred because T4rII simultaneously infected all bacteria, such that no survivors could benefit from suicide. Growth curves are means of four replicates (control treatment eight replicates) and shaded areas represent s.e.

3. Results

(a) The benefit of host suicide

In liquid-shaken medium, we found that very low numbers of T4rII were sufficient to eradicate *E. coli* HK97 cultures (figure 1a). By contrast, *E. coli* λ exhibited unhindered growth that was not significantly different from an uninfected control culture (max. doubling time 24.65 ± 6.92 min; mean $\pm$ 95% CI) at MOI (the phages-to-bacteria ratio) < 1 (at MOI = 10^{-6} : 22.91 ± 3.39 , Tukey's HSD test, $p > 0.99$; 10^{-5} : 22.98 ± 6.23 , $p > 0.99$; 10^{-4} : 21.69 ± 2.34 , $p = 0.94$; 10^{-3} : 21.91 ± 3.38 , $p = 0.96$; 10^{-2} : 22.57 ± 7.30 , $p = 0.99$; 0.1: 26.93 ± 13.95 , $p = 0.98$; figure 1b). This demonstrates that host suicide clears the infection, thereby saving the clonal population from extinction. While at MOI = 1, growth was significantly delayed (44.74 ± 14.24 , $p < 0.0001$), we observed no growth at MOI > 1 , a pattern compatible with population extinction occurring when all bacterial cells commit suicide upon simultaneous infection with T4rII (figure 1b).

(b) The cost of host suicide

In the absence of T4rII and taking marker effects into account, we found no evidence that the frequency of *E. coli* λ

Figure 2. (a) In co-cultures where *Escherichia coli* λ was very common (>95%), suicide could eradicate T4rII (MOI = 0.02) infections. In one case (95.7%), co-cultures entered a growth-arrest phase during which *E. coli* λ significantly increased in frequency suggesting that suicide could be delayed or circumvented for unknown reasons. Co-cultures in which *E. coli* λ was less common (<95%) went extinct, showing that suicide cannot stop infection in well-mixed populations with a susceptible strain. Growth curves are means of four replicates, shaded areas represent standard errors, and colours indicate the type of marker plasmid used for *E. coli* λ (blue circles, pGFP; orange triangles, pDsRed). (b) In those cases where suicide could eradicate T4rII, *E. coli* HK97 successfully hitchhiked along with *E. coli* λ, but never experienced significant fitness benefits (*t*-tests, from left to right: $t_6 = 0$, $p = 1.00$; $t_6 = -13.38$, $p < 0.0001$; $t_6 = 1.43$, $p = 0.20$; $t_6 = -1.04$, $p = 0.34$; $t_6 = 0.73$, $p = 0.49$; $t_6 = 0.56$, $p = 0.59$; $t_6 = 1.39$, $p = 0.21$). This indicates that *E. coli* λ suicide bears negligible costs.

significantly changed during overnight competition against *E. coli* HK97 (average m of *E. coli* λ $\pm$ 95% CI = 0.11 ± 0.32 , $t_{46} = 0.95$, $p = 0.35$). This result indicates that intrinsic costs of maintaining the Rex system are negligible.

Our competition assays in the presence of T4rII also revealed no evidence for significant extrinsic costs. When *E. coli* λ was sufficiently frequent in a culture (>95%), its suicide cleared T4rII infections (figure 2a) and *E. coli* HK97 successfully hitchhiked along but never significantly increased in frequency (figure 2b). At an *E. coli* λ frequency of approximately 95 per cent, cultures entered a growth-arrest phase. In one of these cases (*E. coli* λ pGFP), *E. coli* λ significantly increased in frequency (figure 2b). This finding is compatible with previous observations that under certain conditions the Rex system induces dormancy instead of suicide [23,24] (see §4). By contrast, co-cultures went extinct when the initial *E. coli* λ frequency was less than 95 per cent, showing that suicide cannot stop infection in well-mixed cultures if the

suicidal strain is not sufficiently common, because T4rII multiplication (burst size 100–300) within *E. coli* HK97 outweighs suicide-based phage eradication.

(c) Relatedness and the evolution of host suicide

The earlier-mentioned analyses in well-mixed cultures (where relatedness $r \approx 0$) show that owing to its cost-free nature, altruistic host suicide can be maintained even with $r = 0$ (i.e. non-suicidal strains cannot invade). On the other hand, the data also show that altruistic host suicide cannot evolve with $r = 0$ because populations go extinct before invasion of the Abi trait is possible. Thus, we predict marginal relatedness ($r < 0$) to be required for successful invasion. Our fitness assays in structured environments confirmed this hypothesis.

In a highly structured environment (0.4% agarose, high relatedness), we found that T4rII infections could always be stopped, even at very high MOIs normally sufficient to eradicate *E. coli* λ in liquid culture. Furthermore, *E. coli* λ significantly outcompeted *E. coli* HK97 under all conditions (figure 3a), whereby the relative fitness of *E. coli* λ significantly increased with phage concentration (ANOVA: $F_{1,72} = 24.66$, $p < 0.0001$) but was independent of the initial frequency of *E. coli* λ ($F_{1,72} = 1.51$, $p = 0.22$). These findings are consistent with the view that phage–bacteria interaction occurred at a local scale, where T4rII managed to drive local *E. coli* HK97 patches to extinction. By contrast, T4rII was eradicated from *E. coli* λ patches owing to suicide, leading to large fitness benefits for surviving *E. coli* λ, even when initially rare.

Crucially, we observed that *E. coli* λ also experienced large fitness benefits when moving from solid (0.4% agarose, reflecting high relatedness) to almost liquid (in 0.025% agarose-shaken medium, reflecting relatedness close to zero) conditions (figure 3b). This shows that even marginal population structure suffices for *E. coli* λ to significantly outcompete *E. coli* HK97. Moreover, our data show that fitness benefits for *E. coli* λ significantly peaked at intermediate agarose concentrations (quadratic regression: $F_{2,37} = 5.59$, $p = 0.008$, figure 3). This pattern indicates that in extremely structured environments not only bacteria but also phage mobility is greatly limited. Consequently, susceptible *E. coli* HK97 performs better in these environments because of a reduced risk of encountering phages.

4. Discussion

We showed that *E. coli* λ suicide upon T4rII infection was highly beneficial as it saved the clonal population from extinction, but entailed no detectable cost for cells committing suicide. This resulted in suicidal behaviour being favoured even when relatedness between committers and survivors approached zero. No detectable costs of suicide could arise owing to a combination of low metabolic maintenance costs of the Rex system, and the fact that infected bacteria are moribund anyway. Because infection causes the rapid death of the host, an infected individual has little or no opportunity to reproduce until its death, such that the commitment of suicide might simply hasten an individual's inevitable doom. Our findings demonstrate that altruistic host suicide can clearly be delimited from other forms of suicidal behaviours in microbes, such as cell lysis to release toxins [25] or nutrients [26], cell sacrifices to overcome immune systems [27] or to form fruiting body stalks [28,29].

Figure 3. (a) On solid agarose (0.4%, high relatedness) *Escherichia coli* λ significantly outcompeted *E. coli* HK97 at all phage concentrations, even when initially rare (*t*-tests, from left to right: $t_7 = 3.5$, $p = 0.0097$; $t_7 = 9.6$, $p < 0.0001$; $t_7 = 10.7$, $p < 0.0001$; $t_7 = 12.8$, $p < 0.0001$; $t_7 = 7.6$, $p < 0.0001$; $t_7 = 4.6$, $p = 0.0024$; $t_7 = 5.5$, $p = 0.0009$; $t_7 = 10.6$, $p < 0.0001$; $t_7 = 17.1$, $p < 0.0001$; $t_7 = 25.1$, $p < 0.0001$). (b) *Escherichia coli* λ also significantly outcompeted *E. coli* HK97 (in a 1:1 mix) when the degree of population structure was reduced to a minimum, showing that only marginal relatedness is required for host suicide to be selectively favoured (*t*-tests, from left to right: $t_7 = 4.8$, $p = 0.0020$; $t_7 = 18.2$, $p < 0.0001$; $t_7 = 14.2$, $p < 0.0001$; $t_7 = 10.9$, $p < 0.0001$; $t_7 = 9.5$, $p < 0.0001$). Colours indicate the type of marker plasmid used for *E. coli* λ (blue circles, pGFP; orange triangles, pDsRed).

In contrast to the altruistic suicide described in our study, these other behaviours require substantial relatedness among interacting individuals because suicidal cells lose direct fitness benefits.

The absence of detectable costs for *E. coli* λ suicide further raises the question, whether this behaviour represents a true altruistic act, which per some definitions implies $c > 0$ [30,31]. There are two ways to deal with this issue. First, it could be argued that, although not detectable with our methods, there are certainly some costs ($c > 0$) involved in Rex maintenance, such that *E. coli* λ suicide indeed represents an altruistic act. For instance, marginal costs of carrying Rex might be masked in our experiment because the two prophages (HK97 and λ) differ in a number of minor aspects

other than Rex. Second, $c = 0$ could be true, in that case *E. coli* λ suicide could be considered as a novel type of cooperative behaviour, which is neither altruistic ($c > 0$) nor mutually beneficial ($c < 0$) [31]. However, we do not think that the proclamation of a novel type of cooperative behaviour is useful, and rather endorse the view that the definition of altruism should be expanded to $c \geq 0$ (in line with Foster [32]).

Our results further reveal that under a narrow range of conditions (probably depending on MOI) the Rex system rather induces dormancy than suicide (figure 2a, *E. coli* λ frequency of approx. 95%). This phenomenon was first described by Slavcey & Hayes [23,24], who showed that a fraction of dormant cells are able to resume growth after approximately 2 days. The proximate basis of this dormancy is unknown. Furthermore, it is also unclear whether this form of dormancy is adaptive, providing direct fitness benefits, or whether it represents a by-product of the Rex system, which has primarily evolved to induce suicide. The latter explanation seems more probable, because the mechanism of cell death is clearly established [8,13] and is triggered across most conditions examined here.

Our fitness-based (ultimate) analyses reveal that bacterial suicide is adaptive and represents a highly efficient host-defence mechanism. Our findings are in close agreement with the work of Fukuyo *et al.* [33], who used an engineered *E. coli*–phage system to demonstrate for the first time that host suicide is adaptive in structured habitats. Our study and the work of Fukuyo *et al.* [33] contrast with previous studies, where evolutionary conclusions were mainly based on proximate (mechanistic) aspects of cell death [8,10,11]. This is problematic because mechanistic approaches elucidating how things work can provide only limited answers on why a given behaviour has evolved [31].

More generally, evolutionary conclusions based on proximate findings resulted in much controversy on whether mechanisms resulting in microbial cell death have indeed evolved for the purpose of committing suicide or whether they represent by-products of selection at other traits [34–36]. While this controversy can never distinctly be resolved for past events, it seems that in the Rex system, *rex*B may have originally been favoured because it protects *E. coli* from the killing mediated by addition modules [37]. In combination with the sensor *RexA*, which is triggered by several unrelated phages [38], the system may have subsequently been coopted into an adaptive suicidal host-defence mechanism. Co-option, which is a common phenomenon in evolution [32], might have also been involved in shaping some of the other Abi systems described in bacteria [9–11]. Moreover, it might be that not all Abi systems involve active suicide (i.e. a cell expressing a trait with the purpose to kill itself). It might rather be that certain mechanisms have evolved to block phage progeny production within the host, which consequently lead to bacterial death. Such traits would however spread under the same conditions as the Rex system examined in our study.

The idea of infected hosts committing suicide in order to prevent disease transmission to healthy relatives has also received considerable attention in higher organisms. While theoretical work convincingly revealed the conditions required for host suicide to evolve [39,40], empirical work has often been criticized [41,42]. In particular, the adaptive significance of host suicide has been challenged because studies involved eusocial Hymenoptera [43–45] or clonal aphids [46], where complex life histories impeded rigorous fitness tests and the exclusion of

alternative explanations. Our findings now reveal that, owing to its low cost, altruistic host suicide might easily evolve in any organism, even when relatedness is low. Possible examples include the observation that infected social insect workers leave the colony to die in isolation (i.e. altruistic self-removal [44,45]), a behaviour that could come at negligible costs because infected individuals might be strongly compromised in carrying out worker tasks anyway.

We thank John W. Little, Ing-Nang Wang, Benjamin Kerr, Bärbel Stecher, Martin Ackermann and Winfried Boos for providing bacteria and phage strains, the flow cytometry laboratory at the ETH Zürich, Anette Schütz and Olin Silander for help with flow cytometry, and Jim Bull, Tim Cooper, Fred Inglis, Laurent Keller, Stu West and an anonymous reviewer for valuable comments. This work was supported by the Competence Center Environment and Sustainability of the ETH (to D.R.), the Swiss National Science Foundation, and the European Commission (to R.K.).

References

- Hamilton WD. 1964 The genetical evolution of social behaviour. *J. Theor. Biol.* **7**, 1–52. (doi:10.1016/0022-5193(64)90038-4)
- Abbot P *et al.* 2011 Inclusive fitness theory and eusociality. *Nature* **471**, E1–E4. (doi:10.1038/nature09831)
- Boomsma JJ. 2007 Kin selection versus sexual selection: why the ends do not meet. *Curr. Biol.* **17**, R673–R683. (doi:10.1016/j.cub.2007.06.033)
- Hughes WOH, Oldroyd BP, Beekman M, Ratnieks FLW. 2008 Ancestral monogamy shows kin selection is key to the evolution of eusociality. *Science* **320**, 1213–1216. (doi:10.1126/science.1156108)
- Cornwallis CK, West SA, Davis KE, Griffin AS. 2010 Promiscuity and the evolutionary transition to complex societies. *Nature* **466**, 969–972. (doi:10.1038/nature09335)
- Lukas D, Clutton-Brock T. 2012 Cooperative breeding and monogamy in mammalian societies. *Proc. R. Soc. B* **279**, 2151–2156. (doi:10.1098/rspb.2011.2468)
- Mollineux IJ. 1991 Host–parasite interactions: recent developments in the genetics of abortive phage infections. *New Biol.* **3**, 230–236.
- Parma DH, Snyder M, Sobolevski S, Nawroz M, Brody E, Gold L. 1992 The Rex system of bacteriophage λ : tolerance and altruistic cell death. *Genes Dev.* **6**, 497–510. (doi:10.1101/gad.6.3.497)
- Snyder L. 1995 Phage-exclusion enzymes: a bonanza of biochemical and cell biology reagents? *Mol. Microbiol.* **15**, 415–420. (doi:10.1111/j.1365-2958.1995.tb02255.x)
- Chopin M-C, Chopin A, Bidnenko E. 2005 Phage abortive infection in lactococci: variations on a theme. *Curr. Opin. Microbiol.* **8**, 473–479. (doi:10.1016/j.mib.2005.06.006)
- Labrie SJ, Samson JE, Moineau S. 2010 Bacteriophage resistance mechanisms. *Nat. Rev. Microbiol.* **8**, 317–327. (doi:10.1038/nrmicro2315)
- Queller DC, Strassmann JE. 2009 Beyond society: the evolution of organismality. *Phil. Trans. R. Soc. B* **364**, 3143–3155. (doi:10.1098/rstb.2009.0095)
- Ptashne M. 2004 *A genetic switch: phage lambda revisited*, 3rd edn. Cold Spring Harbor, NY: CSH Laboratory Press.
- Benzer S. 1955 Fine structure of a genetic region in bacteriophage. *Proc. Natl Acad. Sci. USA* **41**, 344–354. (doi:10.1073/pnas.41.6.344)
- Chao L, Levin BR. 1981 Structured habitats and the evolution of anticompetitor toxins in bacteria. *Proc. Natl Acad. Sci. USA* **78**, 6324–6328.
- Kümmerli R, Griffin AS, West SA, Buckling A, Harrison F. 2009 Viscous medium promotes cooperation in the pathogenic bacterium *Pseudomonas aeruginosa*. *Proc. R. Soc. B* **276**, 3531–3538. (doi:10.1098/rspb.2009.0861)
- West SA, Griffin AS, Gardner A, Diggle SP. 2006 Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* **4**, 597–607. (doi:10.1038/nrmicro1461)
- Kümmerli R, van den Berg P, Griffin AS, West SA, Gardner A. 2010 Repression of competition promotes cooperation: experimental evidence from bacteria. *J. Evol. Biol.* **23**, 699–706. (doi:10.1111/j.1420-9101.2010.01936.x)
- R Development Core Team 2010 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Zwietering MH, Jongenburger I, Rombouts FM, van't Ried K. 1990 Modeling of the bacterial growth curve. *Appl. Environ. Microbiol.* **56**, 1875–1881.
- Hartl DL, Clark AG. 1997 *Principles of population genetics*, 3rd edn. Sunderland, MA: Sinauer Associates, Inc.
- Orr HA. 2009 Fitness and its role in evolutionary genetics. *Nat. Rev. Genet.* **10**, 531–539. (doi:10.1038/nrg2603)
- Slavcev RA, Hayes S. 2002 Rex-centric mutualism. *J. Bacteriol.* **184**, 857–858. (doi:10.1128/JB.184.3.857-858.2002)
- Slavcev RA, Hayes S. 2003 Stationary phase-like properties of the bacteriophage λ Rex exclusion phenotype. *Mol. Genet. Genomics* **269**, 40–48. (doi:10.1007/s00438-002-0787-x)
- Cascales E, Buchanan SK, Duché D, Kleantous C, Llobes R, Postle K, Riley M, Slatin S, Cavard D. 2007 Colicin biology. *Microbiol. Mol. Biol. Rev.* **71**, 158–229. (doi:10.1128/MMBR.00036-06)
- Durand PM, Rashidi A, Michod RE. 2011 How an organism dies affects the fitness of its neighbors. *Am. Nat.* **177**, 224–232. (doi:10.1086/657686)
- Ackermann M, Stecher B, Freed NE, Songhet P, Hardt W-D, Doebeli M. 2008 Self-destructive cooperation mediated by phenotypic noise. *Nature* **454**, 987–990. (doi:10.1038/nature07067)
- Strassmann J, Zhu Y, Queller DC. 2000 Altruism and social cheating in the social amoeba *Dictyostelium discoideum*. *Nature* **408**, 965–967. (doi:10.1038/35050087)
- Kraemer SA, Velicer GJ. 2011 Endemic social diversity within natural kin groups of a cooperative bacterium. *Proc. Natl Acad. Sci. USA* **108**, 10 823–10 830. (doi:10.1073/pnas.1100307108)
- Lehmann L, Keller L. 2006 The evolution of cooperation and altruism: a general framework and a classification of models. *J. Evol. Biol.* **19**, 1365–1376. (doi:10.1111/j.1420-9101.2006.01119.x)
- West SA, Griffin AS, Gardner A. 2007 Social semantics: altruism, cooperation, mutualism, strong reciprocity and group selection. *J. Evol. Biol.* **20**, 415–432. (doi:10.1111/j.1420-9101.2006.01258.x)
- Foster K. 2011 The sociobiology of molecular systems. *Nat. Rev. Genet.* **12**, 193–203. (doi:10.1038/nrg2903)
- Fukuyo M, Sasaki A, Kobayashi I. 2012 Success of a suicidal defense strategy against infection in a structured habitat. *Sci. Rep.* **2**, 238. (doi:10.1038/srep00238)
- Gardner A, Kümmerli R. 2008 Social evolution: this microbe will self-destruct. *Curr. Biol.* **18**, R1021–R1023. (doi:10.1016/j.cub.2008.09.003)
- Nedelcu AM, Driscoll WW, Durand PM, Herron MD, Rashidi A. 2011 On the paradigm of altruistic suicide in the unicellular world. *Evolution* **65**, 3–20. (doi:10.1111/j.1558-5646.2010.01103.x)
- Reece SE, Pollitt LC, Colegrave N, Gardner A. 2011 The meaning of death: evolution and ecology of apoptosis in protozoan parasites. *PLoS Pathog.* **7**, e1002320. (doi:10.1371/journal.ppat.1002320)
- Engelberg-Kulka H, Rechtes M, Narasimhan S, Schoulaker-Schwarz R, Klemens Y, Aizenman E, Glaser G. 1998 *rexB* of bacteriophage λ is an anti-cell death gene. *Proc. Natl Acad. Sci. USA* **95**, 15 481–15 486. (doi:10.1073/pnas.95.26.15481)
- Court D, Oppenheim AB. 1983 Phage λ 's accessory genes in Lambda II. In *Lambda II* (eds RW Hendrix, JW Roberts, FW Stahl, RA Weisberg), pp. 251–277. Cold Spring Harbor, NY: CSH Laboratory Press.
- Smith Trail DR. 1980 Behavioral interactions between parasites and hosts: host suicide and the

- evolution of complex life cycles. *Am. Nat.* **116**, 77–92. (doi:10.1086/283612)
40. Debarre F, Lion S, van Baalen M, Gandon S. 2012 Evolution of host life-history traits in a spatially structured host-parasite system. *Am. Nat.* **179**, 52–63. (doi:10.1086/663199)
 41. Tomlinson I, Latta B. 1987 Adaptive and non-adaptive suicide in aphids. *Nature* **330**, 701. (doi:10.1038/330701a0)
 42. Poulin R. 1995 'Adaptive' changes in the behaviour of parasitized animals: a critical review. *Int. J. Parasitol.* **25**, 1371–1383. (doi:10.1016/0020-7519(95)00100-X)
 43. Müller CB, Schmid-Hempel R. 1992 To die for host or parasite? *Anim. Behav.* **44**, 177–179. (doi:10.1016/S0003-3472(05)80770-5)
 44. Heinze J, Walter B. 2010 Moribund ants leave their nests to die in social isolation. *Curr. Biol.* **20**, 249–252. (doi:10.1016/j.cub.2009.12.031)
 45. Rueppell O, Hayworth MK, Ross NP. 2010 Altruistic self-removal of health-compromised honey bee workers from their hive. *J. Evol. Biol.* **23**, 1538–1546. (doi:10.1111/j.1420-9101.2010.02022.x)
 46. McAllister MK, Roitberg BD. 1987 Adaptive suicidal behaviour in pea aphids. *Nature* **328**, 797–799. (doi:10.1038/328797b0)