TECHNICAL NOTE

Tae-Soo Park · Young-Ho Kim

Modification of the Weaver-Dunn procedure

Accepted: 30 December 1998

We report a modification of the Weaver-Dunn procedure for the treatment of acromioclavicular (AC) dislocations (Rockwood type III–VI lesions).

Surgical technique

After exposing the acromion and distal clavicle, subperiosteal flaps are raised from the distal 2.0 to 3.0 cm of the clavicle. The remnant of the AC ligament, joint capsule and periosteum of the medial aspect of the acromion and distal clavicle are dissected and

carefully preserved. The soft tissues attached to the distal clavicle are divided into two flaps. The meniscal remnant is removed; approximately 1.5 cm of distal clavicle is removed taking care to preserve the adjacent soft tissue attachments to the clavicle and acromion. The coracoacromial (CA) ligament is mobilized from the acromion using #2 Ethibond; two to three modified Kessler sutures are passed through the free cut end. Either a 5.0 mm wide Mersilene tape or strong braided sutures are passed around the coracoid process from medial to lateral behind the coracoid attachment of the CA ligament. Two small drill holes are made in the dorsal surface of the cut end of the clavicle, through these are passed the Kessler sutures in the CA ligament which is then sutured firmly into the cut end of the clavicle while it is held reduced. The Mersilene tape or braided sutures are then passed through a separate drill hole in the clavicle and tied firmly to hold the clavicle in the reduced position.

The CA ligament repair can be reinforced using the remains of the AC ligament and joint capsule. The deltoid is reattached and the wound is closed.

Fig. 1 If the length of the superoposterior flap of the AC ligament is adequate, this portion is sutured to remnants of the torn AC ligament attached to the medial aspect of the acromion. Antero-medially and posteriorly, the CA ligament is augmented by the antero-inferior flap of the AC ligament

Fig. 2 If the length of the superoposterior flap of the AC ligament is inadequate, the CA ligament is augmented by the supero-posterior and antero-inferior flaps of the AC ligament attached to the cortices of the remaining clavicle

T.-S. Park (✉) · Y.-H. Kim
Department of Orthopaedic Surgery, Hanyang University, Kuri
Hospital, College of Medicine, 249-1, Kyomoon-Dong, Kuri City,
Kyunggi-Do, 471-020, Korea