Long Term Angiographical Follow-up of Cerebral Aneurysms after Coil Embolization

Y. KAKU, K. HAYASHI, M. SAWADA, N. SAKAI

Department of Neurosurgery, Gifu University School of Medicine, Japan

Key words: follow-up, endovascular occlusion, GDC, recanalization

Summary

We evaluated long-term angiographical follow-up of cerebral aneurysms treated with detachable platinum coils with special reference to the long-term morphological outcomes of incompletely obliterated aneurysms.

Serial long-term follow-up cerebral angiograms (>1 year) were obtained in 47 cases out of 134 cases treated with GDCs or IDCs from 1994 to 1999. In 47 patients, initial angiographical results demonstrated ten complete aneurysmal occlusion, seven aneurysms with a small neck remnant, twenty aneurysms with body filling and ten aneurysms with both of neck remnant and body filling.

In the ten completely obliterated aneurysms, follow-up angiograms revealed no aneurysmal recanalization. In seven aneurysms with neck remnant, three remained unchanged, four showed enlargement of contrast filling in the part of the neck of the aneurysm. In twenty aneurysms with body filling, five had developed into complete obliteration, three remained unchanged, 11 had recanalization, and one aneurysm displayed regrowth. In ten aneurysms

with both of neck remnant and body filling, two remained unchanged, five had recanalization, and three aneurysms displayed regrowth.

Long-term angiographical follow-up results are less satisfactory in cases involving incompletely obliterated lesions. A higher incidence of recanalizations were promoted in cases with neck remnant and/or body filling. Strict follow-up angiography is mandatory when a complete obliteration is not achieved.

Introduction

Endovascular embolization using GDCs or IDCs for complicated intracranial aneurysms has become widely accepted as an alternative to direct surgery. Although remarkable advances have been made in the endovascular treatment of aneurysms, the long-term outcome of this type of treatment is not well defined.

Follow up studies of these patients are necessary to document continued aneurysm exclusion from the cerebral circulation.

In this study, we evaluated long-term angio-

Follow-up results 1

neck remnant 7 cases
(VER : $32.17 \pm 12.58\%$)

unchanged 3 cases
(VER : $28.97 \pm 9.4\%$)

enlarged 4 cases
(VER : $34.58 \pm 14.04\%$)

Figure 1 Long-term angiographical follow-up results of cerebral aneurysms with neck remnant following GDC embolization.

graphical follow up of cerebral aneurysms treated with GDCs with special reference to the long-term morphological outcomes of incompletely obliterated aneurysms.

Clinical materials

134 cases were treated with GDCs or IDCs from 1994 to 1999. Serial long-term follow up cerebral angiograms (>1 year) were obtained

in 47 cases. The remaining 87 cases without long-term angiographical follow up were not included in this study.

In 47 patients; 7 males and 40 females with the ages ranged from 32 to 84 years (mean 61.7 years); 23 aneurysms presented with subarachnoid hemorrhage, 19 aneurysms were found incidentally and 5 aneurysms exhibited mass effect. 31 aneurysms were located in the anterior circulation and 16 in the posterior circulation.

Figure 2 Serial angiograms demonstrating the process of enlargement of the neck remnant following GDC embolization A) Vertebral angiogram showing a small basilar tip aneurysm with ill-defined neck. B) Vertebral angiogram obtained immediately after endovascular occlusion demonstrating a small neck remnant. C) Vertebral angiogram obtained 12 months after GDC embolization, demonstrating enlargement of the neck remnant.

Figure 3 Long-term angiographical follow-up results of cerebral aneurysms with body fillings following GDC embolization.

20 aneurysms were small in size ($<10\text{mm}$) with well defined neck, 11 were small sized with ill-defined neck, 8 were large ($10\text{-}25\text{mm}$) and 8 were giant ($>25\text{mm}$).

The morphological results of initial treatment were categorized as follows; complete occlusion defined as no contrast filling of the dome or neck of the aneurysm, neck remnant defined as some contrast filling in the part of the neck of aneurysm, body filling defined as some contrast filling of the dome of the aneurysm, both of neck remnant and body filling.

Postembolization follow-up cerebral angiograms were obtained within 6 months and repeated annually, if a complete obliteration was not confirmed.

The volume embolization ratio (VER) of each aneurysm was calculated as follows ; total coil volume / aneurysm volume. Detail of the calculation of VER has been reported by Satoh et al⁶.

Results

Initial morphological results: Complete aneurysmal occlusion was achieved in 10 patients (21.3%). A small neck remnant was de-

tected in 7 cases (14.9%), a body filling in 20 cases (42.5%) and both of them were detected in 10 cases (21.3%). The average of VER was 37.3% in small aneurysms with small necks, 37.0% in small aneurysms with ill-defined necks, 22.0% in large aneurysms, 11.9% in giant aneurysms.

Follow-up angiographical results: No aneurysmal recanalization was observed in 10 completely obliterated aneurysms (average VER: 37.4%). In 7 aneurysms with neck remnant, 3 (average VER 29.0%) remained unchanged, 4 (average VER 34.6%) showed enlargement of contrast filling in the part of the neck of the aneurysm (figure 1). Figure 2 shows a typical case of enlargement of neck remnant following GDC embolization of a basilar tip aneurysm.

In 20 aneurysms with body filling, 5 (average VER 26.5%) had developed into complete obliteration, 3 (average VER 33.2%) remained unchanged, 11 (average VER 31.1%) had recanalization, and 1 (VER: 25.8%) aneurysm displayed regrowth (figure 3). Figure 4 shows a typical case of progressive thrombosis following GDC embolization of a small IC aneurysm with small neck, and figure 5 demonstrates a

Figure 4 Serial angiograms demonstrating the process of progressive thrombosis following GDC embolization of a small IC aneurysm. A) Left carotid angiogram showing a small IC aneurysm. B) Left carotid angiogram obtained immediately after endovascular occlusion demonstrating incomplete obliteration of the aneurysm with contrast filling of the dome of the aneurysm. C) Left carotid angiogram obtained 12 months after GDC embolization, demonstrating complete occlusion of the aneurysm.

typical case of aneurysmal recanalization of a large IC aneurysm.

In 10 aneurysms with both of neck remnant and body filling, 2 (average VER 16.3%) remained unchanged, 5 (average VER 20.1%)

had recanalization, and 3 aneurysms (average VER: 24.9%) displayed regrowth (figure 6).

There was no statistical correlation between VER and long-term morphological results in each group.

Figure 5 Serial angiograms demonstrating the process of aneurysmal recanalization of a large IC aneurysm following GDC embolization. A) Left carotid angiogram obtained immediately after endovascular occlusion demonstrating an almost total occlusion of a left IC large aneurysm with some contrast filling of the neck and the body of the aneurysm. B) Left carotid angiogram obtained 12 months after GDC embolization, demonstrating recanalization of the aneurysm.

Figure 6 Long-term angiographical follow-up results of cerebral aneurysms with both of neck remnant and body filling following GDC embolization.

Discussion

Endovascular embolization using GDCs or IDCs for complicated intracranial aneurysms has become widely accepted as an alternative to direct surgery. Published reports of early clinical and angiographical results have been promising^{1,5}, but the long term efficacy of the GDC or IDC methods remains to be undetermined. Incomplete endovascular occlusion of aneurysm leaves the patient at risk for future expansion and future subarachnoid hemorrhage^{3,4}. Hayakawa et al evaluated the anatomical evolution of neck remnants in aneurysms treated with GDCs. They reported that 25% of aneurysms with neck remnant exhibited progressive thrombosis, 26% remained unchanged and 49% displayed recanalization on follow up angiography². Especially in large or giant aneurysms, 88% of aneurysm with neck remnant displayed aneurysmal recanalization. As with their follow up results, 51.1% of incompletely obliterated aneurysms showed aneurysmal recanalization or regrowth in our series.

The size of an aneurysm neck correlates well with the initial morphological results of the endovascular treatment for cerebral aneurysms⁷. In the cases with wide-necked aneurysm,

neointimal formation across the aneurysm neck was rarely completed, furthermore, looser packing with GDCs may prevent neo-intimal and neo-endothelial proliferation across the neck of the aneurysm. Therefore, the coils and the unorganized intra-aneurysmal thrombus were directly exposed to the blood circulation. Subsequently, the coils can be pushed by the arterial pressure and become compacted toward the aneurysm dome with further expansion of the aneurysm.

With this limitation in mind, patients need to be very carefully chosen for GDC embolization and strict follow-up angiography is mandatory when a complete embolization is not achieved.

Conclusions

GDC embolization is a safe treatment for cerebral aneurysms with lower incidence of peri-procedural morbidity, whereas follow-up results are less satisfactory in cases involving incompletely obliterated lesions. Higher incidence of recanalization was promoted in cases with neck remnant and/or body filling. Further refinements of GDC technology is necessary to improve present morphological and clinical outcomes.

References

- 1 Guglielmi G, Viñuela F et Al: Electrothrombosis of saccular aneurysms via endovascular approach. Part 2: Preliminary clinical experience. *J Neurosurg* 75: 8-14, 1991.
- 2 Hayakawa M, Murayama Y et Al: Natural history of the neck remnant of a cerebral aneurysm treated with the Guglielmi detachable coil system. *J Neurosurg* 93: 561-568, 2000.
- 3 Hodgson TJ, Carroll T, Jellinek DA: Subarachnoid hemorrhage due to late recurrence of a previously unruptured aneurysm after complete endovascular occlusion. *Am J Neuroradiol* 19: 1939-1941, 1998.
- 4 Mericle RA, Wakhloo AK et Al: Delayed aneurysm regrowth and recanalization after Guglielmi detachable coil treatment. *J Neurosurg* 89: 142-145, 1998.
- 5 Murayama Y, Viñuela F et Al: Embolization of incidental cerebral aneurysms by using the Guglielmi detachable coil system. *J Neurosurg* 90: 207-214, 1999.
- 6 Satoh K, Satomi J et Al: Measurement of volume ratio to predict coil compaction, on aneurismal embolization. *Interventional Neuroradiology* 4 (Suppl 1): 179-182, 1998.
- 7 Zubillaga AF, Guglielmi G et Al: Endovascular occlusion of intracranial aneurysms with electrically detachable coils: Correlation of aneurysm neck size and treatment results. *Am J Neuroradiol* 15: 821-827, 1994.

Yasuhiko Kaku, M.D.
Department of Neurosurgery
Gifu University School of Medicine
40 Tsukasamachi, Gifu 500-8705, Japan
E-mail : kaku@cc.gifu-u.ac.jp