Published in final edited form as:

Clin Exp Allergy. 2013 May ; 43(5): 527–534. doi:10.1111/cea.12086.

Glutathione S-transferase P1 Ile105Val polymorphism modulates allergen-induced airway inflammation in human atopic asthmatics in vivo

Aimee Hoskins, RN, BSN, Pingsheng Wu, PhD, Sara Reiss, MLI, and Ryszard Dworski, MD, PhD

Division of Allergy, Pulmonary and Critical Care Medicine, Vanderbilt University School of Medicine, Nashville, Tennessee, USA

Abstract

Background—Glutathione S-transferase P1 is a Phase II cytoprotective and detoxifying enzyme that is widely expressed in human airways. The glutathione S-transferase P1 Ile105Val polymorphism has been linked with atopic disorders and asthma. Yet, little is known about the regulation of allergic inflammation by glutathione S-transferase P1 in human asthmatics.

Objective—To establish the effect of the glutathione S-transferase P1 Ile105Val polymorphism on allergen-induced airway inflammation and oxidant stress, and nonspecific bronchial hyperresponsiveness to methacholine and reactivity to specific allergen in mild human atopic asthmatics *in vivo*.

Methods—Five Val¹⁰⁵/Val¹⁰⁵, twelve Val¹⁰⁵/Ile¹⁰⁵, and twenty Ile¹⁰⁵/Ile¹⁰⁵ mild atopic asthmatics underwent methacholine challenge, inhaled allergen challenge, and endobronchial allergen provocation through a bronchoscope. A panel of inflammatory cytokines and chemokines, F₂-isoprostanes and isofuranes, markers of oxidative stress, thromboxane B₂, and immunoglobulin E were measured in bronchoalveolar lavage fluid at baseline and 24 hrs after allergen instillation.

Results—Asthmatics with glutathione S-transferase P1 Val¹⁰⁵/Val¹⁰⁵ compared to asthmatics with the glutathione S-transferase P1 Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ had greater generation of acute phase cytokines (TNF-α, IL-6, CXCL8), IL-12, CCL11, thromboxane B₂, and immunoglobulin E at 24 hrs after local allergen challenge. The GSTP1 genotype had no effect on airway hyperresponsiveness to methacholine and the reactivity to specific allergen.

Conclusion—The glutathione S-transferase P1 Ile105Val polymorphism markedly modifies allergen-provoked airway inflammation in atopic asthmatics *in vivo*. Modulation of the biochemical milieu in response to allergen provides a mechanistic explanation for regulatory effects of glutathione S-transferase P1 polymorphism on airway pathophysiology, and may guide improvement of future therapeutic methods in human atopic asthmatics. These findings must be confirmed in a larger study population of asthmatics with various ethnicities.

Keywords

Glutathione S-transferase P1; Ile105Val polymorphism; atopic asthma; allergen challenge; methacholine challenge; allergic inflammation; oxidant stress

Corresponding author and reprint requests: Ryszard Dworski, MD, PhD, Vanderbilt University School of Medicine, Division of Allergy, Pulmonary and Critical Care Medicine, T-1218 Medical Center North, Phone: 615 322 3412, Fax: 615 343 7448, ryszard.dworski@vanderbilt.edu.

Disclosure of potential conflict of interest: The authors have declared that they have no potential conflict of interest or commercial associations as per journal policy.

Introduction

Glutathione *S*-transferase P1 (GSTP1) is a Phase II cytoprotective and detoxifying enzyme belonging to a large family of GSTs [1]. GSTP1 is widely expressed in human airways, predominantly in alveolar macrophages and the epithelial cells and, and provides the majority of GST activity in the lung [1,2]. The major physiological role of GSTP1 is elimination of electrophiles by conjugating them with glutathione [1]. However, GSTP1 has also antioxidant properties due to selenium-independent peroxidase activity [1] and modulates major cellular signalling and gene transcription pathways including c-Jun N-terminal kinase (JNK) [3], tumour necrosis factor (TNF)-receptor-associated factor 2 (TRAF2) [4], and peroxiredoxin-1 [5].

Asthma is a heterogeneous airway disorder linked to airway inflammation and oxidative stress [6,7]. The loss of GSTP1 activity by gene deletion results in an exaggerated response to allergic sensitization and challenge in a mouse model of asthma [8]. Paradoxically, antigen-induced inflammation inhibits expression of GSTP1 in the lungs of OVA-sensitized mice [9]. Moreover, decreased expression of GSTP1 in nasal epithelium has also been found in asthmatic children [7].

A large number of GST genes have been recognized as polymorphic and the relevance of some genetic variants of GST to human pathology has been documented [1]. A single nucleotide polymorphism at position 313 in GSTP1 that converts an adenine to a guanine (A→G) is both functional and frequently associated with various diseases. GSTP1 variants containing Val¹⁰⁵ have a higher efficiency for diol epoxides of polycyclic aromatic hydrocarbons but are less effective for 1-chloro-2,4-dinitrobenzene than Ile¹⁰⁵ containing GSTP1 [10]. The Ile to Val substitution in codon 5 of exon 5 (Ile¹⁰⁵→Val¹⁰⁵) lies in close proximity to the catalytic centre of the enzyme and presumably accounts for the change in activity of the enzyme [1].

The link between human asthma and the GSTP1 polymorphism was originally reported by British investigators in a case-controlled study that showed a decreased risk of asthma and atopy among individuals with the GSTP1 Val¹⁰⁵/Val¹⁰⁵ genotype as compared to GSTP1 Ile¹⁰⁵/Ile¹⁰⁵ individuals [11]. Subjects with GSTP1 Ile¹⁰⁵/Val¹⁰⁵ variant had an intermediate risk of asthma [11]. A recent meta-analysis of 17 investigations on GSTP1 Ile¹⁰⁵Val and asthma, although suggested a weak protective effect of the Val¹⁰⁵, showed a tremendous heterogeneity of the data [12]. In fact, some studies suggest deteriorating effect of GSTP1 Val¹⁰⁵ on asthma [13,14].

We hypothesized that the Ile¹⁰⁵→Val¹⁰⁵ polymorphism of GSTP1 modulates airway inflammation and reactivity in human atopic asthmatics. The aim of our study was to determine the effect of the Ile¹⁰⁵→Val¹⁰⁵ polymorphism of GSTP1 on allergen-induced airway inflammation and oxidant stress, and nonspecific bronchial hyperresponsiveness to methacholine and reactivity to specific allergen in human atopic asthmatics *in vivo*. We employed segmental and inhaled allergen challenges that are established models of exacerbation in atopic asthmatics and provide important tools for studying the pathophysiology of allergic asthma *in vivo* [15].

Methods

Patients

Thirty seven mild atopic asthmatics by National Asthma Education and Prevention Program guidelines [16] were recruited from the Middle Tennessee area. None of the volunteers had a

history of tobacco smoking, other lung disorders or a respiratory infection in the 6 weeks preceding the study. Patients had positive skin test with aeroallergens, methacholine challenge, and inhaled allergen provocation. All medications were discontinued at least 48 hrs prior to procedures. Pregnant women were excluded by urine HCG testing. Volunteers consented verbally and in writing to the protocol that was approved by the Vanderbilt University Committee for the Protection of Human Subjects (IRB number: 051158). An Investigational New Drug agreement with the U.S. Food and Drug Administration for the use of allergens for inhaled and bronchoscopic provocations in humans was obtained.

Study protocol

Volunteers underwent bronchoscopy with baseline (before allergen challenge) bronchoalveolar lavage (BAL) in the lingua of the left upper lobe followed by segmental allergen challenge (SAC) in a subsegment of the right middle lobe. The allergen challenged subsegment was lavaged 24 hrs later. At least 3 weeks after the bronchoscopy studies asthmatics underwent methacholine challenge followed by an inhaled allergen provocation the next day. The same allergen was used for both the inhaled and segmental allergen challenge.

Allergy skin test

Prick skin testing was performed with diluent and histamine controls in parallel with a set of standardized aeroallergens (*Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, cat hair, Bermuda grass, Kentucky bluegrass, fescue meadow grass, orchard grass, redtop grass, ryegrass, sweet vernal grass, timothy grass, short ragweed) (Greer Laboratories, Lenoir, NC) according to practice guidelines [17].

Bronchoscopy studies

Subjects were fasted overnight. Midazolam 1–2 mg and fentanyl 50–100 mcg were administered intravenously if patients wished conscious sedation. Topical lidocaine (7 mg/kg) was used for airway anaesthesia. BAL was performed using three 50 ml aliquots of normal saline and immediately suctioned back. SAC was done using two incremental doses of the allergen to which the volunteers had positive skin test, each dose in a 5 ml aliquot of normal saline. The first dose of the allergen was 10-fold higher than the concentration that gave 2+ skin reaction on intradermal test (5–8 mm wheal, 20–30 mm erythema). The subsegment was watched for 2 minutes. None of the volunteers developed visible local inflammatory reaction after the first dose of allergen. Thus, in each subject the second dose of the same allergen was instilled at concentration 100-fold higher than the 2+ skin reaction, and the bronchoscope was then removed [18].

Methacholine challenge

The methacholine (Provocholine, Methapharm Inc., Brantford, Ontario, Canada) challenge was performed according to the American Thoracic Society guidelines [19] using a Salter dosimeter (Salter Labs, Arvin, CA) and Flow Screen computerized spirometer (VIASYS Healthcare GmbH, Hoechberg, Germany).

Inhaled allergen challenge

Allergen inhalation challenge was accomplished as described by Cockcroft and colleagues [20]. The greatest fall in FEV₁ between 0–3 hrs post allergen was taken as the early airway response (EAR), and the maximum FEV₁ fall at 3–8 hrs was the late airway response (LAR). Allergen aerosols were generated using a Wright nebulizer connected to a 2-way Hans Rudolph valve mouthpiece (Roxon, Montreal, Canada) attached to a wall oxygen source at 50 psi. Spirometry was performed using Flow Screen computerized spirometer.

Laboratory methods

Genomic DNA was extracted from blood leukocytes with a Wizard Genomic DNA Purification Kit (Promega, Madison, WI). Probes used for the ABI assay (TaqMan® SNP Genotyping Assay, Applied Biosystems by Life Technologies, Carlsbad, CA) were labelled at their 5' end with either VIC® or 6-carboxy-fluorescein reporter dyes. The reaction components include: 2.5µl Taqman Universal PCR Master Mix and 0.125µl or 0.250µl assay mix (Applied Biosystems), and approximately 5ng of genomic DNA in a total volume of 10µl per single tube reaction. PCR amplification was performed in a thermal cycler (Techne TC-412, Burlington, NJ) with an initial step of 95°C for 10 min followed by 50 cycles of 92°C for 15 sec and 60°C for 1 min. After amplification, the fluorescence of each sample was read on the ABI 7900HT (DNA Resources Core, Vanderbilt University) and analyzed with the Sequence Detection Software (Applied Biosystems).

BAL cells were counted in Reichert Brightline Metallized Phase Hemacytometer. F₂-isoprostanes, isofuranes, and thromboxane B₂ (TXB₂) were analyzed by stable isotope dilution method in conjunction with gas chromatography/negative ion chemical ionization/mass spectrometry as previously described [21]. Interleukin(IL)-4, IL-5, IL-10, IL-12p70, IL-13, interferon (IFN)-γ, Tumour Necrosis Factor (TNF)-α, and IL-8 (CXCL8) were measured using human Plex Multi-Spot assay (Meso Scale Discovery, Gaithersburg, MD). IL-3, IL-6, CCL5 (RANTES), CCL11 (eotaxin), and immunoglobulin E (IgE) were analyzed by Cytometric Bead Array (BD Biosciences, San Jose, CA).

Statistical analysis

Patients' baseline characteristics were compared by using the chi-square test for categorical variables, and the Mann-Whitney U test for continuous variables. Data are presented as proportions for categorical variables and mean ± standard deviation (SD) for continuous variables. There was no difference in percentages of cells, concentrations of mediators and markers of oxidative stress between Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ both before and after allergen challenge, therefore a recessive model was used to compare GSTP1 genotype Val¹⁰⁵/Val¹⁰⁵ with Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ combined. In addition, percentages of cells, concentrations of mediators and markers of oxidative stress between genotypes were compared at baseline and after allergen challenge separately by using Mann-Whitney U tests. The effect of GSTP1 genotype was further assessed by comparing the change in outcome variables after allergen challenge between groups using general linear models (GLM). Baseline value of outcome variable was adjusted as a model covariate as well as other baseline covariates including age, gender, and race. Residuals were assessed graphically for normality and transformation on the dependent variable was done to correct non-normal residuals if needed. Significance was accepted when $p < 0.05$. R-software version 2.11.1 and SPSS version 19 (SPSS Inc., Chicago, IL, USA) were used for data analyses.

Results

Subjects

The study population consisted of 37 mild atopic asthmatics (25 women and 12 men). Thirty three patients were Caucasians and 4 African Americans. All patients had a normal baseline spirometry and the vast majority of them did not use any daily asthma medications. Four patients provided history of past inhaled corticosteroid use. Only two individuals took daily 100mg Fluticasone/50mg Salmeterol by inhalation. Discontinuation of the inhaled corticosteroids and beta-agonist prior to provocative challenges was not associated with any adverse consequences as judged by subjective and objective (spirometric) criteria. Volunteers were asked to continue the same treatment regimen throughout the study that

typically consisted of an inhaled short acting beta-agonist as needed, during the entire period of the study. Table 1 provides additional demographic data concerning the volunteers.

Effect on inflammatory airway cells

Asthmatics with Val¹⁰⁵/Val¹⁰⁵ had higher percentage of neutrophils in BAL at baseline compared to patients with Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ GSTP1 genotypes. The difference was not significant at 24 hrs after allergen provocation. Other cell types were not affected by GSTP1 polymorphism (Table 2). In addition, analysis using a recessive model showed no difference in inflammatory cells between Val¹⁰⁵/Val¹⁰⁵ and Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ GSTP1 genotypes after adjusting for baseline cell percentages, age, gender and race, at 24 hrs after allergen challenge.

Effect on allergen-induced inflammatory mediators, oxidant stress, and IgE

Asthmatics with all three GSTP1 SNP combinations at position 105 had similar baseline concentrations of cytokines and TXB₂ in BAL. However, instillation of specific allergen provoked a greater production of IL-6 (p=0.03), IL-12 (p=0.02), CXCL8 (p=0.03), and TNF-α (p=0.01) in the airways of asthmatics with the Val¹⁰⁵/Val¹⁰⁵ compared to asthmatics with Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ GSTP1 genotypes (Figure 1). GSTP1 polymorphism had no effect on allergen-induced generation of other Th2 cytokines and CCL5 (Table 1 online supplement).

In addition, Val¹⁰⁵/Val¹⁰⁵ asthmatics compared to Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ patients generated greater amounts of the cyclooxygenase metabolite, TXB₂ after instillation of specific allergen (p=0.02) (Figure 1), although this effect disappeared after adjusting for baseline TXB₂ level, age, gender and race.

Levels of F₂-isoprostanes and isofuranes, specific markers of oxidative stress *in vivo* [21], were also higher in asthmatics with Val¹⁰⁵/Val¹⁰⁵ compared to patients with GSTP1 Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ although the difference was not quite statistically significant (Table 1 online supplement).

The concentrations of total IgE in BAL fluid following allergen challenge were significantly higher in patients with GSTP1 Val¹⁰⁵/Val¹⁰⁵ compared to Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ asthmatics (Figure 1).

Except of TXB₂, the effect of GSTP1 polymorphism on allergen-induced generation of the measured mediators and IgE was essentially consistent after adjusting for baseline concentrations and the age, gender, and ethnicity of volunteers.

Effect on airway hyperresponsiveness to methacholine and reactivity to specific allergen

The threshold dose of methacholine causing 20% decrease in FEV₁ (Met-PC₂₀) did not differ among asthmatics with various GSTP1 genotypes. Likewise, the concentrations of allergen provoking 20% decrease in FEV₁ during the early asthmatic response (All-PD₂₀) were similar among the groups (Table 3). The study did not show a relationship between the GSTP1 polymorphism and LAR.

Discussion

Our study demonstrates greater generation of acute phase cytokines (TNF-α, IL-6, CXCL8), IL-12, and TXB₂ at 24 hrs after allergen challenge in atopic asthmatics with the homozygous GSTP1 Val¹⁰⁵/Val¹⁰⁵ genotype compared to asthmatics with the GSTP1 Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ genotypes. Allergen-induced levels of F₂-isoprostanes and

isofuranes, markers of oxidative stress [21], were also higher in individuals with GSTP1 Val¹⁰⁵/Val¹⁰⁵ although the difference narrowly missed statistical significance. To our knowledge these data are novel.

Our findings are likely to be relevant to the pathomechanism of human asthma. GSTP1 is mainly expressed in alveolar macrophages and airway epithelial cells that greatly contribute to airway pathobiology in asthma [2]. TNF- α , CXCL8, and IL-6 enhance allergic inflammation and airway remodelling [6]. IL-12 affects expression of the atopic phenotype by supporting differentiation of naïve CD4⁺ helper T cells toward the Th1 subset [6]. TXB₂ is a potent bronchoconstrictor as well as a modulator of inflammation and remodelling [22]. Oxidant stress plays a major pathophysiological role in asthma [7], and evidence of greater airway oxidant stress has been previously reported in asthmatic children with GSTP1 Val¹⁰⁵/Val¹⁰⁵ compared to GSTP1 Ile¹⁰⁵/Ile¹⁰⁵ [23]. Similarly to our study in atopic asthma, an increased production of IgE has been found in diisocyanate-induced asthmatics carrying the GSTP1 Val¹⁰⁵/Val¹⁰⁵ genotype [24].

GSTP1 polymorphism was not associated with nonspecific airway hyperresponsiveness to methacholine or airway reactivity to specific allergen. Our data concur with Imboden and colleagues who found no relationship between GSTP1 genotypes and sensitivity to methacholine [25] but disagree with Fryer and colleagues who have reported a protective role of GSTP1 Val¹⁰⁵/Val¹⁰⁵ on nonspecific bronchial hyperresponsiveness [11]. Importantly, all our volunteers were only mild atopic asthmatics with well controlled disease whereas patients in the latter study were phenotypically inhomogeneous. Nevertheless, studying larger sample size might be unnecessary to reveal modifying effect of the GSTP1 polymorphism on airway reactivity in atopic asthma.

The mechanism underlying the regulation of allergen-provoked airway inflammation in atopic asthma is unknown. Polymorphism in codon 5 alters conjugation of the protein with xenobiotics which may affect detoxifying functions of GSTP1 [1]. However, GSTP1 is also capable of interacting with several proteins involved in cell signalling and gene expression such as JNK1 [3], and mitogen-activated protein (MAPK) [26]. Nonetheless, the hypothesis that GSTP1 regulates signalling and gene expression in atopic asthma needs further investigation.

Currently, the pathophysiologic role of GSTP1 polymorphism in atopic asthma is not completely understood. Although mainly attributed to regulation of oxidant stress and detoxification of xenobiotics [1], our study clearly demonstrates more complex modulatory function of GSTP1 on allergen-induced airway inflammation in atopic asthmatics *in vivo*. Clinical and epidemiological data concerning regulation of asthma by the GSTP1 variants in humans are conflicting and support both protective and deteriorating role of the Val¹⁰⁵ allele on the disease [12], perhaps due to inconsistent study populations that included patients with various asthma phenotypes. Our patients represented exclusively carefully characterized atopic asthmatics. Interestingly, Kamada and colleagues found a link between the homozygous Val¹⁰⁵ genotype and asthma among atopic children [14]. In addition to allergens, GSTP1 regulates airway response to a variety of non-specific environmental factors affecting asthma such as ozone, nitrogen dioxide, tobacco smoke, and air pollution [12, 27]. Thus, future studies should demonstrate how GSTP1 modulates airway response to allergen accompanied by diverse environmental factors because complex exposures are more likely in natural surroundings. For example, Gilliland and colleagues showed augmentation of ragweed-induced generation of IgE, histamine and IL-4 by second-hand smoke [28] and diesel exhaust particles [29] in atopics with Ile¹⁰⁵/Ile¹⁰⁵ compared to Val¹⁰⁵/Ile¹⁰⁵ after nasal challenge, although the study population did not include any subjects with the Val¹⁰⁵/Val¹⁰⁵. However, there are also studies showing detrimental role of

GSTP1 Val¹⁰⁵/Val¹⁰⁵ rather than Ile¹⁰⁵/Ile¹⁰⁵ on allergen sensitization and asthma especially in children [26,30].

A major limitation of our investigation is a small number of participants, especially asthmatics with GSTP1 Val¹⁰⁵/Val¹⁰⁵ who predominantly represented the White ethnicity, although a pooled and race-stratified analysis revealed similar outcomes in our study. Nonetheless, the possibility that regulation of the allergen-induced airway inflammation by GSTP1 in atopic asthma is race-dependent should be addressed in future studies.

In summary, our study provides new data and a novel perspective on the regulatory role of GSTP1 polymorphism on allergen-induced airway inflammation in human atopic asthmatics *in vivo*. Our results suggest upregulation of allergen-induced airway inflammation and oxidative stress in adult atopic asthmatics with the homozygous Val¹⁰⁵ GSTP1 variant. Better understanding of the relationship between environmental and genetic factors and the pathomechanism of asthma may provide important guidance for improvement of therapeutic methods for atopic asthma in humans.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The study was supported by grants from the NIH (K23 HL080030, M01 RR-00095, and NIH P30 ES000267). The authors thank Dr. Ginger Milne, PhD, Stephanie Sanchez and Holly Borntrager for assistance with measurement of eicosanoids in BAL fluid.

References

1. Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu Rev Pharmacol Toxicol.* 2005; 45:51–88. [PubMed: 15822171]
2. Anttila S, Hirvonen A, Vainio H, Husgafvel-Pursiainen K, Hayes JD, Ketterer B. Immunohistochemical localization of glutathione S-Transferases in human lung. *Cancer Res.* 1993; 53:5643–8. [PubMed: 8242618]
3. Adler V, Yin Z, Fuchs SY, Benezra M, Rosario L, Tew KD, et al. Regulation of JNK signaling by GSTp. *EMBO J.* 1999; 18:1321–34. [PubMed: 10064598]
4. Wu Y, Fan Y, Xue B, Luo L, Shen J, Zhang S, et al. Human glutathione S-transferase P1-1 interacts with TRAF2 and regulates TRAF2-ASK1 signals. *Oncogene.* 2006; 25:5787–800. [PubMed: 16636664]
5. Ralat LA, Manevich Y, Fisher AB, Colman RF. Direct evidence for the formation of a complex between l-cysteine peroxiredoxin and glutathione S-transferase π with activity changes in both enzymes. *Biochemistry.* 2006; 45:360–72. [PubMed: 16401067]
6. Holgate ST. Pathophysiology of asthma: what has our current understanding taught us about new therapeutic approaches. *J Allergy Clin Immunol.* 2011; 128:495–505. [PubMed: 21807404]
7. Dworski R. Oxidant stress and asthma. *Thorax.* 2000; 55:S51–S53. [PubMed: 10992559]
8. Zhou J, Wolf CR, Henderson CJ, Cai Y, Board PG, Foster PS, et al. Glutathione transferase P1. An endogenous inhibitor of allergic responses in a mouse model of asthma. *Am J Respir Crit Care Med.* 2008; 178:1202–10. [PubMed: 18787219]
9. Schroer KT, Gibson AM, Sivaprasad U, Bass SA, Ericksen MB, Willis-Karp M, et al. Downregulation of glutathione S-transferase pi in asthma contributes to enhanced oxidative stress. *J Allergy Clin Immunol.* 2011; 128:539–48. [PubMed: 21570714]
10. Hu X, Xia H, Srivastava SK, Herzog C, Awasthi YC, Ji X, et al. Activity of four allelic forms of glutathione S-transferase hGSTP1-1 for diol epoxides of polycyclic aromatic hydrocarbons. *Biochem Biophys Res Comm.* 1997; 238:397–402. [PubMed: 9299520]

11. Fryer AA, Bianco A, Hepple M, Jones PW, Strange RC, Spiteri MA. Polymorphism at the glutathione S-transferase GSTP1 locus. A new marker for bronchial hyperresponsiveness and asthma. *Am J Respir Crit Care Med*. 2000; 161:1437–42. [PubMed: 10806136]
12. Minelli C, Granell R, Newson R, Rose-Zerilli MJ, Torrent M, Ring SM, et al. Glutathione-S-transferase genes and asthma phenotypes: a human epidemiology (HuGE) systemic review and meta-analysis including unpublished data. *Int J Epidemiol*. 2010; 39:539–62.
13. Schroer KT, Myers JMB, Ryan PH, LeMasters GK, Bernstein DI, Villareal M, et al. Associations between multiple environmental exposures and glutathione S-transferase P1 on persistent wheezing in a birth cohort. *J Pediatr*. 2009; 154:401–8. [PubMed: 18950799]
14. Kamada F, Mashimo Y, Inoue H, Shao C, Hirota T, Doi S, et al. The GSTP1 gene is a susceptibility gene for childhood asthma and the GSTM1 gene is a modifier of the GSTP1 gene. *Intern Arch Allergy Immunol*. 2007; 144:275–86.
15. Busse WW, Wanner A, Adams K, Reynolds HY, Castro M, Chowdhury B, et al. Investigative bronchoprovocation and bronchoscopy in airway diseases. *Am J Respir Crit Care Med*. 2005; 172:807–16. [PubMed: 16020805]
16. National Asthma Education and Prevention Program. Experts Panel Report 3: Guidelines for the Diagnosis and Management of Asthma. Bethesda, MD: National Institute of Health, National Heart, Lung, and Blood Institute; 2007. Pub no. 08-5846
17. American College of Allergy, Asthma and Immunology, American Academy of Allergy, Asthma and Immunology, and the Joint Council of Allergy, Asthma and Immunology. Practice Parameters for Allergy Diagnostic Testing. *Ann Allergy Asthma Immunol*. 1995; 75:543–625. [PubMed: 8521115]
18. Dworski R, Simon HU, Hoskins A, Yousefi S. Eosinophil and neutrophil extracellular DNA traps in human allergic asthmatic airways. *J Allergy Clin Immunol*. 2011; 127:1260–6. [PubMed: 21315435]
19. American Thoracic Society. Guidelines for methacholine and exercise challenge testing. *Am J Respir Crit Care Med*. 2000; 161:309–29. [PubMed: 10619836]
20. Cockcroft DW, Davis BE, Boulet L-P, Deschesnes F, Gauvreau GM, O'Byrne PM, et al. The links between allergen skin test sensitivity, airway responsiveness and airway response to allergen. *Allergy*. 2005; 60:56–9. [PubMed: 15575931]
21. Roberts LJ, Morrow JD. Measurement of F₂-isoprostanes as an index of oxidative stress in vivo. *Free Radic Biol Med*. 2000; 28:505–13. [PubMed: 10719231]
22. Boyce JA. Eicosanoids in asthma, allergic inflammation, and host defense. *Curr Mol Med*. 2008; 8:335–49. [PubMed: 18691060]
23. Ercan H, Birben E, Dizdar EA, Keskin O, Karaasian C, Soyer OU, et al. Oxidative stress and genetic and epidemiologic determinants of oxidant injury in childhood asthma. *J Allergy Clin Immunol*. 2006; 118:1097–104. [PubMed: 17088135]
24. Piirila P, Wikman K, Luukkonen R, Kaaria K, Rosenberg C, Nordman H, et al. Glutathione S-transferase genotypes and allergic responses to diisocyanate exposure. *Pharmacogenetics*. 2001; 11:437–45. [PubMed: 11470996]
25. Imboden M, Rochat T, Brutsche M, Schindler C, Downs SH, Gerbase MW, et al. Glutathione S-transferase genotype increases risk of progression from bronchial hyperresponsiveness to asthma in adults. *Thorax*. 2008; 63:322–8. [PubMed: 18057098]
26. Ip YT, Davis RJ. Signal transduction by the c-Jun N-terminal kinase (JNK)-from inflammation to development. *Curr Opin Cell Biol*. 1998; 10:205–19. [PubMed: 9561845]
27. Melén E, Nyberg F, Lindgren CM, Berglind N, Zucchelli M, Nordling E, et al. Interactions between glutathione S-transferase P1, Tumor necrosis factor, and traffic-related air pollution for development of childhood allergic disease. *Environ Health Perspectives*. 2008; 116:1077–84.
28. Gilliland FD, Li Y-F, Gong H Jr, Diaz-Sanchez D. Glutathione S-Transferase M1 and P1 prevent aggravation of allergic responses by secondhand smoke. *Am Rev Respir Crit Care Med*. 2006; 174:1335–41.
29. Gilliland, Li Y-F, Saxon A, Diaz-Sanchez D. Effect of glutathione-S-transferase M1 and P1 genotypes on xenobiotic enhancement of allergic responses: randomized, placebo-controlled crossover study. *Lancet*. 2004; 363:119–25. [PubMed: 14726165]

30. Lee YL, Lin YC, Wang JY, Hsiue TR, Guo YL. Glutathione S-transferase P1 gene polymorphism and air pollution as interactive factors for childhood asthma. *Clin Exp Allergy*. 2004; 34:1707–13. [PubMed: 15544594]

Figure 1.

Box plots comparing baseline (Bas) and post-allergen (All) BAL concentrations of TNF- α (A), IL-6 (B), IL-8 (C), IL-12 (D), TXB₂ (E) and IgE (E) in atopic asthmatics with GSTP1 Val/Val vs. combined group of patients with Val/Ile and Ile/Ile (Mann-Whitney U test).

Table 1

Demographic findings, history of atopy and asthma, asthma medication use, and allergens employed for challenges (p values calculated using one way ANOVA or Fisher's exact test for interval and nominal values, respectively).

	Val ¹⁰⁵ /Val ¹⁰⁵	Val ¹⁰⁵ /Ile ¹⁰⁵	Ile ¹⁰⁵ /Ile ¹⁰⁵	p
	Count or mean±SD, range			
Gender, M/F	2/3	3/9	7/13	<u>0.78</u>
Age	29.7±7.5 (20–37)	30±7.4 (23–46)	28±6.6 (20–43)	<u>0.60</u>
Race				
White	4	12	17	<u>0.32</u>
African American	1	0	3	
Family history of atopy and asthma	4	10	17	<u>0.96</u>
Baseline spirometry (%predicted)				
FVC	101±15.6 (83–123)	110.5±11 (92–128)	101±14 (70–141)	<u>0.68</u>
FEV1	105±9.9 (89–122)	100±14.5 (79–126)	92±12.6 (73–116)	<u>0.80</u>
FEV1/FVC	91.5±9.6 (79–106)	90±8.3 (75–104)	80±9.6 (67–107)	<u>0.80</u>
Drugs (daily or as needed)				
Short-acting β_2 -agonist	5	12	20	NA *
Long-acting β_2 -agonist	1	1	0	<u>0.18</u>
Inhaled corticosteroids	1	2	3	<u>0.96</u>
Nasal corticosteroids	2	2	0	<u>0.03</u>
Leukotriene modifier	1	2	2	<u>0.78</u>
Antihistamine	3	7	6	<u>0.21</u>
Allergens (type/dose) used for segmental allergen challenge				
Dust mite	<u>1/55 AU</u>	<u>4/0.55–5.5 AU, mean 3.85 AU</u>	<u>5/0.55–505 AU, mean 129 AU</u>	<u>0.59</u>
Cat	<u>1/55 AU</u>	<u>6/0.55–550 AU, mean 111.1 AU</u>	<u>10/0.055–5.5, mean 2.3 AU</u>	
Ragweed	<u>1/0.55 AU</u>	<u>0</u>	<u>2/.055–.0055 AU, mean 0.03 AU</u>	
Bermuda	<u>0</u>	<u>1/0.5 AU</u>	<u>1</u>	
Bluegrass	<u>1/0.055 AU</u>	<u>1/550 AU</u>	<u>0</u>	
Rye grass	<u>1/0.055 AU</u>	<u>0</u>	<u>1/105 AU</u>	
Timothy grass	<u>0</u>	<u>0</u>	<u>1/0.055 AU</u>	

* Fisher exact test could not be performed as everyone in the cohort used short-acting beta agonist.

Table 2

Percentages of cells in BAL fluid at baseline (Bas) and after allergen challenge (All) (mean $\pm$ SD, range, p values are obtained using recessive models with Val¹⁰⁵/Val¹⁰⁵ vs. Val¹⁰⁵/Ile¹⁰⁵ and Ile¹⁰⁵/Ile¹⁰⁵ combined).

Variable		Val ¹⁰⁵ /Val ¹⁰⁵	Val ¹⁰⁵ /Ile ¹⁰⁵ + Ile ¹⁰⁵ /Ile ¹⁰⁵	p values *
Macrophages	Bas	76.2 $\pm$ 15.9 (56–89)	86.6 $\pm$ 7.7 (64–98)	0.15
	All	28.2 $\pm$ 10.1 (14–42)	49.5 $\pm$ 26.0 (2–98)	0.09
Eosinophils	Bas	2.6 $\pm$ 3.1 (0–8)	1.7 $\pm$ 2.7 (0–11)	0.31
	All	30.8 $\pm$ 31.9 (3–69)	13.9 $\pm$ 19.6 (1–80)	0.18
Neutrophils	Bas	5.6 $\pm$ 2.7 (2–9)	2.0 $\pm$ 2.3 (0–11)	0.008
	All	35.4 $\pm$ 27.5 (5–68)	27.9 $\pm$ 25.7 (0–75)	0.65
Lymphocytes	Bas	13.0 $\pm$ 13.6 (4–37)	7.9 $\pm$ 7.8 (0–28)	0.75
	All	11.8 $\pm$ 7.6 (5–21)	7.9 $\pm$ 7.2 (0–32)	0.21

* P values were obtained with non parametric Mann-Whitney U test

Table 3

Provocative concentrations of methacholine and allergen (dilution fold of the allergen stock solution) causing 20% decrease in FEV₁ (mean $\pm$ SD, range, p values Val¹⁰⁵/Val¹⁰⁵ vs. Val¹⁰⁵/Ile¹⁰⁵ + Ile¹⁰⁵/Ile¹⁰⁵).

	Val ¹⁰⁵ /Val ¹⁰⁵	Val ¹⁰⁵ /Ile ¹⁰⁵ + Ile ¹⁰⁵ /Ile ¹⁰⁵	p value
Methacholine PC₂₀	4.8 $\pm$ 5.9 (0.24–12.4)	3.35 $\pm$ 4.3 (0.09–20)	0.9
Allergen PC₂₀	132.8 $\pm$ 122 (4–256)	69.8 $\pm$ 113.1 (4–512)	0.3