Mat Soc Med. 2012 Jun; 24(2): 91-93

Received: March 17th, 2012

Accepted: May 12th, 2012

CONFLICT OF INTEREST: none declared.
© AVICENA 2012

doi: 10.5455/msm.2012.24.91-93

Total Fats, Saturated Fatty Acids, Processed Foods and Acute Coronary Syndrome in Transitional Albania

Iris Mone, Anyla Bullo

Clinical and Biochemistry Laboratory Service, Faculty of Medicine, Tirana University, Tirana, Albania

Corresponding author: Iris Mone, MD. Address: Rr. "Dibres", No. 371, Tirana, Albania. Telephone: 00355692149301; E-mail: iris_mone@yahoo.com.

ORIGINAL PAPER

SUMMARY

Background: We aimed to assess the association of acute coronary syndrome (ACS) with selected food groups pertinent to non-Mediterranean prototype in Albania, a transitional post-communist country in Southeast Europe. Methods: We conducted a case-control study in Tirana in 2003-2006 including 467 non-fatal consecutive ACS patients (370 men aged 59.1 ± 8.7 years, 97 women aged 63.3 ± 7.1 years; 88% response) and a population-based control group (469 men aged 53.1 ± 10.4 years, 268 women aged 54.0 ± 10.9 years; 69% response). A semi-quantitative food frequency questionnaire including 105 food items was administered to all participants based on which the daily calorie intake for selected food groups (meat products, overall oils and fats, sweets, and junk food) was calculated. General linear model was used to assess the association of food groups with ACS. Results: Mean age-adjusted values of meat products, overall oils and fats, sweets and junk food were all considerably higher in cases than controls in both sexes. Cases had significantly higher mean "non-Mediterranean" diet scores (consisting of junk food, sweets, oils and fats except olive oil) than controls (10.3% vs. 5.9% in men and 15.2% vs. 8.3% in women, $P < 0.01$ for both). Conclusions: In this Albanian population, intake of total fats, in particular saturated fatty acids was associated with a higher risk of ACS in both sexes. Furthermore, the consumption of processed foods was associated with considerable excess coronary risk which points to serious health implications for the Albanian adult population.

Key words: total fats, saturated fatty acids, risk of Acute coronary syndrome.

1. INTRODUCTION

In Albania, anecdotal evidence indicates that deviation from the traditional Mediterranean diet has taken place in all regions of the country in the past two decades in line with an emergent western lifestyle including a sharp change of dietary patterns (1). Thus, it is reported that many elements of the traditional Mediterranean diet in Albania are substituted by processed foods higher in salt and saturated fats (1, 2). This process of nutritional transition has been paralleled with an increase in the rates of cardiovascular disease in Albania in the past decades (3, 4). To date, however, there are no individually-based data informing about the quantity and quality of food consumption in the Albanian adult population. Therefore, information on the exact consumption and share of processed foods is scant for Albania.

In this context, we undertook a population-based case-control study in Tirana in order to assess the association of acute coronary syndrome (ACS) with selected food groups mainly not from the Mediterranean prototype

including overall oils and fats (except olive oil), sweets and junk food.

2. METHODS

A population-based case-control study of ACS was conducted in 2003-06 among 35-74 year-old residents of Tirana (5). Cases included 467 non-fatal consecutive ACS patients aged 35-74 years, hospitalized at the University Hospital Center (370 men aged 59.1 ± 8.7 years and 97 women aged 63.3 ± 7.1 years; 88% response). The definition of ACS was based on combinations of clinical signs and symptoms, electrocardiographic and echocardiographic criteria (5, 6). The population-based sample comprised an age- and sex-stratified random sample drawn from the adult population of the Tirana municipality. Overall, 469 men aged 53.1 ± 10.4 years and 268 women aged 54.0 ± 10.9 years were recruited (69% response) [5,6].

The data collection included administration of a semi-quantitative food frequency questionnaire (FFQ) and a structured questionnaire including data on socio-demo-

Food groups	Men (N=734)					Women (N=282)				
	Cases (N=361)		Controls (N=373)		p*	Cases N=(90)		Controls N=(192)		p*
	Mean	95%CI	Mean	95%CI		Mean	95%CI	Mean	95%CI	
Butter, margarine, oils, fats (on bread and/or added into food)	79	75-83	58	54-62	<0.01	102	92-111	58	52-65	<0.01
Sweets (cakes, ice-creams)	176	171-182	114	108-120	<0.01	212	198-226	153	144-162	<0.01
Meat and meat products	572	558-586	398	384-412	<0.01	480	454-506	350	333-367	<0.01
Junk food	38	35-41	30	27-33	<0.01	58	49-68	32	25-38	<0.01
“Non-Mediterranean” diet score [junk food + sweets + oils and fats (except olive oil)]	283	274-292	177	169-186	<0.01	364	343-385	215	201-229	<0.01
“Non-Mediterranean” diet score / total calorie intake (percentage of total energy intake)	10.3	10.0- 10.6	5.9	5.6-6.2	<0.01	15.2	14.5- 16.0	8.3	7.9-8.8	<0.01

Table 1. Mean age-adjusted calorie intake (Kcal/day) for selected food groups by Acute Coronary Syndrome status and sex *
Comparison of means by use of the General Linear Model.

graphic and socioeconomic characteristics, conventional coronary risk factors and behavioral factors (5,6).

The FFQ consisted of 105 food items. Participants were asked to indicate how often, on average, they had eaten specified amounts of each food item in the past 12 months. We used nine categories to assess the average frequency of intake of each food item: <1/month, 1-3/month, 1/week, 2-4/week, 5-6/week, 1/day, 2-3/day, 4-5/day, and >6/day. Microdiet, Version 2 (Downlee Systems Limited, UK, 2005) was employed to calculate for each food item the daily calorie intake. The respective values for all 105 food items were added up in order to get a summary score (total daily calorie intake expressed in Kcal) for each participant. For the current analysis, the daily calorie intake (in Kcal) of each food group was calculated.

The study was approved by the Albanian Committee of Medical Ethics. All participants provided written informed consent.

The statistical analysis included 451 ACS patients (361 men and 90 women) and 565 controls (373 men and 192 women) for whom data on nutrient intake were available. Mean age-adjusted values by ACS status, their 95% confidence intervals (CIs) and p-values for selected food groups (meat products, overall oils and fats, sweets, and junk food) were calculated separately in men and women by use of the General Linear Model.

3. RESULTS

The Table presents mean age-adjusted values of selected food groups by ACS status separately in men and women. The analysis indicated that mean age-adjusted values of all food groups presented in the Table were substantially and significantly higher in cases than controls in both sexes (for meat products: 572 Kcal/day in male cases vs. 398 Kcal/day in male controls, and 480 Kcal/day vs. 350 Kcal/day, respectively, in women, $P<0.01$ for both sexes; for overall oils and fats: 79 Kcal/day in male cases vs. 58 Kcal/day in male controls, and 102 Kcal/day vs. 58 Kcal/day, respectively, in women, $P<0.01$ for both sexes; for sweets: 176 Kcal/day in male cases vs. 114 Kcal/day in male controls, and 212 Kcal/day vs. 153 Kcal/day, respectively, in women, $P<0.01$ for both sexes; for junk food: 38 Kcal/day in male cases vs. 30 Kcal/day in male

controls, and 58 Kcal/day vs. 32 Kcal/day, respectively, in women, $P<0.01$ for both sexes).

An overall score referred to as “non-Mediterranean” diet score (consisting of junk food, sweets, oils and fats except olive oil) was calculated for each participant (Table). ACS patients had significantly higher mean “non-Mediterranean” diet scores than controls (10.3% vs. 5.9% in men and 15.2% vs. 8.3% in women, $P<0.01$ for both).

4. DISCUSSION

A salient finding of our study was the significantly higher consumption of “non-Mediterranean” food items (including junk food, sweets, oils and fats except olive oil) in ACS patients than in controls. In this population, there was evidence of a strong positive association of ACS with “non-Mediterranean” diet score expressed both in Kcal/day and as a percentage of total daily calorie intake. In contrast, a Mediterranean prototype diet that reflected a higher consumption of fish, olive oil and fresh fruit and vegetables as well as traditional Albanian dishes was strongly protective (data not shown).

The consumption of processed foods was associated in this Albanian adult population with considerable excess coronary risk which points to serious health implications. This marker of change is an important indicator of the process and extent of transition in Albania that will require targeted public health interventions to avoid its negative consequences, particularly for cardiovascular health and metabolic disorders (1, 2).

This study has some potential limitations given the response rate in the control group (5,6) and the use of FFQ for assessment of dietary patterns. On the face of it, there is no reason to assume differential reporting of dietary patterns in ACS patients as compared to their control counterparts.

Yet, we cannot dismiss the possibility that the acute coronary event may have influenced the reporting of nutrient intake in ACS patients as compared to the population-based control group. However, the FFQs are widely employed for measuring the dietary intake in epidemiological studies of this nature (7).

In conclusion, in this Albanian population, intake of total fats, in particular saturated fatty acids, processed

foods (referred to as “junk” foods), sweets and meat products were all associated with a higher risk of ACS in both sexes.

REFERENCES

1. Ministry of Health of the Republic of Albania. Public Health and Health Promotion Strategy: Towards a healthy country with healthy people. Tirana, Albania, 2004.
2. Shapo L, Pomerleau J, McKee M, Coker R, Ylli A. Body weight patterns in a country in transition: a population-based survey in Tirana city, Albania. *Public Health Nut.* 2003; 6: 471-477.
3. Institute of Statistics of Albania. Health indicators for years 1994-1998. Tirana, INSTAT, 2001.
4. Institute of Statistics of Albania. Causes of deaths for the year 2000. Tirana, INSTAT, 2001.
5. Burazeri G, Goda A, Sulo G, Stefa J, Roshi E, Kark JD. Conventional risk factors and acute coronary syndrome during a period of transition: A population-based case-control study in Tirana, Albania. *Croat Med J.* 2007; 48: 225-233.
6. Burazeri G, Goda A, Kark JD. Religious observance and acute coronary syndrome in predominantly Muslim Albania: a population-based case-control study in Tirana. *Ann Epidemiol.* 2008; 18: 937-945.
7. Willett WC. Food-frequency methods. In: Willett WC (ed.). *Nutritional Epidemiology*, 2nd edition. New York: Oxford University Press, 1998: 74-100.

European Association of Science Editors (EASE)

The European Association of Science Editors (EASE) is an internationally oriented community of individuals from diverse backgrounds, linguistic traditions and professional experience who share an interest in science communication and editing.

[Request new password](#)

[Home](#)
[About us](#)
[Membership](#)
[Publications](#)
[Services](#)
[EASE Events](#)
[Other Events](#)
[Payments](#)

Welcome to EASE

Editing in the Digital World, Tallinn Conference

Registration for Tallinn remains open throughout May. Don't forget that EASE members enjoy significant discounts so ensure your membership is up to date. [See Programme.](#)

We have received many interesting abstracts for **posters** - the latest from Ida Raffaelli.

Tom Babor's abstract on the use of bibliometrics to monitor a field of research is now available for parallel session H.

A new abstract has been added for the parallel session on Digital tools for detecting misconduct. Rachael Lammey of CrossRef will describe both Crosscheck and the new programme, Crossmark.

Parallel session C: unfortunately, Daphne Lees is unable to attend. Eva Baranyiová will speak in her place.

Workshops

There are still some places left at the optional workshops, so if you are already registered and would like to take advantage of doing a workshop while you are at the conference, please email the **EASE secretary** for more information.

EASE Forum

Don't forget that EASE has an interactive **Forum** for discussing various matters pertaining to editing, so if you have a tricky question or want to share a recent

Forthcoming events

Santa Fe Science Writing Workshop
Monday, 7 May, 2012
United States

In the Spotlight

Izet Masic
Bosnia and Herzegovina

Izet Masic is Professor of Family Medicine at the Medical Faculty of the University of Sarajevo and Professor of Social Medicine, Health Care Organization and Health Economics at the Faculty for Health Sciences at the University of Zenica in Bosnia & Herzegovina. He is also Professor of...

[Read more](#)

Jobs

Associate professor of public health
Paris
France
.....
Scientific Editor