Published in final edited form as:

Bioorg Med Chem Lett. 2013 May 15; 23(10): 3034–3038. doi:10.1016/j.bmcl.2013.03.019.

Synthesis and structure-activity relationships of 2-amino-3-carboxy-4-phenylthiophenes as novel atypical protein kinase C inhibitors

Paul M. Titchenell^a, H. D. Hollis Showalter^{b,c}, Jean-François Pons^d, Alistair J. Barber^e, Yafei Jin^b, and David A. Antonetti^f

^aDepartment of Cellular and Molecular Physiology, Penn State University College of Medicine, Hershey, PA 17033, USA

^bVahlteich Medicinal Chemistry Core, University of Michigan, Ann Arbor, MI 48105, USA

^cDepartment of Medicinal Chemistry, University of Michigan, Ann Arbor, MI 48105, USA

^dEvotec (UK) Ltd., 114 Milton Park, Abingdon, OX14 4SA, United Kingdom

^eDepartment of Ophthalmology, Penn State University College of Medicine, Hershey, PA 17033, USA

^fDepartments of Ophthalmology and Visual Sciences and Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI 48105, USA

Abstract

Recent evidence suggests atypical protein kinase C (aPKC) isoforms are required for both TNF- and VEGF-induced breakdown of the blood-retinal barrier (BRB) and endothelial permeability to 70kDa dextran or albumin. A chemical library screen revealed a series of novel small molecule phenylthiophene based inhibitors of aPKC isoforms that effectively block permeability in cell culture and *in vivo*. In an effort to further elucidate the structural requirements of this series of inhibitors, we detail in this study a structure-activity relationship (SAR) built on screening hit **1**, which expands on our initial pharmacophore model. The biological activity of our analogues was evaluated in models of *bona fide* aPKC-dependent signaling including NFκB driven-gene transcription as a marker for an inflammatory response and VEGF/TNF-induced vascular endothelial permeability. The EC₅₀ for the most efficacious inhibitors (**6**, **32**) was in the low nanomolar range in these two cellular assays. Our study demonstrates the key structural elements that confer inhibitory activity and highlights the requirement for electron-donating moieties off the C-4 aryl moiety of the 2-amino-3-carboxy-4-phenylthiophene backbone. These studies suggest that this class has potential for further development into small molecule aPKC inhibitors with therapeutic efficacy in a host of diseases involving increased vascular permeability and inflammation.

© 2013 Elsevier Ltd. All rights reserved.

Correspondence to: David A. Antonetti.

Supplementary Material

Details of the chemical syntheses and assays performed can be found in Supplemental Materials.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

phenylthiophenes; vascular permeability; macular edema; inflammation; protein kinase C; vascular endothelial growth factor; tumor necrosis factor

Retinal vascular permeability and subsequent macular edema (ME) contribute to the loss of vision in a host of blinding diseases. Breakdown of the BRB in response to elevated VEGF or inflammatory cytokines contributes to the pathology of diseases such as diabetic retinopathy, uveitis, and retinal vein occlusions and is associated with vision loss.¹ Biological therapies targeting and binding VEGF have demonstrated excellent results in reducing macular edema and preserving visual function in recent clinical trials for many patients.² Recently, aPKC has been identified as a common signaling molecule that mediates VEGF, TNF, CCL2 and thrombin induced permeability and BRB dysfunction.^{3–6} Drugs targeting aPKC may provide an effective means to control vascular permeability and prevent edema induced by multiple vasoactive cytokines.

Atypical PKC isoforms are a distinct subfamily of the PKC isoforms that do not require Ca^{2+} and diacylglycerol (DAG) for activation. The aPKC isoforms consist of two isozymes, aPKC ζ and aPKC ι (also called λ in mouse), both containing a unique regulatory C1 domain, a Phox and Bem 1 (PB1) protein-protein interacting motif, and a pseudosubstrate domain in the amine terminus with the kinase domain residing within the carboxy terminus.⁷ Genetic loss-of-function experiments for aPKC ι reveal this kinase is intimately involved in the formation of cellular polarity evident from early embryonic lethality.⁸ Importantly, once polarity has been established, depletion of aPKC does not appear to induce cell polarity defects or disorganization of the junctional complex.⁹ Gene deletion of aPKC ζ exhibits only minor defects in immune system development and alterations in NF κ B signaling during the innate immune response.^{10,11} Numerous studies implicate aberrant aPKC signaling associated with diseases involving inflammation and proliferation; therefore, aPKC isoforms have emerged as novel therapeutic targets. Animal studies targeting aPKC isoforms have demonstrated effectiveness in models of type II diabetes, cancer, and inflammation.^{12–14} Hence, targeting aPKC may be an effective treatment for a variety of diseases including macular edema and inflammation.

Screening a Chembridge small molecule library, we identified a series of phenylthiophenes as potent and non-toxic aPKC inhibitors that are effective at preventing both VEGF and TNF-induced retinal permeability in the rodent retina without inducing measurable cell death or observable retinal pathology.^{5,15} These hits exhibit a non-competitive kinetic mechanism of action and maintain selectivity in both kinase screens and in cellular based assays supporting phenylthiophenes as a viable chemotype for aPKC inhibition.¹⁵ Several recent publications have attempted to design small molecule inhibitors of atypical PKCs and some progress has been made with isoform specificity and potency. One report describes a series of allosteric aPKC ζ inhibitors that exhibits micromolar efficacy in the prevention of TNF-induced NF κ B activity.¹³ However this class of inhibitors still exhibit robust activation of PDK1, an upstream activator of aPKC ζ ,¹⁷ which may lead to a less than optimal cellular response as is evident from their efficacy in cell-based assays compared to the phenylthiophene-based inhibitors.

In this manuscript, we report on medicinal chemistry efforts to further characterize and elucidate the inhibitory activity of phenylthiophene based aPKC inhibitors. Based on the screening hit **1** (Table 1), we synthesized a series of analogues to develop and explore structure-activity relationships (SAR) against an aPKC. A subset of the most potent

compounds was then evaluated in cellular assays to determine their efficacy as inhibitors of retinal endothelial permeability and NF κ B activation.

The heterocyclic 2-aminothiophene 3-carboxylic ethyl ester analogues (**1**, **25–28**, **31–39**, Table 2) were prepared in two steps by the Gewald reaction¹⁸ from aryl-substituted acetophenones and ethyl cyanoacetate (Scheme 1a). To make substructure B derivatives shown in Table 1, a two-step sequence of *N*-2 Boc protection and saponification of **1** gave acid **51**, which was further derivatized to intermediate esters **53–66** or amides **67–70** (Table S1, Supplementary Materials) using standard coupling conditions.¹⁹ Boc deprotection with trifluoroacetic acid then provided target analogues **6–20**. This same two-step sequence was applied to selected ethyl esters with different C-4 aromatic substitution patterns (Table 2) to make a subseries of benzyl ester derivatives (Scheme 1b). Analogues **24a** and **24b** (Table 1) were made (Methods A and C, respectively, Supplementary Materials) to compare the kinase activity of **1** with its C-3 nitrile and carboxylic acid congeners.

To make C-4 anilino congeners (**29**, **30**; Table 2) of **1** we started with nitro precursor **94** (Scheme 2). The *N*-2 amino function was bis-Boc protected followed by iron/acetic acid reduction to **96**. Standard alkylation conditions provided *N,N*-dimethyl intermediate **97**. Boc deprotection of **96** and **97** afforded the target compounds.

To explore the effect of modifying the 2-NH₂ on kinase activity, a small series of derivatives was synthesized (Scheme 3). Deamination of **1** under Sandmeyer conditions provided the C-2 protio analogue **2**, whereas simple alkylation gave the mono- and dimethylamino derivatives **3** and **4**, respectively. The urea **5** was prepared in two steps by reaction of **1** with trichloroacetyl isocyanate followed by trichloroacetyl cleavage with ammonia.²⁰

Isosteric replacement of the central thiophene ring of **1** to provide the pyrazole (**100**) and isoxazole (**101**) congeners was achieved by utilizing synthetic procedures for closely related compounds. Thus, condensation of enol ether **99** with hydrazine²¹ or hydroxylamine²² gave the respective target compounds (Scheme 4).

All compounds were rigorously purified by flash chromatography or crystallization, and their structural assignments were supported by diagnostic peaks in the ¹H NMR spectra and by mass spectrometry. Several of the compounds of Tables 1 and 2 are known, but were made to help delineate an SAR for this series. References to prior preparations for each known compound are cited.

SAR for phenylthiophene analogues in an isolated kinase assay

We initially screened the phenylthiophene analogues at, 30 μ M against aPKC ζ in an *in vitro* kinase assay.

Our initial SAR efforts were directed toward exploring variations at the C-2 and C-3 positions of a series of initial phenylthiophene hits, which we had previously characterized.⁴ Two subseries of analogues (**A** and **B**, Table 1) were synthesized with modifications made at R² and R³. For A, it is clear that C-2 amine substitution is required with maximal potency associated with a primary amine (**1** vs **3** vs **4**). Modification to R² = H (**2**) or urea (**5**) essentially abolishes activity

Having established NH₂ as an optimal substituent for R², we then studied a number of ester variations around the C-3 position of the phenylthiophene core (subseries B, Table 1). After confirming earlier results showing significant inhibitory activity (100%) for the ethyl (**1**) and the 2-propyl (**6**) esters,⁴ we synthesized a series of additional ester analogues (**7–19**) with R³ moieties representing a range of physical, electronic and steric properties. The installation of

solubilizing groups (**7**, **8**) as well as phenyl (**9**) decreased inhibitory activity towards aPKC ζ compared to **1**. However, simple homologation of phenyl to benzyl (**10**) restored full inhibition so additional benzyl esters were investigated. Analogues with a range of electron donating and withdrawing substituents at open positions on the phenyl ring were synthesized (**11** – **18**) as well as a single example of a heterocycle (**19**) (Table 1). Here, substituents off the 2- and 3-positions of the phenyl ring (**14**, **15**, **17**) were well tolerated regardless of the electronics whereas 4-position (**16**, **18**) were not. Noteworthy is the remarkable difference in activity due to positioning of fluorine (**15** vs **16**). This, along with the 4-cyano analogue (**18**), suggests that strongly electron-withdrawing moieties at the 4-position are unfavorable, which is reinforced by data for the 4-pyridyl congener (**19**). The mildly electron-donating methyl substituent contributes to modest inhibition regardless of positioning (**11** vs **12** vs **13**). Overall, the ester SAR suggests that the C-3 position of the thiophene core can tolerate lipophilic ester functionality, but with some electronic and geometric constraints.

Selected amide congeners (**20** – **23**, Table 1) were also prepared to evaluate the effect of an ester to amide modification at C-3. XR³ substituents of varying size and lipophilicity showed significantly lower activity suggesting a possible detrimental effect due to a different orientation of the amide substituent relative to ester. Interestingly, a nitrile substituent at R³ (**24a**) demonstrated significant inhibitory activity (87%), whereas the C-3 carboxylic acid analogue (**24b**) exhibited only moderate activity (65% inhibition).

Table 2 expands upon our initial SAR survey by looking at a number of analogues with variable substitution (Z) off the C-4 aryl moiety while retaining optimized substituents established for C-2 and C-3 as shown in Table 1. Analogues with a C-3 ethyl or benzyl ester moiety are shown in subseries **C** and **D**, respectively. A series of electron donating and withdrawing substituents were evaluated at variable positions on the C-4 aryl substituent.

Robust inhibition of aPKC ζ was observed for ethyl ester analogues (subseries **C**) having electron-donating groups in the 3'- and 4'-positions on the phenyl ring (**25** – **34**). Furthermore, this inhibition was generally maintained for corresponding benzyl ester analogues (subseries **D**) where pairwise comparisons can be made (**1** vs **10**, **26** vs **41**, **28** vs **43**, **32** vs **44**, **33** vs **45**). Conversely, electron withdrawing Z-groups for ethyl ester (**35** – **39**) or benzyl ester (**46** – **50**) analogues generally significantly decreased inhibitory activity. In summary, the data of Table 2 suggest that kinase inhibition within the 2-amino-3-carboxy-4-phenylthiophene core is optimized by electron-donating Z-substituents off the C-4 aryl moiety.

The SAR established in Tables 1 and 2 suggests that a generally electron-rich heterocycle is required for optimal kinase activity. To test this hypothesis, we made two congeners of **1** in which the thiophene core was replaced by electron deficient heterocycles pyrazole (**100**) and isoxazole (**101**). Neither of these central core modifications was effective at inhibiting aPKC ζ (30% and 22% inhibition, respectively, at 30 μ M) (Table S2, Supplemental Materials).

Having completed our initial screening, we determined IC₅₀ values for several of our most potent compounds from Tables 1 and 2. These range from IC₅₀ = 1–6 μ M with electron-donating groups at 3'- and 4'-positions on the phenyl ring being the most potent and with analogues **29**, **30**, and **32** demonstrating the highest potency (Table 3).

Evaluation of phenylthiophene inhibitors in cell-based assays

Following IC₅₀ determinations, we evaluated the compounds of Table 3 in two cell-based assays. A single dose of 0.1 μ M was used in an initial screen to determine the ability of a

compound to prevent TNF-induced NF κ B transcriptional activation in a stable cell line (HEK293) expressing a NF κ B reporter system. This assay is a surrogate for downstream signaling of aPKC ζ as this kinase has been shown to regulate NF κ B activation in multiple cell types as evidenced from an animal knockout model.³⁶ Importantly, this assay has been used previously to determine aPKC ζ inhibitor cellular efficacy.¹³ A robust increase of NF κ B by TNF was demonstrated and complete inhibition of TNF-induced NF κ B activation was verified using an I κ B kinase inhibitor VII (IKK-I) (Figure S1, Supplemental Materials). Using this assay, six analogues (**6**, **10**, **14**, **29**, **30**, **32**) were tested and all demonstrated significant inhibition of TNF-induced NF κ B activity and were further analyzed using dose-response curves to determine EC₅₀ values as depicted for compound **6** (Figure 1A). Each compound demonstrated low nanomolar potency toward suppressing NF κ B activity (Table 3), demonstrating that these analogues exhibit potent anti-inflammatory effects. Importantly, this assay is a *bona fide* marker of aPKC activity and downstream signaling, and cellular data correlates with *in vitro* enzyme data supporting aPKC as the cellular target.

We further evaluated compounds of Table 3 in an assay to assess their capability to prevent retinal endothelial permeability, a hallmark of ME. This is a permeability assay that was performed using retinal endothelial cells and monitoring the flux of 70kDa RITC-dextran across an endothelial monolayer. This assay has been used extensively to monitor the permeabilizing effects of growth factors and inflammatory cytokines implicated in ocular disease³⁷ and requires aPKC ι .¹⁵ Dose response curves were generated with **6**, **14**, **29**, **30** and **32** with the results shown in Table 3 along with a representative dose response curve for **6** (Figure 1B). Four compounds, (**6**, **29**, **30**, **32**) showed good efficacy in preventing TNF/VEGF-induced endothelial permeability, with compound **6** exhibiting an EC₅₀ of 1 nM.

The cell based efficacy data is supported with a favorable specificity profile for our better candidates. Importantly, **6** has been shown to exhibit a high degree of specificity towards the aPKC isoforms as compared to its closest AGC family members and does not block VEGF-induced Akt activation or Erk activation in endothelial cells.¹⁵ Compounds **6**, **14**, **30** and **32**-display low micromolar to high nanomolar potency against aPKC isoforms without discrimination for PKC ζ or PKC ι (Table S3, Supplemental Materials). A preliminary *in silico* calculation of physico-chemical parameters (Table S4, Supplemental Materials) for selected compounds of Table 3 suggests favorable biopharmaceutical properties for this class compounds but clearly further stability, pharmacokinetic and toxicity analyses are required. cLogP values for compounds **6**, **14**, **30** and **32** are 4.256, 5.165, 4.459, and 3.854. In addition, topological polar surface area (TPSA) for compounds **6**, **14**, **30** and **32** are 70.78, 80.00, 55.55, and 70.78, respectively. Overall, the enzyme and cellular data support this class of phenylthiophenes as potent and specific inhibitors of aPKC isoforms with significant efficacy in preventing vascular permeability and NF κ B-induced gene transcription.

Previously, we reported a group of phenylthiophene based molecules as the first biologically active aPKC inhibitors with high enzyme specificity and a non-competitive mode of action with nanomolar efficacy in cells.¹⁵ In this paper, we have expanded the screening for aPKC inhibitors by developing an SAR to further delineate structural requirements and refine a pharmacophore for this new class of aPKC inhibitors. The addition of electron donating groups (Z) to the C-4 aryl substituent, principally in the 3'- and 4'-positions, combined with a C-2 primary amine and C-3 ester moiety confer robust inhibitory activity for this class of compounds. The reason for the discrepancy between kinase potency (IC₅₀ low micromolar) and cellular potency (EC₅₀ low nanomolar) is unclear but does not appear to be due to off-target effects as **6** has been shown to possess excellent selectivity in a panel of 20 AGC kinases and this inhibitor class has been demonstrated to not affect canonical VEGF signaling cascades.¹⁵

Our series of phenylthiophenes are the first compounds to exhibit low nanomolar activity against aPKC activity in cells with biomarkers of both NF κ B dependent gene expression and TNF/VEGF-induced vascular permeability. Our data suggest that the phenylthiophene pharmacophore can be further optimized to provide therapeutic agents with the potential to treat diseases, such as macular edema, that involve vascular permeability induced by growth factors such as VEGF and inflammatory cytokines such as TNF. More broadly we believe that expansion of this class can lead to agents that target acute inflammatory responses.

Future SAR will focus on molecular modifications that further expand upon the current series to optimize drug-like properties while maintaining aPKC selectivity and improving potency. These will entail modifications of the phenylthiophene core, which will include exploration of the C-5 position. Our current studies provide a good starting point to achieve this goal.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by research grants from JDRF, NIH (R01 EY012021), and support from Research to Prevent Blindness (DAA). HDHS and YJ acknowledge generous support by the University of Michigan College of Pharmacy Ella and Hans Vahlteich Research Fund.

References and notes

1. Pan PS, Vasko RC, Lapera SA, Johnson VA, Sellers RP, Lin CC, Pan CM, Davis MR, Ardi VC, McAlpine SR. *Bioorg Med Chem.* 2009; 17:5806. [PubMed: 19643615]
2. Nguyen QD, Brown DM, Marcus DM, Boyer DS, Patel S, Feiner L, Gibson A, Sy J, Rundle AC, Hopkins JJ, Rubio RG, Ehrlich JS. *Ophthalmology.* 2012; 119:789. [PubMed: 22330964]
3. Dimitrijevic OB, Stamatovic SM, Keep RF, Andjelkovic AV. *Journal of cerebral blood flow and metabolism: official journal of the International Society of Cerebral Blood Flow and Metabolism.* 2006; 26:797. [PubMed: 16192992]
4. Titchenell PM, Lin CM, Keil JM, Sundstrom JM, Smith CD, Antonetti DA. *Biochem J.* 2012; 446:455. [PubMed: 22721706]
5. Aveleira CA, Lin CM, Abcouwer SF, Ambrosio AF, Antonetti DA. *Diabetes.* 2010; 59:2872. [PubMed: 20693346]
6. Minshall RD, Vandenbroucke EE, Holinstat M, Place AT, Tirupathi C, Vogel SM, van Nieuw Amerongen GP, Mehta D, Malik AB. *Microvasc Res.* 2010; 80:240. [PubMed: 20417648]
7. Steinberg SF. *Physiol Rev.* 2008; 88:1341. [PubMed: 18923184]
8. Moscat J, Rennert P, Diaz-Meco MT. *Cell Death and Differentiation.* 2006; 13:702. [PubMed: 16322752]
9. Suzuki A, Yamanaka T, Hirose T, Manabe N, Mizuno K, Shimizu M, Akimoto K, Izumi Y, Ohnishi T, Ohno S. *J Cell Biol.* 2001; 152:1183. [PubMed: 11257119]
10. Leitges M, Sanz L, Martin P, Duran A, Braun U, Garcia JF, Camacho F, Diaz-Meco MT, Rennert PD, Moscat J. *Mol Cell.* 2001; 8:771. [PubMed: 11684013]
11. Bandyopadhyay G, Standaert ML, Sajan MP, Kanoh Y, Miura A, Braun U, Kruse F, Leitges M, Farese RV. *Molecular Endocrinology.* 2004; 18:373. [PubMed: 14615604]
12. Sajan MP, Nimal S, Mastorides S, Acevedo-Duncan M, Kahn CR, Fields AP, Braun U, Leitges M, Farese RV. *Metabolism: Clinical and Experimental.* 2012; 61:459. [PubMed: 22225955]
13. Frohner W, Lopez-Garcia LA, Neimanis S, Weber N, Navratil J, Maurer F, Stroba A, Zhang H, Biondi RM, Engel M. *Journal of Medicinal Chemistry.* 2011; 54:6714. [PubMed: 21863889]
14. Murray NR, Kalari KR, Fields AP. *Journal of Cellular Physiology.* 2011; 226:879. [PubMed: 20945390]

15. Titchenell PM, Lin CM, Keil JM, Sundstrom JM, Smith CD, Antonetti DA. The Biochemical journal. 2012; 446:455. [PubMed: 22721706]
16. Baell JB, Holloway GA. J Med Chem. 2010; 53:2719. [PubMed: 20131845]
17. Balendran A, Biondi RM, Cheung PC, Casamayor A, Deak M, Alessi DR. J Biol Chem. 2000; 275:20806. [PubMed: 10764742]
18. Buchstaller HP, Siebert CD, Lyssy RH, Frank I, Duran A, Gottschlich R, Noe CR. Monatsh Chem. 2001; 132:279.
19. Montalbetti CAGN, Falque V. Tetrahedron. 2005; 61:10827.
20. Janetka JW, Almeida L, Ashwell S, Brassil PJ, Daly K, Deng C, Gero T, Glynn RE, Horn CL, Ioannidis S, Lyne P, Newcombe NJ, Oza VB, Pass M, Springer SK, Su M, Toader D, Vasbinder MM, Yu D, Yu Y, Zabudoff SD. Bioorg Med Chem Lett. 2008; 18:4242. [PubMed: 18547806]
21. Velicelebi, G.; Stauderman, KA.; King, F.; Pei, Y.; Whitten, JP. PCT Int Appl WO. 2009076454A2..
22. Breinlinger, EC.; Cusack, KP.; Hobson, AD.; Li, B.; Gordon, TD.; Stoffel, RH.; Wallace, GA.; Gronsgaard, P.; Wang, L. PCT Int Appl WO. 2009011850A2..
23. Shvedov VI, Ryzhkova VK, Grinev AN. Khim Geterotsikl Soedin. 1967:239.
24. Chen, D.; Lee, KC.; Terrell, LR. PCT Int Appl WO. 2011075559A1..
25. Moorman, AR.; Romagnoli, R.; Baraldi, PG. PCT Int Appl WO. 2002083083A2..
26. Nara, H.; Kaieda, A.; Sato, K.; Terauchi, J. PCT Int Appl WO. 2007049820A1..
27. Cravo, D.; Hallakou-Bozec, S.; Bolze, S.; Lepifre, F.; Faveriel, L.; Durand, J-D.; Charon, C. PCT Int Appl WO. 2011080277A1..
28. Stein, RL.; Case, A.; Yeh, L-A.; Cuny, G.; Duval, E. PCT Int Appl WO. 2006060702A1..
29. Duval E, Case A, Stein RL, Cuny GD. Bioorg Med Chem Lett. 2005; 15:1885. [PubMed: 15780627]
30. Kathiravan MK, Shishoo CJ, Chitre TS, Mahadik KR, Jain KS. Synth Commun. 2007; 37:4273.
31. Tormyshev VM, Trukhin DV, Rogozhnikova OY, Mikhulina TV, Troitskaya TI, Flinn A. Synlett. 2006:2559.
32. Hodge, CN.; Janzen, WP.; Williams, KP.; Cheatham, LA. PCT Int Appl WO. 2005033102A2..
33. Wang, Y.; Yang, T. PCT Int Appl WO. 2012100732A1..
34. Scammells, PJ.; Aurelio, L.; Christopoulos, A.; Sexton, PM.; Valant, C.; Flynn, BL.; Linden, JM. PCT Int Appl WO. 2009049362A1..
35. Aurelio L, Figler H, Flynn BL, Linden J, Scammells PJ. Bioorg Med Chem. 2008; 16:1319. [PubMed: 17980606]
36. Diaz-Meco MT, Moscat J. Immunological Reviews. 2012; 246:154. [PubMed: 22435553]
37. Antonetti DA, Wolpert EB. Methods Mol Med. 2003; 89:365. [PubMed: 12958433]
38. Huth JR, Mendoza R, Olejniczak ET, Johnson RW, Cothron DA, Liu Y, Lerner CG, Chen J, Hajduk PJ. Journal of the American Chemical Society. 2005; 127:217. [PubMed: 15631471]
39. Mansuy D. Journal of hepatology. 1997; 26(Suppl 2):22. [PubMed: 9204406]
40. Fouda HG, Avery MJ, Dalvie D, Falkner FC, Melvin LS, Ronfeld RA. Drug Metab Dispos. 1997; 25:140. [PubMed: 9029043]
41. Medower C, Wen L, Johnson WW. Chem Res Toxicol. 2008; 21:1570. [PubMed: 18672911]
42. Dalvie DK, Kalgutkar AS, Khojasteh-Bakht SC, Obach RS, O'Donnell JP. Chem Res Toxicol. 2002; 15:269. [PubMed: 11896674]
43. Pereillo JM, Maftouh M, Andrieu A, Uzabiaga MF, Fedeli O, Savi P, Pascal M, Herbert JM, Maffrand JP, Picard C. Drug Metab Dispos. 2002; 30:1288. [PubMed: 12386137]

Figure 1.

Examples of cell-based efficacy of phenylthiophenes

A) HEK293pNFkB cells were treated where indicated with 5 ng/ml TNF for 5 h. A 3 h pre-treatment of a dose-response of **6** was performed where indicated and a Bright-Glo luciferase reaction was performed (SM methods); $n > 5$. B) Permeability assays were performed (SM methods) following a dose-response of compound **6** and the addition of VEGF/TNF for 1.5 h; $n = 7$ combined from multiple independent experiments. Statistics were performed using One way ANOVA with Neuman-Keuls posthoc test in GraphPad.

Scheme 1^a

^a Reagents and conditions: (i) methyl cyanoacetate, NH_4OAc , AcOH , toluene, reflux, 18–48 h; (ii) sulfur powder, Et_2NH , EtOH , 50 °C, 3 h (iii) Boc_2O , DMAP, pyridine, 55 °C, 3 h; (iv) KOH , aq. EtOH , reflux, 5 h; (v) for **52**: R^3OH , CDI, DCM or R^3OH , DCC, THF or R^3OH , Ph_3P , DEAD, THF; (vi) TFA, DCM; (vii) for **52**: R^3NH_2 , EDC, HOBt, NEt_3 , DMF, 24 h.

Scheme 2^a

^a Reagents and conditions: (i) Boc_2O , DMAP, pyridine, 55 °C, 5 h; (ii) for **95**: Fe powder, AcOH, 60 °C, 1 h; (iii) MeI, K_2CO_3 , acetone, 60 °C; (iv) TFA, DCM, rt, 16 h; (v) EtOH, 130 °C, 48 h.

Scheme 3^a

^a Reagents and conditions: (i) *t*-BuONO, CuCl₂, EtOH, 1 h, then aq. NH₄Cl; (ii) MeI, K₂CO₃, acetone 60 °C, 16 h; (iii) MeI, NaH, DMF, rt, 1 h; (iv) trichloroacetyl isocyanate, THF, 0 °C, 3 h; (v) NH₃, MeOH, 0 °C, 1 h.

Scheme 4^a

^a Reagents and conditions: (i) ethyl cyanoacetate, NaH, THF, then Me_2SO_4 , NaHCO_3 , aq. dioxane; (ii) for **100**: $\text{NH}_2\text{NH}_2 \cdot \text{HCl}$, NEt_3 , EtOH, reflux; for **101**: $\text{NH}_2\text{OH} \cdot \text{HCl}$, NEt_3 , EtOH, reflux.

Table 1

Initial phenylthiophene SAR vs. aPKC ζ^a

<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>A</p> </div> <div style="text-align: center;"> <p>B</p> </div> </div>			
No.	Str.	R ² or XR ³	Inh (%) ^a
1 ²³	A	NH ₂	100
2	A	H	3
3	A	NHMe	32
4	A	NMe ₂	60
5	A	NHCONH ₂	14
6	B	O- <i>i</i> -Pr	100
7	B		57
8	B		49
9	B	OPh	48
10	B	OCH ₂ Ph	100
11	B	OCH ₂ Ph-2-Me	64
12	B	OCH ₂ Ph-3-Me	67
13	B	OCH ₂ Ph-4-Me	50
14	B	OCH ₂ Ph-3-OMe	100
15	B	OCH ₂ Ph-2-F	100
16	B	OCH ₂ Ph-4-F	2
17	B	OCH ₂ Ph-3-CN	70
18	B	OCH ₂ Ph-4-CN	10
19	B	OCH ₂ -4-pyridyl	53
20	B	NHEt	50
21	B		55
22	B	NHCH ₂ Ph	43
23	B	NHCH ₂ -4-pyridyl	3
24a ²⁴	B	COXR ³ = CN	87
24b	B	COXR ³ = CO ₂ H	65

^aInhibition determined from ADP Quest assay with 30 μ M of inhibitor using aPKC ζ (500 ng/ml). Values are the mean of at least n = 5 and were repeated in independent experiments. Standard error of the mean <20%. The experimental conditions are reported in the Supplementary Material.

Str. = chemical structure.

Table 2

4-Aryl SAR of phenylthiophenes vs. aPKC ζ ^a

Z	No.	Str.	Inh (%) ^a	No.	Str.	Inh
3',4'-(OMe) ₂	1	C	100	10	D	100
3'-Me	25 ^{25, 26}	C	77	40	D	68
4'-Me	26 ^{24, 26-28, 30}	C	80	41	D	60
3'-OMe	27 ^{24, 26-29}	C	82	42	D	57
4'-OMe	28 ^{23, 28-31}	C	100	43	D	86
4'-NH ₂	29 ^{31, 32}	C	100	----	D	n.d.
4'-NMe ₂	30	C	100	----	D	n.d.
3'-NHAc	31 ³²	C	89	----	D	n.d.
3',4'-OCH ₂ O-	32 ³²	C	100	44	D	85
2'-F,4'-OMe	33	C	100	45	D	78
3'-F,4'-OMe	34	C	100		D	n.d.
3'-CF ₃	35 ³³⁻³⁵	C	30	46	D	29
4'-CF ₃	36 ³²	C	16	47	D	21
2'-F	37	C	62	48	D	57
3'-F	38 ³²	C	38	49	D	53
3'-CN	39 ^{26, 31}	C	32	50	D	25

^aSee footnote a, Table 1.

Table 3

Cellular assays for selected phenylthiophenes

No.	aPKC kinase assay		TNF-induced NFκB activation		TNF/VEGF-induced endothelial permeability EC ₅₀ (μM) ^c
	Inh. (%) ^a	IC ₅₀ (μM) ^a	Inh. (%) ^b	EC ₅₀ (μM) ^b	
6	100	6	90	0.003	0.001
10	100	6	64	0.002	n.d.
14	100	5	80	0.001	>0.1
29	100	2	72	0.001	0.27
30	100	1	37	0.001	0.28
32	100	2	34	0.002	0.02
33	100	4	65	0.01	n.d.

^aInhibition determined from ADP Quest assay with 30 μM (reported in Tables 1 & 2) or a 5 point dose-response curve to determine IC₅₀ against aPKCζ (500 ng/ml). Values are the mean of at least n 5 and were repeated in independent experiments. Standard error of the mean <20%.

^bInhibition determined from TNF-induced NFκB activity assay with 0.1 μM or a 5 point dose-response curve to determine EC₅₀. Values are the mean of at least n>4 and were repeated in independent experiments. Standard error of the mean <35%.

^cEC₅₀ determined from VEGF/TNF-induced retinal endothelial permeability assay using a 5 point dose response curve. Values are the mean of at least n 8 and were repeated in independent experiments. Standard error of the mean <30%. All experimental conditions are reported in the supplementary methods. IC₅₀ and EC₅₀ values were calculated using a variable slope dose-response curve in GraphPad (SM methods).