ORIGINAL ARTICLE

Open Access

Facile synthesis of symmetrical bis(benzhydryl) ethers using *p*-toluenesulfonyl chloride under solvent-free conditions

Goutam Brahmachari* and Bubun Banerjee

Abstract

Background: The benzhydryl ether moiety is widely distributed in nature and constitutes a key structural motif in numerous molecules of significant biological potential and of prospective clinical uses. Solvent-free and cost-effective facile synthesis of symmetrical bis(benzhydryl)ethers is, thus, much desirable.

Results: A simple and efficient method for the facile synthesis of symmetrical bis(benzhydryl)ethers directly from the corresponding benzhydrols has been developed using a catalytic amount of *p*-toluenesulfonyl chloride (5 mol%) at an oil bath temperature of 110°C under solvent-free conditions.

Conclusions: Operational simplicity, low reagent loading, high product yields, short reaction time, and solvent-free conditions are the notable advantages of the present method.

Keywords: Bis(benzhydryl)ethers, Benzhydrols, *p*-Toluenesulfonyl chloride, Solvent-free

Background

The benzhydryl ether moiety is abundant in a number of naturally occurring and biologically active compounds as well as molecules of potential clinical uses [1-8]; this motif was also found as a partial structure in a few new chemical entities showing therapeutic activity as well [9]. A number of reports are available describing the synthesis of molecules bearing this structural motif, which were shown to exhibit various pharmacological potentials such as non-nucleoside reverse transcriptase inhibition [10], anti-plasmodial and anti-trypanosomal action [11], monoamine uptake inhibition, anti-depressant and anti-parkinsonian activity [12,13], and anti-histaminic [14] and anti-spasmodic [15] action. Naturally occurring symmetrical bis(benzhydryl)ethers are also known to show promising therapeutic potentials including significant anti-platelet aggregation efficacy [16]. Very recently, application of such ether substructures in the total syntheses of a number of natural products has nicely been reviewed by Pitsinos et al. [17]. Although there are a good number of reports on the synthetic methodology of

diaryl ethers, there are only two such reports so far on bis(benzhydryl)ethers in the literature [18-20]; symmetrical bis(benzhydryl)ethers were conventionally synthesized from corresponding benzhydrols using 100% sulfuric acid in large excess [18-20] and *p*-toluenesulfonic acid in equivalent amount [21]. Both of these earlier methods require the use of strong acids in relatively large excess. Under this purview, we have been motivated to undertake systematic planning to develop a convenient and efficient protocol for the conversion of benzhydrols into their bis(benzhydryl)ether derivatives.

In continuation of our effort to develop green and solvent-free synthetic methodologies for organic transformations [22-28], we wish to report in this communication a convenient and straightforward protocol for the efficient synthesis of symmetrical bis(benzhydryl)ethers in excellent yields using a catalytic amount of *p*-toluenesulfonyl chloride under solvent-free conditions (Scheme 1). The process is very simple, cost-effective, and environmentally benign.

Methods

Infrared spectra were recorded using a Shimadzu (FT-IR 8400S) Fourier transform infrared (FT-IR) spectrophotometer (Shimadzu, Kyoto, Japan) using KBr disc. ¹H and ¹³C nuclear magnetic resonance (NMR) spectra were

* Correspondence: brahmg2001@yahoo.co.in
Laboratory of Natural Products and Organic Synthesis, Department of Chemistry, Visva-Bharati University, Santiniketan, West Bengal 731 235, India

Scheme 1 Synthesis of symmetrical bis(benzhydryl)ethers.

obtained at 400 and 100 MHz, respectively, using a Bruker DRX400 spectrometer (Bruker Instruments, Billerica, MA, USA) and CDCl₃ as the solvent. Mass spectra (time-of-flight mass spectrometry (TOF-MS)) were measured on a Q-ToF MicroTM mass spectrometer (Waters MS Technologies, Manchester, UK). Elemental analyses were performed with an Elementar Vario EL III Carlo Erba 1108 micro-analyzer instrument (Carlo Erba Reagenti SpA, Rodano, Italy). Melting point was recorded on a Sunvic melting point apparatus (Sunvic, Glasgow, UK) and is uncorrected. Column chromatography was carried out over silica gel (60 to 120 mesh, Merck & Co., Inc., Whitehouse Station, NJ, USA), and thin layer chromatography (TLC) was performed using silica gel 60 F₂₅₄ (Merck) plates.

Results and discussion

Firstly, we carried out the synthesis of bis(benzhydryl) ether **1** from benzhydryl as our model reaction in order to optimize the best suited reaction conditions (Figure 1); it was observed (Table 1) that the alcohol in the presence of *p*-TsCl (5 mol%) afforded the best result with 86% isolated yield at 110°C within a short period of time (15 min) under solvent-free conditions.

Figure 1 Optimization of the reaction conditions.

A number of benzhydryl derivatives containing mono- and di-chloro, mono-bromo, di-fluoro, mono-methoxy, and mono-methyl phenyl groups were then screened for studying the generality as well as the efficacy of this present procedure (Figure 2; Table 2). All the entries find an easy and efficient route to their symmetrical bis (benzhydryl)ether derivatives in the presence of *p*-TsCl under solvent-free conditions (Figure 2) within 8 to 15 min affording excellent yields (85% to 92%). The workup of the reaction mixtures is simple and highly convenient. Each product has been characterized by detailed spectral analyses including FT-IR, ¹H NMR, ¹³C NMR, and TOF-MS. In addition, the molecular structure of bis(bis-

Table 1 Optimization of the reaction conditions following Figure 1

Entry	<i>p</i> -TsCl (mol%)	Solvent	Temperature (°C)	Time (min)	Yield (%) ^a
1	0	-	110	120	No reaction
2	10	-	90	25	82
3	10	-	110	15	87
4	5	-	90	35	80
5	5	-	110	15	86
6	5	-	Rt	240	Trace
7	3	-	110	75	28
8	10	CH ₂ Cl ₂	Rt	1,050	19
9	5	CH ₂ Cl ₂	Rt	1,050	Trace
10	10	THF	Rt	600	Trace
11	10	CH ₃ CN	Rt	720	17
12	10	CH ₂ Cl ₂	Reflux	300	36
13	50	-	110	5	43 (tosylate: 47)
14	100	-	110	5	7 (tosylate: 91)

^aIsolated yield; RT, room temperature; *p*-TsCl, *p*-toluenesulfonyl chloride; Tosylate, benzhydryl *p*-toluenesulfonate.

Figure 2 Synthesis of symmetrical bis(benzhydryl)ethers using *p*-TsCl as reagent under solvent-free conditions.

phenylmethyl)ether (Table 2, entry 1) has unambiguously been confirmed from X-ray crystallographic analysis [29-33] (Figure 3).^a

We propose the following mechanistic pathway for the reaction (Scheme 2). *p*-TsCl reacts rapidly with an equivalent amount of diarylmethanol to generate HCl (and a tosylate derivative as side product) *in situ* that eventually catalyzes the etherification following a catalytic cycle. The corresponding tosylate derivative remains intact as side product. To ensure the fact, we have checked the reaction with 50 and 100 mol% *p*-TsCl in two separate entries (entries 13 and 14; Table 1) where benzhydryl *p*-toluenesulfonate was isolated as 47% and 91% yields, respectively. In addition, we have carried out the reaction with benzhydrol separately using dry HCl gas (passed for a while into the reaction vessel) and concentrated hydrochloric acid; it has been found that dry HCl could also act as an efficient catalyst producing the corresponding ether derivative **1** with 82% yield in 30 min at 110°C, while concentrated HCl (45 mg added to 1 mmol of benzhydrol) required much more time (45 min) producing less amount of yield (62%) at the same reaction temperature. These experimental observations support our proposed mechanistic pathway as well.

Experimental

General procedure for the synthesis of symmetrical bis(benzhydryl)ethers (entries 1 to 7)

An oven-dried screw cap test tube was charged with a magnetic stir bar, benzhydrol (1 mmol), and *p*-toluenesulfonyl chloride (5 mol%). The tube was then evacuated and back-filled with nitrogen. The evacuation/backfill sequence was repeated two additional times. The tube was placed in a preheated oil bath at 110°C, and the reaction mixture was stirred vigorously. The progress of the reaction was monitored by TLC, and on completion, the reaction mixture was cooled to room temperature. The reaction mixture was extracted with dried ethyl acetate (10 ml), and the extract was then concentrated under reduced pressure; the residue was purified via column chromatography using silica gel (60

to 120 mesh) and petrol ether-ethyl acetate mixture. The structure of each purified symmetrical bis(benzhydryl)ethers was confirmed by analytical as well as spectral studies including FT-IR, ¹H NMR, ¹³C NMR, and TOF-MS. Respective physical and spectral properties of bis(diarylmethyl)ethers are described below.

The spectral and analytical data of all the compounds including all new entries are given below (see also Additional file 1):

- *Bis(bis-phenylmethyl)ether (1)*: white solid, 86% yield, m.p. 106°C to 107°C (Lit. 105°C to 107°C [34], 107°C [18]). IR ($\nu_{\max}$, KBr) cm⁻¹: 3,057, 3,028, 2,953, 1,595, 1,489, 1,445, 1,250, 1,163, 1,098, 1,072, 1,029, 6,98. ¹H NMR (CDCl₃, 200 MHz, δ): 7.40 to 7.23 (m, 20H, Ar H), 5.41 (s, 2H, CH). ¹³C NMR (CDCl₃, 100 MHz, δ): 142.28, 128.45, 127.51, 127.33, 80.05. TOF-MS: 373.44 ([M + Na]⁺). Anal. found: C, 89.13; H, 6.28. C₂₆H₂₂O requires C, 89.11; H, 6.33%
- *Bis[[1-(4-methylphenyl)-1-phenyl]methyl]ether (2)*: yellowish white, semi solid, 89% yield. IR ($\nu_{\max}$, KBr) cm⁻¹: 3,060, 3,025, 2,923, 2,852, 1,655, 1,460, 1,277, 1,124, 1,071, 824, 810, 699. ¹H NMR (CDCl₃, 400 MHz, δ): 7.6 (d, 4H, Ar H, *J* = 7.6 Hz), 7.53 to 7.45 (m, 8H, Ar H), 7.43 to 7.41 (m, 2H, Ar H), 7.34 (d, 4H, Ar H, *J* = 7.6 Hz), 5.63 (s, 2H, CH), 2.53 (s, 6H, CH₃). ¹³C NMR (CDCl₃, 100 MHz, δ): 142.82, 142.70, 139.59, 139.48, 137.26, 137.22, 129.33, 129.30, 128.57, 128.54, 127.53, 127.49, 127.46, 127.41, 127.34, 79.96, 21.37. TOF-MS: 401.05 ([M + Na]⁺). Anal. found: C, 89.89; H, 6.90. C₂₈H₂₆O requires C, 89.85; H, 6.92%
- *Bis[[1-(4-chlorophenyl)-1-phenyl]methyl]ether (3)*: white semi solid, 85% yield. IR ($\nu_{\max}$, KBr) cm⁻¹: 3,063, 3,029, 2,925, 2,854, 1,595, 1,490, 1,449, 1,259, 1,185, 1,086, 1,057, 843, 811, 700. ¹H NMR (CDCl₃, 400 MHz, δ): 7.31 to 7.30 (m, 8H, Ar H), 7.28 to 7.25 (m, 10H, Ar H), 5.33 (s, 2H, CH). ¹³C NMR (CDCl₃, 100 MHz, δ): 141.49, 141.39, 140.65, 140.54, 133.37, 133.30, 128.69, 128.63, 128.60, 128.56, 128.48, 127.87, 127.81, 127.22, 127.13, 79.53. TOF-

Table 2 Synthesis of symmetrical bis(benzhydryl)ethers using *p*-TsCl as reagent under solvent-free conditions following Figure 2

Entry	Alcohol	Product	Time (min)	Yield (%) ^a	Melting point (°C)	
					Found	Reported
1		1	15	86	106 to 107	105 to 107 [34]
2		2	15	89	Semisolid	Present work
3		3	10	85	Semisolid	Present work
4		4	10	92	Semisolid	Present work
5		5	10	88	125 to 127	126 to 127 [34,35]
6		6	8	91	88 to 90	Present work
7		7	12	90	Semisolid	Present work

^aIsolated yield.

MS: 441.94 ([M + Na]⁺). Anal. found: C, 74.45; H, 4.83. C₂₆H₂₀Cl₂O requires C, 74.47; H, 4.81%

- *Bis*[[1-(4-bromophenyl)-1-phenyl]methyl]ether (4): white semi solid, 92% yield. IR (ν_{max}, KBr) cm⁻¹: 3,085, 3,062, 3,028, 2,924, 2,854, 1,602, 1,590, 1,486, 1,454, 1,290, 1,185, 1,107, 1,070, 1,028, 847, 793, 700. ¹H NMR (CDCl₃, 400 MHz, δ): 7.33 (dd, 4H, Ar H, *J* = 8.4, 5.2 Hz), 7.21 to 7.15 (m, 10H, Ar H), 7.12 (dd, 4H, Ar H, *J* = 8.4, 3.2 Hz), 5.23 (s, 2H, CH). ¹³C NMR (CDCl₃, 100 MHz, δ): 141.43, 141.33, 141.20, 141.08, 131.68, 131.62, 128.94,

128.86, 128.70, 128.65, 127.94, 127.87, 127.26,

127.17, 121.60, 121.52, 79.61. TOF-MS: 528.74 ([M + Na]⁺). Anal. found: C, 61.49; H, 3.93. C₂₆H₁₈Br₂O requires C, 61.44; H, 3.97%

- *Bis*[bis(4-chlorophenyl)methyl]ether (5): white solid, 88% yield, m.p. 125°C to 127°C (Lit. 126°C to 127°C) [35,36]. IR (ν_{max}, KBr) cm⁻¹: 3,031, 2,924, 1,594, 1,491, 1,410, 1,290, 1,188, 1,089, 1,013, 854, 824, 735, 726. ¹H NMR (CDCl₃, 200 MHz, δ): 7.31 (d, 8H, Ar H, *J* = 8.6 Hz), 7.23 (d, 8H, Ar H, *J* = 8.6 Hz), 5.29 (s, 2H, CH). ¹³C NMR (CDCl₃, 75 MHz,

Figure 3 Diagram and packing arrangement. (a) ORTEP diagram of compound **1** (CCDC 840259). (b) The packing arrangement of molecules viewed down the a-axis.

δ): 139.72, 133.71, 128.82, 128.36, 78.97. TOF-MS: 509.12 ($[M + Na]^+$). Anal. found: C, 63.94; H, 3.69; $C_{26}H_{18}Cl_4O$ requires C, 63.96; H, 3.72%

- *Bis[bis[4-fluorophenyl]methyl]ether* (**6**): white solid, 91% yield, m.p. 88°C to 90°C. IR (ν_{max} , KBr) cm^{-1} : 3,069, 3,057, 2,925, 1,603, 1,507, 1,422, 1,408, 1,298, 1,225, 1,178, 1,155, 1,101, 1,029, 859, 837, 818. 1H NMR ($CDCl_3$, 400 MHz, δ): 7.19 to 7.16 (m, 8H, Ar H), 6.94 to 6.88 (m, 8H, Ar H), 5.22 (s, 2H, CH). ^{13}C NMR ($CDCl_3$, 100 MHz, δ): 163.52, 161.07, 137.51, 137.48, 128.82, 128.74, 115.59, 115.38, 78.91. TOF-MS: 445.98 ($[M + Na]^+$). Anal. found: C, 73.89; H, 4.28. $C_{26}H_{18}F_4O$ requires C, 73.93; H, 4.30%
- *Bis[[1-(4-methoxyphenyl)-1-phenyl]methyl]ether* (**7**): colorless liquid, 90% yield. IR (ν_{max} , KBr) cm^{-1} : 3,062, 3,029, 2,953, 2,932, 2,906, 2,835, 1,510, 1,494, 1,451, 1,249, 1,171, 1,111, 1,080, 849, 819, 698. 1H NMR ($CDCl_3$, 400 MHz, δ): 7.35 (d, 4H, Ar H, J =

7.6 Hz), 7.32 to 7.28 (m, 4H, Ar H), 7.27 to 7.24 (m, 6H, Ar H), 6.84 (d, 4H, Ar H, J = 8.4 Hz), 5.34 (s, 2H, CH), 3.77 (s, 6H, OCH_3). ^{13}C NMR ($CDCl_3$, 100 MHz, δ): 158.97, 158.93, 142.72, 142.52, 134.53, 134.32, 128.67, 128.60, 128.35, 128.32, 127.30, 127.24, 127.17, 127.09, 113.80, 113.77, 79.43, 79.40, 55.26. TOF-MS: 432.99 ($[M + Na]^+$). Anal. found: C, 81.95; H, 6.37. $C_{28}H_{26}O_3$ requires C, 81.92; H, 6.38%

Conclusions

In conclusion, we have developed a very simple and highly efficient solvent-free protocol for the synthesis of symmetrical bis(benzhydryl)ethers using inexpensive *p*-toluenesulfonyl chloride as reagent. The significant features of this environmentally benign and cost-effective straightforward protocol for direct conversion of benzhydrols into symmetrical bis(benzhydryl)ethers include operational simplicity, low reagent loading, high

Scheme 2 Proposed mechanistic pathway for etherification.

product yields, short reaction time, and solvent-free conditions.

Endnote

^aThe molecular structure of the product, bis(bis-phenylmethyl)ether (**1**), was determined by means of X-ray crystallographic studies. CCDC 840259 (**1**) contains the supplementary crystallographic data for this article. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/data_request/cif.

Additional file

Additional file 1: Title: Supplementary information Description: A document showing the general experimental details and procedures for the synthesis of symmetrical bis(benzhydryl)ethers. Copies of ¹H- and ¹³C-NMR spectra of all the entries (**1** to **7**) are also supplied.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

The authors are thankful to I.I.C.B., Kolkata and Chemistry Department, Kalyani University, India for the spectral measurements. B.B. is grateful to the UGC, New Delhi for awarding him a Senior Research Fellowship. G.B. is

thankful to the CSIR, New Delhi for financial support (No. 02(0110)/12/EMR-II dated 01.11.2012). The authors are grateful to Dr. Vivek K. Gupta, Post-Graduate Department of Physics, University of Jammu, Jammu Tawi 180 006, India for collecting the X-ray data.

This study was conducted in memory of Santosh Kr. Brahmachari.

Received: 30 November 2012 Accepted: 26 January 2013

Published: 18 February 2013

References

1. Brahmachari G (2010) Handbook of pharmaceutical natural products, 1st edn. Wiley-VCH, Weinheim
2. Li X, Upton TG, Gibb CLD, Gibb BC (2003) Resorcinarenes as templates: a general strategy for the synthesis of large macrocycles. *J Am Chem Soc* 125:650–651
3. Ley SV, Thomas AW (2003) Modern synthetic methods for copper-mediated C(aryl)-O, C(aryl)-N, and C(aryl)-S bond formation. *Angew Chem Int Ed* 42:5400–5449
4. Tlili A, Monnier F, Taillefer M (2010) Selective one-pot access to symmetrical or unsymmetrical diaryl ethers by copper-catalyzed double arylation of a simple oxygen source. *Chem Eur J* 16:12299–12302
5. Tan ES, Miyakawa M, Bunzow JR, Grandy DK, Scanlan TS (2007) Exploring the structure-activity relationship of the ethylamine portion of 3-iodothyronamine for rat and mouse trace amine-associated receptor 1. *J Med Chem* 50:2787–2798
6. Nicolaou KC, Boddy CNC, Brase S, Winssinger N (1999) Chemistry, biology and medicine of the glycopeptide antibiotics. *Angew Chem Int Ed* 38:2097–2152
7. Harris CM, Kopecka H, Harris TM (1983) Vancomycin: structure and transformation to CDP-I. *J Am Chem Soc* 105:6915–6922
8. Barna JCJ, Williams DH, Stone DJM, Leung TWC, Doddrell DM (1984) Structure elucidation of the teicoplanin antibiotics. *J Am Chem Soc* 106:4895–4902
9. Hayashi M (1997) Drug Data Report 704 (JP 97077745). J. R. Prous Science, Barcelona

10. Su D, Lim JJ, Tinney E, Wan B, Young MB, Anderson KD, Rudd D, Munshi V, Bahnck C, Felock PJ, Lu M, Lai M, Touch S, Moyer G, DiStefano DJ, Flynn JA, Liang Y, Sanchez R, Perlow-Poehnelt R, Miller M, Vacca JP, Williams TM, Anthony NJ (2009) Biaryl ethers as novel non-nucleoside reverse transcriptase inhibitors with improved potency against key mutant viruses. *J Med Chem* 52:7163–7169
11. Weis R, Schlapper C, Brun CR, Kaiser M, Seebacher W (2006) Antiplasmodial and antitrypanosomal activity of new esters and ethers of 4-dialkylaminobicyclo[2.2.2]octan-2-ols. *Eur J Pharm Sci* 28:361–368
12. Van Der Zee P, Hespe W (1978) A comparison of the inhibitory effects of aromatic substituted benzhydryl ethers on the uptake of catecholamines and serotonin into synaptosomal preparations of the rat brain. *Neuropharmacol* 17:483–490
13. Nilsson JL, Wägermark J, Dahlbom R (1969) Potential antiparkinsonism agents. Quinuclidinyl benzhydryl ethers. *J Med Chem* 12:1103–1105
14. McGavack TH, Schulman PM, Boyd LJ (1948) A clinical investigation of beta-morpholino-ethyl benzhydryl ether hydrochloride (linadryl) as an antihistamine agent. *J Allergy* 19:141–145
15. Loew ER, Kaiser ME (1945) Alleviation of anaphylactic shock in guinea pigs with synthetic benzhydryl alkamine ethers. *Exp Biol Med* 58:235–237
16. Pyo MK, Jin JL, Koo YK, Yun-Choi S (2004) Phenolic and furan type compounds isolated from *Gastrodia elata* and their anti-platelet effects. *Arch Pharm Res* 27:381–385
17. Pitsinos EN, Vidali VP, Couladouros EA (2011) Diaryl ether formation in the synthesis of natural products. *Eur J Org Chem* 7:1207–1222
18. Pratt EF, Draper JD (1949) Reaction rates by distillation. I. The etherification of phenylcarbinols and the transesterification of their ethers. *J Am Chem Soc* 71:2846–2849
19. Welch CM, Smith HA (1950) The properties of benzhydrol in sulfuric acid solution. *J Am Chem Soc* 72:4748–4750
20. Smith HA, Thompson RG (1955) Preparation and properties of substituted benzhydryl carbonium ions. *J Am Chem Soc* 77:1778–1783
21. Toda F, Takumi H, Akehi M (1990) Efficient solid-state reactions of alcohols: dehydration, rearrangement, and substitution. *J Chem Soc Chem Commun* 1270–1271
22. Brahmachari G, Laskar S (2010) A very simple and highly efficient procedure for *N*-formylation of primary and secondary amines at room temperature under solvent-free conditions. *Tetrahedron Lett* 51:2319–2322
23. Brahmachari G, Laskar S, Sarkar S (2010) Metal acetate/metal oxide in acetic acid: an efficient reagent for the chemoselective *N*-acetylation of amines. *J Chem Res* 34:288–295
24. Brahmachari G, Laskar S, Sarkar S (2010) A green approach to chemoselective *N*-acetylation of amines using catalytic amount of zinc acetate in acetic acid under microwave irradiation. *Indian J Chem* 49B:1274–1281
25. Brahmachari G, Das S (2012) Bismuth nitrate-catalyzed multicomponent reaction for efficient and one-pot synthesis of densely functionalized piperidine scaffolds at room temperature. *Tetrahedron Lett* 53:1479–1484
26. Brahmachari G, Banerjee B (2012) A comparison between catalyst-free and $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$ -catalyzed Strecker reactions for the rapid and solvent-free one-pot synthesis of racemic α -aminonitrile derivatives. *Asian J Org Chem* 1:251–258
27. Brahmachari G, Das S (2013) One-pot synthesis of 3-[(*N*-alkylanilino)(aryl)methyl] indoles via a transition metal assisted three-component condensation at room temperature. *J Het Chem*, in press
28. Brahmachari G, Das S (2013) A simple and straightforward method for one-pot synthesis of 2,4,5-triarylimidazoles using titanium dioxide as an eco-friendly and recyclable catalyst under solvent-free conditions. *Indian J Chem Sec B*, 52B:387–393
29. Sheldrick GM (1997) SHELXS97, Program for the solution of crystal structures. University of Göttingen, Göttingen
30. Farrugia LJ (1997) ORTEP-3 for windows—a version of ORTEP-III with a graphical user interface (GUI). *J Appl Cryst* 30:565–566
31. Farrugia LJ (1999) WinGX suite for small-molecule single-crystal crystallography. *J Appl Cryst* 32:837–838
32. Nardelli M (1995) PARST95—An update to PARST. A system of Fortran routines for calculating molecular structure parameters from the results of the crystal structure analysis. *J Appl Cryst* 28:659
33. Spek AL (2009) Structure validation in chemical crystallography. *Acta Cryst D65*:148–155
34. Koloves M, Froussios C (1984) *O*-diphenylmethylation of alcohols and carboxylic acids using diphenylmethyl diphenyl phosphate as alkylating agent. *Tetrahedron Lett* 25:3909–3912
35. Grummitt O, Buck AC (1945) Di-(*p*, *p*-dichlorobenzohydryl) ether. *J Am Chem Soc* 67:693
36. Welch CM, Smith HA (1953) Reactions of carboxylic acids in sulfuric acid. *J Am Chem Soc* 75:1412–1415

doi:10.1186/2191-2858-3-1

Cite this article as: Brahmachari and Banerjee: Facile synthesis of symmetrical bis(benzhydryl)ethers using *p*-toluenesulfonyl chloride under solvent-free conditions. *Organic and Medicinal Chemistry Letters* 2013 3:1.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com