Published in final edited form as:

Org Lett. 2013 March 1; 15(5): 992–995. doi:10.1021/ol303431h.

Highly Efficient Cu(I)-Catalyzed Oxidation of Alcohols to Ketones and Aldehydes with Diaziridinone

Yingguang Zhu, Baoguo Zhao, and Yian Shi

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523, United States

Yian Shi: yian@lamar.colostate.edu

Abstract

A novel and efficient Cu(I)-catalyzed oxidation of alcohols has been achieved with di-*tert*-butyldiaziridinone as oxidant under mild conditions. A wide variety of primary and secondary alcohols with various functional groups can be oxidized to aldehydes and ketones in high yields. The reaction proceeds under neutral conditions making it compatible with acid or base-sensitive substrates, and it is amenable to gram scale.

The oxidation of alcohols to the corresponding ketones or aldehydes is one of the most fundamental and widely utilized transformations in organic synthesis.¹ Various effective reagents and methods have been developed, including chromium reagents,² manganese(IV) oxide,³ activated DMSO,⁴ hypervalent iodine reagents,⁵ ruthenium reagents,⁶ osmium(VIII) oxide,⁷ and metal-^{8–16} or TEMPO-catalyzed^{17,18} oxidation, etc. Due to the importance of this transformation in organic synthesis, the development of new, efficient, safe, and mild oxidation processes is still highly desirable and valuable.

We have previously reported that readily prepared di-*tert*-butyldiaziridinone (**1**)^{19,20} is a highly effective nitrogen source for the Pd(0)-²¹ and Cu(I)-²² catalyzed diamination of olefins. In our efforts to further explore the reactivity of diaziridinone and expand its synthetic utility, we have found that alcohols can be efficiently oxidized to the corresponding carbonyl compounds with di-*tert*-butyldiaziridinone (**1**) in the presence of Cu(I) catalyst under mild reaction conditions. Herein we wish to report our preliminary studies on this subject.

Our initial studies were carried out with 2,2-dimethyl-1-phenyl-1-propanol (**2a**) as test substrate. Several copper catalysts were first screened in CDCl₃ at room temperature (Table 1, entries 1–5). High conversions were obtained with CuCl and CuBr (Table 1, entries 3 and 4). In the case of CuBr, ketone **3a** was isolated in 87% yield. The yield decreased to 53% when P(*t*Bu)₃ was added (Table 1, entry 6). Among the solvents examined (Table 1, entries 7–11), CH₃CN gave the highest yield (94%) for ketone **3a** (Table 1, entry 9). No reaction occurred in the absence of the Cu catalyst (Table 1, entry 12) or di-*tert*-butyldiaziridinone

Correspondence to: Yian Shi, yian@lamar.colostate.edu.

Supporting Information Available. Experimental procedures, characterization data, and NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(**1**) under air (Table 1, entry 13), suggesting that both Cu catalyst and di-*tert*-butyldiaziridinone (**1**) are required for the reaction.

With the optimal reaction conditions in hand, the generality for the oxidation was subsequently investigated with various secondary alcohols. As shown in Table 2, high yields were obtained for a wide range of alcohols. Various acyclic and cyclic secondary benzylic alcohols with alkyl, cyclopropyl, vinyl, allyl, alkynyl, and thioether groups were effectively oxidized to ketones in 73–99% yields (Table 2, entries 1–13). As illustrated in the oxidation of benzhydrol (**2c**) (Table 2, entry 3), the method is amenable to gram scale. The oxidation can also be extended to a variety of non-benzylic secondary alcohols, giving the corresponding ketones in 74–99% yields (Table 2, entries 14–28). Acyclic aliphatic alcohol **2o** was oxidized to ketone **3o** in 99% yield (Table 2, entry 15). Cyclic alcohols with variable ring sizes and substituents were found to be effective substrates (Table 2, entries 16–28). Acid-sensitive groups such as silyl ethers and ketal can be tolerated (Table 2, entries 20 and 27). The mildness of the reaction conditions was further demonstrated in the oxidation of (1*R*,2*S*)-*trans*-2-phenyl-1-cyclohexanol (**2bb**) (Table 2, entry 28). Ketone **3bb** was obtained in 98% yield, and no racemization of the α -stereocenter occurred during the oxidation.

Primary alcohols were also found to be effective substrates, but generally less reactive than secondary alcohols. To shorten the reaction time, the reaction temperature was raised to 60 °C. Under these reaction conditions, various benzylic alcohols with different substituents on the phenyl ring can be efficiently oxidized to the corresponding aldehydes in 63–96% yields (Table 3, entries 1–10). Heterocyclic primary alcohols were smoothly oxidized to give the aldehydes in good yields (Table 3, entries 11 and 12). In addition, allylic and aliphatic alcohols were also effective substrates, giving the corresponding aldehydes in 70–90% yields (Table 3, entries 13–16).

While a precise understanding of the reaction mechanism awaits further study, a plausible catalytic pathway is proposed in Scheme 1. The CuBr first reductively cleaves the N–N bond of di-*tert*-butyldiaziridinone (**1**) to form four-membered Cu(III) species **A** and/or Cu(II) nitrogen radical **B**,^{22e,f} which reacts with alcohol **2** to generate alkoxy Cu(III) species **C**. The β -H elimination of intermediate **C** affords ketone **3** and Cu(III) species **D**, which undergoes reductive elimination to form urea and regenerate the CuBr catalyst.

In summary, we developed a novel oxidation of alcohols with CuBr as catalyst and di-*tert*-butyldiaziridinone (**1**) as oxidant. Various primary and secondary alcohols can be efficiently oxidized under mild conditions to the corresponding aldehydes and ketones in high yields. The present process exhibits the following favorable features: (1) the procedure is effective for a wide range of substrates bearing various functional groups, such as, alkenyl, alkynyl, thioether, silyl ether, amide, carbamate, ketal, ester, and heterocycles; (2) the reaction proceeds under neutral and mild conditions (no acid or base required) making it compatible with acid or base-sensitive substrates; (3) the method is amenable to gram scale with no special precautions to exclude air or moisture. Further development of other reaction processes with diaziridinones and related compounds is currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful for the generous financial support from the General Medical Sciences of the National Institutes of Health (GM083944-05).

REFERENCES

1. For leading books, see: Hudlický M. *Oxidations in Organic Chemistry*. 1990 Washington, DC American Chemical Society; Larock RC. *Comprehensive Organic Transformations* (2nd ed.). 1999 New York Wiley-VCH:1234. Arends IWCE, Sheldon RA, Bäckvall J-E. In *modern oxidation methods* (2nd ed.). 2010 Weinheim Wiley-VCH:147.
2. For leading references, see: Bowden K, Heilbron IM, Jones ERH, Weedon BCL. *J. Chem. Soc.* 1946;39. Poos GI, Arth GE, Beyler RE, Sarett LH. *J. Am. Chem. Soc.* 1953; 75:422. Coates WM, Corrigan JR. *Chem. Ind. (London)*. 1969:1594. Corey EJ, Suggs JW. *Tetrahedron Lett.* 1975; 16:2647. Piancatelli G, Scettri A, D'Auria M. *Synthesis*. 1982:245. Luzzio FA, Guziec FS Jr. *Org. Prep. Proced. Int.* 1988; 20:533.
3. For leading references, see: Harfenist M, Bavley A, Lazier WA. *J. Org. Chem.* 1954; 19:1608. Brink M. *Synthesis*. 1975:253. Fatiadi AJ. *Synthesis*. 1976:65. Fatiadi AJ. *Synthesis*. 1976:133. Taylor RJK, Reid M, Foot J, Raw SA. *Acc. Chem. Res.* 2005; 38:851. [PubMed: 16285708]
4. For leading references, see: Pfitzner KE, Moffatt JG. *J. Am. Chem. Soc.* 1963; 85:3027. Mancuso AJ, Huang S-L, Swern D. *J. Org. Chem.* 1978; 43:2480. Mancuso AJ, Swern D. *Synthesis*. 1981:165. Tidwell TT. *Synthesis*. 1990:857.
5. For leading references, see: Dess DB, Martin JC. *J. Org. Chem.* 1983; 48:4155. Dess DB, Martin JC. *J. Am. Chem. Soc.* 1991; 113:7277. Frigerio M, Santagostino M. *Tetrahedron Lett.* 1994; 35:8019. Meyer SD, Schreiber SL. *J. Org. Chem.* 1994; 59:7549. Corey EJ, Palani A. *Tetrahedron Lett.* 1995; 36:3485. Nicolaou KC, Zhong Y-L, Baran PS. *J. Am. Chem. Soc.* 2000; 122:7596. Uyanik M, Ishihara K. *Chem. Commun.* 2009:2086.
6. For leading references, see: Berkowitz LM, Rylander PN. *J. Am. Chem. Soc.* 1958; 80:6682. Tomioka H, Takai K, Oshima K, Nozaki H. *Tetrahedron Lett.* 1981; 22:1605.
7. For a leading reference, see: Maione AM, Romeo A. *Synthesis*. 1984:955.
8. For leading references on copper-catalyzed oxidation of alcohols, see: Semmelhack MF, Schmid CR, Cortés DA, Chou CS. *J. Am. Chem. Soc.* 1984; 106:3374. Markó IE, Giles PR, Tsukazaki M, Brown SM, Urch CJ. *Science*. 1996; 274:2044. [PubMed: 8953027] Ansari IA, Gree R. *Org. Lett.* 2002; 4:1507. [PubMed: 11975615] Gamez P, Arends IWCE, Reedijk J, Sheldon RA. *Chem. Commun.* 2003:2414. Markó IE, Gautier A, Dumeunier R, Doda K, Philippart F, Brown SM, Urch CJ. *Angew. Chem. Int. Ed.* 2004; 43:1588. Jiang N, Ragauskas AJ. *J. Org. Chem.* 2006; 71:7087. [PubMed: 16930071] Onomura O, Arimoto H, Matsumura Y, Demizu Y. *Tetrahedron Lett.* 2007; 48:8668. Liu C, Han J, Wang J. *Synlett.* 2007:643. Figiel PJ, Kopylovich MN, Lasri J, Guedes da Silva MFC, Fraústo da Silva JJR, Pombeiro AJL. *Chem. Commun.* 2010:2766. Hoover JM, Stahl SS. *J. Am. Chem. Soc.* 2011; 133:16901. [PubMed: 21861488] Han C, Yu M, Sun W, Yao X. *Synlett.* 2011:2363. Babu SG, Priyadarsini PA, Karvembu R. *Appl. Catal. A: Gen.* 2011; 392:218.
9. For leading references on palladium-catalyzed oxidation of alcohols, see: Peterson KP, Larock RC. *J. Org. Chem.* 1998; 63:3185. ten Brink G-J, Arends IWCE, Sheldon RA. *Science*. 2000; 287:1636. [PubMed: 10698735] Jensen DR, Pugsley JS, Sigman MS. *J. Am. Chem. Soc.* 2001; 123:7475. [PubMed: 11472200] Ferreira EM, Stoltz BM. *J. Am. Chem. Soc.* 2001; 123:7725. [PubMed: 11481006] Stahl SS, Thorman JL, Nelson RC, Kozee MA. *J. Am. Chem. Soc.* 2001; 123:7188. [PubMed: 11459511] Liu J, Wang F, Sun K, Xu X. *Catal. Commun.* 2008; 9:386. Dileep R, Bhat BR. *Appl. Organometal. Chem.* 2010; 24:663.
10. For leading references on ruthenium-catalyzed oxidation of alcohols, see: Griffith WP, Ley SV, Whitcombe GP, White AD. *J. Chem. Soc. Chem. Commun.* 1987:1625. Morris PE, Kiely DE. *J. Org. Chem.* 1987; 52:1149. Ley SV, Norman J, Griffith WP, Marsden SP. *Synthesis*. 1994:639. Fung W-H, Yu W-Y, Che C-M. *J. Org. Chem.* 1998; 63:2873. Csajnyik G, Éll AH, Fadini L, Pugin B, Bäckvall J-E. *J. Org. Chem.* 2002; 67:1657. [PubMed: 11871899] Yamaguchi K, Mizuno N. *Angew. Chem. Int. Ed.* 2002; 41:4538. Gonsalvi L, Arends IWCE, Sheldon RA. *Org. Lett.* 2002; 4:1659. [PubMed: 12000267] Zhan B-Z, White MA, Sham T-K, Pincock JA, Doucet RJ, Rao KVR, Robertson KN, Cameron TS. *J. Am. Chem. Soc.* 2003; 125:2195. [PubMed: 12590547]
11. For leading references on gold-catalyzed oxidation of alcohols, see: Tsunoyama H, Sakurai H, Negishi Y, Tsukuda T. *J. Am. Chem. Soc.* 2005; 127:9374. [PubMed: 15984857] Guan B, Xing D, Cai G, Wan X, Yu N, Fang Z, Yang L, Shi Z. *J. Am. Chem. Soc.* 2005; 127:18004. [PubMed: 16366542] Ni J, Yu W-J, He L, Sun H, Cao Y, He H-Y, Fan K-N. *Green Chem.* 2009; 11:756.

12. For leading references on tungsten-catalyzed oxidation of alcohols, see: Bortolini O, Conte V, Furia FD, Modena G. *J. Org. Chem.* 1986; 51:2661. Sato K, Aoki M, Takagi J, Noyori R. *J. Am. Chem. Soc.* 1997; 119:12386. Sloboda-Rozner D, Alsters PL, Neumann R. *J. Am. Chem. Soc.* 2003; 125:5280. [PubMed: 12720432]
13. For leading references on manganese-catalyzed oxidation of alcohols, see: Son Y-C, Makwana VD, Howell AR, Suib SL. *Angew. Chem. Int. Ed.* 2001; 40:4280. Bagherzadeh M. *Tetrahedron Lett.* 2003; 44:8943. Mardani HR, Golchoubian H. *Tetrahedron Lett.* 2006; 47:2349. Kwong H-K, Lo P-K, Lau K-C, Lau T-C. *Chem. Commun.* 2011:4273.
14. For leading references on iron-catalyzed oxidation of alcohols, see: Pearson AJ, Kwak Y. *Tetrahedron Lett.* 2005; 46:5417. Shi F, Tse MK, Pohl M-M, Brückner A, Zhang S, Beller M. *Angew. Chem. Int. Ed.* 2007; 46:8866. Schröder K, Junge K, Bitterlich B, Beller M. *Top. Organomet. Chem.* 2011; 33:83. Kunisu T, Oguma T, Katsuki T. *J. Am. Chem. Soc.* 2011; 133:12937. [PubMed: 21780821]
15. For leading references on cobalt-catalyzed oxidation of alcohols, see: Tovrog BS, Diamond SE, Mares F, Szalkiewicz A. *J. Am. Chem. Soc.* 1981; 103:3522. Iwahama T, Yoshino Y, Keitoku T, Sakaguchi S, Ishii Y. *J. Org. Chem.* 2000; 65:6502. [PubMed: 11052094] Das S, Punniyamurthy T. *Tetrahedron Lett.* 2003; 44:6033. Gunasekaran N, Jerome P, Ng SW, Tiekink ERT, Karvembu R. *J. Mol. Catal. A: Chem.* 2012; 353–354:156.
16. For leading references on vanadium-catalyzed oxidation of alcohols, see: Li C, Zheng P, Li J, Zhang H, Cui Y, Shao Q, Ji X, Zhang J, Zhao P, Xu Y. *Angew. Chem. Int. Ed.* 2003; 42:5063. Pawar VD, Bettigeri S, Weng S-S, Kao J-Q, Chen C-T. *J. Am. Chem. Soc.* 2006; 128:6308. [PubMed: 16683782] Bagherzadeh M, Amini M. *J. Coord. Chem.* 2010; 63:3849. Alagiri K, Prabhu KR. *Tetrahedron.* 2011; 67:8544.
17. For leading reviews on TEMPO-catalyzed oxidation of alcohols, see: Adam W, Saha-Möller CR, Ganeshpure PA. *Chem. Rev.* 2001; 101:3499. [PubMed: 11840992] Sheldon RA, Arends IWCE. *Adv. Synth. Catal.* 2004; 346:1051.
18. For leading references on TEMPO-catalyzed oxidation of alcohols, see: Anelli PL, Biffi C, Montanari F, Quici S. *J. Org. Chem.* 1987; 52:2559. Einhorn J, Einhorn C, Ratajczak F, Pierre J-L. *J. Org. Chem.* 1996; 61:7452. [PubMed: 11667673] Rychnovsky SD, Vaidyanathan R. *J. Org. Chem.* 1999; 64:310. [PubMed: 11674122] Bolm C, Magnus AS, Hildebrand JP. *Org. Lett.* 2000; 2:1173. [PubMed: 10804582] Liu R, Liang X, Dong C, Hu X. *J. Am. Chem. Soc.* 2004; 126:4112. [PubMed: 15053593] Shibuya M, Osada Y, Sasano Y, Tomizawa M, Iwabuchi Y. *J. Am. Chem. Soc.* 2011; 133:6497. [PubMed: 21473575]
19. For a leading review on diaziridinones, see: Heine HW, Hassner A. *The Chemistry of Heterocyclic Compounds*. 1983 New York John Wiley & Sons, Inc:547.
20. For the preparation of di-*tert*-butyldiaziridinone (**1**), see: Greene FD, Stowell JC, Bergmark WR. *J. Org. Chem.* 1969; 34:2254. Du H, Zhao B, Shi Y. *Org. Synth.* 2009; 86:315.
21. For Pd(0)-catalyzed diamination of olefins, see: Du H, Zhao B, Shi Y. *J. Am. Chem. Soc.* 2007; 129:762. [PubMed: 17243803] Du H, Yuan W, Zhao B, Shi Y. *J. Am. Chem. Soc.* 2007; 129:11688. [PubMed: 17803307] Du H, Yuan W, Zhao B, Shi Y. *J. Am. Chem. Soc.* 2007; 129:7496. [PubMed: 17521192] Du H, Zhao B, Shi Y. *J. Am. Chem. Soc.* 2008; 130:8590. [PubMed: 18549207] Zhao B, Du H, Cui S, Shi Y. *J. Am. Chem. Soc.* 2010; 132:3523. [PubMed: 20166669]
22. For Cu(I)-catalyzed diamination of olefins, see: Yuan W, Du H, Zhao B, Shi Y. *Org. Lett.* 2007; 9:2589. [PubMed: 17536815] Du H, Zhao B, Yuan W, Shi Y. *Org. Lett.* 2008; 10:4231. [PubMed: 18763785] Wen Y, Zhao B, Shi Y. *Org. Lett.* 2009; 11:2365. [PubMed: 19408926] Zhao B, Du H, Shi Y. *J. Org. Chem.* 2009; 74:8392. [PubMed: 19827794] Zhao B, Peng X, Cui S, Shi Y. *J. Am. Chem. Soc.* 2010; 132:11009. [PubMed: 20698659] Zhao B, Peng X, Zhu Y, Ramirez TA, Cornwall RG, Shi Y. *J. Am. Chem. Soc.* 2011; 133:20890. [PubMed: 22081888]

Scheme 1.
Proposed Catalytic Cycle for the Oxidation

Table 1

Studies on Reaction Conditions^a

entry	catalyst	solvent	time (h)	conv (%) ^b	yield (%) ^c
1	CuCN	CDCl ₃	4	0	–
2	CuI	CDCl ₃	4	47	38
3	CuCl	CDCl ₃	4	93	78
4	CuBr	CDCl ₃	4	97	87
5	CuBr ₂	CDCl ₃	4	57	49
6	CuBr·P(<i>t</i> Bu) ₃ (1:1)	CDCl ₃	4	65	53
7	CuBr	CH ₂ Cl ₂	4	99	82
8	CuBr	DCE	4	99	80
9	CuBr	CH₃CN	2	100	94
10	CuBr	DMF	4	42	29
11	CuBr	THF	4	89	79
12 ^d	none	CH ₃ CN	24	0	–
13 ^e	CuBr	CH ₃ CN	24	0	–

^a All reactions were carried out with alcohol **2a** (0.30 mmol), di-*tert*-butyldiaziridinone (**1**) (0.33 mmol), and Cu catalyst (0.030 mmol) in solvent (0.6 mL) at rt under Ar unless otherwise stated.

^b The conversion was based on alcohol **2a** and determined by the ¹H NMR.

^c Isolated yield.

^d The reaction was carried out at rt or 60 °C under Ar.

^e The reaction was carried out at 60 °C under air in the absence of **1**.

Table 2

Oxidation of Secondary Alcohols^a

$ \begin{array}{c} \text{O} \\ \parallel \\ \text{}^t\text{BuN}-\text{N}^t\text{Bu} \\ \mathbf{1} \\ \text{CuBr (10 mol \%), CH}_3\text{CN, rt} \\ \xrightarrow{\hspace{1cm}} \\ \begin{array}{ccc} \text{R}^1-\text{CH}(\text{OH})-\text{R}^2 & & \text{R}^1-\text{C}(=\text{O})-\text{R}^2 \\ \mathbf{2} & & \mathbf{3} \end{array} \end{array} $				
entry	substrate 2	product 3	time (h)	yield (%) ^b
1	2a , R = ^t Bu	3a	2	95
2	2b , R = Me	3b	10	97
3 ^c	2c , R = Ph	3c	12	99
4	2d , R = cyclopropyl	3d	3	99
5	2e , R = 2-phenylvinyl	3e	0.2	99
6	2f , R = vinyl	3f	5	87
7	2g , R = allyl	3g	1.5	74
8 ^d			4.5	99
	2h	3h		
9	2i , n = 1	3i	6	97
10	2j , n = 2	3j	3	98
11	2k , n = 3	3k	0.5	91
12	2l , X = O	3l	12	98
13 ^e	2m , X = S	3m	5	73
14 ^d			4.5	86
	2n	3n		

$ \begin{array}{c} \text{O} \\ \parallel \\ \text{'BuN-N'Bu} \\ \mathbf{1} \\ \xrightarrow{\text{CuBr (10 mol \%), CH}_3\text{CN, rt}} \\ \text{R}^1\text{-CH(OH)-R}^2 \quad \mathbf{2} \quad \longrightarrow \quad \text{R}^1\text{-C(=O)-R}^2 \quad \mathbf{3} \end{array} $				
entry	substrate 2	product 3	time (h)	yield (%) ^b
15	$n\text{C}_{10}\text{H}_{21}$ 2o	$n\text{C}_{10}\text{H}_{21}$ 3o	12	99
16 ^f	2p , $n = 1$	3p	12	80
17	2q , $n = 3$	3q	12	85
18 ^g	2r , $n = 7$	3r	6	99
	2s , R = Ph	3s	10	99
20	2t , R = OSiEt ₃	3t	12	93
21 ^h	2u	3u	12	90
22 ^d	2v	3v	5	91
23 ⁱ	2w	3w	8	74
24 ^h	2x	3x	8	83
25	2y	3y	3	90

$ \begin{array}{c} \text{O} \\ \parallel \\ \text{'BuN}-\text{N'Bu} \\ \mathbf{1} \\ \xrightarrow{\text{CuBr (10 mol \%), CH}_3\text{CN, rt}} \\ \text{R}^1-\text{CH}(\text{OH})-\text{R}^2 \quad \mathbf{2} \quad \longrightarrow \quad \text{R}^1-\text{C}(=\text{O})-\text{R}^2 \quad \mathbf{3} \end{array} $				
entry	substrate 2	product 3	time (h)	yield (%) ^b
26	2z	3z	8	92
27 ^j	2aa	3aa	4	87
28	2bb (>99% ee)	3bb (>99% ee)	11	98

^a All reactions were carried out with alcohol **2** (0.60 mmol), di-*tert*-butyldiaziridinone (**1**) (0.66 mmol), and CuBr (0.060 mmol) in CH₃CN (1.2 mL) at rt under Ar unless otherwise stated.

^b Isolated yield.

^c The reaction was carried out on 10 mmol scale of **2c** under open air.

^d The reaction was carried out at 60 °C.

^e The reaction was carried out with 0.90 mmol of **1** at 60 °C.

^f The reaction was carried out on 0.90 mmol scale of **2p**.

^g The reaction was carried out in CH₃CN (2.0 mL) at 60 °C.

^h The reaction was carried out in CDCl₃ (1.5 mL) at 60 °C.

ⁱ The reaction was carried out with 1.14 mmol of **1** in CDCl₃ (1.5 mL) at 60 °C.

^j The reaction was carried out in CH₃CN (2.4 mL) at 60 °C.

Table 3

Oxidation of Primary Alcohols^a

$ \begin{array}{c} \text{O} \\ \parallel \\ \text{}^t\text{BuN}-\text{N}^t\text{Bu} \\ \mathbf{1} \\ \xrightarrow{\text{CuBr (10 mol \%), CH}_3\text{CN, 60 }^\circ\text{C}} \\ \text{R}-\text{CH}_2\text{OH} \quad \mathbf{4} \quad \quad \quad \text{R}-\text{CHO} \quad \mathbf{5} \end{array} $				
entry	substrate 4	product 5	time (h)	yield (%) ^b
1	4a , X = H	5a	5	83
2	4b , X = <i>o</i> -OMe	5b	3	90
3	4c , X = <i>m</i> -OMe	5c	3	89
4	4d , X = <i>p</i> -OMe	5d	3	90
5	4e , X = <i>p</i> -SMe	5e	5	63
6	4f , X = <i>p</i> - ^{<i>i</i>} Pr	5f	3	91
7	4g , X = <i>p</i> -Br	5g	3	82
8	4h , X = <i>p</i> -CO ₂ Me	5h	3	94
9	 4i	 5i	2	92
10	 4j	 5j	5	96
11	 4k	 5k	4	90
12	 4l	 5l	4	80
13 ^c	 4m	 5m	12	87
14 ^d	 4n	 5n	4	70
15	 4o	 5o	10	83

entry	substrate 4	product 5	time (h)	yield (%) ^b
16 ^d	<i>n</i> -C ₁₅ H ₃₁ -OH 4p	<i>n</i> -C ₁₅ H ₃₁ -CHO 5p	4	90

^a All reactions were carried out with alcohol **4** (0.60 mmol), di-*tert*-butyldiaziridinone (**1**) (0.66 mmol), and CuBr (0.060 mmol) in CH₃CN (1.2 mL) at 60 °C under Ar unless otherwise stated.

^b Isolated yield.

^c The reaction was carried out with 0.72 mmol of **1**.

^d The reaction was carried out in CDCl₃ (1.2 mL).