Published in final edited form as:

Biochem Biophys Res Commun. 2013 April 19; 433(4): 508–512. doi:10.1016/j.bbrc.2013.03.016.

The Regulation of Apoptosis by the Downstream Regulatory Element Antagonistic Modulator/Potassium Channel Interacting Protein 3 (DREAM/KChIP3) Through Interactions with Hexokinase I

Theodore A. Craig^{1,2}, Pradeep L. Ramachandran^{1,2}, H. Robert Bergen III³, Jewel L. Podratz⁴, Anthony J. Windebank⁴, and Rajiv Kumar^{1,2,3}

¹Division of Nephrology and Hypertension, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA

²Department of Medicine, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA

³Department of Biochemistry and Molecular Biology, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA

⁴Department of Neurology, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA

Abstract

The EF-hand protein, DREAM/KChIP3 (henceforth referred to as DREAM), regulates apoptosis by incompletely understood mechanisms. We demonstrate that in the presence of Ca^{2+} , DREAM interacts with hexokinase I, a protein known to bind mitochondria and regulate apoptosis. A mutant DREAM protein construct incapable of binding Ca^{2+} does not associate with hexokinase I. The amino-terminal portion of DREAM is required for binding to hexokinase I, as a DREAM construct lacking the first 94 amino terminal residues fails to bind hexokinase I. Expression of DREAM in neuroblastoma cells enhances cisplatin mediated caspase-3 activity. Simultaneous expression of hexokinase I in such cells reduces DREAM-stimulated apoptosis. DREAM overexpression in neuroblastoma cells reduces hexokinase I localization on isolated mitochondria. The interaction of DREAM with hexokinase I may be important in the regulation of neuronal apoptosis.

Keywords

Apoptosis; DREAM; calsenilin; KChIP3; hexokinase I; neuronal calcium sensor proteins; EF-hand proteins; calcium

© 2013 Elsevier Inc. All rights reserved.

Correspondence: Rajiv Kumar, MS 1-120, Mayo Clinic, 200 First street SW, Rochester, MN 55905, United States. Fax: +1 507 538 3954. rkumar@mayo.edu.

TAC and PLR contribute equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

INTRODUCTION

Cell number and tissue viability are regulated by apoptosis [1,2,3,4,5]. Altered apoptosis is noted in several diseases including cancer and neurodegenerative diseases such as Alzheimer's, Huntington's and Parkinson's diseases [1,6,7]. Understanding mechanisms by which apoptosis is regulated is important in the prevention and treatment of various neoplastic and neurodegenerative disorders.

The neuronal EF-hand protein, downstream regulatory element antagonistic modulator (DREAM/Calsenilin/KChIP3) regulates diverse metabolic processes [8,9,10,11]. DREAM represses endogenous opioid production by binding to regulatory elements in the *prodynorphin* (*pdyn*) and *c-fos* genes [8] and binds to Ca^{2+} -activated Kv4 potassium channels in the brain and heart [12]. *Dream*^{-/-} mice display reduced responses to pain, and improved contextual fear conditioning [13,14]. DREAM regulates apoptosis [15,16] by altering intracellular Ca^{2+} signaling [16], changing presenilin 1 and 2 activity [9,15,17] and by binding to peroxiredoxin 3, an antioxidant enzyme [18]. DREAM binds to the carboxyl-terminal portion of presenilin to regulate γ -secretase function [9]. DREAM regulates the levels of a proteolytic products of presenilin 2 [9] and reverses the presenilin-mediated enhancement of calcium signaling [19]. Indeed, *dream*^{-/-} mice have approximately 50% less A β 40 and A β 42 in the cerebellum [20]. DREAM concentrations are increased in the cortex of brains of patients with Alzheimer's disease and in the neocortex and the hippocampus of brains of mice overexpressing a Swedish mutant β -amyloid precursor protein [15]. DREAM is also a substrate for caspase-3 [10].

To gain insights into how DREAM alters apoptosis in neuronal cells, we investigated whether DREAM interacts with other previously identified regulators of apoptosis and regulates their activity. We report that DREAM interacts with brain hexokinase I in a Ca^{2+} -dependent manner, and demonstrate the functional relevance of the interaction.

MATERIALS AND METHODS

Expression and purification of DREAM protein constructs

Full length (FL) DREAM (amino acids 1–256), sFL-DREAM (amino acids 95–256), and calcium insensitive Mut-1234-DREAM (E103A, D110A, D139A, D141A, D175A, N177A, D223A, N225A), were expressed as N-terminal glutathione S-transferase (GST) fusion products in *Escherichia coli* BL21 cells and purified as previously described [21,22].

Synthesis of DREAM expression vectors

DREAM pcDNA3.1 sense and anti-sense constructs were prepared by inserting the *Eco*RI cleaved DREAM fragments obtained from the chimeric DREAM pCR2.1 plasmid into the *Eco*RI site of pcDNA3.1(–) (Life Technologies) as described earlier [23].

Expression and purification of full length hexokinase I protein

Full length human hexokinase I with a carboxyl-terminal 10× His tag (GenBank accession NM_000188.2) was expressed in the pET28a(+) vector and *Escherichia coli* Rosetta 2 (DE3) host (EMD-Novagen, Billerica, MA). The cDNA (Origene) was amplified using the following oligonucleotides (5' → 3') for PCR: 5PCI1hHK1 : 5'GAGAACATGTCCATGATCGCCGCGCAGCTCCTGGCCTATTACTTCAC3' ; 3HK110xHisXh: 5'GAGACTCGAGTTAGTGATGGTGATGATGGTGATGGTGATGGCTGCTTGCC TCTGTGCGTAACCGCACGCCACGGCCGTG3'. The PCR amplified products were cleaved by *Scl*I and *Xho*I and ligated into *Nco*I/*Xho*I cleaved, dephosphorylated pET28a(+).

Transformed *E. coli* Rosetta 2 (DE3) were grown in 2xYT medium with 40 µg/mL kanamycin, and 20 µg/mL chloramphenicol, initially at 37 °C to an OD 600nm of ~1. The temperature was reduced (16 °C), and protein expression was induced for 23 hours with 0.2 mM *iso*-propylthiogalactoside. Cells were lysed and protein was purified on Ni-NTA Superflow resin (Qiagen, Valencia, CA).

Synthesis of a pCMV6-AC-hexokinase I-mRFP plasmid

The hexokinase I gene was cloned in to the pCMV6-AC-mRFP vector to create a vector which expressed hexokinase I with a c-terminal monomer RFP (mRFP) fusion product. The hexokinase I cDNA (OriGene, Rockville, MD) was amplified by PCR using the following primers: 5' → 3': 5_3gfpHK*Asc*I: GAGAGGCGCGCCATG ATCGCCGCGCAGCTCCTGGCCTAT and 3_5gfpHK*Xho*I: TCTCCTCGAGGCTGC TTGCCTCTGTGCGTAACCGC. The product was treated with restriction enzymes *Asc*I and *Xho*I (New England Biolabs, Ipswich, MA) and ligated into the pCMV6-AC-mRFP vector treated with similar restriction enzymes.

DREAM affinity capture assay

This was performed as described earlier [22]. Homogenates of rat brain (10% weight/volume) were prepared in 10 mL buffer containing 50 mM Tris-HCl (pH 7.5), 50 mM NaCl, 2 mM dithiothreitol (DTT), mini cOmplete, EDTA-free, protease inhibitor cocktail (Roche Diagnostics, Indianapolis, IN) and either 2 mM EDTA (buffer A), or 5 mM CaCl₂ (buffer B), respectively. Homogenates in Ca²⁺ or EDTA buffer were applied to columns prepared by incubating GST-DREAM or GST with glutathione sepharose resin (Miltényi Biotec, Auburn, CA). The resins were extensively washed with Ca²⁺ or EGTA buffer containing 2 M NaCl. Residual proteins bound to the column were eluted with Ca²⁺ or EGTA buffer containing 50 mM reduced glutathione. Fractions were run on an SDS-PAGE gels and were stained with 0.02% PhastGel Blue R. Mass spectrometric identification of proteins bound to GST-DREAM or GST was carried out as described in Ramachandran *et al.* [22]. Scaffold (version Scaffold_2_00_06, Proteome Software Inc., Portland, OR) was used to validate MS/MS based peptide and protein identification.

Assessment of DREAM hexokinase I interactions

A protein capture assay was conducted to test the binding of rat brain hexokinase I to full length, short full length, and a calcium insensitive mutant GST-tagged DREAM constructs in the presence of either EDTA or Ca²⁺ as described above [22]. GST bound to glutathione resin and resin alone, were used as controls. Resins (125 µL of each, equilibrated with 2.5 mL of either Buffer A (50 mM Tris-HCl pH 7.5, 50 mM NaCl, 2 mM DTT, 5 mM CaCl₂) or Buffer B (50 mM Tris-HCl pH 7.5, 50 mM NaCl, 2 mM DTT, 2 mM EDTA) were prepared. Rat brain homogenate was added to appropriate resins in the presence of EDTA or Ca²⁺ buffers. Resins were washed with either Buffer A or Buffer B, and bound proteins were eluted with either Buffers A or B containing 50 mM glutathione. Eluates were electrophoresed on a SDS gel, transferred to a PVDF membrane and analyzed by western analysis with a rabbit anti-HK I antibody (Cell Signaling Technology, Danvers, MA) and an appropriate secondary antibody (peroxidase labeled goat anti-rabbit antibody, DAKO).

Determination of hexokinase I activity

Hexokinase I activity [24] was measured at 22 °C in 96-well plates by assaying the change in absorbance of NADPH in the presence or absence of added buffer or DREAM (6–12:1 molar ratio, DREAM:hexokinase I) with Ca²⁺ or EGTA present. To 80 µl assay buffer (90 mM tricine-KOH, pH 8.1, 5 mM glucose, 0.25 mM NADP⁺ (Sigma-Aldrich, St. Louis, MO), 10 mM MgCl₂, 5 mM ATP, disodium salt, and glucose-6-phosphate dehydrogenase

were added Ca^{2+} or EDTA (1 mM and 1.25 mM final concentrations), and 20 μL recombinant hexokinase I (20, 40 or 80 ng diluted in 0.1 M tricine pH 8.1). The absorbance of the reaction mixture was measured for 5 minutes at 340 nm with a SpectraMax M2 microplate reader (Molecular Devices).

Biolayer interferometry (BLI) measurements to assess the binding of DREAM to hexokinase I

Biotinylated DREAM was synthesized by adding a 10-fold molar excess of sulfosuccinimidobiotin (Thermo Scientific) to 40 μM DREAM in 50 mM Na^+/K^+ phosphate buffer, pH 7.5, 150 mM NaCl for 30 minutes at room temperature. Biotinylated DREAM was dialyzed against 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, and 1 mM DTT buffer. Streptavidin biosensors (Pall Forte Bio, Menlo Park, CA), were hydrated in H_2O for 10 minutes, and then in either Ca^{2+} or EGTA containing buffer for 30 seconds. The biosensor was submerged in a 4 μL drop containing 2 μL , 30 μM biotinylated DREAM and 2 μL of fresh dialysis buffer with either 2 mM CaCl_2 or 2 mM EGTA for 120 seconds. A new baseline was acquired for 60 seconds. The biosensor tip with bound-DREAM was submerged in a 4 μL drop containing 2 μL of 2 μM hexokinase I, and 2 μL of dialysis buffer containing 2 mM CaCl_2 or EGTA. The response was recorded for 240 seconds. The response to dissociation was recorded by submerging the biosensor tip in to Buffer A with either 2 mM CaCl_2 or EGTA for 240 seconds. Kinetic constants were derived by using BLItz pro software.

Measurement of Apoptosis in Neuroblastoma N1E-115 cells

N1E-115 mouse neuroblastoma cells were grown in DMEM supplemented with 10% FBS and penicillin-streptomycin. Cells in two 24-well plates were transiently transfected using Lipofectamine with each of the following chimeric plasmids: 1) Sense DREAM cDNA in pcDNA3.1 vector, and hexokinase I cDNA in pCMV6-RFP vector; 2) pcDNA3.1 blank vector and hexokinase I cDNA in a pCMV6-RFP vector, 3) Sense DREAM in pcDNA3.1 vector and a blank cDNA pCMV6-RFP vector, 4) pcDNA3.1 blank cDNA vector and pCMV6-RFP blank cDNA vector. Cells were cultured for 48 hours and transferred in equal numbers to a 96-well plate. After 24 hours, cells were treated with 5 $\mu\text{g}/\text{mL}$ of cisplatin. Caspase-3 activity was measured after 24 hours treatment with cisplatin using a Caspase-3 Fluorescence assay kit (Cayman Chemicals, Ann Arbor, MI). Plates were read using a SpectraMax M2 plate reader with excitation and emission wavelengths set to 485 nm and 535 nm, respectively.

Isolation of mitochondria and assessment of hexokinase-1 bound to mitochondria

N1E-115 neuroblastoma cells were cultured in 60 mm dishes ($n = 18$) and DMEM, 10% FBS. Cells were transfected with chimeric DREAM-pcDNA3.1 plasmid ($n = 9$), or with pcDNA3.1 plasmid ($n = 9$) using Lipofectamine and grown for 24 hours post-transfection. Cells were scraped from the dishes and centrifuged at $1000 \times g$ for 5 minutes. Cell pellets were washed with 1 mL mitochondrial isolation buffer (MIB, 10 mM Tris-HCl, pH 7.4, 0.32 mM sucrose, and 1 mM EDTA). After centrifuging at $1000 \times g \sim 7$ volumes MIB was added to the pellet. The pellet was resuspended, and homogenized using a DUAL PTFE pestle and glass tube (Kimble Chase, Vineland NJ). The cell lysate was centrifuged at $1000 \times g$ for five minutes. The resultant supernatant was centrifuged at $18,000 \times g$ for 15 minutes. 1 mL of MIB was added to the pellet which was centrifuged for 10 minutes at $18,000 \times g$. The final mitochondrial pellet was lysed by with H_2O . Protein content was determined by the Bio-Rad protein assay. Thirty μg of protein from each lysate, and 20 ng and 30 ng standards of recombinant hexokinase I were electrophoresed on 10% Bis-Tris SDS-PAGE gels (Life Technologies), and transferred and immuno-blotted as described earlier. Hexokinase I bands were imaged and quantitated using the NIH Image program.

RESULTS AND DISCUSSION

Hexokinase I is captured by a GST- DREAM affinity column in the presence of calcium but not in the presence of EDTA

As shown in Figure 1, following exposure of rat brain homogenates to GST-DREAM glutathione Sepharose in the presence of Ca^{2+} , a high molecular weight protein band was captured that on subsequent analysis by mass spectrometry was identified as hexokinase I. In the presence of EDTA (and absence of Ca^{2+}) no hexokinase I related peptides were identified on mass spectrometry. The results demonstrated that Ca^{2+} is required for the interaction of DREAM with hexokinase I.

The interaction of DREAM with hexokinase I is dependent on the presence Ca^{2+} -binding domains of DREAM

As shown in Figure 2, when rat brain homogenates were bound to a GST-full-length DREAM resin in the presence of Ca^{2+} (Lane 1), bound hexokinase I was readily detected. In the presence of EDTA (Lane 2) bound hexokinase I was not detected. Appropriate controls with GST-resin or resin alone (Lanes 7 and 8, and lanes 9 and 10) showed no hexokinase I binding. To confirm the importance of Ca^{2+} -binding in DREAM-hexokinase I interactions, we examined the binding of hexokinase I derived from rat brain homogenates to a GST-EFMut1234 DREAM resin (Lanes 5 and 6). No binding of hexokinase I was observed, confirming that in the absence of Ca^{2+} -binding and associated conformational to DREAM, interactions with hexokinase I do not occur.

Further assessment of the influence of Ca^{2+} on DREAM-hexokinase I interactions was performed by bio-layer interferometry. As shown in Table 1, the K_D for the interaction between DREAM-hexokinase I in the presence of Ca^{2+} was 1.57×10^{-6} M whereas in the presence of EDTA the K_D was 5.72×10^{-6} M.

The interaction of DREAM with hexokinase I is dependent on the presence of the NH_2 -terminus of DREAM and the presence of Ca^{2+} -binding domains of DREAM

Neuronal calcium-sensor proteins, including DREAM, often bind other proteins via their NH_2 - terminal domains. To test whether the NH_2 -terminus of DREAM was required for binding to hexokinase I, we examined the interaction of brain-derived hexokinase I with a GST-sFL-DREAM resin. The bound DREAM construct lacks the first 94 residues of DREAM. As noted in Figure 2, (Lanes 3 and 4), no hexokinase I is bound to the GST-sFL-DREAM resin either in the presence or absence of Ca^{2+} .

The influence of DREAM on hexokinase I function

As shown in Table 2, the addition of DREAM did not influence the activity of hexokinase I. To assess the influence of DREAM on hexokinase I-modulated apoptosis, we transfected neuroblastoma cells with DREAM and-induced apoptosis by the addition of cisplatin and assessed the influence of hexokinase I on this process. Caspase-3 in cisplatin treated, blank vector-transfected neuroblastoma cells (Figure 3, column 1) is increased by transfection of cells with a DREAM expression plasmid (Figure 3, column 2, $p < 0.005$). Transfection of neuroblastoma cells with a hexokinase I expression plasmid does not reduce caspase-3 activity when compared to blank vector-transfected cells (Figure 3, column 4 versus column 1). However, following transfection of neuroblastoma cells with a DREAM expression plasmid, the increased caspase-3 noted in such cells is reduced by co-transfection with a hexokinase I expression plasmid (Figure 3, column 3 versus column 4, $p < 0.05$).

DREAM alters hexokinase I localization

To determine how DREAM alters hexokinase I function, we assessed the amount of hexokinase I on mitochondrial membranes in the presence and absence of DREAM overexpression. Hexokinase I amount was assessed using immunoblotting and a specific antibody directed against the protein. As shown in Figure 4, overexpression of DREAM reduced the amount of hexokinase I localized to the surface of mitochondria.

The regulation of apoptosis is important in understanding normal development and the pathogenesis of diseases such as cancer and neurodegenerative diseases [1,3,5,6,7,25,26]. Apoptosis is controlled by multiple pathways and proteins such as the B-cell lymphoma-2 (Bcl-2) proteins. Many of the effectors of apoptosis alter mitochondrial permeability and the release of cytochrome c, which in turn influences caspase activity [4,26,27,28,29,30,31,32,33,34]. The VDAC plays a key role in determining mitochondrial permeability [33,35,36,37,38,39]. Its activity is regulated by hexokinase I and II. The binding of hexokinase I and II to the VDAC decreases channel conductance, cytochrome c release and apoptosis [28,40,41]. Overexpression of VDAC1 is noted in Alzheimer's hippocampi, and increases in hippocampal pVDAC1 as well as a decrease in HK1, pAkt and pGSK3 expression are found in a mouse model of Alzheimer's disease [42,43].

EF-hand proteins such as calbindin D_{28K} [44] and DREAM regulate apoptosis [9,10,15,17,19]. DREAM increases the rate of apoptosis in neural cells [15,16,17,18] and is expressed in increased amounts in the hippocampus of patients with Alzheimer's disease and in animal models of Alzheimer's disease [15,17,20]. DREAM interacts with the carboxyl-terminal region of presenilin-2 and alters presenilin regulation of γ -secretase activity [9,10]. In addition, DREAM modulates intracellular Ca²⁺ concentrations and opposes some of the effects of presenilin on intracellular Ca²⁺ concentrations [19].

The data presented in this report suggests a heretofore unrecognized pathway by which DREAM influences apoptosis through mitochondrial hexokinase I binding and localization. The role of other KChIPs in hexokinase binding was not investigated. It is likely (although not completely definitive) that DREAM binds only hexokinase I, as no peptides from hexokinase II were conclusively identified. Since the amino acid sequences of the hexokinase I and II are 72% identical, and mass spectrometry may not identify segments that are divergent, we cannot be completely sure that DREAM does not bind hexokinase II. The amino acids in hexokinase I required for the interaction with DREAM have not been identified. It is unknown if DREAM displaces hexokinase I from VDAC. The interaction between DREAM and hexokinase I could conceivably be an important target for modulating apoptosis. Targeted down-regulation of DREAM may allow binding of HK1 to VDAC preventing apoptosis and neurodegeneration.

CONCLUSION

DREAM alters apoptosis in neural cells by binding hexokinase I, and reducing mitochondrial hexokinase I localization.

Acknowledgments

Supported by grants from the NIH (AR-058003 and AR-60869), and from the Marion and Ralph Falk Medical Trust.

Abbreviations

Ca²⁺ calcium ion

DREAM	downstream regulatory element antagonist modulator
KChIP3	potassium channel interacting protein
EDTA	ethylenediaminetetraacetic acid
EGTA	ethylene glycol-bis(2-aminoethylether)- <i>N,N,N',N'</i> -tetraacetic acid

REFERENCES

1. Breiden DE, Rao RV, Mehlen P. Cell death in the nervous system. *Nature*. 2006; 443:796–802. [PubMed: 17051206]
2. Crawford ED, Wells JA. Caspase substrates and cellular remodeling. *Annual review of biochemistry*. 2011; 80:1055–1087.
3. Evan GI, Vousden KH. Proliferation, cell cycle and apoptosis in cancer. *Nature*. 2001; 411:342–348. [PubMed: 11357141]
4. Hammerman PS, Fox CJ, Thompson CB. Beginnings of a signal-transduction pathway for bioenergetic control of cell survival. *Trends in biochemical sciences*. 2004; 29:586–592. [PubMed: 15501677]
5. Jiang X, Wang X. Cytochrome C-mediated apoptosis. *Annual review of biochemistry*. 2004; 73:87–106.
6. Baehrecke EH. How death shapes life during development, *Nature reviews. Molecular cell biology*. 2002; 3:779–787. [PubMed: 12360194]
7. Mathew R, Karantza-Wadsworth V, White E. Role of autophagy in cancer, *Nature reviews. Cancer*. 2007; 7:961–967. [PubMed: 17972889]
8. Carrion AM, Link WA, Ledo F, Mellstrom B, Naranjo JR. DREAM is a Ca²⁺-regulated transcriptional repressor. *Nature*. 1999; 398:80–84. [PubMed: 10078534]
9. Buxbaum JD, Choi EK, Luo Y, Lilliehook C, Crowley AC, Merriam DE, Wasco W. Calsenilin: a calcium-binding protein that interacts with the presenilins and regulates the levels of a presenilin fragment. *Nature medicine*. 1998; 4:1177–1181.
10. Choi EK, Zaidi NF, Miller JS, Crowley AC, Merriam DE, Lilliehook C, Buxbaum JD, Wasco W. Calsenilin is a substrate for caspase-3 that preferentially interacts with the familial Alzheimer's disease-associated C-terminal fragment of presenilin 2. *The Journal of biological chemistry*. 2001; 276:19197–19204. [PubMed: 11278424]
11. Hammond PI, Craig TA, Kumar R, Brimijoin S. Regional and cellular distribution of DREAM in adult rat brain consistent with multiple sensory processing roles. *Brain Res Mol Brain Res*. 2003; 111:104–110. [PubMed: 12654510]
12. Shibata R, Misonou H, Campomanes CR, Anderson AE, Schrader LA, Doliveira LC, Carroll KI, Sweatt JD, Rhodes KJ, Trimmer JS. A fundamental role for KChIPs in determining the molecular properties and trafficking of Kv4.2 potassium channels. *The Journal of biological chemistry*. 2003; 278:36445–36454. [PubMed: 12829703]
13. Alexander JC, McDermott CM, Tunur T, Rands V, Stelly C, Karhson D, Bowlby MR, An WF, Sweatt JD, Schrader LA. The role of calsenilin/DREAM/KChIP3 in contextual fear conditioning. *Learning & memory*. 2009; 16:167–177. [PubMed: 19223600]
14. Cheng HY, Pitcher GM, Laviolette SR, Whishaw IQ, Tong KI, Kockeritz LK, Wada T, Joza NA, Crackower M, Goncalves J, Sarosi I, Woodgett JR, Oliveira-dos-Santos AJ, Ikura M, van der Kooy D, Salter MW, Penninger JM. DREAM is a critical transcriptional repressor for pain modulation. *Cell*. 2002; 108:31–43. [PubMed: 11792319]
15. Jo DG, Lee JY, Hong YM, Song S, Mook-Jung I, Koh JY, Jung YK. Induction of pro-apoptotic calsenilin/DREAM/KChIP3 in Alzheimer's disease and cultured neurons after amyloid-beta exposure. *Journal of neurochemistry*. 2004; 88:604–611. [PubMed: 14720210]
16. Lilliehook C, Chan S, Choi EK, Zaidi NF, Wasco W, Mattson MP, Buxbaum JD. Calsenilin enhances apoptosis by altering endoplasmic reticulum calcium signaling. *Molecular and cellular neurosciences*. 2002; 19:552–559. [PubMed: 11988022]

17. Jo DG, Chang JW, Hong HS, Mook-Jung I, Jung YK. Contribution of presenilin/gamma-secretase to calsenilin-mediated apoptosis. *Biochemical and biophysical research communications*. 2003; 305:62–66. [PubMed: 12732196]
18. Rivas M, Aurrekoetxea K, Mellstrom B, Naranjo JR. Redox signaling regulates transcriptional activity of the Ca²⁺-dependent repressor DREAM. *Antioxidants & redox signaling*. 2011; 14:1237–1243. [PubMed: 20618065]
19. Leissring MA, Yamasaki TR, Wasco W, Buxbaum JD, Parker I, LaFerla FM. Calsenilin reverses presenilin-mediated enhancement of calcium signaling. *Proceedings of the National Academy of Sciences of the United States of America*. 2000; 97:8590–8593. [PubMed: 10900016]
20. Lilliehook C, Bozdagi O, Yao J, Gomez-Ramirez M, Zaidi NF, Wasco W, Gandy S, Santucci AC, Haroutunian V, Huntley GW, Buxbaum JD. Altered Abeta formation and long-term potentiation in a calsenilin knock-out. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2003; 23:9097–9106. [PubMed: 14534243]
21. Craig TA, Benson LM, Venyaminov SY, Klimtchuk ES, Bajzer Z, Prendergast FG, Naylor S, Kumar R. The metal-binding properties of DREAM: evidence for calcium-mediated changes in DREAM structure. *The Journal of biological chemistry*. 2002; 277:10955–10966. [PubMed: 11788589]
22. Ramachandran PL, Craig TA, Atanasova EA, Cui G, Owen BA, Bergen HR 3rd, Mer G, Kumar R. The Potassium Channel Interacting Protein 3 (DREAM/KChIP3) Heterodimerizes with and Regulates Calmodulin Function. *The Journal of biological chemistry*. 2012; 287:39439–39448. [PubMed: 23019329]
23. Craig TA, Benson LM, Venyaminov SY, Klimtchuk ES, Bajzer Z, Prendergast FG, Naylor S, Kumar R. The metal-binding properties of DREAM: evidence for calcium-mediated changes in DREAM structure. *J Biol Chem*. 2002; 277:10955–10966. [PubMed: 11788589]
24. Scheer WD, Lehmann HP, Beeler MF. An improved assay for hexokinase activity in human tissue homogenates. *Analytical biochemistry*. 1978; 91:451–463. [PubMed: 9762131]
25. Fischer SJ, McDonald ES, Gross L, Windebank AJ. Alterations in cell cycle regulation underlie cisplatin induced apoptosis of dorsal root ganglion neurons in vivo. *Neurobiology of disease*. 2001; 8:1027–1035. [PubMed: 11741398]
26. Leist M, Jaattela M. Four deaths and a funeral: from caspases to alternative mechanisms, *Nature reviews. Molecular cell biology*. 2001; 2:589–598. [PubMed: 11483992]
27. Adrain C, Creagh EM, Martin SJ. Apoptosis-associated release of Smac/DIABLO from mitochondria requires active caspases and is blocked by Bcl-2. *The EMBO journal*. 2001; 20:6627–6636. [PubMed: 11726499]
28. Azoulay-Zohar H, Israelson A, Abu-Hamad S, Shoshan-Barmatz V. In self-defence: hexokinase promotes voltage-dependent anion channel closure and prevents mitochondria-mediated apoptotic cell death. *Biochem J*. 2004; 377:347–355. [PubMed: 14561215]
29. Gottlob K, Majewski N, Kennedy S, Kandel E, Robey RB, Hay N. Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes Dev*. 2001; 15:1406–1418. [PubMed: 11390360]
30. Green DR, Galluzzi L, Kroemer G. Mitochondria and the autophagy-inflammation-cell death axis in organismal aging. *Science*. 2011; 333:1109–1112. [PubMed: 21868666]
31. Lin MT, Beal MF. Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature*. 2006; 443:787–795. [PubMed: 17051205]
32. Ow YP, Green DR, Hao Z, Mak TW. Cytochrome c: functions beyond respiration, *Nature reviews. Molecular cell biology*. 2008; 9:532–542. [PubMed: 18568041]
33. Shoshan-Barmatz V, Keinan N, Abu-Hamad S, Tyomkin D, Aram L. Apoptosis is regulated by the VDAC1 N-terminal region and by VDAC oligomerization: release of cytochrome c, AIF and Smac/Diablo. *Biochim Biophys Acta*. 2010; 1797:1281–1291. [PubMed: 20214874]
34. Zamzami N, Kroemer G. The mitochondrion in apoptosis: how Pandora's box opens, *Nature reviews. Molecular cell biology*. 2001; 2:67–71. [PubMed: 11413468]
35. Colombini M. VDAC: the channel at the interface between mitochondria and the cytosol. *Molecular and cellular biochemistry*. 2004; 256–257:107–115.

36. De Stefani D, Bononi A, Romagnoli A, Messina A, De Pinto V, Pinton P, Rizzuto R. VDAC1 selectively transfers apoptotic Ca^{2+} signals to mitochondria. *Cell death and differentiation*. 2012; 19:267–273. [PubMed: 21720385]
37. Granville DJ, Gottlieb RA. The mitochondrial voltage-dependent anion channel (VDAC) as a therapeutic target for initiating cell death. *Current medicinal chemistry*. 2003; 10:1527–1533. [PubMed: 12871124]
38. Shoshan-Barmatz V, Mizrahi D. VDAC1: from structure to cancer therapy. *Frontiers in oncology*. 2012; 2:164. [PubMed: 23233904]
39. Thomsen MB, Wang C, Ozgen N, Wang HG, Rosen MR, Pitt GS. Accessory subunit KChIP2 modulates the cardiac L-type calcium current. *Circulation research*. 2009; 104:1382–1389. [PubMed: 19461043]
40. Abu-Hamad S, Zaid H, Israelson A, Nahon E, Shoshan-Barmatz V. Hexokinase-I protection against apoptotic cell death is mediated via interaction with the voltage-dependent anion channel-1: mapping the site of binding. *J Biol Chem*. 2008; 283:13482–13490. [PubMed: 18308720]
41. Pastorino JG, Hoek JB. Hexokinase II: the integration of energy metabolism and control of apoptosis. *Current medicinal chemistry*. 2003; 10:1535–1551. [PubMed: 12871125]
42. Cuadrado-Tejedor M, Vilarino M, Cabodevilla F, Del Rio J, Frechilla D, Perez-Mediavilla A. Perez-Mediavilla, Enhanced expression of the voltage-dependent anion channel 1 (VDAC1) in Alzheimer's disease transgenic mice: an insight into the pathogenic effects of amyloid-beta. *Journal of Alzheimer's disease : JAD*. 2011; 23:195–206.
43. Ferrer I. Altered mitochondria, energy metabolism, voltage-dependent anion channel, and lipid rafts converge to exhaust neurons in Alzheimer's disease. *Journal of bioenergetics and biomembranes*. 2009; 41:425–431. [PubMed: 19798558]
44. Kojetin DJ, Venters RA, Kordys DR, Thompson RJ, Kumar R, Cavanagh J. Structure, binding interface and hydrophobic transitions of Ca^{2+} -loaded calbindin-D(28K). *Nat Struct Mol Biol*. 2006; 13:641–647. [PubMed: 16799559]

HIGHLIGHTS

- DREAM increases apoptosis in neural cells.
- DREAM binds hexokinase I, in a calcium-dependent manner.
- DREAM reduces mitochondrial HK I concentrations.
- DREAM-dependent apoptosis is reduced by overexpression of hexokinase I.
- DREAM and hexokinase I interact to regulate apoptosis in cells.

Figure 1.

Total number of unique hexokinase I peptides identified from MS/MS analysis of pull-down experiment with rat brain homogenates applied to a GST-full-length DREAM (amino acids 1–256) resin in the presence or absence of Ca^{2+} .

Figure 2.

Western blot analysis of rat brain homogenate proteins bound to various constructs of GST-DREAM immobilized on glutathione resin in the presence or absence of Ca^{2+} . The lane designation are as follows: 1) GST-FL DREAM (1–256), Ca^{2+} present, 2) GST-FL DREAM, EDTA present, 3) GST-sFL DREAM (95–256), Ca^{2+} present, 4) GST-sFL DREAM, EDTA present, 5) GST-EFMut1234 DREAM, Ca^{2+} present, 6) GST-EFMUT1234 DREAM, EDTA present, 7) GST, Ca^{2+} present, 8) GST, EDTA present, 9) Resin, Ca^{2+} present, 10) Resin, EDTA present.

Figure 3.

Caspase-3 assay of cisplatin-induced apoptosis in N1E-115 neuroblastoma cells transfected with DREAM and/or hexokinase I expression plasmids. Column 1. Cells transfected with blank vectors, pcDNA3.1 and pCMV6. Column 2. Cells transfected with sense DREAM-pcDNA3.1 vector and blank pCMV6 vector. Column 3. Cells transfected with DREAM-pcDNA3.1 vector plus hexokinase I-RFP vector. Column 4. Cells transfected with hexokinase I-RFP vector and pcDNA3.1 blank vector. The cells were treated with 5 μ g/ml cisplatin for 24 hours. Number of unique observations = 5. * $P < 0.05$, *** $P < 0.005$

Figure 4.

Hexokinase I bound to isolated mitochondria. Hexokinase I bound (ng) to isolated mitochondria from N1E-115 neuroblastoma cells transfected with either DREAM-pcDNA3.1 or pcDNA3.1. Mean $\pm$ standard error of mean. $P < 0.005$, $n = 9$.

The association and dissociation constants of DREAM and hexokinase I as assessed by bio-layer interferometry.

Table 1

	K_D (M)	k_a (1/Ms)	k_a Error	k_d (1/s)	k_d Error
Ca²⁺ Condition	1.57 × 10 ⁻⁶	6.28 × 10 ³	5.36 × 10 ²	9.85 × 10 ⁻³	1.52 × 10 ⁻⁴
EGTA Condition	5.72 × 10 ⁻⁶	2.14 × 10 ³	1.07 × 10 ³	1.23 × 10 ⁻²	3.11 × 10 ⁻⁴

Table 2

The influence of DREAM on hexokinase I glycolytic activity (mean $\pm$ standard deviation).

Buffer alone + Ca ²⁺ n = 63	DREAM + buffer + Ca ²⁺ n = 63	Buffer alone + EGTA n = 64	DREAM + buffer + EGTA n = 64
8.15 $\pm$ 9.11	8.75 $\pm$ 9.57	8.35 $\pm$ 9.00	9.24 $\pm$ 10.22