Published in final edited form as:

Org Lett. 2013 April 5; 15(7): 1568–1571. doi:10.1021/ol4003459.

Chiral Brønsted Acid Catalyzed Enantioselective Synthesis of *anti*-Homopropargyl Alcohols via Kinetic Resolution-Aldehyde Allenylboration Using Racemic Allenylboronates

Andy S. Tsai, Ming Chen, and William R. Roush

Department of Chemistry, Scripps Florida, Jupiter, Florida 33458, United States

William R. Roush: roush@scripps.edu

Abstract

A chiral phosphoric acid catalyzed kinetic resolution/allenylboration of racemic allenylboronates with aldehydes is described. Allenylboration of aldehydes with 2.8 equivalents of allenylboronate ($\pm$)-1 in the presence of 10 mol% of catalyst (*R*)-2 provided *anti*-homopropargyl alcohols **3** in 83–93% yield with 9:1 to 20:1 diastereoselectivity and 73–95% e.e. The catalyst enables the kinetic resolution of the racemic allenylboronate ($\pm$)-1 to set the methyl stereocenter and biases the facial attack of the aldehyde to set the stereochemistry of the hydroxyl group in **3**.

Asymmetric crotylation of aldehydes with crotylboron reagents is a well established method to generate stereodefined acyclic molecules bearing an olefinic handle.¹ Similarly, allenylboration of 3-substituted allenylboron reagents provides homopropargyl alcohols with an alkyne moiety which are equally valuable in polyketide natural product syntheses.^{2,3} Due to the closed transition states involved in allenylboration reactions, the axial chirality of 3-substituted allenylboron reagents is directly transferred to the stereochemistry of the propargylic position in the product (e.g. the position that bears the methyl group in **3** and **4**, Scheme 1).^{3,4} The stereochemistry of the hydroxyl group in the homopropargyl products is controlled by the facial approach of the allenylboronate to the aldehyde.

As shown in Scheme 1, allenylboration of an aldehyde with (*M*)-**1** can provide either *anti*-homopropargyl alcohol *ent*-**3** or the *syn* adduct **4** via the two competing transition states **TS-1** and **TS-2**. These two transition states, arising from opposing diastereofacial approaches of the reagent to the aldehyde substrates, often have similar energy and poor *anti/syn* diastereoselectivity is generally observed.^{3b–c,3h,4} Consequently, in order to obtain homopropargyl alcohol products with high diastereo- and enantioselectivity, single enantiomer 3-substituted allenylboron reagents are required as well as a means to control the diastereofacial selectivity of the addition of the reagent to the aldehyde substrate.

Correspondence to: William R. Roush, roush@scripps.edu.

Supporting Information Available: Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

We recently demonstrated that the chiral, nonracemic phosphoric acid catalyst **2**⁵ can bias the two competing transition states (e.g. **TS-1** and **TS-2**) such that either *syn*- or *anti*-products can be obtained with good to excellent diastereoselectivity and with very high enantioselectivity.⁶ As illustrated in Scheme 2, allenylboration of aldehydes with (*M*)-**1** in the presence of 5 mol% of catalyst (*S*)-**2** gave *anti*-homopropargyl alcohols *ent*-**3** in 83–98% yield and excellent diastereo- (>50:1 ds) and enantioselectivity (>98% ee). We infer that this transformation is stereochemically matched. On the other hand, mismatched allenylborations of aldehydes with (*M*)-**1** in the presence of the enantiomeric catalyst (*R*)-**2** provided the *syn*-adducts **4** with 9:1 ds and >98% ee. Importantly, the reaction of (*M*)-**1** and aldehydes in the presence of (*S*)-**2** proceeded at a faster rate than the reaction with the (*M*)-**1**/(*R*)-**2** pairing.⁶ These differing rates form the basis of a possible kinetic resolution of a racemic allenylboronate, such as ($\pm$)-**1**,⁷ in reactions catalyzed by chiral phosphoric acids such as **2**, and would eliminate the need to synthesize enantioenriched allenylboron reagents for use in aldehyde allenylboration reactions.^{2b,8} Thus, a single enantiomer phosphoric acid catalyst might be able *both* to select the more reactive allenylboronate enantiomer from a racemic mixture *and* direct its addition to the aldehyde with high diastereo facial selectivity. Accordingly, we are pleased to report that the chiral, nonracemic Brønsted acid (*R*)-**2** catalyzes the allenylboration of aldehydes with racemic allene ($\pm$)-**1** with kinetic resolution to obtain *anti*-homopropargyl alcohols **3** with high diastereo- and enantioselectivity.

Initial studies of the chiral phosphoric acid catalyzed kinetic resolution-allenylboration reactions of ($\pm$)-**1** were performed with hydrocinnamaldehyde as the substrate (Table 1). Treatment of hydrocinnamaldehyde with 2.1 equiv. of allenylboronate ($\pm$)-**1** and 5 mol% of catalyst (*R*)-**2** at ambient temperature provided enantiomerically enriched (71% ee) *anti*-homopropargyl alcohol **3a** in 92% yield with 10:1 diastereoselectivity (Table 1, entry 2). The enantiomeric excess of **3a** was determined by using the Mosher ester analysis.⁹ To confirm that the reaction indeed proceeded via the kinetic resolution pathway, the remaining allenylboronate was isolated and subjected to the uncatalyzed allenylboration with hydrocinnamaldehyde. The homopropargyl alcohol product obtained from the reaction is a 3.5:1 mixture of *anti*- and *syn*-diastereomers *ent*-**3a** and *ent*-**4a**. The enantiomeric excess of both diastereomers obtained from this experiment was 60% ee.

When the allenylboration reaction of ($\pm$)-**1** and hydrocinnamaldehyde was performed in toluene at –30 °C in the presence of 5 mol% of catalyst (*R*)-**2**, *anti*-homopropargyl alcohol **3a** was obtained with 20:1 diastereoselectivity and 78% ee (entry 3). A solvent screen showed that while dichloromethane was a competent solvent (entry 4), reactions performed in toluene provided better diastereoselectivity with nearly equivalent enantioselectivity. Coordinating solvents, such as diethyl ether, were detrimental (entry 5). Increasing the equivalents of allenylboronate ($\pm$)-**1** used in the reaction to 2.8 equiv resulted in improved enantioselectivity (84% ee; entry 6). Finally, by increasing the catalyst loading to 10 mol% and by performing the reaction in toluene at –50 °C for 16 h, the allenylboration reaction provided homopropargyl alcohol **3a** in 93% yield with 20:1 diastereoselectivity and 90% ee (entry 7).

The conditions developed for the kinetic resolution-allenylboration of hydrocinnamaldehyde were then applied to the reactions of a variety of representative achiral aldehydes; the results of these experiments are summarized in Figure 1. Unhindered aliphatic and α,β -unsaturated aldehydes provided homopropargyl alcohols **3a–c** with excellent diastereo- and enantioselectivities. The homopropargyl alcohols derived from a range of substituted aromatic aldehydes were obtained in 86–93% yield with 9–15:1 diastereoselectivities and 95% ee (**3f–h**). High diastereoselectivities were obtained for the reactions leading to **3d** and **3e**, however the enantioselectivities were only moderate in these cases (73–77% ee).

To further evaluate the synthetic utility of this kinetic resolution-allenylboration procedure, allenylboration reaction with the chiral, nonracemic aldehyde **5** was explored. When the allenylboration of aldehyde **5** with 2.8 equiv. of racemic **1** was performed in the presence of 10 mol% of acid catalyst (*S*)-**2**, an 8:1 mixture of homopropargyl alcohols **6** and **7** was obtained, favoring the *anti*, *anti*-homopropargyl alcohol **6**. This result demonstrates an exceedingly simple means to generate the *anti*, *anti*-stereotriad from the racemic allenylboronate ($\pm$)-**1**.¹⁰

Finally, racemic allenylboronates ($\pm$)-**8** and ($\pm$)-**9** were tested in this kinetic resolution-allenylboration reaction sequence (Scheme 4). By using the conditions determined to be optimal in our studies of the kinetic resolution-allenylboration reactions of ($\pm$)-**1**, the allenylboration of hydrocinnamaldehyde with allene ($\pm$)-**8** provided a 6.5:1 mixture of homopropargyl alcohols **10** and **11** with a terminal alkyne unit in 85% yield and 73% ee. Similarly, the allenylboration of hydrocinnamaldehyde with allene ($\pm$)-**9** (2.8 equiv) and 10 mol % of catalyst (*R*)-**2** at -10°C afforded homopropargyl alcohols **12** and **13** in 78% yield with 10:1 diastereoselectivity and 67% ee. These results suggest that further optimization may be required in order to achieve high diastereo- and enantioselectivity in the kinetic resolution-allenylboration reactions of substrates like ($\pm$)-**8** and ($\pm$)-**9**.

In summary, we developed a chiral Brønsted acid catalyzed kinetic resolution-allenylboration reaction of racemic allenylboronate ($\pm$)-**1**. When performed in the presence of 10 mol% of acid (*R*)-**2**, the kinetic resolution-aldehyde allenylboration reaction of ($\pm$)-**1** provided *anti*-homopropargyl alcohols **3** in 83–93% yield with 9:1 to 20:1 diastereoselectivity and 73–95% ee. The kinetic resolution of racemic allene ($\pm$)-**1** and its controlled diastereofacial addition to a chiral aldehyde was exemplified by the chiral Brønsted acid catalyzed allenylboration of **5**, which provided the *anti*, *anti*-homopropargyl alcohol **6** with 8:1 selectivity. Synthetic applications of this methodology will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support provided by the National Institutes of Health (GM038436 and CA162504) is gratefully acknowledged. We thank Eli Lilly for a Graduate Fellowship to M. Chen.

References

- (a) Roush, WR. Comprehensive Organic Synthesis. Trost, BM.; Fleming, I., editors. Vol. 2. Pergamon Press; New York: 1991. p. 1(b) Denmark SE, Fu J. Chem Rev. 2003; 103:2763. [PubMed: 12914480] (c) Lachance H, Hall DG. Org React. 2008; 73:1.(d) Yus M, Gonzles-Gmez JC, Foubelo F. Chem Rev. 2011; 111:7774. [PubMed: 21923136]
- For recent reviews on asymmetric propargylation of carbonyl compounds, see: Ding CH, Hou XL. Chem Rev. 2011; 111:1914. [PubMed: 21344874] Marshall JA. J Org Chem. 2007; 72:8153. [PubMed: 17595141] For selected asymmetric aldehyde propargylation reactions where only one stereocenter is set, see: Haruta R, Ishiguro M, Ikeda N, Yamamoto H. J Am Chem Soc. 1982; 104:7667. Corey EJ, Yu CM, Lee DH. J Am Chem Soc. 1990; 112:878. Hernandez E, Burgos CH, Alicea E, Soderquist JA. Org Lett. 2006; 8:4089. [PubMed: 16928081] Shi SL, Oisaki K, Kanai M, Shibasaki M. J Am Chem Soc. 2010; 132:6638. [PubMed: 20420398] Fandrick DR, Fandrick KR, Reeves JT, et al. J Am Chem Soc. 2010; 132:7600. [PubMed: 20481453] Barnett DS, Schaus SE. Org Lett. 2011; 13:4020. [PubMed: 21732609] Keck GE, Krishnamurthy D, Chen X. Tetrahedron Lett. 1994; 35:8323. Yu CM, Yoon SK, Choi HS, Baek K. Chem Commun. 1997:763. Denmark SE,

- Wynn TJ. *Am Chem Soc.* 2001; 123:6199. Even DA, Sweeney ZK, Rovis T, Tedrow JS. *J Am Chem Soc.* 2001; 123:12095. [PubMed: 11724622] Chen J, Captain B, Takenaka N. *Org Lett.* 2011; 13:1654. [PubMed: 21366250] Woo SK, Geary LM, Krische MJ. *Angew Chem Int Ed.* 2012; 51:7830. Jain P, Wang H, Houk KN, Antilla JC. *Angew Chem, Int Ed.* 2012; 51:1391. Reddy LR. *Org Lett.* 2012; 14:1142. [PubMed: 22273041] Haddad TD, Hirayama LC, Buckley JJ, Singaram B. *J Org Chem.* 2012; 77:889. [PubMed: 22148263] Hirayama LC, Daddad TD, Oliver AG, Singaram B. *J Org Chem.* 2012; 77:4342. [PubMed: 22497595] Wan H, Jain P, Antilla JC, Houk KN. *J Org Chem.* 2013; 78:1208. [PubMed: 23298338]
3. For selected asymmetric propargylations of carbonyl compounds where two stereocenters are set, see: Marshall JA. *Chem Rev.* 1996; 96:31. [PubMed: 11848743] Ito H, Sasaki Y, Sawamura M. *J Am Chem Soc.* 2008; 130:15774. [PubMed: 18973297] Matsumoto Y, Naito M, Uozumi Y, Hayashi T. *J Chem Soc, Chem Commun.* 1993:1468. Han JW, Tokunaga N, Hayashi T. *J Am Chem Soc.* 2001; 123:12915. [PubMed: 11749556] Marshall JA, Adams ND. *J Org Chem.* 1997; 62:8976. Marshall HA, Maxson K. *J Org Chem.* 2000; 65:630. [PubMed: 10813988] Brawn RA, Panek JS. *Org Lett.* 2007; 9:2689. [PubMed: 17559219] Shimizu M, Kurahashi T, Kitagawa H, Hiyama T. *Org Lett.* 2003; 5:225. [PubMed: 12529146] Marshall JA, Wang XJ. *J Org Chem.* 1992; 57:1242. Marshall JA, Perkins J. *J Org Chem.* 1994; 59:3509. Marshall JA, Adams ND. *J Org Chem.* 1999; 64:5201. Marshall JA, Grant CM. *J Org Chem.* 1999; 64:696. [PubMed: 11674133] Marshall JA, Chobanian HR, Yanik MM. *Org Lett.* 2001; 3:3369. [PubMed: 11594836] Geary LM, Woo SK, Leung JC, Krische MJ. *Angew Chem, Int Ed.* 2012; 51:2972.
4. Sasaki Y, Sawamura M, Ito H. *Chem Lett.* 2011; 40:1044.
5. For recent reviews of Brønsted acid catalysts, see: Akiyama T. *Chem Rev.* 2007; 107:5744. [PubMed: 17983247] Rueping M, Kuenkel A, Atodiresei I. *Chem Soc Rev.* 2011; 40:4539. [PubMed: 21614342] Rueping M, Nachtsheim BJ, Ieawsuwan W, Atodiresei I. *Angew Chem Int Ed.* 2011; 50:6706. Giacalone F, Gruttadauria M, Agrigento P, Noto R. *Chem Soc Rev.* 2012; 41:2406. [PubMed: 22167174]
6. Chen M, Roush WR. *J Am Chem Soc.* 2012; 134:10947. [PubMed: 22731887]
7. For a recent study of kinetic resolution in the enantioselective hydroboration of racemic allenylboronate ($\pm$)-**1** with diisopinocampheylborane, see: Han JL, Chen M, Roush WR. *Org Lett.* 2012; 14:3028. [PubMed: 22646839]
8. The synthesis of these reagents have generally been derived from the S_N2' substitution of enantioenriched propargyl alcohol derivatives or enantioselective hydrometalation of enynes amongst other methods.; see refs. 2a and 3.
9. (a) Dale JA, Mosher HS. *J Am Chem Soc.* 1973; 95:512. (b) Ohtani I, Kusumi T, Kashman Y, Kakisawa H. *J Am Chem Soc.* 1991; 113:4092.
10. (a) Hoffmann RW. *Angew Chem, Int Ed.* 1987; 26:489. (b) Hoffmann RW, Dahmann G, Andersen MW. *Synthesis.* 1994:629. (c) Marshall JA, Perkins J, Wolf MA. *J Org Chem.* 1995; 60:5556. (d) Chemler SR, Roush WR. *J Org Chem.* 1998; 63:3800. (e) Chemler SR, Roush WR. *J Org Chem.* 2003; 68:1319. [PubMed: 12585871] (f) Chen M, Roush WR. *J Am Chem Soc.* 2012; 134:3925. [PubMed: 22332989] (g) Chen M, Roush WR. *Org Lett.* 2012; 14:426. [PubMed: 22148850] (h) Chen M, Roush WR. *Org Lett.* 2012; 14:1880. [PubMed: 22409510]

Figure 1.

Kinetic Resolution/Allenylboration Reactions of Representative Aldehydes with Allenylboronate (±)-**1**^{a-d}

^aReactions were performed with 0.2 mmol of aldehyde at a concentration of 0.2 M. ^bProduct diastereoselectivities were determined by ¹H NMR analysis of crude reaction products. ^cCombined isolated yield of both diastereomers. ^dEnantiomeric purity of the major product was determined by Mosher ester analysis.⁹

Scheme 1.
Aldehyde Allenylboration Reactions with Allenylboronate (*M*)-1

Scheme 2. Proposed Kinetic Resolution/Aldehyde Allenylboration with Racemic Allenylboronate **(±)-1**

Scheme 3.
Allenylboration of Chiral Aldehyde **5**

Scheme 4. Kinetic Resolution/Allenylboration Reactions with Racemic Allenylboronates **8** and **9**

Table 1

Kinetic Resolution/Allenylboration Reactions of Hydrocinnamaldehyde Using Racemic Allenylboronate ($\pm$)-**1** and Chiral Acid (*R*)-**2**.^a

entry	solvent	temp (°C)	time (h)	(<i>R</i>)- 2 (mol %)	($\pm$)- 1 (equiv)	<i>d</i> _s ^b	yield ^c	% ee ^d
1	Toluene	55	24	0	1.1	3.5:1	90%	-
2	Toluene	23	24	5%	2.1	10:1	92%	71
3	Toluene	-30	24	5%	2.1	20:1	92%	78
4	CH ₂ Cl ₂	-30	24	5%	2.1	11:1	89%	80
5	Et ₂ O	-30	38	5%	2.1	6:1	63%	49
6	Toluene	-30	24	5%	2.8	20:1	89%	84
7	Toluene	-50	16	10%	2.8	20:1	93%	90

^aReactions were performed with 0.15 mmol of aldehyde at a concentration of 0.2 M.

^bProduct diastereoselectivities were determined by ¹H-NMR analysis of crude reaction mixture.

^cCombined isolated yield of both diastereomers.

^dEnantioselectivity of the major product was determined by Mosher ester analysis.⁹