Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading

doi:10.1016/S2221-1691(13)60102-1

© 2013 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Seroepidemiology of bluetongue disease in small ruminants of north-east of Iran

Vahid Najamezhad^{1*}, Mahin Rajae²¹Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran²Veterinary Council of Khorasan Razavi, Mashad, Iran

PEER REVIEW

Peer reviewer

Dr Rouz-Ali Batavani, Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Tel: +98 9141414153

Fax: +98 441 2777099

E-mail: r.batavani@urmia.ac.ir

Comments

This is a good study with a significant number of serum samples compared with similar works in other regions. The results are interesting and suggested that bluetongue can be one of the causes of abortion in small ruminants of the region.

Details on Page 494

ABSTRACT

Objective: To estimate the prevalence and distribution of bluetongue virus antibody in sheep and goats in 25 townships of Khorasan Razavi. Bluetongue is an infectious, non-contagious, arthropod born viral disease of ruminants and has been reported from most of the tropical and subtropical regions of the world. **Methods:** A total number of 1034 serum samples from sheep and goats were collected and transmitted to Serological Laboratory of Veterinary Council of Khorasan Razavi. Serums were screened for the presence of group-specific bluetongue virus antibody using competitive Enzyme Linked Immuno Sorbent Assay (c-ELISA). **Results:** The seropositivity of sheep and goats for bluetongue was found to be 89.2%. The highest prevalence rate was seen in Taybad, Khalilabad and Torbat-jam (100%) and the least prevalence rate was seen in Jovein (55%). **Conclusions:** The results showed that the majority of animals in the north-east of Iran are infected with bluetongue virus. High correlation between abortion history and seropositivity emphasize the economical importance of bluetongue virus in the sheep herds of the region.

KEYWORDS

Khorasan Razavi, Seroepidemiology, Bluetongue, Sheep, Goat

1. Introduction

Bluetongue is an infectious, non-contagious, arthropod borne, viral disease of domesticated and wild ruminants. Bluetongue virus (BTV) is the type specific of the genus *Orbivirus* in the family Reoviridae. Twenty-four antigenically distinct serotypes of BTV have been identified worldwide[1]. The BTV are transmitted between ruminants by the bite of certain vector species of *Culicoides* midges[2]. Venereal transmission between ruminants has been recorded but occurs very rarely and it is not considered to

be of epidemiological significance[3]. Among the animals, clinical signs occur most often in sheep and certain species of deer with significant morbidity[4], although cattle and goat may be affected[5]. Affected sheep may have erosion and/or ulcer on the mucous membranes and inflammation of the coronary band. Other clinical signs include fever, depression, excessive salivation, nasal discharge, facial edema, hyperaemia, lameness and death[6]. Older animals tend to be more susceptible to disease than younger ones and the severity of clinical signs seem to vary with the breed of the animal, the serotype and strain of the infecting

*Corresponding author: Vahid Najamezhad, Department of Clinical Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Tel: +98 9153023733

E-mail: v.najamezhad@urmia.ac.ir

Foundation Project: Supported by Veterinary Council of Khorasan Razavi. Grant No.: 05/09/38671.

Article history:

Received 7 May 2013

Received in revised form 12 May, 2nd revised form 16 May, 3rd revised form 20 May 2013

Accepted 10 Jun 2013

Available online 28 Jun 2013

virus and with certain rather ill-defined interactions with the environment[7,8]. Bluetongue typically occurs when susceptible animal species are introduced into areas with circulating virulent BTV strains, or when virulent BTV strains extend their range to previously unexposed populations of ruminants[9]. Some strains of the virus can result in mortality rates as high as 70% in highly susceptible sheep[1,10]. The exact worldwide economic losses due to bluetongue have not been expressed, but it estimated as 3 billion US dollar a year due to death, abortions, weight loss, reduced milk yield, meat efficiency, export restrictions for live animals, their semen and some products such as fetal bovine serum[11]. The costs of preventive and control measures should also be taken into the account[12,13]. Various techniques have been used to detect antibodies against BTV. Agar gel immuno diffusion and competitive enzyme linked immuno sorbent assay (c-ELISA) are recommended as prescribed tests for International Trade in the OIE Manual of Standards for Diagnostic Test and Vaccines[1].

Khorasan–Razavi province is located at the north–east Iran between latitudes 30°24′–38°17′ N and between longitudes 55°17′–61°15′ E. It has more than 8 000 000 small ruminants which represents an important source of income for rural areas. At present, no information is available concerning on the seroprevalence of bluetongue disease in the sheep and goats in Khorasan Razavi province. Thus, the present study was conducted to clarify different aspects of the bluetongue disease in the sheep and goats population of the region.

2. Materials and methods

A total number of 1034 serum samples were collected from apparently healthy sheep and goats of different age and sex from different herds of 22 townships of Khorasan Razavi. All the serum samples were submitted to Serological Laboratory of Veterinary Council of Khorasan Razavi and stored at –20 °C until examined. Serums were screened for the presence of group-specific BTV antibody by competitive enzyme linked immunosorbent assay[14], using the c-ELISA IDEXX Bluetongue Competition® assay (IDEXX BT, Netherland). The optic density of each sample was read by an ELISA micro plate reader (BioTek, power wave XS2, USA) at 450 nm. Results are expressed as percentage of negativity (PN) compared with the kit control and designated as positive, doubtful or negative according to the cut-off values recommended by the manufacturer (PN ≤ 70 is positive; 70 < PN < 80 is doubtful; PN ≥ 80 is negative).

2.1. Statistical analysis

Statistical analyses were performed using SPSS software version 11.5 (SPSS Inc., Chicago, IL, USA). Chi-square test and fisher exact tests were applied to analyze the association between seroprevalence status, age, sex and abortion.

quantitative variables were subjected to two independent *t*-tests. Values of *P* < 0.05 were considered to be significant.

3. Results

The results of seroprevalence bluetongue among sheep and goats in Khorasan Razavi are summarized in Table 1. In the present study, the seropositivity of sheep and goats for bluetongue was found to be 89.2% (922 out of 1034). The rate of positivity in sheep and goats was 90.0% and 87.6% respectively. Comparison of age in positive and negative groups showed that the age of positive group was significantly more than that of negative group (*P* < 0.05). Regarding abortion history in the herd, the results showed that ewes with abortion history had 92.9% and those without abortion history had 85.1% seropositivity. Goats with and without abortion history had 91.0% and 82.4% seropositivity, respectively (Table 2). Concerning the disease agent with abortion history, in Chi-square test, *P* value for sheep and goats was 0.001 and 0.012, respectively. Sex didn't affect the rate of seropositivity (*P* = 0.75).

Table 1

The result of ELISA for bluetongue antibodies in small ruminants of different areas of north–east of Iran.

City	Sheep		Goat		Seropositivity %
	Positive	Negative	Positive	Negative	
Kalat	12	2	–	–	85.7
Kashmar	20	1	15	1	94.6
Khaf	30	2	30	1	95.2
Mashad	41	9	18	2	84.3
Ghuchan	10	1	12	4	81.5
Mahvelat	11	2	7	–	90.0
Jovein	11	9	–	–	55.0
Roshtkhar	28	2	5	–	94.3
Kalat	19	5	15	7	73.9
Sarakhs	56	1	–	–	98.2
Taybad	23	–	15	–	100.0
Bajestan	13	–	13	1	96.3
Khalil–abad	15	–	–	–	100.0
Torbat–heydarie	28	1	25	2	94.6
Bardaskan	30	5	12	2	85.7
Chenaran	15	5	15	6	73.2
Dargaz	23	5	10	3	80.5
Sabzevar	44	6	20	5	85.3
Gonabad	10	2	13	4	79.3
Nishaboor	62	7	40	4	90.3
Fariman	50	2	14	3	92.8
Torbat–jam	52	–	40	–	100.0
Total	603	69	319	45	89.2

Table 2

The percentage of bluetongue antibody positive sheep and goats with and without abortion.

Species			Abortion		Total	
			yes	no		
Sheep	ELIZA	Pos	Count	392	211	603
			% within abortion	92.9	85.1	90.0
		Neg	Count	30	37	67
			% within abortion	7.1	14.9	10.0
	Total	Count	422	248	670	
		% within abortion	100.0	100.0	100.0	
	Pos	Count	202	117	319	
		% within abortion	91.0	82.4	87.6	
Goat	ELIZA	Neg	Count	20	25	45
			% within abortion	9.0	17.6	12.4
	Total	Count	222	142	364	
		% within abortion	100.0	100.0	100.0	

4. Discussion

In the present study, the seropositivity of sheep and goat for bluetongue was found to be 89.2% (922 out of 1034) which is higher than the reported seroprevalence of 6.57% in south-east of Iran^[15], 51.6% in Isfahan^[16], 34.7% in west Azerbaijan^[17] and 73.5% in Shiraz^[18]. Other studies have shown that bluetongue disease occurs worldwide and lead to significant losses in the small ruminant population^[10,17]. In Thailand, a serological study was carried out and demonstrated 39.4% and 73.0% seropositive rates in indigenous sheep and goats respectively^[19]. Seroprevalence rates up to 90% were detected in some regions of European Turkey^[20]. Bluetongue was confirmed in sheep herds located in Germany. The overall herd seroprevalences were estimated at 37.5% in 2006 and 41.5% in 2007^[21]. Since vaccination against BT is not done in Iran, antibody indicated direct exposure to the *Orbivirus*. The distribution of infected animals in Khorasan Razavi province showed the highest prevalence rate in cities near Afghanistan border. Unfortunately no information is available regarding bluetongue disease in Afghanistan. Movements of infected animals have been thought to be a source of infection of blue tongue in new areas^[22]. Despite the high seroprivalence of bluetongue in Khorasan Razavi there is no clinical report of the disease in this region. Clinical recognition of the disease depends largely on the presence of highly susceptible sheep breeds, which invariably act as indicator^[1,23].

It is reported that BTV can cause 25% abortion and 50% decrease in fertility in sheep^[24]. In the present study there was a significant relation between seroprevalence rate and

abortion history. Hence it emphasizes the importance of the BTV in the herd.

There are differences in age susceptibility to clinical disease which, inexplicably vary with different outbreaks. With Australian serotypes, disease occurs in sheep only at the age of 3 years or older^[25]. Seroprevalence increases with age, probably a reflection of increased duration of exposure^[10]. In the present study the seroprivalence rates were increased with increase of age in sheep and goats. It could be a reason for the disease to be endemic, because newborn lambs could be protected by maternal immunity. Also, Taylor and Mellor (1994) reported that after bluetongue epidemic in Turkey, the disease became endemic and probability of infection was much less in sheep up to 2 years old^[20]. However, in a seroprevalence study in the south-east of Iran the results showed that seroprevalence rates were decreased with the increasing of age in sheep^[15].

This study showed that the majority of animals in the north-east of Iran were infected with BTV. High correlation between abortion history and seropositivity, emphasized the economical importance of BTV in the sheep herds of the region.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This study was supported by a grant (No: 05/09/38671) from the Veterinary Council of Khorasan Razavi. The authors gratefully acknowledge the help of Rahim Mohammadi who proofread the manuscript.

Comments

Background

There is some information about the sero-epidemiological statues of bluetongue disease in various parts of the world and Iran including West Azerbaijan (northwest of Iran), Shiraz (central part of Iran) and Kerman (Southeast of Iran). But there is no information of that in northeast of Iran.

Research frontiers

In this study the authors have studied seroepidemiological statue of bluetongue disease in Khorasan Razavi province (northeast of Iran). In this study, the relationships between abortion, age, sex and seropositivity of bluetongue disease have been investigated.

Related reports

There are some similar seroepidemiological reports of bluetongue disease from other regions of Iran which is mentioned in the manuscript. The results of similar studies in other countries are mentioned too.

Innovations and breakthroughs

This is the first report that describes the seroprevalence rates of BTV in sheep and goat flocks in northeast of Iran.

Applications

The result of this study can be very important for other neighbor countries. According to high sero-positivity of bluetongue disease in the region, a good control program of bluetongue disease should be investigated.

Peer review

This is a good study with a significant number of serum samples compared with similar works in other regions. The results are interesting and suggested that bluetongue can be one of the causes of abortion in small ruminants of the region.

References

- [1] Breard E, Hamblin C, Hammoui S, Sailleau C, Dauphin G, Zientara S. The epidemiology and diagnosis of bluetongue with particular reference to Corsica. *Res Vet Sci* 2004; **77**: 1–8.
- [2] Kaufmann C, Steinmann IC, Hegglin D, Schaffner F, Mathis A. Spatio-temporal occurrence of *Culicoides* biting midges in the climatic regions of Switzerland, along with large scale species identification by MALDI-TOF mass spectrometry. *Parasit Vectors* 2012; **5**: 1–10.
- [3] Steukers L, Bertels G, Cay AB, Nauwynck HJ. Schmallenberg virus: emergence of an *Orthobunyavirus* among ruminant in Western Europe. *Vlaams Diergen Tijds* 2012; **81**: 119–127.
- [4] Coetzee P, Stokstad M, Venter EH, Myrmel M, Vuuren MV. Bluetongue: a historical and epidemiological perspective with the emphasis on South Africa. *Virol J* 2012; **9**(198): 1–9.
- [5] Baldet T, Baylis M, Belli G. *Vector of bluetongue virus*. Italy: 3th International symposium on bluetongue; 2003: 26–29.
- [6] Sperlova A, Zendulkova D. Bluetongue: a review. *Vet Med–Czech* 2011; **56**(9): 430–452.
- [7] Miller MM, Brown J, Cornish T, Johnson G, Mecham JO, Reeves WK, et al. Investigation of abluetongue disease epizootic caused by bluetongue virus serotype 17 in sheep in Wyoming. *J Am Vet Med Assoc* 2010; **237**(8): 955–959.
- [8] Mellor PS, Leake CJ. Climatic and geographical influences on arboviral infections and vectors. *OIE Sci Tech Rev* 2000; **19**: 41–54.
- [9] Zientara S, MacLachlan N, Calistri P, Sanchez-Vizcaino J, Savini G. Bluetongue vaccination in Europe. *Expert Rev Vaccin* 2010; **9**: 989–991.
- [10] Radostits OM, Gay CC, Blood DC, Hinchclif KW. *Veterinary medicine: a textbook of the diseases of cattle, sheep, pigs, goats and horses*. London: WB Saunders Company; 2007.
- [11] Venter EH, Gerdes T, Wright I, Terblanche J. An investigation into the possibility of bluetongue virus transmission by transfer of infected ovine embryos. *Onderstepoort J Vet Res* 2011; **78**(1): 1–7.
- [12] Blancou J, Pearson J. Bioterrorism and infectious animal diseases. *Comp Immunol Microbiol Infect Dis* 2003; **26**: 431–443.
- [13] Zendulkova D, Pospisil Z. Bluetongue and its possible impact on production and quality of food of animal origin (in Czech). Brno: Scientific Report for the Veterinary Committee for Food Safety; 2007, p. 18.
- [14] OIE WAoDot. Bluetongue and epizootic haemorrhagic disease. In: *OIE Terrestrial Manual*. Paris; 2009: 11–12.
- [15] Mozaffari A, Khalili M. The first survey for antibody against bluetongue virus in sheep flocks in southeast of Iran. *Asian Pac J Trop Biomed* 2012; S1808–S1810.
- [16] No`man V, Kargar Moakhar R, Shah Moradi AH, Heydari MR, Tabatabaei J. *A seroepidemiological survey for bluetongue virus antibody in sheep and goats of Isfahan province, Iran*. 1st ed. Isfahan, Iran: Isfahan Agricultural and Natural Resources Research Center; 2006.
- [17] Shoorijeh S, Ramin A, MacLachlan N, Osburn B, Tamadon A, Behzadi M, et al. High seroprevalence of bluetongue virus infection in sheep flocks in West Azerbaijan, Iran. *Comp Immunol Microbiol Infect Dis* 2010; **33**(3): 243–247.
- [18] Mohammadi A, Tanzifi P, Nemati Y. Seroepidemiology of bluetongue disease and risk factors in small ruminants of Shiraz suburb, Fars province, Iran. *Trop Biomed* 2012; **29**(4): 632–637.
- [19] Apiwatnakorn B, Bura P, Pasavorakul O. Serological study for bluetongue in Thailand. In: George TD, Kagao P, editors. *First Southeast Asia and pacific regional bluetongue symposium*. Kunming, P. R. China; 1996: 20–22.
- [20] Taylor WP, Mellor PS. Bluetongue virus distribution in Turkey 1978–1981. *Epidemiol Infect* 1994; **112**: 623–633.
- [21] Franz J, Conraths JM, Gethmann C, Thomas CM, Beer M, Hoffmann B. Epidemiology of bluetongue virus serotype 8. *Emerg Infect Dis* 2009; **15**: 23–30.
- [22] Niedbalski W. Monitoring studies of bluetongue disease in ruminants imported to Poland from EU. *Pol J Vet Sci* 2010; **13**(2): 333–336.
- [23] Erasmus BJ, Potgieter AC. The history of bluetongue. In: Mettrons P, Baylis M, Mellor PS, editors. *Bluetongue*. London: Elsevier, Academic Press; 2009: 7–21.
- [24] Toussaint JF, Sailleau C, Mast J, Houdart P, Czaplicki G, Demeestere L, et al. Bluetongue in Belgium. *Emerg Infect Dis* 2007; **13**: 614–616.
- [25] Caporale M, Wash R, Pini A, Savini G, Franchi P, Golder M, et al. Determinants of bluetongue virus virulence in murine models of disease. *J Virol* 2011; **85**(21): 11479–11489.