Full-Genome Characterization of a G8P[8] Rotavirus That Emerged among Children with Diarrhea in Croatia in 2006

Roberto Delogu,^a Alessandra Lo Presti,^b Franco Maria Ruggeri,^c Eleonora Cella,^b Marta Giovanetti,^b Massimo Ciccozzi,^b Suncanica Ljubin-Sternak,^d Suzana Bukovski-Simonoski,^e Amarela Lukic-Grlic,^f Giovanni Ianaro,^a Lucia Fiore^a

National Center for Immunobiologicals Research and Evaluation,^a Department of Infectious, Parasitic and Immune-Mediated Diseases,^b and Department of Veterinary Public Health and Food Safety,^c Istituto Superiore di Sanità, Rome, Italy; Croatian National Institute of Public Health and School of Medicine, University of Zagreb, Zagreb, Croatia^d; University Hospital for Infectious Diseases Dr. Fran Mihaljevic, Zagreb, Croatia^e; Zagreb University Children's Hospital and School of Medicine, University of Zagreb, Zagreb, Croatia^f

The whole genome of a G8P[8] rotavirus from the 2006 epidemic in Croatia was sequenced and showed a Wa-like genotype constellation. Its VP7 gene clustered with DS-1-like G8 African rotaviruses and a G8P[4] German strain. Remaining genes clustered with contemporary Belgian G1P[8] rotaviruses, suggesting reassortment between human G8 and G1P[8] rotaviruses in Croatia or other European countries.

Rotavirus A (RVA) causes acute gastroenteritis in humans and animals globally, killing 450,000 children per year, mainly in developing countries (1).

The RVA genome includes 11 double-stranded RNA segments encoding six viral proteins (VP1 to VP4, VP6, VP7) and six non-structural proteins (NSP1 to NSP6) (2). Rotavirus outer layer protein VP7 and spike protein VP4 are neutralization antigens, determining virus genotypes G (glycoprotein; VP7) and P (protease sensitive; VP4). At least 27 G and 37 P genotypes have been identified among human, animal, and avian RVAs (3–5).

Five genotypes (G1P[8], G2P[4], G3P[8], G4P[8], and G9P[8]) represent over 90% of global human RVAs (6), which may explain the efficacy of both the monovalent (P[8], G1) Rotarix and the pentavalent (P[8], G1 to G4) RotaTeq vaccines against rotavirus diarrhea worldwide (7).

Serotypes other than G1 to G4 and strains evolving by mutation or genomic reassortment between human and/or animal RVAs represent an emerging threat to children (8–10). Millions of people are exposed to animal rotaviruses worldwide, and several uncommon human genotypes (G6, G8, G12, P3[9], P5A[3]) closely resemble animal RVA strains (11).

Molecular epidemiology and viral typing can help monitor the emergence of novel RVAs, confirming the efficacy of current vaccines against unconventional genotypes.

Reassortment events can involve any double-stranded RNA segments, and G and P typing may be insufficient to investigate RVA origin and interspecies transmission. Genotyping based on sequencing of all 11 genome segments (3, 4) is very helpful, distinguishing 9 VP1 (R), 9 VP2 (C), 8 VP3 (M), 35 VP4 (P), 16 VP6 (I), 27 VP7 (G), 16 NSP1 (A), 9 NSP2 (N), 12 NSP3 (T), 14 NSP4 (E), and 11 NSP5 (H) genotypes.

Bovine-like G8 rotavirus was first reported in an Indonesian child (12), and further G8 RVA human cases occurred worldwide, including in industrialized countries (13–21). Using full-genome sequencing, G8P[8] African RVAs were shown to involve reassortment between at least four human, swine, and bovine strains (22).

This paper reports the whole-genome characterization of a G8P[8] human rotavirus strain that emerged in Croatia in 2006 (23).

Cases showed severe gastroenteritis and included children <5

years old from Croatian hospitals in 2005 and 2006. Stools were screened by use of the RotaMax assay, and RVA was characterized by reverse transcription-PCR (RT-PCR) (23). Agarose gel-purified amplicons were sequenced using RT-PCR primers (8, 18, 24, 25), assembling consensus sequences for all gene segments (18, 22, 26).

Twelve data sets were built and included all NCBI sequences (<http://www.ncbi.nlm.nih.gov/pubmed>) showing >89% similarity with query sequences (between 116 for VP2 and 127 for NSP4).

Sequences were aligned using ClustalX software (27) and edited by Bioedit software. The ModelTest v3.0 (28) with the hierarchical likelihood ratio test was used to select the best-fit models for sequence data analysis.

Bayesian phylogenetic trees were constructed for the VP7, VP4, and NSP4 data sets (29), using the GTR+I+G nucleotide substitution model for VP7 and the HKY+G model for VP4/NSP4.

Markov chain Monte Carlo searches were made (50×10^6 generations, tree sampling every 5,000 generation, and 10% burn-in fraction; clade statistical support followed >0.90 posterior probability).

Maximum-likelihood (ML) phylogenetic trees of NSP1 to NSP3, NSP5 and NSP6, VP1 to VP3, and VP6 were generated with the PAUP* v4.0 package (30), using the GTR+I+G model for the VP6, NSP1, and VP1 data sets; the HKY+G model for the NSP6/NSP5 data set; and the TrN+I+G for the NSP2, NSP3, VP2, and VP3 data sets. Statistical robustness and the reliability of the branching order within trees were confirmed by bootstrap analysis (1,000 replicates), with bootstrap values of >70% considered clade support.

Received 9 February 2013 Accepted 14 February 2013

Published ahead of print 20 February 2013

Address correspondence to Franco Maria Ruggeri, franco.ruggeri@iss.it.

R.D. and A.L.P. contributed equally to this article.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JCM.00396-13>.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.
[doi:10.1128/JCM.00396-13](https://doi.org/10.1128/JCM.00396-13)

FIG 2 Bayesian phylogenetic analysis of human rotavirus NSP4 nucleotide sequences. The tree was rooted by using the midpoint rooting method. Branch lengths were estimated with the best-fitting nucleotide substitution model according to a hierarchical likelihood ratio test. For significant statistical support, posterior probabilities of >0.90 are reported. Several clusters are replaced by a triangle for simplicity, and the size of the triangle is an indication of the number of sequences that it represents. The number of strains for each place of isolation is indicated in parentheses. The phylogenetic cluster containing CR2006 is boxed, and the CR2006 strain is in boldface type. For each strain, the following data are given: species of origin/place of isolation/year/strain name. The number of substitutions per site is indicated by the scale bar.

Initial sequencing of genes encoding VP4 and NSP4 for 10/31 G8P[8] RVAs showed nearly complete nucleotide sequence identity, suggesting circulation of a single G8P[8] strain in Croatia in 2005 and 2006. The consensus sequence of the VP7 gene of the Croatian strain from 2006 (CR2006) showed

>98% nucleotide sequence identity with the sequences of other human G8 strains whose sequences are in GenBank, particularly sequences from neighboring Slovenia recovered in 2006 and from Africa recovered in 2000 to 2004. CR2006 VP4 and NSP4 recalled human G3P[8] strains reported in Italy in

following data are given: species of origin/place of isolation/year/strain name. The number of substitutions per site is indicated by the scale bar. (A) Phylogenetic tree based on VP7 gene nucleotide sequence; (B) detail of the phylogenetic relationship between RV strains of the G8 genotype; (C) phylogenetic tree based on VP4 gene nucleotide sequence. Kor, South Korea; Jap, Japan; Switz, Switzerland; Slo, Slovenia; Ger, Germany; Philip, The Philippines; Bangl, Bangladesh; Bel, Belgium; GBR, Great Britain; Bo, bovine; Si, simian; Ov, ovine; Po, porcine; La, lapine (rabbit).

2005 (31). To investigate the Croatian G8P[8] origin, full-genome sequencing was performed, using the RotaC tool (<http://rotac.regatools.be>) (32). CR2006 presented a genotype G8P[8]-I1-R1-C1-M1-A1-N1-T1-E1-H1.

Bayesian analysis identified CR2006 VP7 in a statistically supported cluster with African strains (Malawi, Tunisia, Cameroon, Ethiopia) and European strains (Slovenia [SI-885/06], Germany [GER1H-09]) (Fig. 1A). In the VP4 and NSP4 trees (Fig. 1C and 2), CR2006 also clustered with European and African RVAs, including human (Hu)/Slovenia/2006/G8P[8]/SI-885/06, Hu/Italy/2005/G3P[8], Hu/Ethiopia/2004/G8P[8]/ARN, and Hu/Tunisia/2000-2004/G8P[8]/ARN.

CR2006 clustered with the Wa-like genotype constellation strains: Hu/Belgium/2005-2009/G1P[8], Hu/Germany/G3G12P[8]/GER126-08, and other global RVAs in NSP1 to NSP6, VP1 to VP3, and VP6 (see Fig. S1 to S9 in the supplemental material).

G8P[8] rotavirus circulated abnormally in Croatia in 2006, reaching 15.8% of children hospitalized with diarrhea (23). Full-genome sequencing of these viruses was performed to investigate links between their emergence and genome characters.

The G8 rotavirus genotype, normally combined with the P[1], P[5], P[11], or P[21] type (11, 33, 34), is typical of bovine strains but has also been sporadically detected in humans since 1978 (35) in association with different P types (13, 16–18, 36–38). Human G8 strains normally exhibit P[4] or P[6] both in developed countries and in Africa, where they are particularly prevalent (15–17, 26). Recently, full-genome sequencing of a human G8P[8] rotavirus in the Democratic Republic of Congo (DRC) in 2003 (18, 22) suggested that this strain originated from reassortment between a bovine G8P[6] ancestor with three or more different rotaviruses, including human Wa- and DS-1-like strains. This strain exhibited a DS-1-like genotype constellation.

Croatian CR2006 G8P[8], however, presented the Wa-like constellation (3) G8-P[8]-I1-R1-C1-M1-A1-N1-T1-E1-H1, with all genes except the VP7 gene, which is related to African and Slovenian human G8 strains, showing a close evolutionary relationship to the prototype G1P[8] human strain Wa. Rotaviruses are usually species specific, and gene reassortment with human strains appears to be necessary for zoonotic adaptation (26, 39–41). Characterization of epidemic CR2006 further supports this view. As VP7 plays a major role in the immune response (42, 43), CR2006 emergence and epidemic spread may have involved reassortment with common rotaviruses preexisting in Croatia.

CR2006 was more frequent in younger children (23) than any other rotavirus genotypes cocirculating in Croatia in 2006. This could be fortuitous or suggest that maternal antibodies protected younger children from CR2006 less than they did from viruses of other serotypes. Maternal immunity fades quickly (44, 45), and it is conceivable that in 2006 older Croatian children were similarly unprotected and hence susceptible to G8 or any other common rotavirus genotypes.

Phylogenetic analyses revealed that, except for the VP7 gene, CR2006 was more closely related to humans G1P[8] strains identified between 2005 and 2009 in Belgium. Therefore, the origin of CR2006 likely involved reassortment of a Wa-like rotavirus circulating in one or more countries of Europe, including Belgium and possibly Croatia, with a cocirculating G8 virus. In 2006, a similar G8P[8] strain was reported and partially sequenced in Slovenia (20), whereas no G8P[8] rotaviruses were detected in other European countries. It is possible that the reassortment event leading to

CR2006 may have occurred in or near Croatia. The low correlation between VP7 of CR2006 and VP7 of G8 DS-1-like rotaviruses characterized in DRC (18) indicates the independent evolution of these G8 reassortants. Conversely, CR2006 shared a similar gene 9 with human G8 strains from northern/central Africa in the early 2000s, suggesting that the G8 gene followed immigration routes from Africa. A closely similar VP7 was found in a G8P[4] human strain identified in Germany 2 years later (21), suggesting that the CR2006 G8 gene participated in additional reassortments with European DS-1-like rotaviruses. The German strain was strictly correlated with the DRC G8 rotaviruses reported earlier, except for VP4 (18), and may represent a further evolution of G8P[8] rotaviruses imported from Africa. The VP7 of all European G8 strains differed markedly from the VP7 of bovine G8 rotaviruses, contrary to a recent human-bovine reassortment event within Europe.

Rotavirus genetic evolution is certainly influenced by immune pressure (46) that may favor the spread of G8 and other uncommon rotavirus G types (19, 20, 47), as occurred for G9 strains that originated from swine (48, 49). International transmission of genotypes is facilitated by globalization and travel, urging worldwide rotavirus strain surveillance to anticipate the emergence of novel strains and animal/human reassortants and monitor the effectiveness of current human rotavirus vaccines.

Nucleotide sequence accession numbers. Sequences were deposited in the GenBank database (accession numbers [JQ988894](https://www.ncbi.nlm.nih.gov/nuclot/JQ988894) to [JQ988904](https://www.ncbi.nlm.nih.gov/nuclot/JQ988904)).

ACKNOWLEDGMENTS

This study was sponsored by the Ministry of Health, Italy (CCM Epidemiologia Molecolare di Rotavirus in Età Pediatrica in Italia; Creazione di una Rete di Sorveglianza per Monitorare la Diffusione e l'Evoluzione di Genotipi Virali [to L.F.] and Strategic Program: Preparedness and Response to Emerging Zoonoses and Exotic Viral Infection through an Integrated Medical and Veterinary Approach [to F.M.R.]), ISS/NIH (Molecular and Antigenic Evolution of Rotavirus Strains of Human and Animal Origin to F.M.R.), and EuroRotaNet (<http://www.eurorota.net>).

REFERENCES

1. Tate JE, Burton AH, Boschi-Pinto C, Steele AD, Duque J, Parashar UD. 2012. 2008 estimate of worldwide rotavirus-associated mortality in children younger than 5 years before the introduction of universal rotavirus vaccination programmes: a systematic review and meta-analysis. *Lancet Infect. Dis.* 12:136–141.
2. Estes MK, Kapikian AZ. 2007. Rotaviruses, p 1917–1974. In Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE (ed), *Fields virology*, vol 2. Kluwer/Lippincott, Williams & Wilkins, Philadelphia, PA.
3. Matthijnssens J, Ciarlet M, Rahman M, Attoui H, Banyai K, Estes MK, Gentsch JR, Iturriza-Gomara M, Kirkwood CD, Martella V, Mertens PP, Nakagomi O, Patton JT, Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K, Desselberger U, Van Ranst M. 2008. Recommendations for the classification of group A rotaviruses using all 11 genomic RNA segments. *Arch. Virol.* 153:1621–1629.
4. Matthijnssens J, Ciarlet M, McDonald SM, Attoui H, Banyai K, Brister JR, Buesa J, Esona MD, Estes MK, Gentsch JR, Iturriza-Gomara M, John R, Kirkwood CD, Martella V, Mertens PP, Nakagomi O, Parreno V, Rahman M, Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K, Patton JT, Desselberger U, Van Ranst M. 2011. Uniformity of rotavirus strain nomenclature proposed by the Rotavirus Classification Working Group (RCWG). *Arch. Virol.* 156:1397–1413.
5. Trojnar E, Sachsenroder J, Twardziok S, Reetz J, Otto PH, John R. 2013. Identification of an avian group A rotavirus containing a novel VP4 gene with a close relationship to those of mammalian rotaviruses. *J. Gen. Virol.* 94:136–142.

6. Santos N, Hoshino Y. 2005. Global distribution of rotavirus serotypes/genotypes and its implication for the development and implementation of an effective rotavirus vaccine. *Rev. Med. Virol.* 15:29–56.
7. Jiang V, Jiang B, Tate J, Parashar UD, Patel MM. 2010. Performance of rotavirus vaccines in developed and developing countries. *Hum. Vaccin.* 6:532–542.
8. Iturriza-Gomara M, Isherwood B, Desselberger U, Gray J. 2001. Reassortment in vivo: driving force for diversity of human rotavirus strains isolated in the United Kingdom between 1995 and 1999. *J. Virol.* 75:3696–3705.
9. Maunula L, Von Bonsdorff CH. 2002. Frequent reassortments may explain the genetic heterogeneity of rotaviruses: analysis of Finnish rotavirus strains. *J. Virol.* 76:11793–11800.
10. Gentsch JR, Laird AR, Bielfelt B, Griffin DD, Banyai K, Ramachandran M, Jain V, Cunliffe NA, Nakagomi O, Kirkwood CD, Fischer TK, Parashar UD, Bresee JS, Jiang B, Glass RI. 2005. Serotype diversity and reassortment between human and animal rotavirus strains: implications for rotavirus vaccine programs. *J. Infect. Dis.* 192(Suppl 1):S146–S159.
11. Martella V, Banyai K, Matthijnssens J, Buonavoglia C, Ciarlet M. 2010. Zoonotic aspects of rotaviruses. *Vet. Microbiol.* 140:246–255.
12. Hasegawa A, Inouye S, Matsuno S, Yamaoka K, Eko R, Suharyono W. 1984. Isolation of human rotaviruses with a distinct RNA electrophoretic pattern from Indonesia. *Microbiol. Immunol.* 28:719–722.
13. Holmes JL, Kirkwood CD, Gerna G, Clemens JD, Rao MR, Naficy AB, Abu-Elyazed R, Savarino SJ, Glass RI, Gentsch JR. 1999. Characterization of unusual G8 rotavirus strains isolated from Egyptian children. *Arch. Virol.* 144:1381–1396.
14. Santos N, Lima RC, Pereira CF, Gouvea V. 1998. Detection of rotavirus types G8 and G10 among Brazilian children with diarrhea. *J. Clin. Microbiol.* 36:2727–2729.
15. Steele AD, Parker SP, Peenze I, Pager CT, Taylor MB, Cubitt WD. 1999. Comparative studies of human rotavirus serotype G8 strains recovered in South Africa and the United Kingdom. *J. Gen. Virol.* 80(Pt 11):3029–3034.
16. Cunliffe NA, Gentsch JR, Kirkwood CD, Gondwe JS, Dove W, Nakagomi O, Nakagomi T, Hoshino Y, Bresee JS, Glass RI, Molyneux ME, Hart CA. 2000. Molecular and serologic characterization of novel serotype G8 human rotavirus strains detected in Blantyre, Malawi. *Virology* 274:309–320.
17. Adah MI, Nagashima S, Wakuda M, Taniguchi K. 2003. Close relationship between G8-serotype bovine and human rotaviruses isolated in Nigeria. *J. Clin. Microbiol.* 41:3945–3950.
18. Matthijnssens J, Rahman M, Yang X, Delbeke T, Arijis I, Kabue JP, Muyembe JJ, Van Ranst M. 2006. G8 rotavirus strains isolated in the Democratic Republic of Congo belong to the DS-1-like genogroup. *J. Clin. Microbiol.* 44:1801–1809.
19. Montenegro FM, Correia JB, Rodrigues Falbo A, Dove W, Nakagomi T, Nakagomi O, Cuevas LE, Cunliffe NA, Hart CA. 2007. Anticipating rotavirus vaccines in Brazil: detection and molecular characterization of emerging rotavirus serotypes G8 and G9 among children with diarrhoea in Recife, Brazil. *J. Med. Virol.* 79:335–340.
20. Steyer A, Poljsak-Prijatelj M, Bufon TL, Marcun-Varda N, Marin J. 2007. Rotavirus genotypes in Slovenia: unexpected detection of G8P[8] and G12P[8] genotypes. *J. Med. Virol.* 79:626–632.
21. Pietsch C, Petersen L, Patzer U, Liebert UG. 2009. Molecular characteristics of German G8P[4] rotavirus strain GER1H-09 suggest that a genotyping and subclassification update is required for G8. *J. Clin. Microbiol.* 47:3569–3576.
22. Matthijnssens J, Ciarlet M, Heiman E, Arijis I, Delbeke T, McDonald SM, Palombo EA, Iturriza-Gomara M, Maes P, Patton JT, Rahman M, Van Ranst M. 2008. Full genome-based classification of rotaviruses reveals a common origin between human Wa-like and porcine rotavirus strains and human DS-1-like and bovine rotavirus strains. *J. Virol.* 82:3204–3219.
23. Tcheremenskaia O, Marucci G, De Petris S, Ruggeri FM, Dovecar D, Sternak SL, Matyasova I, Dhimolea MK, Mladenova Z, Fiore L. 2007. Molecular epidemiology of rotavirus in central and southeastern Europe. *J. Clin. Microbiol.* 45:2197–2204.
24. Das BK, Gentsch JR, Cicirello HG, Woods PA, Gupta A, Ramachandran M, Kumar R, Bhan MK, Glass RI. 1994. Characterization of rotavirus strains from newborns in New Delhi, India. *J. Clin. Microbiol.* 32:1820–1822.
25. Gentsch JR, Glass RI, Woods P, Gouvea V, Gorziglia M, Flores J, Das BK, Bhan MK. 1992. Identification of group A rotavirus gene 4 types by polymerase chain reaction. *J. Clin. Microbiol.* 30:1365–1373.
26. Esona MD, Geyer A, Page N, Trabelsi A, Fodha I, Aminu M, Agbaya VA, Tsion B, Kerin TK, Armah GE, Steele AD, Glass RI, Gentsch JR. 2009. Genomic characterization of human rotavirus G8 strains from the African rotavirus network: relationship to animal rotaviruses. *J. Med. Virol.* 81:937–951.
27. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673–4680.
28. Posada D, Buckley TR. 2004. Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Syst. Biol.* 53:793–808.
29. Zehender G, Ebranati E, Lai A, Santoro MM, Alteri C, Giuliani M, Palamara G, Perno CF, Galli M, Lo Presti A, Ciccozzi M. 2010. Population dynamics of HIV-1 subtype B in a cohort of men-having-sex-with-men in Rome, Italy. *J. Acquir. Immune Defic. Syndr.* 55:156–160.
30. Ciccozzi M, Babakir-Mina M, Lo Presti A, Marcuccilli F, Perno CF, Ciotti M. 2011. Phylogenesis and clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *Open Virol. J.* 5:22–26.
31. De Grazia S, Martella V, Colomba C, Cascio A, Arista S, Giammanco GM. 2009. Genetic characterization of G3 rotaviruses detected in Italian children in the years 1993–2005. *J. Med. Virol.* 81:2089–2095.
32. Maes P, Matthijnssens J, Rahman M, Van Ranst M. 2009. RotaC: a web-based tool for the complete genome classification of group A rotaviruses. *BMC Microbiol.* 9:238. doi:10.1186/1471-2180-9-238.
33. Dhama K, Chauhan RS, Mahendran M, Malik SV. 2009. Rotavirus diarrhea in bovines and other domestic animals. *Vet. Res. Commun.* 33:1–23.
34. Monini M, Cappuccini F, Battista P, Falcone E, Lavazza A, Ruggeri FM. 2008. Molecular characterization of bovine rotavirus strains circulating in northern Italy, 2003–2005. *Vet. Microbiol.* 129:384–389.
35. Matsuno S, Hasegawa A, Mukoyama A, Inouye S. 1985. A candidate for a new serotype of human rotavirus. *J. Virol.* 54:623–624.
36. Palombo EA, Clark R, Bishop RF. 2000. Characterisation of a “European-like” serotype G8 human rotavirus isolated in Australia. *J. Med. Virol.* 60:56–62.
37. Armah GE, Pager CT, Asmah RH, Anto FR, Oduro AR, Binka F, Steele D. 2001. Prevalence of unusual human rotavirus strains in Ghanaian children. *J. Med. Virol.* 63:67–71.
38. Volotao EM, Soares CC, Maranhao AG, Rocha LN, Hoshino Y, Santos N. 2006. Rotavirus surveillance in the city of Rio de Janeiro-Brazil during 2000–2004: detection of unusual strains with G8P[4] or G10P[9] specificities. *J. Med. Virol.* 78:263–272.
39. Rahman M, Matthijnssens J, Yang X, Delbeke T, Arijis I, Taniguchi K, Iturriza-Gomara M, Iftekhharuddin N, Azim T, Van Ranst M. 2007. Evolutionary history and global spread of the emerging g12 human rotaviruses. *J. Virol.* 81:2382–2390.
40. Steyer A, Poljsak-Prijatelj M, Barlic-Maganja D, Marin J. 2008. Human, porcine and bovine rotaviruses in Slovenia: evidence of interspecies transmission and genome reassortment. *J. Gen. Virol.* 89:1690–1698.
41. Browning GF, Snodgrass DR, Nakagomi O, Kaga E, Sarasini A, Gerna G. 1992. Human and bovine serotype G8 rotaviruses may be derived by reassortment. *Arch. Virol.* 125:121–128.
42. Ward R. 2009. Mechanisms of protection against rotavirus infection and disease. *Pediatr. Infect. Dis. J.* 28:S57–S59.
43. Offit PA, Shaw RD, Greenberg HB. 1986. Passive protection against rotavirus-induced diarrhea by monoclonal antibodies to surface proteins vp3 and vp7. *J. Virol.* 58:700–703.
44. Dennehy PH, Cortese MM, Begue RE, Jaeger JL, Roberts NE, Zhang R, Rhodes P, Gentsch J, Ward R, Bernstein DI, Vitek C, Bresee JS, Staat MA. 2006. A case-control study to determine risk factors for hospitalization for rotavirus gastroenteritis in U.S. children. *Pediatr. Infect. Dis. J.* 25:1123–1131.
45. Clemens J, Rao M, Ahmed F, Ward R, Huda S, Chakraborty J, Yunus M, Khan MR, Ali M, Kay B, van Loon F, Sack D. 1993. Breast-feeding and the risk of life-threatening rotavirus diarrhea: prevention or postponement? *Pediatrics* 92:680–685.
46. Kirkwood CD. 2010. Genetic and antigenic diversity of human rotaviruses: potential impact on vaccination programs. *J. Infect. Dis.* 202(Suppl):S43–S48.

47. Esona MD, Geyer A, Banyai K, Page N, Aminu M, Armah GE, Hull J, Steele DA, Glass RI, Gentsch JR. 2009. Novel human rotavirus genotype G5P[7] from child with diarrhea, Cameroon. *Emerg. Infect. Dis.* 15:83–86.
48. Ramachandran M, Gentsch JR, Parashar UD, Jin S, Woods PA, Holmes JL, Kirkwood CD, Bishop RF, Greenberg HB, Urasawa S, Gerna G, Coulson BS, Taniguchi K, Bresee JS, Glass RI. 1998. Detection and characterization of novel rotavirus strains in the United States. *J. Clin. Microbiol.* 36:3223–3229.
49. Santos N, Lima RC, Nozawa CM, Linhares RE, Gouvea V. 1999. Detection of porcine rotavirus type G9 and of a mixture of types G1 and G5 associated with Wa-like VP4 specificity: evidence for natural human-porcine genetic reassortment. *J. Clin. Microbiol.* 37:2734–2736.