Complaints of micturition, defecation and sexual function in cauda equina syndrome due to lumbar disk herniation: a systematic review

N. S. Korse · W. C. H. Jacobs · H. W. Elzevier ·
C. L. A. M. Vleggeert-Lankamp

Received: 23 May 2012 / Revised: 29 September 2012 / Accepted: 20 November 2012 / Published online: 13 December 2012
© Springer-Verlag Berlin Heidelberg 2012

Abstract

Purpose Cauda equina syndrome (CES) is a rare complication of lumbar disk herniation. Although micturition, defecation and/or sexual function are by definition affected, little seems to be known about long-term outcome. Aim of this study is to review current literature on outcome of micturition, defecation and sexual function in CES due to lumbar disk herniation.

Methods A literature search was done in Pubmed, Embase and Web of Science using a sensitive search string combination. Studies were selected by predefined selection criteria and risk of bias was assessed using a Cochrane checklist adjusted for this purpose.

Results Fifteen studies were included. Risk of bias varied with six studies showing low risk. Mean minimal follow-up time was 17.0 months (range 3–24 months). All studies evaluated micturition and reported dysfunction at follow-up in 42.5 % (range 13.3–90.0 %). Defecation and sexual function were evaluated in eight and nine studies, respectively, and reported to be 49.6 (range 10.5–90.0 %) and 44.3 % (range 10.0–76.6 %), respectively. Only two studies assessed sexual function in all patients at follow-up.

Conclusion This review offers an insight into the extent of micturition dysfunction, defecation dysfunction and sexual dysfunction (SD) in CES after decompression. Our

findings show that dysfunction is extremely common, even at long-term follow-up. A condition as invalidating as CES requires proper patient information and the outcomes presented here may help in providing those data. Bias in included studies, lack of universal definitions and incomplete follow-up results qualify these data as the best we momentarily have, but still subject to improvement. Since SD seems to be severely underreported, we recommend further research to explore the extent of this problem, as well as the use of questionnaires in future clinical (prospective) studies to accomplish a more patient-based approach.

Keywords Cauda equina syndrome · Lumbar disk herniation · Micturition · Defecation · Sexual dysfunction · Pelvic floor

Introduction

Cauda equina syndrome (CES) is an uncommon neurologic condition caused by compression of the cauda equina. The first notion of CES in English literature was taken by Mixter and Barr [1]. They described a spectrum of neurological and urological complaints in patients with a lumbar hernia and attributed this to a severe compression of the cauda equina, urging for emergency decompression as opposed to a more expectative treatment in uncomplicated hernia nuclei pulposi (HNP). Although CES can be instigated by any pathological process compressing the cauda equina, the lumbar disk is the most prevalent structure: 45 % of described CES cases in literature are attributed to the disk [2]. In patients operated on because of HNP, the incidence of CES is 1–10 % [3, 4]. Because CES is an indication for surgery in HNP, the incidence of CES among general HNP patients is probably lower [5].

N. S. Korse (✉) · W. C. H. Jacobs ·
C. L. A. M. Vleggeert-Lankamp
Department of Neurosurgery, Leiden University Medical Center
(LUMC), Postzone J11-R-83, Postbus 9600, RC 2300 Leiden,
The Netherlands
e-mail: n.s.korse@lumc.nl

H. W. Elzevier
Department of Urology, Leiden University Medical Center
(LUMC), Leiden, The Netherlands

Exact definitions of CES have always been a topic of ambiguity, although most authors agree that micturition dysfunction should be present [6–9]. After reviewing more than a hundred articles, Fraser et al. [2] stated that at least one or more of the following should be present to diagnose CES: (1) micturition and/or defecation dysfunction, (2) reduced sensation in the saddle area, and (3) sexual dysfunction (SD) with possible neurologic deficit in the lower limb (motor/sensory loss, reflex change). Even though the relevance of micturition dysfunction, defecation dysfunction and SD is clear from this definition, little seems to be known about the long-term outcome of these functions. This poses a problem for the clinician who needs to provide his patient with an accurate prognosis and thereby obtain true informed consent for therapy.

Solid information about long-term recovery of micturition, defecation and sexual function in CES is not abundant. One factor is probably the shame that revolves around discussing problems of micturition, defecation and sexual function. Secondly, the extensive focus in literature on the timing of decompression, pushes the actual hard figures on long-term outcome more to the background. For the surgeon confronted with a CES patient, however, questions do not arise around the best timing of surgery, since it is generally accepted that decompression should be done as soon as possible. Questions do arise around the long-term prognosis. More specifically, the prognoses of micturition, defecation and sexual function, which are so inevitably affected in CES, need to be clarified. Individual studies at best provide the clinician with uncertain estimates based on few patients.

The aim of this study, therefore, is to review current literature on the outcome of micturition, defecation and sexual function in CES due to HNP.

Materials and methods

Data searches and study selection

In January 2012, the electronic databases Pubmed, Embase and Web of Science were searched using the search strategies as shown in Fig. 1.

Two of the authors (N.S.K., W.C.H.J.) separately screened the articles by title, abstract or by full article, when necessary, to select the studies that met the pre-defined selection criteria. Selection criteria were stated as followed:

- the article was published in English, Dutch, French, German or Spanish;
- the study included patients diagnosed with CES due to HNP;

PUBMED (149 results)
 ("lumbar disk herniation" OR "lumbar disc herniation" OR "lumbar herniated disk" OR "lumbar herniated disc" OR "lumbar disectomy" OR "lumbar disectomy" OR "lumbar disk surgery" OR "lumbar disc surgery" OR "lumbar disc" OR "lumbar disk" OR "lumbar disk prolapse" OR "lumbar disc prolapse" OR "prolapsed intervertebral disk" OR "prolapsed intervertebral disc" OR "discogenic compression") AND ("cauda equina syndrome" OR "cauda equina compression" OR "cauda syndrome" OR "cauda equina"[Majr])

WEB OF SCIENCE (155 results)
 Same strategy as used in Pubmed, adjusted for this database.

EMBASE (316 results)
 Same strategy as used in Pubmed, adjusted for this database.

Fig. 1 Search strategy (as performed 22.01.2012)

- the study reported the following disease characteristics: HNP diagnosed by means of MRI, CT or X contrast RSG;
- the study reported the following patient characteristics: function of micturition, function of defecation and/or sexual function at base line (e.g., pre-operative) and at follow-up (e.g., post-operative), with a follow-up period of at least 2 weeks;
- the study included/evaluated a (primary) treatment, excluding reoperations;
- the study was a case study (with a minimum of ten patients), cohort study or randomized controlled trial. Systematic reviews or meta-analysis were not included;
- the article was published fully in a peer-reviewed journal.

Any discrepancy in selection between the two reviewers was resolved in open discussion. Reference screening and citation tracking were performed on the identified articles and as a final check, the meta-analyses found in the first search were studied to make sure no relevant articles were missed.

Quality assessment

The methodological quality of these studies was assessed by two independent reviewers (H.W.E., C.L.A.M.V.L.), using an adjusted version of the checklist for cohort studies of the Dutch Cochrane Center [10]. When there was no consensus about the assessment, a third reviewer (N.S.K.) was consulted.

The items reviewed in the assessment were: definition of patient group; selection bias; definition of type of surgery (laminectomy/partial excision of vertebral arches); method for assessing outcome (urodynamic/grading/descriptive) and loss to follow-up. A point was given for each of the following items: well-defined patient group [information was given about patients age (mean or range), and about presence/absence of saddle anesthesia, radicular

complaints and micturition dysfunction at presentation], absence of selection bias and absence of attrition bias (attrition bias: loss to follow-up >20 %). Studies were divided into different groups by risk of bias, with the maximum of three points indicating the lowest risk of bias.

Statistical analyses

Statistical analyses were done in STATA version 11.0 [11]. Mean prevalence and range of micturition, defecation and sexual function were calculated. To evaluate the effect of risk of bias and the time of follow-up, as well as pre-operative proportion of micturition dysfunction, defecation dysfunction and saddle hypo/anesthesia on the amount of dysfunction of micturition, defecation and sexual function at follow-up, a meta-regression was performed. Risk of bias was dichotomized in low risk of bias (three points) and medium to high risk of bias (two points or one point). Follow-up was dichotomized in 12 months or less and more than 12 months. Backward elimination was used by repeating the test after removing the least significant factors. A Monte-Carlo permutation test was performed to correct for multiple testing and a probability value of <0.05 was considered statistically significant [12]. Sensitivity analyses were performed for influence of shorter follow-up and low risk of bias studies.

For studies that also included patients without pre-operative dysfunction, we corrected post-operative prevalence of dysfunction for pre-operative prevalence of dysfunction, in order to make comparison between studies fair.

Results

Characteristics of included studies and risk of bias

Through our search, 620 articles were identified, of which 527 original articles were left after removing duplicates (Fig. 2). Selection procedure and subsequent citation tracking resulted in 16 reports on 15 different studies that met all criteria [5, 13–27]. One study was reported in two publications [19, 20]. Reasons to exclude articles were among others small patient numbers [28, 29], inclusion of HNP patients instead of CES patients exclusively [30–37], inclusion of hemi CES instead of total CES [38], no imaging done [3, 4, 39–41], no evaluation of a primary treatment [42–44], follow-up of at least 2 weeks not guaranteed [45–49] and no adequate report on post-operative functions [50].

Fifteen studies, published from 1956 to 2011, were included with 464 patients (Table 1). All were retrospective studies with relatively small sample sizes (range 14–54) and a rather young patient population (mean

Fig. 2 Flow chart of literature search

43.5 years). The mean minimal time of follow-up was 17.0 months after surgery.

Except for three patients who were treated conservatively [5, 17, 24], all patients underwent surgery. Four authors did not give exact information about type of surgery [5, 22–24]. Outcome was descriptive in eight studies [13, 14, 16, 17, 21, 22, 26, 27], graded in six studies [5, 15, 18, 20, 23, 25] and both urodynamic and descriptive in one study [24].

After consultation of the third reviewer, consensus about risk of bias score was achieved in all cases. Six studies were assessed to have a low risk of bias (Table 2). Two studies showed a high risk with both selection bias and either a poor-defined patient group or attrition bias. The remaining seven studies had an intermediate risk of bias.

Clinical presentation of CES

Included studies varied in definition of CES and thus in patient characteristics at presentation (Table 3). Micturition dysfunction was regarded as an important element of CES by most authors and all articles assessed micturition at presentation. Exact definition of micturition dysfunction varied across studies. Mean prevalence of micturition dysfunction at presentation was 88.9 %. Eight authors reviewed function of defecation at presentation, resulting in a prevalence of dysfunction in 47.1 %. Most authors agreed that saddle anesthesia—or to a lesser extent, saddle hypoesthesia—is one of the hallmarks of the classic presentation of CES. In included studies, saddle anesthesia/hypoesthesia was found in 80.8 % of patients at presentation. Pre-operative SD was only assessed in a total of three patients, in three studies [5, 16, 22]. Sciatica, whether unilateral or bilateral, was present in 95.5 % of patients.

Table 1 Characteristics of included studies

Study	N (464)	Mean age in years (range)	Men in %	Mode of surgery (number of patients)	Minimal time of follow-up in months
Jennett [5]	25	NR (20–72)	60.0	NR (24), no surgery (1)	12
Spännare [27]	30	NR (20–70)	46.7	Complete laminectomy with discectomy (17), partial excision of two adjacent vertebral arches bilateral (7), or unilateral (6)	24
O’Laoire [18]	29	40.6 (23–69)	58.6	Laminectomy with discectomy (29)	12
Hellström [24]	17	43 (33–63)	58.8	NR (16), no surgery (1)	24
Kostuik [17]	31	40 (23–61)	54.8	Wide bilateral laminectomy (30), no surgery (1)	24
Gleave [25]	33	40.6 (23–67)	NR	Laminectomy (29), fenestration (4)	24
Shapiro [13]	14	43 (22–67)	64.3	Laminectomy with discectomy (14)	6
Kennedy [26]	19	55 (31–76)	63.2	Bilateral laminectomy (19)	22
Buchner [19]	22	42 (22–67)	59.1	Bilateral hemilaminectomy (12), bilateral laminectomy (9)	12
Shapiro [14]	44	44 (22–67)	61.4	Laminectomy with discectomy (44) of which 1 unilateral microdiscectomy approach	12
Hussain [21]	20	45 (33–67)	50.0	Bilateral laminectomy with discectomy (20)	10
McCarthy [22]	54	41 (24–67)	54.8	NR	25
Qureshi [16]	33	43 (30–79)	57.6	Interlaminar discectomy (15) of which 6 microdiscectomy approach, laminectomy with discectomy (12), hemilaminectomy (6)	3
Crocker [23]	43	NR	NR	NR	24
Dhatt [15]	50	48 (25–85)	66.0	Wide laminectomy with discectomy (50)	12
Mean (corrected)	–	43.5	55.7	–	17.0

NR not reported

Table 2 Risk of bias assessment

	Study	Score on risk of bias scale	Well-defined patient group	Absence of selection bias	Absence of attrition bias
	Jennett [5]	***	*	*	*
	Spännare [27]	***	*	*	*
	O’Laoire [18]	***	*	*	*
	Hellström [24]	**	*	—	*
	Kostuik [17]	**	*	—	*
	Gleave [25]	***	*	*	*
	Shapiro [13]	**	—	*	*
	Kennedy [26]	**	*	—	*
	Buchner [19]	***	*	*	*
	Shapiro [14]	*	*	—	#
	Hussain [21]	**	*	*	#
	McCarthy [22]	**	*	*	—
# No information was provided about exact loss to follow-up	Qureshi [16]	**	*	*	—
Asterisks represent the number of points gained	Crocker [23]	*	—	—	*
	Dhatt [15]	***	*	*	*

Outcome: micturition

Outcome of micturition was evaluated in a total of 409 patients (Table 4). At an average mean minimal follow-up time of 17.0 months after surgery, mean prevalence of micturition dysfunction is 42.5 % (range 13.3–90.0). The corrected mean prevalence of micturition dysfunction is 45.1 % (range 13.3–90.0).

Micturition dysfunction is often described as incontinence [5, 13, 14, 16, 18, 19, 25] or by the presence of urologic complaints [21, 22, 24] or disturbances [15, 23, 27]. Also mentioned are the absence of a ‘normal voiding pattern’ [17] and ‘any residual deficit that was regarded as a physical or psychological impairment’ [26]. In one study [24], in addition to clinical assessment, extensive urodynamics tests were performed to measure micturition outcome, but the author concluded that the complaints patients gave were not always consistent with these functional outcomes.

Variation of prevalence across studies is large. The highest rates of dysfunction were found by Hussain, Dhatt (both 90 %) and Jennett (76.9 %); these three studies all showed a low to intermediate risk of bias. Lowest prevalences were found by Kostuik and Gleave (13.3 and 21.2 %, respectively), which also demonstrated a low to intermediate risk of bias at 24 months post-operative.

Outcome: defecation

Outcome of defecation function was evaluated in a total of 238 patients [5, 14–16, 18, 22, 24, 26]. At an average mean minimal follow-up time of 17.0 months after surgery,

mean prevalence of defecation dysfunction is 49.6 %. Prevalences range from 10.5 [26] to 90.0 % [15]. Interestingly, the studies that evaluated post-operative function of defecation are not the same studies that evaluated this function pre-operatively.

Different definitions of defecation dysfunction are used: ‘a patulous anal sphincter leading usually to fecal incontinence’ or ‘constipation with defective anal sensation’ [5], complaints of ‘bowel disturbance’ [15, 22], ‘abnormal sphincter tone’ [24], being ‘grossly incontinent of stool’ [14], no control of flatus and being ‘occasional incontinence of feces’ [18], ‘poor fecal continence and no control of flatulence’ [26] and a state different from ‘never or rarely leaked from bowels’ [16].

Outcome: sexual function

Outcome of sexual function was reviewed in a total of 201 patients. Only two studies [15, 17] reviewed sexual function in every patient seen at follow-up. Seven studies [5, 13, 14, 18, 22, 24, 26] recorded sexual function in a selection of included patients, more often in men than in women. At an average mean minimal follow-up time of 17.0 months after surgery, mean prevalence of SD is 44.3 % (range 10.0–76.6).

Many different varieties of SD are mentioned, even within the same study: impotence [5, 13, 14, 17, 22, 24], decreased potency [14, 22, 24], more difficulties to obtain orgasm [13], less intense orgasm [13, 14], anorgasm [13, 22], decreased [17, 22] or absent [14] penile/vaginal sensation, incontinence during intercourse [14, 17], dyspareunia [22], absent bulbocavernosus reflex [26], and the very general terms SD [15] and abnormal intercourse [18].

Table 3 Characteristics of patients (at presentation)

Study	Score on risk of bias scale	Micturition dysfunction in % (N)	Defecation dysfunction in % (N)	Saddle hypo/anesthesia in % (N)	Sciatica in % (N)	Definitions for micturition dysfunction at presentation
Jennett [5]	***	92.0 (23/25)	92.0 (23/25)	96.0 (24/25)	100.0 (25/25)	“Some degree of paralysis of the anal and vesical sphincters”
Spännare [27]	***	100.0 (30/30)	–	70.0 (21/30)	100.0 (30/30)	“Some disorders of micturition”
O’Laoire [18]	***	100.0 (29/29)	–	51.7 (15/29)	82.8 (24/29)	“Impaired sphincter function”, i.e. “catheterization required to treat urinary retention or micturition by bladder compression, either by breath-holding or by manually compressing the abdomen”
Hellström [24]	**	70.6 (12/17)	–	94.1 (16/17)	100.0 (17/17)	“Retention of urine”, urodynamics
Kostuik [17]	**	100.0 (31/31)	54.8 (17/31)	54.8 (17/31)	100.0 (31/31)	“Urinary retention”
Gleave [25]	***	100.0 (33/33)	–	72.7 (24/33)	90.1 (30/33)	“Urinary retention with overflow incontinence”, i.e. “inability to void, or bladder enlargement with dribbling incontinence”
Shapiro [13]	**	92.9 (13/14)	–	–	100.0 (14/14)	“Urine incontinence”
Kennedy [26]	**	100.0 (19/19)	78.9 (15/19)	100.0 (19/19)	94.7 (18/19)	“Symptoms of urinary dysfunction”
Buchner [19]	***	100.0 (22/22)	36.4 (8/22)	68.2 (15/22)	100.0 (22/22)	“Urinary retention with overflow incontinence requiring catheterization”
Shapiro [14]	*	100.0 (44/44)	–	100.0 (44/44)	100.0 (44/44)	“Urologic problems of retention, incontinence or both”
Hussain [21]	**	100.0 (20/20)	70.0 (14/20)	100.0 (20/20)	100 (20/20)	“Lack of bladder control”
McCarthy [22]	**	59.5 (25/42)	50.0 (21/42)	76.2 (32/42)	90.4 (38/42)	“Urinary retention”
Qureshi [16]	**	90.9 (30/33)	30.3 (10/33)	81.8 (27/33)	84.8 (28/33)	“Urological dysfunction”
Crocker [23]	*	48.8 (21/43)	–	81.4 (35/43)	100.0 (43/43)	“Sphincter distension”
Dhatt [15]	***	100.0 (50/50)	12.0 (6/50)	90.0 (45/50)	–	“Minor or major bladder dysfunction”
Mean	–	88.9 (402/452)	47.1 (114/242)	80.8 (354/438)	95.5 (384/402)	–

Asterisks represent the number of points gained

Table 4 Outcome of micturition, defecation and sexual function at follow-up

Study	Score on risk of bias scale	Micturition dysfunction in % (N)	Corrected # micturition dysfunction in % (N)	Defecation dysfunction in % (N)	Corrected # defecation dysfunction in % (N)	Sexual dysfunction (SD) in % (N)
Jennett [5]	***	76.9 (10/13)	Idem	84.6 (11/13)	Idem	25.0 (1/4)
Spännare [27]	***	33.3 (10/30)	Idem	–	–	–
O’Laoire [18]	***	37.9 (11/29)	Idem	37.9 (11/29)	Idem	35.3 (6/17)
Hellström [24]	**	41.2 (7/17)	58.3 (7/12)	43.8 (7/16)	Idem	20.0 (2/10)
Kostuik [17]	**	13.3 (4/30)	Idem	–	–	26.6 (8/30)
Gleave [25]	***	21.2 (7/33)	Idem	–	–	–
Shapiro [13]	**	28.6 (4/14)	30.8 (4/13)	–	–	75.0 (6/8)
Kennedy [26]	**	26.3 (5/19)	Idem	10.5 (2/19)	13.3 (2/15)	10.0 (1/10)
Buchner [19]	***	22.7 (5/22)	Idem	–	–	–
Shapiro [14]	*	36.4 (16/44)	Idem	20.5 (9/44)	Idem	76.6 (23/30)
Hussain [21]	**	90.0 (18/20)	Idem	–	–	–
McCarthy [22]	**	33.3 (14/42)	56.0 (14/25)	59.5 (25/42)	119.0 (25/21)	57.1 (24/42)
Qureshi [16]	**	44.0 (11/25)	Idem	32.0 (8/25)	80.0 (8/10)	–
Crocker [23]	*	33.3 (7/21)	Idem	–	–	–
Dhatt [15]	***	90.0 (45/50)	Idem	90.0 (45/50)	750.0 (45/6)	36.0 (18/50)
Mean in % (range)	–	42.5 (13.3–90.0)	45.1 (13.3–90.0)	49.6 (10.5–90.0)	76.6 (13.3–750.0)	44.3 (10.0–76.6)

Corrected for prevalence of dysfunction at presentation

Asterisks represent the number of points gained

Sensitivity analysis

We looked more closely at the relationship between follow-up time and dysfunction (Table 5). Eight studies had a minimal follow-up of 12 months or less (average 9.9 months, range 3–12) and seven studies of more than 12 months (average 23.9 months, range 22–25). In the group with follow-up time 12 months or less, prevalence of micturition dysfunction was higher than in the group with follow-up time more than 12 months (55.3 vs. 28.1 %). The difference was statistically significant ($p = 0.043$). Regarding post-operative defecation dysfunction, we found

the group with follow-up time 12 months or less again showing a higher prevalence of dysfunction than the group with follow-up time more than 12 months (52.2 vs. 44.2 %), this difference was not statistically significant.

For sexual function, the difference was statistically significant ($p = 0.007$), with the group of follow-up time with 12 months or less showing a higher prevalence of dysfunction than the group with follow-up time with more than 12 months (47.5 vs. 38.0 %).

Reviewing the effect of bias on outcome showed that low risk of bias studies reported higher dysfunction of micturition and defecation at follow-up (49.7 and 72.8 %,

Table 5 Effect of follow-up time and risk of bias on post-operative dysfunction

	Studies with minimal follow-up time ≤12 months	Studies with minimal follow-up time >12 months	Adjusted # <i>p</i> value	Low risk of bias studies (***)	Medium to high risk of bias studies (*/**)	Adjusted # <i>p</i> value
Post-operative micturition dysfunction in % (N)	55.3 (120/217)	28.1 (54/192)	0.043	49.7 (88/177)	37.1 (86/232)	Not significant
Post-operative defecation dysfunction in % (N)	52.2 (84/161)	44.2 (34/77)	Not significant	72.8 (67/92)	34.9 (51/146)	0.017
Post-operative sexual dysfunction (SD) in % (N)	47.5 (48/101)	38.0 (35/92)	0.007	35.2 (25/71)	49.2 (64/130)	0.031

Adjusted for multiple testing

respectively) than the intermediate and high risk of bias studies (37.1 and 34.9 %, respectively) (Table 5). The difference for defecation was statistically significant ($p = 0.017$). Regarding sexual function, less dysfunction was seen in the studies with low risk of bias than in the medium–high risk of bias studies (35.2 vs. 49.2 %), with the difference being statistically significant ($p = 0.031$).

Relation of mode of presentation and reported post-operative dysfunction

Since we wished to know if there is any correlation between pre-operative dysfunction and post-operative dysfunction, e.g., whether patients presented with defecation dysfunction show a higher prevalence of post-operative sexual dysfunction, we evaluated the effect of micturition dysfunction, defecation dysfunction and saddle hypo/anesthesia at presentation on post-operative micturition, defecation and sexual function. However, none of these symptoms were found to be statistically prognostic factors for the outcome of micturition, defecation and sexual function.

Discussion

Fifteen studies reviewing outcome of micturition, defecation and sexual function in CES secondary to radiologically confirmed HNP were found, with a minimal mean follow-up time of 17.0 months. All reviewed micturition dysfunction (mean prevalence 42.5 %), eight studies reviewed defecation dysfunction (49.6 %) and nine studies SD (44.3 %). Percentages of dysfunction varied largely across studies.

In studies with a shorter follow-up time, more dysfunction was reported than in studies with a longer follow-up, with a statistically significant difference for micturition and sexual function ($p = 0.043$ and $p = 0.007$, respectively). None of the presenting symptoms analyzed were found to be statistically significant prognostic factors for post-operative micturition, defecation or sexual function.

Lower risk of bias studies reported more dysfunction of micturition and defecation (but less SD; $p = 0.031$), with the difference in defecation dysfunction being statistically significant ($p = 0.017$). We came across several other meta-analyses on various subjects in literature supporting our finding that studies with more risk of bias showed a greater treatment effect [51–53].

Differences in definition and in reviewing outcome

Reviewing literature on outcome for CES involves combining data from various studies. Differences in definitions

of CES pose difficulties in comparing outcome between studies.

Different definitions of dysfunction across studies (Table 3) may cause bias when analyzing results together. The obvious solution to this problem is to create a clear and workable term for dysfunction; however, this may be more difficult than it sounds. On the one hand, we have the physician, looking at dysfunction as organ failure, based on urodynamics and other physical tests, and on the other hand, there is the patient, experiencing dysfunction by complaints and problems in daily life. These two perspectives do not necessarily agree: e.g., in his study McCarthy measured reduced rectal tone in 21 patients, but found complaints of defecation in 25 patients. When a patient is suffering from defecation complaints due to CES, there is a problem of dysfunction, even without measurable dysfunction of normal rectal tone. In addition, the only study that evaluated micturition objectively by means of urodynamics [24], concluded that the complaints of patients were not always consistent with those urodynamic tests. Therefore, we would like to mark complaints as dysfunction even in the absence of aberrant test results, since it is quite common to not find any dysfunction by objective tests in these cases. The use of standardized questionnaires to ask for complaints, as was done in the study of Kennedy, therefore seems an elegant way to address dysfunction.

Sexual function

At an average mean minimal follow-up time of 17.0 months after surgery, almost half of patients experience SD. Sexual function may not only be directly, but also indirectly affected by CES, which is illustrated by the statement of some patients that incontinence of bladder or bowel often causes distress and a great sense of shame during sexual activity [13, 17].

The assessment of sexual function in included studies seems quite arbitrary and is done more often in men than in women. Perhaps inability to obtain or maintain erection are perceived as more basic problems than reduced sensation for women, or the participating clinicians, more often men, find it easier to discuss sexual function with the same sex. In any case, SD seems a topic difficult to discuss for both doctor and patient. Research on this subject, e.g., in the form of questionnaires for clinicians, can provide information about the place sexual problems have in current therapy, and more specifically, ideas to make the treatment of SD more common practice.

Overall, literature evaluating sexual function after CES is scarce [54]. More is written about the effect on sexual function after HNP [31, 55, 56] and spinal cord injury [57–59], some with suggestions for treatment [60].

Overview of literature

This is the first systematic review done on the outcome of micturition, defecation and sexual function in CES. Reviews have been written about these functions separately, but none of these systematically reviewed outcomes combining these three functions. As stated before, this is partly due to traditional focus of literature on timing of decompression. Two large meta-analyses found statistically significant differences in recovery of, among others, micturition and defecation in favor of early decompression [6, 61]. Smaller studies reported similar findings [13, 40, 41, 50]. In one study, the differences were statistically significant [14]. The most recent study that was included in this review evaluated outcome after delayed decompression and stated that the high prevalences of post-operative dysfunction that were found, could possibly have been prevented by early decompression [15]. One of the included studies [16], together with a study from 2009 [49] did not find differences between outcomes. Other authors suggest emergency surgery is more important for some cases of CES than for others [62–65]. However, no doubt arises that CES is an absolute indication for emergency decompression and surgery should be undertaken as soon as possible to obtain better recovery of functions [34, 41].

Clinical interpretation and conclusion

This review offers an insight into the extent of micturition dysfunction, defecation dysfunction and SD in CES after decompression. Since the discussion about timing of decompression has already been researched extensively in literature, instead we looked at the actual outcomes. Our findings show that dysfunction of micturition, defecation and/or sexual function is extremely common with about half of the patients affected in at least one of these domains at an average mean minimal follow-up time of 17 months after surgery. So, even though CES patients get decompression as soon as possible with the aim to restore function, a lot of patients still suffer from dysfunction long after surgery, something for which ideally therapy should be provided. Since micturition, defecation and sexual function are closely related and may affect each other, we believe that already existing individual therapies are best to be combined. A condition as invalidating as CES requires good patient information and the outcomes presented in this review may help in providing those data. Bias in included studies, lack of universal definitions and incomplete follow-up qualify these data as the best we momentarily have, but still subject to improvement. Since SD seems to be severely underreported, we recommend further research to explore the extent of this problem, as well as the use of questionnaires next to urodynamic tests in future

clinical (prospective) studies to accomplish a more patient-based approach.

Conflict of interest None.

References

1. Mixter WJ, Barr JS (1934) Rupture of the intervertebral disk with involvement of the spinal canal. *N Engl J Med* 211(5):210–225
2. Fraser S, Roberts L, Murphy E (2009) Cauda equina syndrome: a literature review of its definition and clinical presentation. *Arch Phys Med Rehabil* 90(11):1964–1968. doi:10.1016/j.apmr.2009.03.021
3. Shephard RH (1959) Diagnosis and prognosis of cauda equina syndrome produced by protrusion of lumbar disk. *Br Med J* 2(5164):1434–1439
4. Aho AJ, Auranen A, Pesonen K (1969) Analysis of cauda equina symptoms in patients with lumbar disk prolapse. Pre-operative and follow-up clinical and cystometric studies. *Acta Chir Scand* 135(5):413–420
5. Jennett WB (1956) A study of 25 cases of compression of the cauda equina by prolapsed intervertebral disks. *J Neurol Neurosurg Psychiatry* 19(2):109–116
6. DeLong WB, Polissar N, Neradilek B (2008) Timing of surgery in cauda equina syndrome with urinary retention: meta-analysis of observational studies. *J Neurosurg Spine* 8(4):305–320. doi:10.3171/spi.2008.8.4.305
7. Radcliff KE, Kepler CK, Delasotta LA, Rihn JA, Harrop JS, Hilibrand AS, Albert TJ, Vaccaro AR (2011) Current management review of thoracolumbar cord syndromes. *Spine J* 11(9):884–892. doi:10.1016/j.spinee.2011.07.022
8. Spector LR, Madigan L, Rhyne A, Darden B, Kim D (2008) Cauda equina syndrome. *J Am Acad Orthop Surg* 16(8):471–479
9. Ma B, Wu H, Jia LS, Yuan W, Shi GD, Shi JG (2009) Cauda equina syndrome: a review of clinical progress. *Chin Med J* 122(10):1214–1222. doi:10.3760/cma.j.issn.0366-6999.2009.10.019
10. Platform-Evidence-Based-Richtlijn-Ontwikkeling (2006) Formulier III voor het beoordelen van een Cohortonderzoek. Dutch Cochrane Center. <http://dcc.cochrane.org/sites/dcc.cochrane.org/files/uploads/cohort.pdf>. Accessed 29 March 2012
11. Harbord RM, Higgins JPT (2009) Meta Regression in Stata. In: Sterne JAC (ed) *Meta-analysis in Stata: an updated collection from the Stata journal*, 1st edn. Stata Press, College Station, pp 70–96
12. Higgins JPT, Thompson SG (2004) Controlling the risk of spurious findings from meta-regression. *Stat Med* 23:1663–1682
13. Shapiro S (1993) Cauda equina syndrome secondary to lumbar disk herniation. *Neurosurgery* 32(5):743–746 discussion 746–747
14. Shapiro S (2000) Medical realities of cauda equina syndrome secondary to lumbar disk herniation. *Spine (Phila Pa 1976)* 25(3):348–351 discussion 352
15. Dhatt S, Tahasildar N, Tripathy SK, Bahadur R, Dhillon M (2011) Outcome of spinal decompression in cauda equina syndrome presenting late in developing countries: case series of 50 cases. *Eur Spine J* 20(12):2235–2239. doi:10.1007/s00586-011-1840-4
16. Qureshi A, Sell P (2007) Cauda equina syndrome treated by surgical decompression: the influence of timing on surgical outcome. *Eur Spine J* 16(12):2143–2151. doi:10.1007/s00586-007-0491-y
17. Kostuik JP, Harrington I, Alexander D, Rand W, Evans D (1986) Cauda equina syndrome and lumbar disk herniation. *J Bone Joint Surg Am* 68(3):386–391

18. O'Laoire SA, Crockard HA, Thomas DG (1981) Prognosis for sphincter recovery after operation for cauda equina compression owing to lumbar disk prolapse. *Br Med J (Clin Res Ed)* 282(6279):1852–1854
19. Buchner M, Schiltenswolf M (2000) Cauda equina syndrome caused by intervertebral lumbar disk prolapse—mid-term results of 22 patients and review of the literature. *Neuro Orthop* 27(1–2):55–64
20. Buchner M, Schiltenswolf M (2002) Cauda equina syndrome caused by intervertebral lumbar disk prolapse: mid-term results of 22 patients and literature review. *Orthopedics* 25(7):727–731
21. Hussain SA, Gullan RW, Chitnavis BP (2003) Cauda equina syndrome: outcome and implications for management. *Br J Neurosurg* 17(2):164–167
22. McCarthy MJH, Aylott CEW, Grevitt MP, Hegarty J (2007) Cauda equina syndrome—factors affecting long-term functional and sphincteric outcome. *Spine* 32(2):207–216. doi:[10.1097/01.brs.0000251750.20508.84](https://doi.org/10.1097/01.brs.0000251750.20508.84)
23. Crocker M, Fraser G, Boyd E, Wilson J, Chitnavis BP, Thomas NW (2008) The value of inter hospital transfer and emergency MRI for suspected cauda equina syndrome: a 2-year retrospective study. *Ann R Coll Surg Engl* 90(6):513–516. doi:[10.1308/003588408x301154](https://doi.org/10.1308/003588408x301154)
24. Hellstrom P, Kortelainen P, Kontturi M (1986) Late urodynamic findings after surgery for cauda equina syndrome caused by a prolapsed lumbar intervertebral disk. *J Urol* 135(2):308–312
25. Gleave JR, MacFarlane R (1990) Prognosis for recovery of bladder function following lumbar central disk prolapse. *Br J Neurosurg* 4(3):205–209
26. Kennedy JG, Soffe KE, McGrath A, Stephens MM, Walsh MG, McManus F (1999) Predictors of outcome in cauda equina syndrome. *Eur Spine J* 8(4):317–322
27. Spannare BJ (1978) Prolapsed lumbar intervertebral disk with partial or total occlusion of the spinal canal. A study of 30 patients with and 28 patients without cauda equina symptoms. *Acta Neurochir (Wien)* 42(3–4):189–198
28. Elgamri A, Sami A, Aqqad A, Hilmani S, Ibahoin K, Naja A, El Kamar A, El Azhari A (2009) Posterior migration of a lumbar disk herniation as a cause of cauda equina syndrome. *J Radiol* 90(6):731–733 French
29. Schaeffer HR (1966) Cauda equina compression resulting from massive lumbar disk extrusion. *Aust N Z J Surg* 35(4):300–306
30. von Wild K (1990) Lumbar intervertebral disk operation in the aged. *Zentralbl Neurochir* 51(1):34–41
31. Akbas NB, Dalbayrak S, Kulcu DG, Yilmaz M, Yilmaz T, Naderi S (2010) Assessment of sexual dysfunction before and after surgery for lumbar disk herniation. *J Neurosurg Spine* 13(5):581–586. doi:[10.3171/2010.5.spine09906](https://doi.org/10.3171/2010.5.spine09906)
32. Fairburn B, Stewart JM (1955) Lumbar disk protrusion as a surgical emergency. *Lancet* 269(6885):319–321
33. Lafuente DJ, Andrew J, Joy A (1985) Sacral sparing with cauda equina compression from central lumbar intervertebral disk prolapse. *J Neurol Neurosurg Psychiatry* 48(6):579–581
34. Todd NV (2005) Cauda equina syndrome: the timing of surgery probably does influence outcome. *Br J Neurosurg* 19(4):301–306. doi:[10.1080/02688690500305324](https://doi.org/10.1080/02688690500305324)
35. Tsai CH, Chou ECL, Chou LW, Chen YJ, Chang CH, Tsou HK, Chen HT (2010) The evaluation of bladder symptoms in patients with lumbar compression disorders who have undergone decompressive surgery. *Spine* 35(17):E849–E854
36. Jegede KA, Ndu A, Grauer JN (2010) Contemporary management of symptomatic lumbar disk herniations. *Orthop Clin North Am* 41(2):217–224
37. Akbar A, Mahar A (2002) Lumbar disk prolapse: management and outcome analysis of 96 surgically treated patients. *J Pak Med Assoc* 52(2):62–65
38. Bartels R, deVries J (1996) Hemi-cauda equina syndrome from herniated lumbar disk: a neurosurgical emergency? *Can J Neurol Sci* 23(4):296–299
39. Choudhury AR, Taylor JC (1980) Cauda equina syndrome in lumbar disk disease. *Acta Orthop Scand* 51(3):493–499
40. Nielsen B, de Nully M, Schmidt K, Hansen RI (1980) A urodynamic study of cauda equina syndrome due to lumbar disk herniation. *Urol Int* 35(3):167–170
41. Dinning TA, Schaeffer HR (1993) Discogenic compression of the cauda equina: a surgical emergency. *Aust N Z J Surg* 63(12):927–934
42. Podnar S, Oblak C, Vodusek DB (2002) Sexual function in men with cauda equina lesions: a clinical and electromyographic study. *J Neurol Neurosurg Psychiatry* 73(6):715–720
43. Podnar S (2006) Bowel dysfunction in patients with cauda equina lesions. *Eur J Neurol* 13(10):1112–1117. doi:[10.1111/j.1468-1331.2006.01423.x](https://doi.org/10.1111/j.1468-1331.2006.01423.x)
44. Podnar S, Trsinar B, Vodusek DB (2006) Bladder dysfunction in patients with cauda equina lesions. *NeuroUrol Urodyn* 25(1):23–31. doi:[10.1002/nau.20188](https://doi.org/10.1002/nau.20188)
45. Scott PJ (1965) Bladder paralysis in cauda equina lesions from disk prolapse. *J Bone Joint Surg Br* 47:224–235
46. Mosdal C, Iversen P, Iversen-Hansen R (1979) Bladder neuropathy in lumbar disk disease. *Acta Neurochir (Wien)* 46(3–4):281–286
47. Jalloh I, Minhas P (2007) Delays in the treatment of cauda equina syndrome due to its variable clinical features in patients presenting to the emergency department. *Emerg Med J* 24(1):33–34. doi:[10.1136/emj.2006.038182](https://doi.org/10.1136/emj.2006.038182)
48. Inui Y, Doita M, Ouchi K, Tsukuda M, Fujita N, Kurosaka M (2004) Clinical and radiologic features of lumbar spinal stenosis and disk herniation with neuropathic bladder. *Spine (Phila Pa 1976)* 29(8):869–873
49. Olivero WC, Wang HA, Hanigan WC, Henderson JP, Tracy PT, Elwood PW, Lister JR, Lyle L (2009) Cauda equina syndrome (CES) from lumbar disk herniations. *J Spinal Disord Tech* 22(3):202–206. doi:[10.1097/BSD.0b013e31817baad8](https://doi.org/10.1097/BSD.0b013e31817baad8)
50. Busse JW, Bhandari M, Schnitker JB, Reddy K, Dunlop RB (2001) Delayed presentation of cauda equina syndrome secondary to lumbar disk herniation: functional outcomes and health-related quality of life. *CJEM* 3(4):285–291
51. Geisler BP, van Dam RM, Gazelle GS, Goehler A (2009) Risk of bias in meta-analysis on erythropoietin-stimulating agents in heart failure. *Heart* 95(15):1278–1279. doi:[10.1136/hrt.2009.172163](https://doi.org/10.1136/hrt.2009.172163) author reply 1279
52. Mangum K, Partna L, Vavrek D (2012) Spinal manipulation for the treatment of hypertension: a systematic qualitative literature review. *J Manip Physiol Ther*. doi:[10.1016/j.jmpt.2012.01.005](https://doi.org/10.1016/j.jmpt.2012.01.005)
53. Hartling L, Bond K, Vandermeer B, Seida J, Dryden DM, Rowe BH (2011) Applying the risk of bias tool in a systematic review of combination long-acting beta-agonists and inhaled corticosteroids for persistent asthma. *PLoS ONE* 6(2):e17242. doi:[10.1371/journal.pone.0017242](https://doi.org/10.1371/journal.pone.0017242)
54. Scheiber-Nogueira MC (2009) Sexual dysfunction in cauda equina syndrome. *Pelvi-Perineologie* 4(3):191–195. doi:[10.1007/s11608-009-0249-6](https://doi.org/10.1007/s11608-009-0249-6)
55. Choy DS (1999) Early relief of erectile dysfunction after laser decompression of herniated lumbar disk. *J Clin Laser Med Surg* 17(1):25–27
56. Orlin JR, Klevmark B (2008) Successful disk surgery after 17 years of erectile dysfunction caused by a “silent” disk protrusion. *Scand J Urol Nephrol* 42(1):91–93. doi:[10.1080/00365590701428467](https://doi.org/10.1080/00365590701428467)
57. Alexander M, Rosen RC (2008) Spinal cord injuries and orgasm: a review. *J Sex Marital Ther* 34(4):308–324. doi:[10.1080/00926230802096341](https://doi.org/10.1080/00926230802096341)

58. Deforge D, Blackmer J, Garritty C et al (2006) Male erectile dysfunction following spinal cord injury: a systematic review. *Spinal Cord* 44(8):465–473
59. Larsen E, Hejgaard N (1984) Sexual dysfunction after spinal cord or cauda equina lesions. *Paraplegia* 22(2):66–74. doi:[10.1038/sc.1984.14](https://doi.org/10.1038/sc.1984.14)
60. Zabihi N, Mourtzinos A, Maher MG, Raz S, Rodriguez LV (2008) The effects of bilateral caudal epidural S2–4 neuromodulation on female sexual function. *Int Urogynecol J Pelvic Floor Dysfunct* 19(5):697–700. doi:[10.1007/s00192-007-0504-8](https://doi.org/10.1007/s00192-007-0504-8)
61. Ahn UM, Ahn NU, Buchowski JM, Garrett ES, Sieber AN, Kostuik JP (2000) Cauda equina syndrome secondary to lumbar disk herniation —A meta-analysis of surgical outcomes. *Spine* 25(12):1515–1522. doi:[10.1097/00007632-200006150-00010](https://doi.org/10.1097/00007632-200006150-00010)
62. Gleave JRW, Macfarlane R (2002) Cauda equina syndrome: what is the relationship between timing of surgery and outcome? *Br J Neurosurg* 16(4):325–328. doi:[10.1080/0268869021000032887](https://doi.org/10.1080/0268869021000032887)
63. Nascone JW, Lauerma WC, Wiesel SW (1999) Cauda equina syndrome: is it a surgical emergency? *University Pennsylvania Orthop J* 12:73–76
64. Gardner A, Gardner E, Morley T (2011) Cauda equina syndrome: a review of the current clinical and medico-legal position. *Eur Spine J* 20(5):690–697. doi:[10.1007/s00586-010-1668-3](https://doi.org/10.1007/s00586-010-1668-3)
65. Tandon PN, Sankaran B (1967) Cauda equina syndrome due to lumbar disk prolapse. *Indian J Orthop* 1:112–119