Published in final edited form as:

Am J Health Behav. 2007 ; 31(3): 238–248. doi:10.5555/ajhb.2007.31.3.238.

Alcohol Consumption, Medical Conditions, and Health Behavior in Older Adults

Derek D. Satre, PhD, Nancy P. Gordon, ScD, and Constance Weisner, DrPH

Derek D. Satre, Assistant Adjunct Professor, Department of Psychiatry, University of California, San Francisco, San Francisco, CA and Adjunct Investigator, Division of Research, Kaiser Permanente Northern California Region, Oakland, CA. Nancy P. Gordon, Investigator, Division of Research, Kaiser Permanente Northern California Region, Oakland, CA. Constance Weisner, Professor, Department of Psychiatry, University of California, San Francisco, CA and Investigator, Division of Research, Kaiser Permanente, Northern California Region, Oakland CA

Abstract

Objectives—To examine associations between drinking patterns, medical conditions, and behavioral health risks among older adults.

Methods—Analyses compared survey participants (health plan members ages 65 to 90, N = 6662) who drank moderately to those who drank over recommended limits or did not drink.

Results—Overlimit drinking was associated with smoking; not trying to eat low-fat foods (in men), and lower BMI (in women). Predictors of not drinking during the prior 12 months included ethnicity, lower education, worse self-reported health, diabetes and heart problems.

Conclusions—Significant relationships exist between health and alcohol consumption patterns, which vary by gender.

Keywords

older adults; alcohol; behavioral health risks; obesity; gender; ethnicity

Older adults compose a large and fast-growing segment of the US population. Yet alcohol consumption in this demographic group has received little study. This is a significant oversight because heavy drinking has been associated with development of multiple medical,^{1,2} functional,³ and psychiatric problems to which older adults are especially vulnerable.^{4,5} Excessive alcohol consumption may be clustered with other behavioral health risks such as smoking and poor diet^{6,7} and obesity,⁸ increasing cumulative risk for medical problems. In contrast, light to moderate drinking may have health benefits.⁹ In studies of heart disease, moderate drinking has been associated with lower mortality than either abstinence or heavy consumption.^{10–12} These studies indicate that alcohol is harmful to health in excess but potentially helpful in moderation and that the relationship of drinking to health is complex.

To prevent serious health problems, recommended drinking limits for older adults have been proposed.^{7,13} Because alcohol can exacerbate medical conditions and has potential for adverse medication interactions, older adults with health problems who drank when they were younger may stop drinking completely.^{14–16} This tendency may explain why studies comparing older adults who drink heavily or moderately to those who abstain completely

have found that abstainers have poorer health.^{17–19} For example, in a large sample of male primary care patients ages 65 and over, patients who reported drinking alcohol in the previous year had better health status (as measured by the SF-36) than did abstainers.¹⁷

Epidemiological studies have found that older women tend to drink smaller quantities of alcohol than older men do, have fewer alcohol-related problems,^{20,21} and are more likely to abstain totally.²² Yet prevalence of drinking over recommended limits is comparable when the limits are set lower for women than for men, as some guidelines have proposed.¹³ For example, in a large (N=5065) multisite primary care study of patients ages 60 and over, 15% of men consumed over 14 drinks per week and 12% of women regularly drank over 7 drinks per week.⁷ These data indicate that although women drink less than men, both men and women may be at risk due to overlimit drinking.

Gender may be associated with changes in drinking patterns in late life as well as other health behaviors.²³ Older women have higher abstinence rates than older men following chemical dependency treatment, based on a study of 92 alcohol dependent patients ages 55 and over in a managed-care outpatient treatment program.²⁴ In that study, women had a later onset of heavy drinking than men had, but were drinking similar amounts of alcohol at the time of treatment entry. In a Canadian population-based survey of older adults, women were more likely than men to take steps to improve their health such as changing diet and using preventive services.²⁵ In a study of 139 primary care patients in San Francisco, women who stopped drinking had higher rates of hypertension and heart problems than did current drinkers, whereas the same relationship was not observed among men.¹⁶ Based on this limited literature, we anticipated that older women in the current study might be more likely to eliminate drinking in the presence of chronic health problems.

There have been few studies of any kind examining ethnic differences in older adult drinking patterns. In one large study, adults 65 and over were included in the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC), a nationally representative sample. Among men, alcohol abuse (based on DSM-IV criteria) was diagnosed in 2.38% of whites, 1.79% of African Americans, 0.00% of Asians, and 3.69% of Hispanics. Among women, alcohol abuse was diagnosed in 0.36% of whites, 0.12% of African Americans, 0.00% of Asians and 0.00% of Hispanics.²⁶ This study examined only substance use disorders and did not report alcohol use patterns (abstinence or quantity/frequency) in the sample. One study comparing small samples of African American and white alcohol-dependent men found that African Americans were more likely to have experienced adverse social consequences due to drinking.²⁷ The authors speculated that once alcohol problems develop, they may become more severe for older minorities due to having fewer financial resources. In addition to cultural and economic factors, ethnic differences in life course trajectories of alcohol use²⁸ may be influenced by higher rates of health problems for which drinking is not recommended, such as diabetes, especially among Hispanic and African American adults.^{29,30} Based on these factors, it was anticipated that ethnic minorities in the present study of older adults might be more likely than whites to report cessation of alcohol consumption.

This current study contributes to the existing literature in several ways. It uses a large, representative health plan population sample to describe patterns of alcohol consumption and abstinence by ethnicity and gender among older adults. Consumption measures correspond to widely used drinking guidelines and thus are relevant to clinical practice. It differentiates those who reported quitting drinking from those who never consumed alcohol as adults, an important yet often overlooked distinction in examining the relationship of alcohol to health.³¹ To our knowledge, it is the first study that correlates alcohol consumption among older adults with chronic medical conditions, health status, and

behavioral health risks. It reports on the relationship of alcohol use to obesity, an important area of investigation in which the existing literature is sparse.

In summary, it was anticipated that individuals who drank above recommended limits or who had ceased drinking would be more likely to report physical and mental health problems than would those who drank moderately (within recommended limits) and that overlimit drinking would be associated with other behavioral health risks. Based on the limited literature described above, it was expected that alcohol consumption would vary by gender and ethnicity. Understanding these patterns has the potential to inform strategies to limit alcohol consumption among older adults.

METHODS

Data for this study came from the 2002 Kaiser Permanente (KP) Adult Member Health Survey, a mailed survey of health plan members in the northern California region. The survey, which was available only in English, included questions about demographic and health-related characteristics. All measures were based on self-report. In April 2002, 8-page questionnaires were mailed to a stratified random sample of 40,000 adult KP Northern California Region Health Plan members aged 20 and over. To ensure that the sample reflected geographic differences in the membership, separate samples of set numbers of men and women from different age-groups were randomly selected for each of the health plan's 18 medical center service populations, with oversampling of members aged 65 and over. Two subsequent survey packet mailings were sent to nonrespondents, with the final wave of questionnaires mailed in mid-July 2002.

Among adults ages 65 and over, response rate was 74.4% for women ($n = 3461/4655$) and 73.6% for men ($n = 3436/4670$) after excluding from the denominator those individuals who were deceased, no longer health plan members, or for whom no current address was available through KP records or the post office. The sample for the current analyses was restricted to adults ages 65 to 90.

Measures

Demographic characteristics—The questionnaire asked age, gender, race/ ethnicity, highest level of education, and marital status.

Alcohol consumption—participants were asked “During the past 12 months, how often have you had a drink containing alcohol?” with response options of almost every day, 5 to 6 times a week, 3 to 4 times a week, 1 to 2 times a week, 2–4 times a month, 1 time a month or less, never in the past 12 months (but, used to drink) or never drank as an adult. Those who reported consuming any alcohol were asked “On days when you had a drink, how many drinks did you usually have?” with examples given of one drink equaling a 12-oz can of beer, 4 oz of wine or 1 oz shot of hard liquor. Average amount of alcohol consumed per week was estimated based on a combination of these 2 items.³² Although no collateral or biochemical verification of participants' alcohol use was available, similar self-report quantity, frequency, and heavy drinking questions have shown moderate to good validity and excellent reliability in an older adult primary care sample.³³ Consistent with prior studies, women were categorized as overlimit drinkers if they reported drinking either more than one drink at a time or more than 7 drinks per week, although men were considered over limit if they consumed either more than 2 drinks at a time or more than 14 drinks per week.⁷

Health conditions—Participants were asked to indicate whether, during the prior 12 months, they had (or took medication for) any of 28 major health conditions. We restricted our analysis to 7 self-reported chronic conditions common in older adults (heart problems,

high blood pressure, high cholesterol, arthritis, diabetes, depression, and anxiety). Participants were also asked if they had a problem with either alcohol or drugs (yes/no) in the prior 12 months.

Participants were asked to rate their overall health status using the question “In general, would you say your health is excellent, very good, good, fair or poor?” This question is predictive of future health care utilization,³⁴ morbidity,³⁵ and mortality.³⁶ The measure was dichotomized (excellent, very good, or good vs fair or poor) in the analyses.^{37, 38} Body mass index (BMI) was calculated using questions asking about height and weight.

Behavioral health risks—Participants were asked if they currently smoked cigarettes (yes/no). They were asked “During the past 12 months, how often did you usually get physical exercise (such as walking, swimming, gardening, golf, tennis, etc.): 5 or more times per week, 3 to 4 times a week, 1 to 2 times a week, 2 to 4 times a month, once a month or less, or never?” As an indicator of a sedentary lifestyle, we dichotomized the measure to compare those who exercised 2 to 4 times per month or less (well under recommended frequency)³⁹ to those who exercised more often. Regarding diet, participants were asked “About how often do you try to eat reduced-fat foods: all the time, most of the time, some of the time, a little of the time, or never?” This variable was dichotomized (a little of the time/never vs some/most/all or the time) as an indicator of having poor dietary practices.

Analyses

Participants were divided into 4 categories of drinking behavior: adult non-drinkers, abstinent for 12 months, moderate drinkers (ie, those who drank under recommended limits), and over-limit drinkers (more than one drink at a time or more than 7 drinks per week for women, more than 2 drinks at a time or more than 14 drinks per week for men).⁷ Because some current guidelines suggest no more than 7 drinks per week for both men and women,⁴⁰ the study replicated the analysis for men using this lower limit, to see whether using a lower drinking cutoff yielded different results regarding the relationship of alcohol consumption to health measures. A subanalysis compared medical conditions of heavy drinkers (over 3 drinks per day for women and over 4 drinks per day for men) to those of current drinkers who consumed less than this amount, controlling for age and ethnicity, using logistic regression. Overall group differences in demographic characteristics and health measures between categories of drinking behavior were examined using Rao-Scott chi-square, with respondent data weighted to reflect the sampling procedures in the survey.

To examine the relationship of medical conditions, self-reported health and behavioral health risks to both drinking cessation (12-month abstinence) and overlimit drinking, separate logistic regression analyses were conducted to test the effect of drinking status on these health variables, controlling for age and ethnicity and BMI (for analysis of medical conditions).

To develop the model predicting drinking cessation, bivariate comparisons were made between current drinkers and those who had been drinkers during adulthood, but had ceased drinking (12 month abstinence). Variables selected for the logistic regression models included those factors significant in the bivariate comparisons: age (measured in 5-year intervals, reference group 65–69), ethnicity (African American, Asian, and Latino, with white non-Hispanic as reference group), lower education (12 years vs at least some college as reference group), perceived health status (fair or poor vs good-excellent as reference group), and the presence of selected health conditions.

Logistic regression was used to examine factors associated with cessation of drinking, among participants who had consumed alcohol as adults, with analyses conducted separately

for men and women. Respondents were assigned a poststratification weighting factor based on the sampling frame so that the weighted data used in the analysis would reflect the age (in 5-year intervals), gender, and geographic distribution of the adult membership at the time of the survey. All analyses were conducted with weighted data using SAS version 9.1 procedures (Proc Surveyfreq and Proc Survelogistic) for analysis of data obtained from a complex survey design, using medical center service population as the stratum variable. These special procedures take into account error introduced by design effects when calculating the variances used for significance testing. For further information on the survey design and weighting procedures, please contact the corresponding author.

We calculated the *c* statistic to report discriminative power of the models for men and women. Pseudo-R-squared was calculated (using unweighted data in order to obtain accurate sample size) to estimate the variance explained by the models.

RESULTS

Sample

The final sample consisted of 6662 individuals ages 65 to 90. This included 3316 men (737 men ages 65 to 69, 737 ages 70 to 74, 1330 ages 75 to 79, 351 ages 80 to 84, and 161 ages 85 to 90) and 3346 women (803 ages 65 to 69, 750 ages 70 to 74, 1329 ages 75 to 79, 330 ages 80 to 84, and 134 ages 85 to 90). Based on weighted data, participants had a mean age of 73.7 (sd = 6.3). Ethnicity data were available for 92% of the sample. Among men, respondents were 82.5% white, 5.0% African American, 5.6% Latino, and 6.9% Asian American. Among women, respondents were 79.4% white, 5.2% African American, 7.2% Latino, and 8.2% Asian American. A small percentage (2 %) reporting Pacific Islander, Native American or “other” were excluded from analyses by ethnicity.

Patterns of Alcohol Consumption

Drinking data were available for 6116 participants (3081 men and 3035 women) with 546 missing (8%) due to nonresponse on one or more drinking items. Those with missing data were older ($P < .001$) and more likely to be female ($P < .001$) and nonwhite ($P < .001$) than were those without missing data.

Results found that 12.1% of men and 16.6% of women drank over recommended limits. Of these, men reported an average of 3.8 (sd = 1.53) drinks and women an average of 2.2 (sd = 0.59) drinks consumed on days when they drank alcohol (not shown). There were significant overall effects on drinking category by age-group, ethnicity, and relationship status (Table 1). Among both men and women, overlimit drinking was lower in the older age-groups (eg, ages 80 to 90 vs 65–74 or 75–79), whereas 12-month abstinence was higher in older age-groups. Whites and Latinos had the highest percentages of overlimit drinkers and Asians the lowest percentages, and African Americans higher percentages of 12-month abstainers than other ethnic groups. Less educated men and women were more likely to report long-term abstinence or 12-month abstinence than were those with higher levels of education.

Medical Conditions and Behavioral Health Risks

Medical conditions and behavioral health risks were examined for men and women. Overall group differences by drinking category were examined using chi square (Table 2). Logistic regression analyses examined odds of having health conditions, fair or poor self-reported health (vs good, very good, or excellent), and behavioral health risks among those who reported overlimit drinking in the prior 12 months vs moderate drinkers, adjusting for age and ethnicity. Medical conditions were also adjusted for BMI. Few significant relationships were found between overlimit drinking and prevalence of major medical conditions.

Contrary to expectation, men who drank over limits had lower prevalence of heart problems (OR = 0.71, CI = 0.51, 0.99), and high cholesterol (OR = 0.71, CI = 0.53, 0.95) than that of moderate drinkers. Women who drank over limits had lower prevalence of heart problems (OR = 0.61, CI = 0.41, 0.93) and high cholesterol (OR = 0.75, CI = 0.57, 0.99) than that of moderate drinkers. When compared with moderate drinking, overlimit drinking was associated with poorer dietary habits (not trying to eat reduced-fat foods) among men (OR = 1.58, CI = 1.21, 2.06) and with current cigarette smoking among women (OR = 2.36, CI = 1.56, 3.57) and men (OR = 2.35, CI = 1.56, 3.53). Among women, overlimit drinking was associated with greater likelihood of being sedentary (OR = 1.49, CI = 1.09, 2.06), but lower likelihood of being overweight (BMI ≥ 30), (OR = 0.61, CI = 0.43, 0.85).

When the relationship of drinking level with medical problems was examined using the lower recommended drinking limit for men (no more than one drink per day), as expected a much larger number of individuals were categorized as overlimit drinkers (32.2% of male study participants, rather than 12.1%). However, only one health correlate changed: overall effect of drinking level on having a lower rate of heart problems became significant ($P = .002$, not shown).

A subanalysis compared health conditions of male heavy drinkers ($n=131$), who consumed over 4 drinks per day, to currently drinking males who drank less than this amount. The small number of heavy drinkers resulted in limited statistical power for this analysis within men and precluded any such comparisons for women ($n=17$) heavy drinkers. The only statistically significant difference was that heavy drinkers were less likely to report having or being treated for high cholesterol than were other drinkers (39.1% vs 22.1%, $P = .001$).

Health Factors Associated With Drinking Cessation

Logistic regression analyses examined odds of having health conditions, fair or poor self-reported health (vs good, very good, or excellent), and behavioral health risks among those who reported moderate drinking vs those who reported no drinking in the prior 12 months vs adjusting for age and ethnicity (see post hoc comparisons reported on Table 2). Medical conditions were also adjusted for BMI. Variables associated with moderate drinking vs 12-month abstinence included lower likelihood of having diabetes (OR = 0.55, CI = 0.41, 0.75), depression (OR = 0.55, CI = 0.36, 0.83), anxiety (OR = 0.45, CI = 0.24, 0.85), fair or poor health (OR = 0.41, CI = 0.31, 0.53), smoking cigarettes (OR = 0.53, CI = 0.35, 0.80), and being sedentary (OR = 0.60, CI = 0.44, 0.83) among men; and heart problems (OR = 0.47, CI = 0.33, 0.66), diabetes (OR = 0.35, CI = 0.24, 0.52), arthritis (OR = 0.73, CI = 0.55, 0.97), depression (OR = 0.50, CI = 0.34, 0.72), fair or poor health (OR = 0.25, CI = 0.19, 0.34), smoking (OR = 0.48, CI = 0.30, 0.77) and being sedentary (OR = 0.37, CI = 0.26, 0.51) among women.

Predictors of Drinking Cessation

In logistic regression analysis, predictors of having quit drinking (12-month abstinence among former alcohol consumers) vs any level of current drinking were examined separately for men and women. To select variables for the model, bivariate chi-square analyses were used to compare demographic factors and health conditions of current drinkers to those of participants who had abstained for 12-months. Individuals who never drank as adults were excluded. Factors significant at the $P < .10$ level were included in the initial model: Age-group, ethnicity (African American, Hispanic, and Asian American vs white), marital status, education, self-reported health, diabetes, and heart problems were in the models for both genders; high blood pressure and high cholesterol were in the initial model for women only. Results of the final model found that among both men and women, drinking cessation was predicted by ethnicity (African American vs white), lower education,

diabetes, and worse self-reported health. Depression was significant among men only. Heart problems were significant among women only (Table 3). Age-group was not significant in either final model.

To estimate overall discriminative power of the model in separating current drinkers from those who had quit drinking, we calculated the *c* statistic, which is equivalent to the area under the ROC curve for the model.⁴¹ For men, *c* = 0.641; for women, *c* = 0.718. Results indicate the probability that the model would correctly assign possible cases involving the dependent variable 64.1% of the time for men and 71.8% of the time for women. For the overall model, pseudo-R-squared = .04 for men, *P* < .001, and .08 for women, *P* < .001.

DISCUSSION

This study examined how health conditions and behaviors, gender, and ethnicity are related to drinking patterns among older adults, including overlimit drinking and drinking cessation (12-month abstinence). Although increased prevalence of medical problems with overlimit drinking was not found, overlimit drinking was associated with other behavioral health risks. Results found an association between health problems and having stopped drinking. Factors independently associated with drinking cessation among both women and men included lower education, ethnicity (African American), and worse self-reported health.

The results help inform understanding of how age influences alcohol consumption. Although not uniform across cohorts,⁴² longitudinal studies have found a decline in quantity of alcohol consumed and an increase in abstinence with age.^{20,21} Medical illness has also been associated with changes in alcohol consumption in middle-aged populations, as we found in our sample.⁴³ Lack of an independent effect of age on drinking cessation in our sample suggests that medical illness may be more important than age in motivating drinking reduction.

Gender Differences

Among both men and women, medical conditions and worse self-rated health were associated with nondrinking in the prior 12 months. Similar to findings in a study of a small sample of older adult primary care patients,¹⁶ heart disease was independently associated with drinking cessation among women but not among men. This may indicate that women with heart problems are more likely to stop drinking than men. Alternatively, it has been suggested that alcohol consumption could benefit women more than men,⁴⁴ particularly for heart problems.

Gender differences were also observed in the relationship of overlimit drinking to poor dietary practices (not trying to eat low-fat foods) and having BMI in the obese range (> 30). Overlimit drinking men had worse results than moderate drinkers on both measures, whereas among women overlimit drinkers were less likely than moderate drinkers to be obese. Previous investigations in younger samples have yielded inconsistent results regarding the relationship of alcohol consumption to obesity,^{45,46} with some recent evidence from a general population sample that quantity of alcohol consumed may increase BMI (measured as a continuous variable) among both women and men.²² In contrast, our findings suggest that among older women, overlimit drinking may not have an adverse impact on rates of obesity.

Clinical Implications

Our results indicate that older men and women who drink over recommended limits may not, in the short term, present in health care settings with worse health than other patients.⁵ However, overlimit drinkers are more likely than moderate drinkers to have behavioral

health risks such as smoking and poor dietary practices, which could be targeted for intervention by medical providers. It is cautioned that because of the correlational study design in a sample with relatively few heavy drinkers, and potential risks associated with overlimit drinking, our findings do not support clinical recommendations for older adults to consume alcohol or to disregard established drinking guidelines.

The relationship among health problems, worse self-reported health, and cessation of alcohol consumption implies that medical illness and the perception of having poor health may motivate elimination of alcohol use. Brief interventions focused on adverse drinking consequences may effectively capitalize on this inclination.^{5,47} In addition, brief interventions may also be useful to reduce drinking in patients with type 2 diabetes and hypertension,⁴⁸ a finding particularly relevant to this sample.

Limitations

Because the survey was conducted only in English, individuals who are non-English speaking or who have very poor reading ability may be underrepresented. African Americans and Hispanics may be less likely to respond to the survey than whites and Pacific Islanders, based on analysis of a previous (1993) Kaiser Member Health Survey year. Health plan members who were very sick and frail or who resided in skilled nursing facilities were unlikely to respond. Nonresponse on the alcohol questions was higher among nonwhites and women, people with lower levels of education, and the oldest study participants. Because these nonrespondents were demographically similar to nondrinkers in the sample, it seems likely that some nondrinkers chose to skip the alcohol questions in the survey. Our results may therefore overestimate the proportion of the total sample that drinks. However, it is not anticipated that these missing data would bias the results regarding the relationships between drinking and health reported in the study.

The current investigation uses retrospective data and provides observational support for an association between health and alcohol consumption, but cannot draw conclusions regarding directionality of these relationships. Further, both the alcohol use and health conditions examined were based on self-report. Although self-report alcohol measures have generally good reliability, and survey participants may be less motivated to underreport drinking than they would in a clinical context, underreporting of alcohol consumption by participants is a potential source of bias. Some respondents who reported no health conditions may have an undiagnosed disorder. However, strengths of the study include its large older adult sample, inclusion of women and minority groups who have been underrepresented in prior studies, and analysis of a privately insured sample with access to health care. The analyses also distinguished former drinkers from those who never drank as adults and included measures of self-reported health, body mass index (BMI), and behavioral health risks.

CONCLUSION

This study reported patterns of alcohol consumption in a large, representative sample of adults ages 65 to 90. Results of this investigation found that those who quit drinking were more likely than current drinkers to report having medical problems, including diabetes, heart problems, and worse self-reported health. Drinking over recommended limits was not associated with medical problems in a one-year period, but was correlated with other behavioral health risks such as smoking and poor dietary practices. Gender differences were observed in the association of alcohol consumption with health and behaviors. Findings have implications for identification of older adults with problematic drinking and the development of strategies to reduce alcohol consumption.

Acknowledgments

This study was supported by the Kaiser Permanente Community Benefit Program, Oakland, Calif. The first author was supported by a grant from the National Institute on Alcohol Abuse and Alcoholism (K23 AA015411). The authors would like to thank Frederic Blow, PhD, and Jennifer Mertens, PhD, for helpful comments on the manuscript; Qing Zheng, BA, for assistance with data analysis; Kevin Delucchi, PhD, for statistical consultation; and Agatha Hinman, BA, for editorial assistance. Findings were presented in part at the annual meeting of the International Society of Addiction Medicine, April 23, 2005, Mar del Plata, Argentina.

REFERENCES

1. Finlayson, RE.; Hurt, RD. Medical consequences of heavy drinking by the elderly. In: Gombert, ESL.; Hegedus, AM.; Zucker, RA., editors. Alcohol Problems and Aging. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; 1998. p. 193-212. Research Monograph No. 33
2. Meyerhoff DJ, Bode C, Nixon SJ, et al. Health risks of chronic moderate and heavy alcohol consumption: how much is too much? Alcohol Clin Exp Res. 2005; 29(7):1334–1340. [PubMed: 16088997]
3. Moore AA, Endo JO, Carter MK. Is there a relationship between excessive drinking and functional impairment in older persons? J Am Geriatr Soc. 2003; 51(1):44–49. [PubMed: 12534844]
4. Ferreira KM, Sloan FA. Excess alcohol consumption and health outcomes: a 6-year follow-up of men over age 50 from the health and retirement study. Addiction. 2002; 97(3):301–310. [PubMed: 11964106]
5. Blow FC, Walton MA, Barry KL, et al. The relationship between alcohol problems and health functioning of older adults in primary care settings. J Am Geriatr Soc. 2000; 48(7):769–774. [PubMed: 10894315]
6. Ma J, Betts NM, Hampl JS. Clustering of lifestyle behaviors: the relationship between cigarette smoking, alcohol consumption, and dietary intake. Am J Health Promot. 2000; 15(2):107–117. [PubMed: 11194694]
7. Adams WL, Barry KL, Fleming MF. Screening for problem drinking in older primary care patients. JAMA. 1996; 276(24):1964–1967. [PubMed: 8971065]
8. Breslow RA, Smothers BA. Drinking patterns and body mass index in never smokers: National Health Interview Survey, 1997–2001. Am J Epidemiol. 2005; 161(4):368–376. [PubMed: 15692081]
9. Stampfer MJ, Kang JH, Chen J, et al. Effects of moderate alcohol consumption on cognitive function in women. N Engl J Med. 2005; 352(3):245–253. [PubMed: 15659724]
10. Britton A, Marmot M. Different measures of alcohol consumption and risk of coronary heart disease and all-cause mortality: 11-year follow-up of the Whitehall II Cohort Study. Addiction. 2004; 99(1):109–116. [PubMed: 14678069]
11. Wannamethee SG, Shaper AG. Type of alcoholic drink and risk of major coronary heart disease events and all-cause mortality. Am J Public Health. 1999; 89(5):685–690. [PubMed: 10224979]
12. Goldberg RJ, Burchfiel CM, Reed DM, et al. A prospective study of the health effects of alcohol consumption in middle-aged and elderly men The Honolulu Heart Program. Circulation. 1994; 89(2):651–659. [PubMed: 8313554]
13. National Institute on Alcohol Abuse and Alcoholism. The physicians' guide to helping patients with alcohol problems. Bethesda, MD: U.S. Department of Health and Human Services [NIH Publication No. 95-3769]; 1995.
14. Krause N. Stress, religiosity, and abstinence from alcohol. Psychol Aging. 1991; 6(1):134–144. [PubMed: 2029362]
15. Moos RH, Brennan PL, Schutte KK, Moos BS. Older adults' health and changes in late-life drinking patterns. Aging Ment Health. 2005; 9(1):49–59. [PubMed: 15841832]
16. Satre DD, Areal PA. Effects of gender, ethnicity, and medical illness on drinking cessation in older primary care patients. J Aging Health. 2005; 17(1):70–84. [PubMed: 15601784]
17. Bridevaux IP, Bradley KA, Bryson CL, et al. Alcohol screening results in elderly male veterans: association with health status and mortality. J Am Geriatr Soc. 2004; 52(9):1510–1517. [PubMed: 15341553]

18. Graham K, Schmidt G. The effects of drinking on health of older adults. *Am J Drug Alcohol Abuse*. 1998; 24(3):465–481. [PubMed: 9741947]
19. Vogeltanz ND, Wilsnack SC, Vickers KS, Kristjanson AP. Sociodemographic characteristics and drinking status as predictors of older women's health. *J Gen Psychol*. 1999; 126(2):135–147. [PubMed: 10368940]
20. Fillmore KM. Women's drinking across the adult life course as compared to men's: a longitudinal and cohort analysis. *Br J Addict*. 1987; 82(7):801–811. [PubMed: 3478069]
21. Moore AA, Gould R, Reuben DB, et al. Longitudinal patterns and predictors of alcohol consumption in the United States. *Am J Public Health*. 2005; 95(3):458–465. [PubMed: 15727977]
22. Breslow RA, Smothers B. Drinking patterns of older Americans: National Health Interview Surveys, 1997–2001. *J Stud Alcohol*. 2004; 65(2):232–240. [PubMed: 15151355]
23. Gallant MP, Dorn GP. Gender and race differences in the predictors of daily health practices among older adults. *Health Educ Res*. 2001; 16(1):21–31. [PubMed: 11252281]
24. Satre DD, Mertens JR, Weisner C. Gender differences in treatment outcomes for alcohol dependence among older adults. *J Stud Alcohol*. 2004; 65(5):638–642. [PubMed: 15536774]
25. Newsom JT, Kaplan MS, Huguet N, McFarland BH. Health behaviors in a representative sample of older Canadians: prevalences, reported change, motivation to change, and perceived barriers. *Gerontologist*. 2004; 44(2):193–205. [PubMed: 15075416]
26. Grant BF, Dawson DA, Stinson FS, et al. The 12-month prevalence and trends in DSM-IV alcohol abuse and dependence: United States, 1991–1992 and 2001–2002. *Drug Alcohol Depend*. 2004; 74(3):223–234. [PubMed: 15194200]
27. Gomberg, ES.; Nelson, BW. Black and white older men: alcohol use and abuse. In: Beresford, TP.; Gomberg, ES., editors. *Alcohol and Aging*. New York: Oxford University Press; 1995. p. 307–323.
28. Jackson, JS.; Williams, DR.; Gomberg, ESL. Aging and alcohol use among African Americans: A life-course perspective. In: Gomberg, ESL.; Hegedus, AM.; Zucker, RA., editors. *Alcohol Problems and Aging*. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; 1998. p. 63–87. Research Monograph No. 33
29. Narayan KM, Boyle JP, Thompson TJ, et al. Lifetime risk for diabetes mellitus in the United States. *JAMA*. 2003; 290(14):1884–1890. [PubMed: 14532317]
30. Brancati FL, Kao WH, Folsom AR, et al. Incident type 2 diabetes mellitus in African American and white adults: the Atherosclerosis Risk in Communities Study. *JAMA*. 2000; 283(17):2253–2259. [PubMed: 10807384]
31. Fillmore KM, Kerr WC, Stockwell T, et al. Moderate alcohol use and reduced mortality risk: systematic error in prospective studies. *Addict Res Theory*. 2006; 14(2):101–132.
32. Armstrong MA, Midanik LT, Klatsky AL. Alcohol consumption and utilization of health services in a health maintenance organization. *Med Care*. 1998; 36(11):1599–1605. [PubMed: 9821947]
33. Bradley KA, McDonell MB, Bush K, et al. The AUDIT alcohol consumption questions: reliability, validity, and responsiveness to change in older male primary care patients. *Alcohol Clin Exp Res*. 1998; 22(8):1842–1849. [PubMed: 9835306]
34. Bierman AS, Bubolz TA, Fisher ES, Wasson JH. How well does a single question about health predict the financial health of Medicare managed care plans? *Eff Clin Pract*. 1999; 2(2):56–62. [PubMed: 10538477]
35. Weisen SF, Frishman WH, Aronson MK, Wassertheil-Smoller S. Self-rated health assessment and development of both cardiovascular and dementing illnesses in an ambulatory elderly population: A report from the Bronx Longitudinal Aging Study. *Heart Dis*. 1999; 1(4):201–205. [PubMed: 11720624]
36. Idler EL, Benyamini Y. Self-rated health and mortality: a review of twenty-seven community studies. *J Health Soc Behav*. 1997; 38(1):21–37. [PubMed: 9097506]
37. McGee DL, Liao Y, Cao G, Cooper RS. Self-reported health status and mortality in a multiethnic US cohort. *Am J Epidemiol*. 1999; 149(1):41–46. [PubMed: 9883792]
38. Morrison RS, Zayas LH, Mulvihill M, et al. Barriers to completion of health care proxies: an examination of ethnic differences. *Arch Intern Med*. 1998; 158(22):2493–2497. [PubMed: 9855388]

39. Centers for Disease Control and Prevention. [Accessed April 6, 2006] Physical activity for everyone: Recommendations: Are there special recommendations for older adults? (on-line). 2006. Available at: http://www.cdc.gov/nccdphp/dnpa/physi-cal/recommendations/older_adults.htm
40. The American Geriatrics Society. [Accessed Sep 26, 2005] Clinical guidelines for alcohol use disorders in older adults (on-line). 2003. Available at: <http://www.americangeriatrics.org/products/positionpapers/alcohol.shtml>
41. Peng C-YJ, Lee KL, Ingersoll GM. An introduction to logistic regression analysis and reporting. *The Journal of Educational Research*. 2002; 96(1):3–14.
42. Levenson MR, Aldwin CM, Spiro A 3rd. Age, cohort and period effects on alcohol consumption and problem drinking: findings from the Normative Aging Study. *J Stud Alcohol*. 1998; 59(6): 712–722. [PubMed: 9811093]
43. Perreira KM, Sloan FA. Life events and alcohol consumption among mature adults: a longitudinal analysis. *J Stud Alcohol*. 2001; 62(4):501–508. [PubMed: 11513228]
44. Green CA, Perrin NA, Polen MR. Gender differences in the relationships between multiple measures of alcohol consumption and physical and mental health. *Alcohol Clin Exp Res*. 2004; 28(5):754–764. [PubMed: 15166650]
45. Liu S, Serdula MK, Williamson DF, et al. A prospective study of alcohol intake and change in body weight among US adults. *Am J Epidemiol*. 1994; 140(10):912–920. [PubMed: 7977278]
46. Wannamethee SG, Shaper AG. Alcohol, body weight, and weight gain in middle-aged men. *Am J Clin Nutr*. 2003; 77(5):1312–1317. [PubMed: 12716687]
47. Mundt MP, French MT, Roebuck MC, et al. Brief physician advice for problem drinking among older adults: an economic analysis of costs and benefits. *J Stud Alcohol*. 2005; 66(3):389–394. [PubMed: 16047528]
48. Fleming M, Brown R, Brown D. The efficacy of a brief alcohol intervention combined with %CDT feedback in patients being treated for type 2 diabetes and/or hypertension. *J Stud Alcohol*. 2004; 65(5):631–637. [PubMed: 15536773]

Table 1

Drinking Patterns of Demographic Subgroups Among Adults Ages 65 to 90

Age	Drinking Pattern				P
	Never Drank as Adult	12-month Abstinent	Moderate Drinker	Overlimit Drinker	
Age					
Men (%)					<.001
65–74 (n = 1377)	12.3	14.3	59.0	14.5	
75–79 (n = 1237)	12.6	16.9	60.7	9.8	
80–90 (n = 467)	18.5	15.9	58.9	6.7	
Women (%)					.002
65–74 (n = 1430)	27.2	9.6	43.3	19.8	
70–79 (n = 1210)	29.8	12.1	43.8	14.3	
80–90 (n = 395)	30.4	13.9	45.6	10.1	
Ethnicity					
Men (%)					<.001
White (n = 2554)	11.6	14.3	61.2	12.9	
African American (n = 143)	12.0	28.9	51.2	7.9	
Latino (n = 180)	15.5	19.1	50.0	15.4	
Asian (n = 204)	36.9	12.2	48.8	2.1	
Women (%)					<.001
White (n = 2470)	24.4	11.3	45.8	18.6	
African American (n = 157)	43.3	16.8	32.8	7.1	
Latino (n = 185)	27.6	9.9	45.8	16.6	
Asian (n = 223)	63.0	5.7	30.0	1.3	
Relationship Status					
Men (%)					.015
In relationship (n = 2437)	13.8	14.3	60.6	11.3	
Single (n = 615)	12.0	17.7	55.0	15.3	
Women (%)					.054
In relationship (n = 1453)	29.9	9.4	43.7	17.0	
Single (n = 1555)	26.9	12.7	44.2	16.3	

Education level	Drinking Pattern				P
	Never Drank as Adult	12-month Abstinent	Moderate Drinker	Overlimit Drinker	
Men					
HS graduate (n = 980)	16.9	19.5	52.7	10.9	<.001
Some college or higher (n = 2080)	11.9	13.3	62.2	12.6	
Women					
HS graduate (n = 1300)	37.3	13.4	36.0	13.3	<.001
Some college or higher (n = 1072)	21.4	9.3	50.1	19.1	

Note. N = 6060 to 6116. "Moderate drinker" equals 1 drink per day for women, 2 drinks per day for men; "over limit" equals > 1 drink per day for women or > 2 drinks per day for men. "Single" includes widowed, divorced, or never married. "In relationship" includes married or in a relationship but not married.

P-values determined by Rao-Scott Chi-Square test.

Prevalence of Medical Conditions, Self-reported Health and Behavioral Health Risks Among Men and Women Ages 65 to 90 in 4 Drinking Pattern Categories

	Men (N=3081)				Women (N=3035)			
	Never Drank as Adult (n=405)	12 month Abstinent (n=482)	Mod erate Drinker (n=841)	Over limit Drinker (n=353)	Never Drank as Adult (n=861)	12 month Abstinent (n=344)	Mod erate Drinker (n=1325)	Over limit Drinker (n=505)
Medical Condition (%)								
Heart problems	25.7	28.9	27.5	21.2 ^b	20.2	25.8 ^a	14.7	8.9 ^b <.001
HTN	53.0	51.8	47.8	51.6	60.4	57.5	49.8	43.2 <.001
High cholesterol	34.9	35.2	39.2	32.8 ^b	33.8	30.5	27.3	20.0 ^b <.001
Arthritis	17.3	16.0	18.2	18.4	30.7	36.0 ^a	28.4	24.5 .012
Diabetes	24.8	22.8 ^a	14.3	12.6	18.3	21.4 ^a	8.2	5.7 <.001
Depression	6.9	10.0 ^a	5.7	5.7	11.0	20.3 ^a	11.5	14.3 <.001
Anxiety	3.0	4.3 ^a	1.9	2.4	5.3	8.0	5.9	5.3 .398
Fair/Poor Health	29.7	30.3 ^a	14.5	16.1	29.5	43.9 ^a	16.0	13.1 <.001
Behavioral Health Risk (%)								
Current smoker	3.7	11.9 ^a	6.8	15.8 ^b	4.1	11.5 ^a	6.0	13.8 ^b <.001
Sedentary	14.6	16.1 ^a	10.3	11.0	21.5	32.7 ^a	14.4	19.4 ^b <.001
Doesn't eat low fat foods	40.7	41.2	44.5	56.0 ^b	40.3	45.9	40.1	44.1 .280
BMI 30	16.8	19.4	18.4	25.6	25.0	27.3	21.9	15.9 ^b .002

Note. P-values indicate overall significance of the difference in distribution of health variables among drinking categories, determined by Rao-Scott Chi-Square test. Post hoc comparisons used logistic regression to examine effect of drinking level (a = 12-month abstinent vs moderate, b = overlimit vs moderate) on health variables, controlling for age and ethnicity. Results marked "a" or "b" are significant at <.05 level.

12-month abstinent" = participants who have not consumed alcohol in the past 12 months. "Moderate drinker" = 1 drink per day for women, 2 drinks per day for men, "over limit" = >1 drink per day for women or >2 drinks per day for men. Sedentary = exercise <once per week. HTN = high blood pressure. BMI = body mass index.

Table 3

Logistic Regression Analysis of Predictors of Having Quit Drinking (12-Month Abstinence) in Men and Women

	OR	95% CI	P
Men (N = 2490)			
Race (African American vs others)	2.2	(1.4, 3.7)	.002
Lower education	1.5	(1.2, 2.0)	<.001
Diabetes	1.6	(1.2, 2.2)	.002
Worse self-reported health	2.2	(1.7, 2.9)	<.001
Depression	1.6	(1.3, 2.4)	.038
Women (N = 2148)			
Race (African American vs white)	1.6	(0.9, 2.9)	.073
Lower education	1.9	(1.5, 2.6)	<.001
Heart problems	1.7	(1.2, 2.5)	.003
Diabetes	2.2	(1.5, 3.3)	<.001
Worse self-reported health	3.4	(2.5, 4.6)	<.001

Note. Analysis compares 12-month abstinence to current drinkers (light or overlimit combined) and excludes individuals who never drank as adults or who had missing data on any of the predictors.

“Worse self-reported health” = fair or poor vs good, very good or excellent. “Lower education” = high school graduate or less.