Published in final edited form as:

Tetrahedron Lett. 2009 April ; 50(15): 1723–1726. doi:10.1016/j.tetlet.2009.01.133.

A new strategy for the synthesis of β -benzylmercaptoethylamine derivatives

Subrata Ghosh and Gregory P. Tochtrop*

Department of Chemistry, Case Western Reserve University, 10900 Euclid Avenue, Cleveland, Ohio, 44106, USA.

Abstract

Here we describe a new experimental approach to the synthesis of the β -benzylmercaptoethylamine functionality and illustrate its synthetic utility in multi-component reactions. Although prevalent in modern organic synthesis, no general methods have been described for this functionality. Using a carefully developed LiOH-water-ethanol reaction mixture we were able to produce a diverse collection of β -benzylmercaptoethylamines containing a range of sensitive functional groups in excellent yields. To further illustrate their utility in molecular library synthesis, we also report the use of β -benzylmercaptoethylamines in five different multi-component reactions.

The β -benzylmercaptoethylamine functionality and its derivatives are widely utilized with applications in organic,¹ inorganic,² medicinal,³ and cosmetic chemistry.⁴ For example, this functionality played a critical role in the synthesis of pantetheine (a cysteamine derivative of pantothenic acid–Vitamin B₅).^{3a} Reisner reported the synthesis of D/L- γ -sulfamyl- α -amino acids as potential anti-metabolites utilizing β -benzylmercaptoethylamine as a key intermediate.^{3b} When Okarvi *et al.* was exploring an alternative of ¹³¹I-hippuran (a renal radiopharmaceutical), they synthesized a number of S-protected derivatives of MAG3 (mercaptoacetyl triglycine) and discovered that the *p*-methoxy derivative of β -benzylmercaptoethylamine was a key starting material.^{3l} Finally, in order to prove the necessity of the disulfide unit present in a series of psammaplin A type antibacterial agents (prepared by a novel combinatorial disulfide exchange strategy), Nicolaou *et al.* used β -benzylmercaptoethylamine as a thioether containing primary amine coupling partner.^{3k}

Despite the wide reaching implications of this functionality, the disparate methods for producing it have afforded varying yields, and a systematic synthetic study is lacking. Some conditions that have been utilized with varying degrees of success include NaOMe-methanol,^{3a} hydrazine hydrate-methanol,^{3b} thiourea-NaOH-ethanol,^{1e} ethyleneimine-ethanol,^{1b} K₂CO₃-ethanol,^{1c,3k} NaOEt-ethanol,^{1d} Na/liq. NH₃,^{1h} NaI-NaOEt,^{2c} TFA-CH₂Cl₂,^{2f} and NaOH-ethanol.³ⁱ These methods suffer from various disadvantages such as long reaction times,^{1b,2f,3i,k} refluxing conditions,^{1b,3a,b} the use of moisture sensitive

© 2009 Elsevier Ltd. All rights reserved.

* Corresponding author. Tel.: +1 2163682351; fax: +1 2163683006; tochtrop@case.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary Material: Experimental procedure, NMR spectra, high resolution mass spectra and compound characterization data are available for all molecules.

solvents and reagents,^{1h} extensive work-up procedures,^{2f} or a two step reaction procedure.^{1e}

During the course of synthesizing a small library of molecules to modulate the secretion of the A β peptide by a neuronal cell line, we became interested in a specific molecule that contained the β -benzylmercaptoethylamine functionality. It was during these studies that we identified the need to develop the method described here.

In our initial consideration of this reaction, and after a careful evaluation of previously reported reaction conditions, we hypothesized that the general reaction (depicted in Scheme 1) could possibly be proceeding via a borderline/S_N1-type reaction mechanism, in contrast to the prototypical S_N2-type pathway. Thus, we envisioned that a careful optimization of both the solvent conditions and the base used would be critical for developing a superior method. Towards the first point, and lending support to our borderline mechanism hypothesis, we were able to identify mixtures of water and ethanol as ideal for smooth conversions. Further lending support to our hypothesis, we found that simple ionic bases further facilitated smooth conversions (Scheme 1).

Here we report a LiOH-water-ethanol reaction system that, with the corresponding benzyl chlorides/bromides and cysteamine hydrochloride, affords excellent yields, rapid reaction times, and a facile workup.⁵ We have subsequently tested this methodology utilizing a wide range of substrates, and we illustrate the utility of these β -benzylmercaptoethylamines in combinatorial chemistry and diversity oriented synthesis through five separate multi-component reactions (MCRs).

Table 1 illustrates a sampling of the reaction conditions we studied. Our initial exploration focused primarily on alcoholic solvents; however we quickly determined that varying amounts of water played an important role in both reaction time and yield. It is likely that the 3:1 ratio of ethanol:water provided a balance between solvent dielectric for the reaction pathway while still allowing the reagents to be soluble in the media. Keeping the ethanol:water ratio fixed, we subsequently screened the effect of various alkali metal and organic bases to find LiOH the most effective.

Using this approach, not only were our yields superior to those previously reported, but we were also able to accomplish the conversions in a minimal amount of time (approximately 40 minutes) and at low temperatures (35°C). For the reactions listed in Table 1 we utilized 3-chlorobenzyl chloride as the representative benzyl halide mainly due to its importance in the molecules we were synthesizing to modulate A β secretion.

After our careful optimization, we applied this methodology to a wide range of substrates summarized in Table 2 using a uniform set of reaction conditions.⁵ Interestingly, we obtained comparatively lower yields for the nitro derivatives. Among all other nitro derivatives, *p*-nitro (entries **12** and **13**, Table 2) gave the lowest yield, further substantiating a borderline/S_N1-type mechanism, where at least a partial charge builds at the benzylic position. Further, the best yields were obtained for the substrates containing an electron donating functional group such as -OCH₃ (entry **3**, Table 2). It is worth pointing out that the reaction was highly compatible with a number of functional groups including -OCH₃, -NO₂, halogens, -CN, -CO₂CH₃, -CH₃, -CF₃. Benzyl chlorides and bromides gave similar yields (compare entries **1** and **2**, **5** and **6**, **12** and **13**; Table 2).

In conjunction with our efforts to synthesize small libraries that modulate A β -peptide excretion we became further interested in exploiting β -benzylmercaptoethylamines in MCRs, as they create significant possibilities for molecular diversity in one step, affording more economical synthetic approaches as compared to linear syntheses.⁶ We explored five

separate MCRs with aims of generating a broad swath of diverse small molecules quickly (Scheme 2). We first explored α -aminophosphonates due, in large part, to their well documented, diverse biological activities.⁷ As a first trial, we synthesized the novel α -aminophosphonate **23** by following a literature procedure utilizing 2-(3-nitrobenzylsulfanyl)-ethylamine as the amine component. The reaction proceeded smoothly, affording a similar yield (70%) to those previously reported utilizing the Zr(IV) catalyst.⁸ Because of the well known therapeutic value of barbiturates,⁹ next we turned our attention towards the synthesis of **24** via a three component reaction using 2-benzylsulfanyl-ethylamine. Our 68% yield is in line with the 72%-90% yields previously reported in the water-based solvent system.¹⁰ Due to the prevalence of the thiazoles across diverse pharmaceutical targets,¹¹ we utilized the methodology of Yavari *et al.*¹² to synthesize **25** via a four component reaction with 2-(3-chlorobenzylsulfanyl)-ethylamine.¹² The reaction proceeded with reasonable yield (65%) after 24 hours. We further explored the synthesis of novel bis-2,3-dihydroquinazolin-4(1*H*)-one derivatives **26** utilizing a three component reaction following a literature procedure.¹³ As compared to precedent, our substrate proved as effective a reagent in the *p*-toluenesulfonic acid catalyzed reaction affording **26** in 50% yield. Finally, since its discovery, the 'Ugi four-component reaction' is one of the best known, and most popular multicomponent reactions. In this reaction an aldehyde/ketone, an amine, a carboxylic acid, and an isocyanide form an α -acylamino carboxamide through a one pot condensation.¹⁴ The final application is illustrated by the formation of **27**, derived from 2-(4-methoxybenzylsulfanyl)-ethylamine, in 72% yield.^{14c}

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (AG015885 & HL053315).

References

- 1(a). Marshall K. Can. J. Chem. 1959; 37:325.(b) Chu SH, Mautner HG. J. Org. Chem. 1961; 26:4498.
(c) Johnston TP, Gallagher A. J. Org. Chem. 1963; 28:1305.(d) Carroll FI, Dickson HM, Wall ME. J. Org. Chem. 1965; 30:33.(e) Aroyan AA, Ovsepyan TR. Armyanskii Khimicheskii Zhurnal. 1968; 21:858.(f) Bewick A, Mellor JM, Owton WM. J. Chem. Soc., Perkins Trans. 1. 1985:1039.(g) Vyas DM, Benigni D, Partyka RA, Doyle TW. J. Org. Chem. 1986; 51:4307.(h) Weintraub PM, Miller FP, Wiech NL. Heterocycles. 1987; 26:1503.(i) Walker MA, Heathcock CH. J. Org. Chem. 1992; 57:5566.(j) Kumar S, Saini R, Sing H. J. Chem. Soc., Perkins Trans. 1. 1992:2011.(k) Akaji K, Kuriyama N, Kiso Y. J. Org. Chem. 1996; 61:3350.(l) Tan DS, Foley MA, Shair MD, Schreiber SL. J. Am. Chem. Soc. 1998; 120:8565.(m) Tan DS, Foley MA, Stockwell BR, Shair MD, Schreiber SL. J. Am. Chem. Soc. 1999; 121:9073.(n) Zhang Z, Martell AE, Motekaitis RJ, Fu L. Tetrahedron Lett. 1999; 40:4615.(o) Petra DGI, Kamer PCJ, Spek AL, Schoemaker HE, van Leeuwen PWNM. J. Org. Chem. 2000; 65:3010. [PubMed: 10814191] (p) Rombouts FJR, Fridkin G, Lubell WD. J. Comb. Chem. 2005; 7:589. [PubMed: 16004503] (q) Angelosante JK, Lewis BJ, Cooper LE, Swanson RA, Daley CJA. Phosphorus, Sulfur, and Silicon. 2006; 181:2263.
- 2(a). Kennard GJ, Deutsch E. Inorg. Chem. 1978; 17:2225.(b) Mirza SA, Pressler MA, Kumar M, Day RO, Maroney MJ. Inorg. Chem. 1993; 32:977.(c) Hay RW, Galyer AL. Transition Met. Chem. 1997; 22:97.(d) Gallo V, Mastrorilli P, Nobile CF, Braunstein P, Englert U. Dalton Trans. 2006:2342. [PubMed: 16688322] (e) Carson EC, Lippard SJ. J. Inorg. Biochem. 2006; 100:1109. [PubMed: 16439023] (f) Smith AL, Day CS, Que L Jr. Zhou Y, Bierbach U. Inorg. Chim. Acta. 2007; 360:2824.
- 3(a). Walton E, Wilson AN, Holly FW, Folkers K. J. Am. Chem. Soc. 1954; 76:1146.(b) Reisner DB. J. Am. Chem. Soc. 1956; 78:5102.(c) Moffatt JG, Khorana HG. J. Am. Chem. Soc. 1961; 83:663.

(d) Tucker H, Coope JF. J. Med. Chem. 1978; 21:769. [PubMed: 29123] (e) Sterk GJ, Van der Goot H, Henk T. Eur. J. Med. Chem. 1986; 21:305. (f) Sami SM, Remers WA, Bradner WT. J. Med. Chem. 1989; 32:703. [PubMed: 2918519] (g) Hardy GW, Lowe LA, Mills G, Sang PY, Simpkin DSA, Follenfant RL, Shankley C, Smith TW. J. Med. Chem. 1989; 32:1108. [PubMed: 2709377] (h) Buschauer A, Lachenmayr F, Schunack W. Pharmazie. 1991; 46:840. [PubMed: 1687832] (i) Oya S, Plössl K, Kung MP, Stevenson DA, Kung HF. Nucl. Med. Biol. 1998; 25:135. [PubMed: 9468028] (j) Zhuang ZP, Kung MP, Mu M, Hou C, Kung HF. Bioconjugate Chem. 1999; 10:159. (k) Nicolaou KC, Hughes R, Pfefferkorn JA, Barluenga S. Chem. Eur. J. 2001; 7:4296. [PubMed: 11686610] (l) Okavi SM, Torfs P, Adriaens P, Verbruggen AM. J. Label. Compd. Radiopharm. 2002; 45:407. (m) Zhang Y, Dai X, Kallmes DF, Pan D. Tetrahedron Lett. 2004; 45:8673. (n) Masip I, Ferrándiz-Huertas C, García-Martínez C, Ferragut JA, Ferrer-Montiel A, Messeguer A. J. Comb. Chem. 2004; 6:135. [PubMed: 14714997] (o) Cleynhens BJ, de Groot TJ, Vanbilloen HP, Kieffer D, Mortelmans L, Bormans GM, Verbruggen AM. Bioorg. Med. Chem. 2005; 13:1053. [PubMed: 15670913] (p) Zhuang ZP, Kung MP, Hou C, Ploessl K, Kung HF. Nucl. Med. Biol. 2005; 32:171. [PubMed: 15721763] (q) Kieffer DM, Vanbilloen HP, Cleynhens BJ, Terwinghe CY, Mortelmans L, Bormans GM, Verbruggen AM. Nucl. Med. Biol. 2006; 33:125. [PubMed: 16459268]

- 4(a). LaPorte G. Parfumerie, Cosmetique, Savons. 1968; 11:516. (b) LaPorte G. American Perfumer and Cosmetics. 1970; 85:47.
5. Typical reaction procedure: LiOH (0.245 g, 10.2 mmol) was dissolved in 5 mL of water and 15 mL of ethanol was added. The resulting solution was added to a flask containing cysteamine hydrochloride (0.568 g, 5 mmol) followed by the dropwise addition of benzyl halides (5 mmol) with continuous stirring. The reaction mixture was stirred for 40 min at 35 °C, and ethanol was removed *in vacuo*. 20 mL of water was subsequently added and the mixture was extracted with dichloromethane (3 × 30 mL), dried over anhydrous Na₂SO₄, concentrated *in vacuo*, and purified via column chromatography over silica gel (60Å, 230 to 400 mesh, SiliCycle) using a mobile phase consisting of a suitable mixture of dichloromethane-methanol (gradient from 2% v/v methanol/dichloromethane to 20% v/v methanol/dichloromethane) to afford the chromatographically pure desired β -benzylmercaptoethylamine derivatives.
- 6(a). Gallop MA, Barrett RW, Dower WJ, Fodor SPA, Gordon EM. J. Med. Chem. 1994; 37:1233. [PubMed: 8176700] (b) Gordon EM, Barrett RW, Dower WJ, Fodor SPA, Gallop MA. J. Med. Chem. 1994; 37:1385. [PubMed: 8182695]
- 7(a). Pratt RF. Science. 1989; 246:917. [PubMed: 2814513] (b) Allen MC, Fuhrer W, Tuck B, Wade R, Wood JM. J. Med. Chem. 1989; 32:1652. [PubMed: 2661820] (c) Bonarska D, Kleszczyńska H, Sarapuk J. Cell. Mol. Biol. Lett. 2002; 7:929. [PubMed: 12378278] (d) Grembecka J, Mucha A, Cierpicki T, Kafarski P. J. Med. Chem. 2003; 46:2641. [PubMed: 12801228] (e) Skropeta D, Schwörer R, Schmidt RR. Bioorg. Med. Chem. Lett. 2003; 13:3351. [PubMed: 12951124]
8. Bhagat S, Chakraborti AK. J. Org. Chem. 2008; 73:6029. [PubMed: 18576690]
- 9(a). Bojarski JT, Mokrosz JL, Barto HJ, Paluchowska MH. Adv. Heterocycl. Chem. 1985; 38:229. (b) Foley LH, Palermo R, Dunten P, Wang P. Bioorg. Med. Chem. Lett. 2001; 11:969. [PubMed: 11327602]
10. Teimouri MB, Abbasi T, Mivehchi H. Tetrahedron. 2008; 64:10425.
- 11(a). Kumar Y, Green R, Borysko KZ, Wise DS, Wotring LL, Townsend LB. J. Med. Chem. 1993; 36:3843. [PubMed: 8254614] (b) Miwatashi S, Arikawa Y, Kotani E, Miyamoto M, Naruo K-I, Kimura H, Tanaka T, Asahi S, Ohkawa S. J. Med. Chem. 2005; 48:5966. [PubMed: 16162000] (c) Papadopoulos C, Geronikaki A, Hadjipavlou-Litina D. Farmaco. 2005; 60:969. [PubMed: 16040029] (d) Pereira R, Gaudon C, Iglesias B, Germain P, Gronemeyer H, de Lera AR. Bioorg. Med. Chem. Lett. 2006; 16:49. [PubMed: 16242326]
12. Yavari I, Hossaini Z, Shirgahi-Talari F, Seyfi S. Synlett. 2008:1631.
13. Baghbanzadeh M, Salehi P, Dabiri M, Kozezhgari G. Synthesis. 2006:344.
- 14(a). Ugi I, Meyer R, Fetzer U, Steinbrückner C. Angew. Chem. 1959; 71:386. Ugi I, Steinbrückner C. Angew. Chem. 1960; 72:267. Marcaccini S, Torroba T. Nat. Protoc. 2007; 2:632. [PubMed: 17406624] and references cited therein.

Scheme 1.

General synthesis of β -benzylmercaptoethylamine and its derivatives using a combination of LiOH-water-ethanol.

α -aminophosphonates

Barbiturates

Thiazoles

Bis-2,3-dihydroquinazolin-4(1*H*)-ones α -acylamino carboxamides **Scheme 2.**

The use of β -benzylmercaptoethylamine and its derivatives in various multicomponent reactions.

Table 1

A systematic evaluation of solvent systems and bases for the formation of β -benzylmercaptoethylamines. The reported yields are for reaction 7 from Table 2. All entries reacted for 40 minutes at 35°C.

Base used	Solvent used	Yield
LiOH	EtOH	72%
LiOH	EtOH-H ₂ O (3:1)	86%
LiOH	H ₂ O	20%
NaOH	EtOH	66%
NaOH	EtOH-H ₂ O (3:1)	78%
KOH	EtOH-H ₂ O (3:1)	75%
CsOH	EtOH-H ₂ O (3:1)	75%
K ₂ CO ₃	EtOH	30%
K ₂ CO ₃	EtOH-H ₂ O (3:1)	50%
Et ₃ N	EtOH	25%
Et ₃ N	DCM	5%

Table 2

Reaction between various benzyl halides and cysteamine hydrochloride using a combination of LiOH-H₂O-EtOH.

Entry	Benzyl halide ^a	Product	Yield ^b (%)
1	Z = H, X = Br	Z = H	90
2	Z = H	Z = H	90
3	Z = 4-OMe	Z = 4-OMe	94
4	Z = 3-OMe	Z = 3-OMe	85
5	Z = 3-F, X = Br	Z = 3-F	83
6	Z = 3-F	Z = 3-F	82
7	Z = 3-Cl	Z = 3-Cl	86
8	Z = 3-Br	Z = 3-Br	85
9	Z = 3-I	Z = 3-I	85
10	Z = 2-NO ₂	Z = 2-NO ₂	78
11	Z = 3-NO ₂	Z = 3-NO ₂	82
12	Z = 4-NO ₂ , X = Br	Z = 4-NO ₂	75
13	Z = 4-NO ₂	Z = 4-NO ₂	75
14	Z = 3-CO ₂ Me	Z = 3-CO ₂ Me	60 ^c
15	Z = 3-CN	Z = 3-CN	80 ^d
16	Z = 3-Me	Z = 3-Me	82
17	Z = 3-CF ₃	Z = 3-CF ₃	80
18	Z = 4- <i>t</i> -Butyl	Z = 4- <i>t</i> -Butyl	82
19	Z = 2,3-di-OMe	Z = 2,3-di-OMe	81
20	Z = 2,6-di-Cl	Z = 2,6-di-Cl	80
21	Z = 3-Cl-5-OMe	Z = 3-Cl-5-OMe	78
22	Z = 2-CN-3-Cl, X = Br	Z = 2-CN-3-Cl	80

^aUnless otherwise stated, X = Cl and reaction time 40 min

^b = Isolated yield.

^cReaction time was 15 min.

^dReaction time was 30 min