Published in final edited form as:

J Agric Food Chem. 2012 July 18; 60(28): 6977–6983. doi:10.1021/jf3006728.

Exploration of the antiplatelet activity profile of betulinic acid on human platelets

Andreas G. Tzacos[†], Vassiliki G. Kontogianni^{†,‡}, Maria Tsoumani^{†,‡}, Eleni Kyriakou[†], John Hwa[‡], Francisco A. Rodrigues^{||}, and Alexandros D. Tselepis^{*,†}

[†]Section of Organic Chemistry and Biochemistry, Department of Chemistry, University of Ioannina, GR-45110, Ioannina, Greece

[‡]Yale School of Medicine, Section of Cardiovascular Medicine, 300 George St, Rm 759 New Haven, CT 06511

^{||}Departamento de Matemática Aplicada e Estatística, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo-Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos, SP, Brazil

Abstract

Betulinic acid, a natural pentacyclic triterpene acid, presents a diverse mode of biological actions including anti-retroviral, antibacterial, antimalarial and anti-inflammatory activities. The potency of betulinic acid as an inhibitor of human platelet activation was evaluated and its antiplatelet profile against *in vitro* platelet aggregation, induced by several platelet agonists (Adenosine Diphosphate, Thrombin Receptor Activator Peptide-14 and Arachidonic Acid), was explored. Flow cytometric analysis was performed to examine the effect of betulinic acid on P-selectin membrane expression and PAC-1 binding to activated platelets. Betulinic acid potently inhibits platelet aggregation and also reduced PAC-1 binding and the membrane expression of P-selectin. Principal component analysis was used to screen, on the chemical property space, for potential common pharmacophores of betulinic acid with approved antithrombotic drugs. A common pharmacophore was defined between the NMR derived structure of betulinic acid and prostacyclin agonists (PGI₂) and the importance of its carboxylate group in its antiplatelet activity was determined. The present results indicate that betulinic acid has potential use as an antithrombotic compound and suggest that the mechanism underlying the antiplatelet effects of betulinic acid is similar to that of the PGI₂ receptor agonists, a hypothesis that reserves further investigation.

Keywords

betulinic acid; platelet aggregation; ADP; antithrombotics

INTRODUCTION

Thrombogenesis is a multicomponent and complex pathophysiological process that requires both humoral and cellular factors (1). Platelets play a crucial role in arterial

*Corresponding Author: A. D. Tselepis, Section of Organic Chemistry and Biochemistry, Department of Chemistry, University of Ioannina, GR-45110, Ioannina, Greece Tel: +30 26510 08365, Fax: +30 26510 08785, atselep@uoi.gr.

[‡]These authors contributed equally.

ASSOCIATED CONTENT

Supporting Information available: NMR (2D spectra, assignment, NOE correlations), list of approved antithrombotic drugs used in the study, details on the hierarchical clustering approach and resulted dendrograms. This material is available free of charge via the Internet at <http://pubs.acs.org>.

thrombosis. Platelet activation leads to shape change, secretion of granular contents and release of arachidonic acid, adhesion to the site of injury, and aggregation (2).

In order to combat a complex pathophysiological process such as thrombogenesis, multi-targeted drugs should be developed. Natural products present an important source of chemicals with such properties (3–7), with approximately half of drugs currently used in the clinic having derived from natural products (8). Natural products have evolved through natural selection to interact with multiple targets and to modulate multiple signal transduction pathways. Furthermore, natural products frequently resemble biosynthetic intermediates or endogenous metabolites, and thus can favourably utilize native active transport mechanisms. Thus, natural products present an important source for identification of multi-target compounds.

A number of studies have suggested that certain bioactive chemicals present in plants may protect against thrombosis (9–11). One such group of compounds is pentacyclic triterpenes from the lupane, oleanane and ursane groups. Pharmacological relevance has increased during the last two decades demonstrating multi-target properties combined with low toxicity (12). One of the most promising multifunctional compounds that targets multiple steps in signal transduction pathways is betulinic acid (Scheme 1) (13–15). Specifically, it has demonstrated a broad range of pharmacological actions including anti-malarial properties (16), topoisomerase inhibitory activity (17), antitumor and anticancer properties (18–20), and anti-inflammatory and anti-retroviral activity (21, 22). To the best of our knowledge, the effect of betulinic acid as an inhibitor of human platelet activation has yet to be explored. This study aims to analyze and dissect the anti-platelet effect of betulinic acid in addition to studying the potential mechanisms involved in this action.

MATERIALS AND METHODS

Chemicals

Betulinic acid and betulin were purchased from Sigma (Steinheim, Germany). DMSO- d_6 (99.8%) was purchased from Deutero (Kastellaun, Germany). Adenosine diphosphate (ADP) was purchased from Chrono-Log Corp. (Havertown, PA, USA). Arachidonic acid (AA) and Thrombin Receptor Activator Peptide-14 (TRAP) were purchased from Sigma Aldrich, St Louis, MO USA. Fluorescently-labeled monoclonal antibodies, PAC-1-FITC, anti-CD62P-PE and anti-CD61-PerCP were obtained from Beckton Dickinson (San Jose, CA, USA).

NMR experiments and modeling of betulinic acid and betulin

NMR experiments were performed at 298K on a Bruker AV-500 spectrometer equipped with a TXI cryoprobe (Bruker BioSpin, Rheinstetten, Germany). Samples were dissolved in 0.5 mL DMSO- d_6 (final concentration 5mM for both betulinic acid and betulin) and transferred to 5mm NMR tubes. The NMR system was controlled by the software TopSpin 2.1. 2D ^1H - ^1H NOESY spectra were acquired using a spectral width of 6502 Hz, acquisition time of 0.079s F_2 and 0.020s F_1 , relaxation delay of 2s, mixing time of 1000 ms ($d_8 = 1\text{s}$). Assignment was determined on the basis of 2D ^1H - ^{13}C HSQC and HMBC spectra. Betulinic acid and betulin were modeled on the basis of NMR derived distance restraints (NOEs) and available X-ray data (23).

Principal component analysis (PCA)

A total of 18 antithrombotic drugs (Figure 3) were selected for analysis. This relatively small dataset allows for identification of individual compounds in the resulting chemical space plot. A set of 8 physicochemical and stereochemical properties for all 18 compounds

were then calculated using chemaxon cheminformatic tools (24): molecular weight (MW), nitrogens (N), oxygens (O), hydrogen bond donors (HBD), hydrogen bond acceptors (HBA), topological polar surface area (tPSA), stereogenic centers (nStereo), solvent accessible / water accessible surface area. To provide a visual representation of the position of each compound in chemical space, we then carried out principal component analysis by considering the Matlab Statistical Toolbox to reduce the 8-dimensional vector corresponding to each compound to a 2-dimensional vector, with minimal loss of information. PCA can be defined as the orthogonal projection of the original data onto a lower dimensional space, called the principal subspace, such that the variance of the projected data is maximized along its first axes (25). This technique can be understood as a rotation of the axes of the original variable coordinate system to new orthogonal axes in order to make the new axes coincide with the directions of maximum variation of the original variables. Indeed, PCA also allows removing the correlations between these features in an optimum fashion. Before implementing the PCA, it is necessary to standardize the original data by subtracting the mean from each feature and dividing by its respective standard deviation. This step is necessary, since the features present different scales, which yields heterogeneous variances. The new transformed data features present zero mean and unitary variance. We have implemented the standardization procedure by considering the Matlab software. In practice, PCA consists initially of finding the eigenvalues and eigenvectors of the sample covariance matrix, obtained from the attribute matrix, i.e. a matrix whose rows represent observations (each antithrombotic drugs) and columns, each of the eight attributes. Subsequently, the eigenvectors are sorted in decreasing order according to their eigenvalues. The multiplication of the original data by the two main eigenvectors gives the data projection. All PCA analyses were performed using Matlab software. Hierarchical clustering has been also performed to identify the clusters of the antithrombotic drugs (see supporting information).

Platelet aggregation in PRP

Platelet aggregation studies in platelet-rich plasma (PRP) prepared from peripheral venous blood of apparently healthy volunteers were performed as we have previously described (26). Briefly, the platelet count of PRP was adjusted to a final platelet concentration of $2.5 \times 10^8/\text{mL}$ with homologous platelet-poor plasma (PPP). Platelet aggregation in the presence of ADP (10 μM), AA (10 μM) and TRAP (10 μM) was measured in aliquots of 0.5mL PRP, in a Chronolog Lumi-Aggregometer (model 560-Ca) at 37°C, with continuous stirring at 1200 rpm. The maximal aggregation, achieved within 3 min after the addition of each agonist, was determined and expressed as a percentage of 100% light transmission calibrated for each specimen (maximal percentage of aggregation). Betulinic acid and betulin obtained as white powder were dissolved in DMSO. The final DMSO concentration in PRP did not exceed 1% (by volume), a concentration that does not influence platelet activation (27). The inhibitory efficacy of betulinic acid was expressed as IC_{50} values (concentration that induces 50% inhibition of platelet aggregation). All aggregation studies were conducted within 3 h of blood draw.

P-selectin membrane expression and PAC-1 binding

The surface expression of CD62P (P-selectin) and the PAC-1 binding to activated platelets was studied by flow cytometry in a FACSCalibur flow cytometer (Becton-Dickinson, San Jose, CA) using a slight modification of a technique previously described (28). Briefly, platelets were incubated in the presence or in the absence of 440 μM betulinic acid or 300 μM betulin with ADP, AA or TRAP (50 mM final concentration for each agonist) for 5 min at 37°C. Platelets were then incubated with PAC-1-FITC and anti-CD62P-PE for 20 min in the dark at room temperature; diluted (1:5, v/v) with 10 mM PBS, pH 7.4 and immediately analyzed by flow cytometry (FACSCalibur, Becton-Dickinson, San Jose, CA, USA) as

previously described (29). Platelets were gated according to staining for the platelet specific antigen, CD61. The gated events were further analyzed in histograms for FL-1 for PAC-1 and FL-2 for the detection of P-selectin, respectively. Analyses included the percentage of positive events facilitated by the evaluation of mean fluorescence intensity (MFI).

RESULTS AND DISCUSSION

The effect of betulinic acid and betulin on platelet aggregation *in vitro* was studied in human platelet-rich plasma (PRP) activated by Adenosine Diphosphate (ADP), Thrombin Receptor Activator Peptide-14 (TRP) and Arachidonic Acid (AA). As shown in Table 1, betulinic acid significantly inhibited platelet aggregation induced by all agonists in a dose-dependent manner, the maximum inhibition being observed at a concentration of 440 μ M. Moreover, betulinic acid is more efficient in inhibiting platelet aggregation induced by AA and TRAP, than ADP, with significantly higher percent (%) inhibition values (Table 1) and lower IC50 values, (210 μ M, 187 μ M and 102 μ M, for ADP, AA and TRAP, respectively). Typical aggregation curves illustrating the dose-dependent inhibitory effect of betulinic acid, are presented in Figure 1A–C. In contrast to betulinic acid, betulin even at a high concentration (300 μ M) similar to the highest concentration of betulinic acid used in the present study, did not affect platelet aggregation by ADP while only a marginal inhibition was observed in platelet aggregation induced by AA and TRAP. It should be noted that we could not use higher concentration than 300 μ M for betulin due to its much lower solubility in DMSO in comparison to betulinic acid.

The above results prompted us to further investigate the inhibitory effect of betulinic acid on platelet activation by studying the conformational change of the integrin receptor α IIb/ β 3 (PAC-1 binding) and the membrane expression of P-selectin. PAC-1 is a monoclonal antibody that binds to the activated form of the integrin receptor α IIb/ β 3 (30). The activation of this integrin leads to its conformational change and the recognition of various ligands, primarily fibrinogen, resulting in platelet aggregation and further activation through α IIb/ β 3-mediated outside-in signaling (26). P-selectin is a major platelet α -granule protein that is highly expressed on the platelet surface during activation and plays significant role in platelet-leukocyte and platelet-endothelial cell interactions (31). As shown in Table 2, betulinic acid at a concentration of 440 μ M significantly inhibits PAC-1 binding and P-selectin expression induced by all agonists, maximal inhibition being observed when TRAP was used as an agonist. By contrast, betulin failed to inhibit PAC-1 binding and P-selectin expression induced by all agonists (Table 2). Representative histograms illustrating the effect of betulinic acid and betulin on PAC-1 binding and P-selectin expression induced by TRAP are shown in Figure 2A–D. The above inhibitory effects of betulinic acid, which are more potent when AA or TRAP are used as agonists compared with ADP, are in accordance with its inhibitory effects on platelet aggregation.

Having defined the potency of betulinic acid in inhibiting platelet activation, induced by 3 different agonists, we next aimed to define pharmacophores responsible for this activity. To investigate a potential overlap in coverage of biologically relevant chemical space between betulinic acid and approved antithrombotic drugs, maps of the chemical space were produced from property spaces and visualized by principal component analysis. A small database of 18 approved antithrombotic drugs (Figure 3) was constructed and grouped in five families according to their mode of action: cyclooxygenase-1 (COX-1) inhibitors, ADP receptor antagonists, prostacyclin (PGI2) IP receptor agonists, thromboxane receptor antagonists, and phosphodiesterase inhibitors. The property spaces describe 8 calculated structural and physicochemical parameters such as size, polarizability, polarity, flexibility, and hydrogen bond capacity. Principal component analysis was utilized to replot the data in a 2-dimensional format representing 84.1% of the original information in the full 8-

dimensional dataset (Figure 3). The two unitless, orthogonal axes represent linear combinations of the original 8 parameters.

Notably, PGI₂ receptor agonists cluster largely in one region of the plot, and betulinic acid belongs also in this cluster (this is in accordance to hierarchical clustering approach considering either the single or the Ward's linkage functions (see supporting information)). In contrast, the target specific ADP receptor antagonists, COX-1 inhibitors, thromboxane TP receptor antagonists and phosphodiesterase inhibitors cover a different part of the chemical space. Analysis of component loadings indicates that, in general, antithrombotic drugs feature higher polar surface area compared to betulinic acid. From the studied drugs betulinic acid is closest, in the defined chemical space, to iloprost that is an analogue of PGI₂. The recorded similarity of betulinic acid to PGI₂ and its analogues, on the chemical property space, by the use of physico-chemical descriptor metrics, prompted us to investigate the potential sampling of common pharmacophores.

PGI₂ is a potent endogenous vasodilator and inhibitor of platelet aggregation. Both actions are mediated by the IP receptor that is a G-protein coupled receptor (32). The PGI₂ synthetic analogues beraprost, iloprost, and treprostinil (Scheme 1) have been successfully used in the clinical treatment of pulmonary arterial hypertension (27). The main structural features of PGI₂ analogues illustrate common features such as a carboxylate group at C1, the α -chain; a hydroxyl group at C11, the ω -chain; and the C15-OH (Scheme 1). Previously, a common pharmacophore was constructed for human IP receptor agonists (33). The pharmacophore deduced from this study indicated the existence of three main structural characteristics: a carboxylate group that is an essential feature for all agonists, a hydrogen bond accepting and/or donating group located in a distance of 8 to 11 Å from the carboxylate group and a spacer among these groups formed by a relative and extended lipophilic area composed by aromatic or aliphatic side chains. This pharmacophore emphasized the 3D orientation of chemical functions necessary to binding to IP receptor and to express their inhibitor activity in platelet aggregation (33). The importance of the conserved carboxylate group for all IP receptor ligands led to the assumption of a hydrogen bond between this carboxylate group and Arg279 (33).

Betulinic acid, similarly to PGI₂, is composed by the following chemotypes: a hydroxyl group at C-3, a carboxylate group at C-28, an alkene group at C-20 and a pentacyclic carbon skeleton. To determine whether betulinic acid chemotypes folds in the pharmacophore model of IP receptor agonists we determined its 3D structure. The 3D architecture of betulinic acid was build on the basis of the recently determined X-ray structure of 3 β -Hydroxylup-20(29)-en-28-yl 1H-imidazole-1-carboxylate (23) and 2D ¹H-¹H NOESY NMR data. As can be seen in Figure 4, in the NMR derived architecture, betulinic acid is composed by a carboxylate group that is located in a distance of 11.2 Å from the hydroxyl group and these two chemotypes are connected by a planar hydrophobic spacer composed by a tetracyclic ring system. Interestingly, these chemical groups of betulinic acid show similar spatial rearrangement to the pharmacophore of IP receptor agonists suggesting that it could also participate in similar ligand binding interaction modes. Thus, it could be suggested that due to this similarity betulinic acid could exert its effects at least partially on the IP receptor. Our results showing that betulinic acid is more efficient in inhibiting platelet aggregation induced by AA and TRAP, than ADP are in accordance with the above suggestion. To further support the above suggestion, we examined the agonist binding features of the ligand binding pocket of the human IP receptor. The agonist binding pocket of hIP became evident after the construction of a model of iloprost bound to the homology modelled hIP receptor (34). Iloprost is a stable high affinity hIP agonist that substitutes a secondary cyclopentane ring in place of the PGI₂ oxolane ring, carries an additional C16-methyl group and a ω -chain triple bond (Scheme 1). From the iloprost-hIP receptor 3D

model the carboxylate group of iloprost was identified as a crucial site for its activity due to its participation in an ionic interaction with the highly conserved Arg279. Mutation of this residue in hIP resulted in a significant decrease in agonist binding affinity (34).

Superposition of our NMR-derived structure of betulinic acid and the previously determined structure of iloprost, in its bound form to the hIP receptor (34), emphasized a high overall structure similarity, suggesting that betulinic acid could be similarly accommodated in the same hIP binding pocket (Figure 4B). Betulinic acid could therefore accommodate successfully the same ligand binding pocket in hIP as iloprost. A direct ionic interaction between the C17-COOH of betulinic acid and R279 could be formed as also extensive hydrophobic interactions of the pentacyclic carbon skeleton of betulinic acid to the receptor similarly to iloprost. On the basis of these observations and in order to probe the potential importance of the carboxylate group of betulinic acid in its activity, we explored the antiplatelet profile of betulin (Scheme 1). Betulin lacks a carboxyl group, and according to our aforementioned hypothesis, should present reduced antiplatelet activity in comparison to betulinic acid. Indeed, betulin had no effect on ADP, Arachidonic acid, TRAP and induced platelet aggregation, even at highest concentrations. Moreover, betulin was not effective in inhibiting PAC-1 binding and P-selectin membrane expression.

In view of development of multitargeted ligands as effective antithrombotic drugs, betulinic acid could be a good lead molecule. From our study a potential pharmacophore overlay between betulinic acid and PGI₂ analogues was identified suggesting a potential common function among these molecules. Through this similarity the importance of the carboxylate group of betulinic acid was suggested and it was experimentally validated. This identified similarity between PGI₂ and betulinic acid is of importance, since PGI₂, besides being an effective antiplatelet inhibitor has been implicated, similarly to betulinic acid, in inhibiting pathways involved in the development of cancer (35–37). Although the detailed mechanism for the antiplatelet activity of betulinic acid remains to be clarified, the present results suggest that betulinic acid could be a useful antithrombotic agent.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

All the MS and NMR data were recorded in the MS and NMR centers of the University of Ioannina. Dr. Scott Gleim (Yale School of Medicine) is greatly acknowledged for providing the coordinates of the three-dimensional model of iloprost bound to the human prostacyclin receptor. Esthir Gkani Foundation (Ioannina, Greece) is acknowledged for economic support.

LITERATURE CITED

1. Fareed J, Hoppensteadt DA, Fareed D, Demir M, Wahi R, Clarke M, Adiguzel C, Bick R. Survival of heparins, oral anticoagulants, and aspirin after the year 2010. *Semin Thromb Hemost.* 2008; 34(1):58–73. [PubMed: 18393143]
2. Gawaz M. Platelets in the onset of atherosclerosis. *Blood Cells Mol Dis.* 2006; 36(2):206–210. [PubMed: 16476558]
3. Newman DJ, Cragg GM, Snader KM. Natural products as sources of new drugs over the period 1981–2002. *J Nat Prod.* 2003; 66(7):1022–1037. [PubMed: 12880330]
4. Ganesan A. The impact of natural products upon modern drug discovery. *Curr Opin Chem Biol.* 2008; 12(3):306–317. [PubMed: 18423384]
5. Janga SC, Tzakos A. Structure and organization of drug-target networks: insights from genomic approaches for drug discovery. *Mol Biosyst.* 2009; 5(12):1536–1548. [PubMed: 19763339]

6. Kyriakou E, Primikyri A, Charisiadis P, Katsoura M, Gerothanassis IP, Stamatis H, Tzakos AG. Unexpected enzyme-catalyzed regioselective acylation of flavonoid aglycones and rapid product screening. *Org Biomol Chem*. 2012; 10(9):1739–1742. [PubMed: 22261720]
7. Vasilopoulou CG, Kontogianni VG, Linardaki ZI, Iatrou G, Lamari FN, Nerantzaki AA, Gerothanassis IP, Tzakos AG, Margarity M. Phytochemical composition of "mountain tea" from *Sideritis clandestina* subsp. *clandestina* and evaluation of its behavioral and oxidant/antioxidant effects on adult mice. *Eur J Nutr*. 2011
8. Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod*. 2007; 70(3):461–477. [PubMed: 17309302]
9. Chen Y, Deuster P. Comparison of quercetin and dihydroquercetin: antioxidant-independent actions on erythrocyte and platelet membrane. *Chem Biol Interact*. 2009; 182(1):7–12. [PubMed: 19555678]
10. Jin YR, Han XH, Zhang YH, Lee JJ, Lim Y, Chung JH, Yun YP. Antiplatelet activity of hesperetin, a bioflavonoid, is mainly mediated by inhibition of PLC-gamma2 phosphorylation and cyclooxygenase-1 activity. *Atherosclerosis*. 2007; 194(1):144–152. [PubMed: 17092506]
11. Poeckel D, Tausch L, Altmann A, Feisst C, Klinkhardt U, Graff J, Harder S, Werz O. Induction of central signalling pathways and select functional effects in human platelets by beta-boswellic acid. *Br J Pharmacol*. 2005; 146(4):514–524. [PubMed: 16086030]
12. Jager S, Trojan H, Kopp T, Laszczyk MN, Scheffler A. Pentacyclic triterpene distribution in various plants - rich sources for a new group of multi-potent plant extracts. *Molecules*. 2009; 14(6):2016–2031. [PubMed: 19513002]
13. Pisha E, Chai H, Lee IS, Chagwedera TE, Farnsworth NR, Cordell GA, Beecher CW, Fong HH, Kinghorn AD, Brown DM, et al. Discovery of betulinic acid as a selective inhibitor of human melanoma that functions by induction of apoptosis. *Nat Med*. 1995; 1(10):1046–1051. [PubMed: 7489361]
14. Eichenmuller M, Hemmerlein B, von Schweinitz D, Kappler R. Betulinic acid induces apoptosis and inhibits hedgehog signalling in rhabdomyosarcoma. *Br J Cancer*. 2003; 103(1):43–51. [PubMed: 20517313]
15. Liby KT, Yore MM, Sporn MB. Triterpenoids and rexinoids as multifunctional agents for the prevention and treatment of cancer. *Nat Rev Cancer*. 2007; 7(5):357–369. [PubMed: 17446857]
16. de Sa MS, Costa JF, Krettli AU, Zalis MG, Maia GL, Sette IM, Camara Cde A, Filho JM, Giulietti-Harley AM, Ribeiro Dos Santos R, Soares MB. Antimalarial activity of betulinic acid and derivatives in vitro against *Plasmodium falciparum* and in vivo in *P. berghei*-infected mice. *Parasitol Res*. 2009; 105(1):275–279. [PubMed: 19367418]
17. Chowdhury AR, Mandal S, Mitra B, Sharma S, Mukhopadhyay S, Majumder HK. Betulinic acid, a potent inhibitor of eukaryotic topoisomerase I: identification of the inhibitory step, the major functional group responsible and development of more potent derivatives. *Med Sci Monit*. 2002; 8(7):BR254–BR265. [PubMed: 12118187]
18. Chintharlapalli S, Papineni S, Ramaiah SK, Safe S. Betulinic acid inhibits prostate cancer growth through inhibition of specificity protein transcription factors. *Cancer Res*. 2007; 67(6):2816–2823. [PubMed: 17363604]
19. Fulda S. Betulinic acid: a natural product with anticancer activity. *Mol Nutr Food Res*. 2009; 53(1):140–146. [PubMed: 19065582]
20. Takada Y, Aggarwal BB. Betulinic acid suppresses carcinogen-induced NF-kappa B activation through inhibition of I kappa B alpha kinase and p65 phosphorylation: abrogation of cyclooxygenase-2 and matrix metalloproteinase-9. *J Immunol*. 2003; 171(6):3278–3286. [PubMed: 12960358]
21. Vijayan V, Antony H, Varma RL. Betulinic acid inhibits endotoxin stimulated phosphorylation cascade and pro-inflammatory prostaglandin E(2) production in human peripheral blood mononuclear cells. *Br J Pharmacol*.
22. Evers M, Poujade C, Soler F, Ribeill Y, James C, Lelievre Y, Gueguen JC, Reisdorf D, Morize I, Pauwels R, De Clercq E, Henin Y, Bousseau A, Mayaux JF, Le Pecq JB, Dereu N. Betulinic acid derivatives: a new class of human immunodeficiency virus type 1 specific inhibitors with a new mode of action. *J Med Chem*. 1996; 39(5):1056–1068. [PubMed: 8676341]

23. Santos RC, Matos Beja A, R SJA, A PJ. 3[beta]-Hydroxylup-20(29)-en-28-yl 1H-imidazole-1-carboxylate. *Acta Cryst.* 2010; E66:o1878–o1879.
24. <http://www.chemaxon.com/>
25. Theodoridis S, Koutroumbas K. Pattern recognition. Academic Press. 2008
26. Mitsios JV, Tambaki AP, Abatzis M, Biris N, Sakarellos-Daitsiotis M, Sakarellos C, Soteriadou K, Goudevenos J, Elisaf M, Tsoukatos D, Tsikaris V, Tselepis AD. Effect of synthetic peptides corresponding to residues 313–332 of the alphaIIb subunit on platelet activation and fibrinogen binding to alphaIIbbeta3. *Eur J Biochem.* 2004; 271(4):855–862. [PubMed: 14764102]
27. Olschewski H, Simonneau G, Galie N, Higenbottam T, Naeije R, Rubin LJ, Nikkho S, Speich R, Hoeper MM, Behr J, Winkler J, Sitbon O, Popov W, Ghofrani HA, Manes A, Kiely DG, Ewert R, Meyer A, Corris PA, Delcroix M, Gomez-Sanchez M, Siedentop H, Seeger W. Inhaled iloprost for severe pulmonary hypertension. *N Engl J Med.* 2002; 347(5):322–329. [PubMed: 12151469]
28. Mitsios JV, Stamos G, Rodis FI, Tsironis LD, Stanica MR, Sakarellos C, Tsoukatos D, Tsikaris V, Tselepis AD. Investigation of the role of adjacent amino acids to the 313–320 sequence of the alphaIIb subunit on platelet activation and fibrinogen binding to alphaIIbbeta3. *Platelets.* 2006; 17(5):277–282. [PubMed: 16928597]
29. Dimitriou AA, Stathopoulos P, Mitsios JV, Sakarellos-Daitsiotis M, Goudevenos J, Tsikaris V, Tselepis AD. Inhibition of platelet activation by peptide analogs of the beta(3)-intracellular domain of platelet integrin alpha(IIb)beta(3) conjugated to the cell-penetrating peptide Tat(48–60). *Platelets.* 2009; 20(8):539–547. [PubMed: 19863457]
30. Biris N, Abatzis M, Mitsios JV, Sakarellos-Daitsiotis M, Sakarellos C, Tsoukatos D, Tselepis AD, Michalis L, Sideris D, Konidou G, Soteriadou K, Tsikaris V. Mapping the binding domains of the alpha(IIb) subunit. A study performed on the activated form of the platelet integrin alpha(IIb)beta(3). *Eur J Biochem.* 2003; 270(18):3760–3767. [PubMed: 12950259]
31. von Hundelshausen P, Weber C. Platelets as immune cells: bridging inflammation and cardiovascular disease. *Circ Res.* 2007; 100(1):27–40. [PubMed: 17204662]
32. Namba T, Oida H, Sugimoto Y, Kakizuka A, Negishi M, Ichikawa A, Narumiya S. cDNA cloning of a mouse prostacyclin receptor. Multiple signaling pathways and expression in thymic medulla. *J Biol Chem.* 1994; 269(13):9986–9992. [PubMed: 7511597]
33. Stoll F, Liesener S, Hohlfeld T, Schror K, Fuchs PL, Holtje HD. Pharmacophore definition and three-dimensional quantitative structure-activity relationship study on structurally diverse prostacyclin receptor agonists. *Mol Pharmacol.* 2002; 62(5):1103–1111. [PubMed: 12391273]
34. Stitham J, Stojanovic A, Merenick BL, O'Hara KA, Hwa J. The unique ligand-binding pocket for the human prostacyclin receptor. Site-directed mutagenesis and molecular modeling. *J Biol Chem.* 2003; 278(6):4250–4257. [PubMed: 12446735]
35. Tennis MA, Van Scoyk M, Heasley LE, Vandervest K, Weiser-Evans M, Freeman S, Keith RL, Simpson P, Nemenoff RA, Winn RA. Prostacyclin inhibits non-small cell lung cancer growth by a frizzled 9-dependent pathway that is blocked by secreted frizzled-related protein 1. *Neoplasia.* 2003; 5(3):244–253. [PubMed: 12534818]
36. Nemenoff R, Meyer AM, Hudish TM, Mozer AB, Snee A, Narumiya S, Stearman RS, Winn RA, Weiser-Evans M, Geraci MW, Keith RL. Prostacyclin prevents murine lung cancer independent of the membrane receptor by activation of peroxisomal proliferator-activated receptor gamma. *Cancer Prev Res (Phila).* 2008; 1(5):349–356. [PubMed: 19138979]
37. Keith RL, Miller YE, Hoshikawa Y, Moore MD, Gesell TL, Gao B, Malkinson AM, Golpon HA, Nemenoff RA, Geraci MW. Manipulation of pulmonary prostacyclin synthase expression prevents murine lung cancer. *Cancer Res.* 2002; 62(3):734–740. [PubMed: 11830527]

Figure 1.
Dose-response curves for betulinic acid demonstrating the inhibition of platelet aggregation induced by ADP (A), Arachidonic acid (AA) (B) and TRAP (C).

Figure 2.

Representative histograms, obtained by flow cytometry analysis, illustrating the effect of betulinic acid (A), (C) and betulin (B), (D) on PAC-1-FITC binding, and CD62P-PE membrane expression on activated with TRAP platelets, respectively.

Figure 3.

Principal component analysis (PCA) comparison of five families of 18 approved antithrombotic drugs representative of five mechanistic groups: adenosine diphosphate receptor inhibitors (orange), prostanoid prostacyclin receptor agonists (blue), COX inhibitors (purple), thromboxane inhibitors (green) and phosphodiesterase inhibitors (red). The first two principal components account for 84.1% of the original information. Clusters of antithrombotic drugs were identified by considering the hierarchical clustering approach (single linkage method) (25). The 4 prostacyclin receptor agonists cluster largely in one region of the plot (grey area).

Figure 4.

(A) NMR derived 3D structure of betulinic acid. The two chemotypes of betulinic acid, the carboxylate group and the hydroxyl group, are highlighted and their van der Waals surfaces are colored in red. (B) Superposition of the NMR derived structure of betulinic acid to the synthetic PGI2 analogue iloprost. Iloprost is shown in its bound form to the hIP receptor as was previously determined (34). Betulinic acid is colored in blue and iloprost in orange.

betulinic acid (1) **betuline (2)** **PGI₂ (3)** **iloprost (4)** **beraprost (5)** **treprostinil (6)****Scheme 1.**

2D structures of betulinic acid and betulin. The 2D structure of prostacyclin and its synthetic analogues (iloprost, beraprost, treprostinil) show common features: the C1-COOH, the α -chain, the C11-OH, the C15-OH and the ω -chain.

Table 1

Effect of betulinic acid and betulin on platelet aggregation induced by ADP, AA and TRAP.

Natural product	Concentration (μ M)	ADP ^a	AA ^b	TRAP ^c
Inhibition, %				
Betulinic acid (1)	440	32 $\pm$ 3.5	86 $\pm$ 11	80 $\pm$ 9
	220	19 $\pm$ 4.2	48 $\pm$ 7.4	60 $\pm$ 2.4
	176	8 $\pm$ 5	39 $\pm$ 2	45 $\pm$ 7
	88	5 $\pm$ 3	31 $\pm$ 2.5	38 $\pm$ 9.5
	63	0	22 $\pm$ 5	27 $\pm$ 5
	44	0	14 $\pm$ 2	16 $\pm$ 4
Betulin (2)	300	0	7 $\pm$ 2	13 $\pm$ 3

^aADP = Adenosin Diphosphate,^bTRAP = Thrombin Receptor Activator Peptide-14,^cAA = Arachidonic Acid.Data represent mean $\pm$ SD values from at least three experiments.

Table 2

Effect of betulinic acid (1) and betulin (2) on P-selectin expression and PAC-1 binding.

Natural product	PAC-1 binding	P-selectin expression					
		Inhibition, %					
		ADP	AA	TRAP	ADP	AA	TRAP
Betulinic acid (1)		30.5 ± 2.4	51.0 ± 4.5	75.9 ± 4	22.6± 2.3	57.0± 2.2	93.9± 2.6
Betulin (2)		3.4 ± 0.3	0.5± 1.4	4.9± 5.2	6.3 ± 0.4	11.0± 2.4	8.8± 3.9

Values represent the inhibitory effect of each natural product at a concentration of 440 µM and 300 µM respectively on P-selectin (CD62P-PE) membrane expression and PAC-1 (PAC-1-FITC) binding induced by ADP, AA and TRAP. Data represent mean ± SD values from at least three experiments.