CORRESPONDENCE

Local Treatment of Chronic Wounds in Patients With Peripheral Vascular Disease, Chronic Venous Insufficiency and Diabetes

by Dr. med. Mike Rüttermann, Dr. med. Andreas Maier-Hasselmann, Brigitte Nink-Grebe, Marion Burckhardt in volume 3/2013

Benefits of Hyperbaric Oxygen Therapy Still Doubtful

Rüttermann et al. were able to determine “the efficacy” from whole-body hyperbaric oxygen therapy (HBOT) on the basis of “good evidence”, demanding an unequivocal recommendation (at least grade B) for its use. In support they cite a meta-analysis by Kränke et al. from 2004 (1).

In 2012 however, Kränke et al. had updated their meta-analysis, reaching a notably different conclusion: “In people with foot ulcers due to diabetes, HBOT significantly improved the ulcers healed in the short term but not the long term and the trials had various flaws in design and/or reporting that means we are not confident in the results” (2).

While Rüttermann et al. refer to “valid data in support of hyperbaric oxygen,” Kränke et al. (2) felt compelled to comment that the relevance of hyperbaric oxygen therapy for persons with chronic wounds, after all, needs proper evaluation by studies adequately powered and designed to minimize all kinds of biases.

In the US, hyperbaric oxygen therapy completely failed in 6259 diabetes patients with foot ulcers and sufficient arterial perfusion, as was shown in the most recent—and by far most comprehensive—publication on the subject (3).

DOI: 10.3238/arztebl.2013.0372a

REFERENCES

1. Kränke P, Bennett MH: Hyperbaric oxygen therapy for chronic wounds. Cochrane Database of Systematic Reviews. Reviews 2004, Issue 1.
2. Kränke P, Bennett MH, Martyn-St James M, Schnabel A, Debus SE: Hyperbaric oxygen therapy for chronic wounds. Cochrane Database of Systematic Reviews 2012, Issue 4. Art. No.: CD004123. DOI: 10.1002/14651858.CD004123.pub3
3. Margolis DJ, Gupta J, Hoffstad O, Papadopoulos M, Glick HA, Thom SR, Mitra N: Lack of effectiveness of hyperbaric oxygen therapy for the treatment of diabetic foot ulcer and prevention of amputation: a cohort study. Diabetes Care 2013, February 19 [ahead of print].
4. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral

vascular disease, chronic venous insufficiency and diabetes. Dtsch Arztebl Int 2013; 110(3): 25–31.

em. Prof. Dr. med. Ernst A. Chantelau
Bremen
chantelau@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Hydrogel Dressings Are Obsolete

I find it impossible to understand that none of the guidelines includes as the basis for hygienic wound care soap baths or showers, which should precede any other measures. Water, even sterile saline solution would not have yielded any evidence. These would counteract any surgical hand scrubbing and any recommendation of hygiene committees for health-care staff to wash their hands after any contact with patients.

What do colleagues do in case of filthy and grossly malodorous wounds, in case the hair on the head is glued together by blood, in case of ingrown stockings, or in case of maggot or insect infestation?

The point of any lavage is to reduce any massive contamination of body cavities and skin by bacteria, fungi, or insects to a minimum. Because of the environment it is impossible to make wounds completely sterile.

By eliminating protein, fat, and detritus, the breeding ground for any pathogens is removed. Further therapies should be undertaken only after this initial step.

It is then of no crucial relevance how the wound is dressed, unless the patient has allergies. Sterile dressings and daily changes of dressings, wherever possible, are what is needed.

For this reason we do not allow hydrogel dressings in our clinic for homeless people. Some 24 hours later, specimens almost always show massive growth of mostly *Staphylococcus aureus*. In patients with venous-lymphatic stasis, compression therapy is essential.

DOI: 10.3238/arztebl.2013.0372b

REFERENCES

1. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral vascular disease, chronic venous insufficiency and diabetes. *Dtsch Arztebl Int* 2013; 110(3): 25–31.

Dr. med. Fred Reiß
Berliner Straße 99
14169 Berlin

Conflict of interest statement

The author declares that no conflict of interest exists.

Topical Honey for Diabetic Foot Ulcers

The data regarding different treatments for wounds are still disappointingly scant. No grade B recommendations are available for wound care products, and grade A seems completely out of the question! A grade B recommendation exists only in favor of hydrogel, hyperbaric oxygenation in ulcers that are beyond treatment in diabetic foot syndrome, and integrated care. In view of the many patients and the vast market in wound care products, this is sobering. I was therefore all the more surprised at the recommendation not to use honey for chronic wounds (the context was really generally that of chronic wounds) because honey resulted in more pain in a single study of venous ulcers. Why was a special scenario (venous ulcers) generalized to include all chronic wounds?

Years of experience with manuka honey in patients with diabetic foot syndrome have shown that pain usually does not constitute a problem in these wounds owing to the comorbid polyneuropathy. I would have welcomed a fact-based recommendation, differentiated by evidence level (which exists for a negative recommendation only for venous ulcers). Numerous small, qualitatively moderate to poor studies of honey treatment in diabetic foot syndrome have shown mostly promising results, which are entirely consistent with my own observations. Obviously, high-quality randomized controlled studies of honey treatment in diabetic foot syndrome are absolutely necessary and desirable. But until these have been conducted and reported, the treatment of the diabetic foot syndrome with honey provides an efficient, cost-effective variant with a poor evidence base. This, however, is also the case for all other wound treatments.

DOI: 10.3238/arztebl.2013.0373a

REFERENCES

1. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral vascular disease, chronic venous insufficiency and diabetes. *Dtsch Arztebl Int* 2013; 110(3): 25–31.

Dr. med. Roland Zerm

Gemeinschaftskrankenhaus Havelhöhe gGmbH, Berlin
roland.zerm@havelhoehe.de

Conflict of interest statement

Dr Zerm has been reimbursed for composing a poster from Neuseelandhaus, a wholesaler of manuka honey.

Wound Care Products Are not Completely Expendable

The authors lucidly explained that the evidence base for evaluating most of the currently available and used products for wound treatment is unsatisfactory. However, should the lack of evidence now lead to a situation where such wound products are regarded as not effective and will therefore not be reimbursed by the health insurers in future, then we run the risk of completely halting the progress achieved over the past decades. We would fail because of our own expectations of such medical products. The expectation is often that using a particular wound product accelerates wound healing that had stagnated before. The products should be seen for what they are—aids that can support some partial aspects of the often complex treatment process—which may include managing wound exudates or avoiding pain—very effectively and therefore greatly improve patients' quality of life (1). Many studies do not include these parameters as criteria for their objectives. Whether the wounds eventually heal or not is primarily a function of the causal treatment and not the wound products. To give an example, in venous ulcers of the lower limb it is essential whether the underlying chronic venous insufficiency (CVI) has been treated. For the treatment of CVI, the evidence is much better (2, 3). Such meta-analyses mention as early as in their introductions that modern, moist wound therapy should be the gold standard for treating such patients.

I fear that the lack of evidence for wound products may be interpreted to mean that such wound products are completely expendable. This conclusion would be erroneous, however, because some of the products can support wound healing very effectively in many respects if they are embedded in causal treatment.

DOI: 10.3238/arztebl.2013.0373b

REFERENCES

1. Wozniak G, Mauckner P, Steinsträsser L, Dissemond J: Standardisierte Wundtherapie. *Gefäßchirurg* 2011; 16: 281–92.
2. Nelson EA, Bell-Syer SE: Compression for preventing recurrence of venous ulcers. *Cochrane Database Syst Rev* 2012; 8: CD002303.
3. O'Meara S, Cullum N, Nelson EA, Dumville JC: Compression for venous leg ulcers. *Cochrane Database Syst Rev* 2012; 11: CD000265.
4. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral vascular disease, chronic venous insufficiency and diabetes. *Dtsch Arztebl Int* 2013; 110(3): 25–31.

Prof. Dr. med. Joachim Dissemond

Universitätsklinikum Essen, Klinik und Poliklinik für Dermatologie, Venerologie und Allergologie
Joachim.dissemond@uk-essen.de

Conflict of interest statement

Professor Dissemond has received consultancy honoraria from KCI, Medoderm, Urgo, Coloplast, and Draco. He has received honoraria from Lohmann & Rauscher, Draco, and Hartmann for authorship in the context of a publication that relates to the subject]. He has received honoraria for preparing scientific further educational events and has received conference participation fees, travel expenses, and hotel

expenses from 3M, B Braun, Coloplast, Draco, KCI, Lohmann & Rauscher, Customed, Systagenix, UCB-Pharma, and Urgo. He has received honoraria for conducting commissioned clinical studies from 3M, Convatec, Customed, Urgo, and Bayer.

Clear Decision Criteria Are Lacking

The guideline presented by the authors meets the formal criteria of the S3 level. The main messages, however, are based on a consensus, which reflects more closely the character of an S1 guideline. So what validity does this guideline have? This is not so much a question for the authors but for the Association of the Scientific Medical Societies in Germany (*Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften*, AWMF) itself. What kind of sense does a guideline make at an S3 level, if of 4998 literature citations analyzed, only 38 randomized controlled trials meet the criteria? Does the AWMF have mechanisms in place that stipulate a certain quantity of evidence as the basis for developing an S3 guideline, or does S3 refer mainly to the guideline's genesis but states nothing about the meaningfulness of its content?

The lack of evidence means that clear criteria for decision making are lacking in the guideline. Regarding wound care products, it says that in existing studies, none of the investigated materials (alginate, foams, acrylate, hydrocolloid, hydrofiber dressings, moist dressings, gauze) were found to have any advantage over other materials for wound healing. In selecting the materials the following criteria should have priority, in accordance with the patient's objectives and the requirements of the wound and economicalness: avoidance of pain, practicality for the patient, condition of wound margin and environment, adhesive strength, absorption and retention of exudate, allergies, and tolerability (expert consensus).

The statement that an advantage was not found for any of the materials under investigation leads to a situation where no one can advise against any of the materials, and the doctor/nurse with wound expertise has to make the selection themselves. In sum, the S3 guideline puts its trust into the qualifications and experience of the wound expert. This underlines the importance of qualified wound experts, whose education is an important aim of the "Chronische Wunden e. V. [chronic wounds reg assoc]" initiative.

DOI: 10.3238/arztebl.2013.0374a

REFERENCES

1. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral vascular disease, chronic venous insufficiency and diabetes. *Dtsch Arztebl Int* 2013; 110(3): 25–31.

Prof. Dr. med. Knut Kröger
Klinik für Gefäßmedizin
HELIOS Klinikum Krefeld GmbH, Krefeld
knut.kroeger@helios-kliniken.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank everyone who read our S3 guideline on local treatment and all those who contributed to the technical-critical discourse with their reader's letters. Professor Dr. Dissemond very aptly describes how wound care products are often credited with far too much importance and that wound healing is ultimately centrally dependent on treating the underlying cause and not on the dressings.

It would not be expedient to have a discussion around the expendability of wound care products. Because of lacking robust study results, the guideline group therefore approved "good clinical practice" recommendations that define clear requirements in terms of wound treatment (among others, see recommendations E28 and E29) and has created a basis for the care of persons with chronic wounds on the basis of systematically consented expert opinion.

Similarly, criteria to be heeded for the use of wound care products were agreed via consensus with the participating specialty societies within the AWMF. Adhering to these criteria should lead to a sensible use of wound care products in the context of patient-oriented treatment.

Lacking evidence should not incite fears but prompt insistence on an assessment of the benefit of wound care products, with end points that are of relevance to patients.

The negative recommendation for medical honey indeed resulted from a study of chronic leg ulcer, and comparable data for diabetic foot syndrome are lacking. On the basis of the existing evidence, and balancing possible strain imposed on the patient with costs, the guideline group decided on a negative recommendation.

As Dr. Zerm correctly states, a positive effect of honey on wound healing has not been documented by high-quality studies so far. This is the case for all wounds that are the subject of our guideline.

We share Dr. Zerm's desire for good evidence, not only for honey treatment.

The recommendation of the guideline, that hyperbaric oxygenation should be used as an additional therapeutic option "for patients with diabetic foot syndrome in whom all possible revascularization measures have been tried without success and the danger remains that the limb may have to be amputated," was approved on the basis of the existing evidence at the time the consensus on the recommendation was reached. When the guideline is updated the recommendation will be adapted to the new, relevant evidence.

We thank Dr. Reiß for his critical comments and share his view that expedient wound treatment requires taking into consideration aspects of hygiene, adherence to the recommendations from the Robert Koch-Institute and individual bodily hygiene as well as, for example, active periodic wound cleaning.

Nowadays, the systematic review of the evidence is seen as the gold standard for guidelines (1, 2). Because of the method used in its compilation, the guideline was

recognized by the Association of the Scientific Medical Societies in Germany (*Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften*, AWMF) as an S3 guideline according to the association's regulations (3), such as described correctly by Professor Dr. Kröger. The difference between S3 guidelines and S1 guidelines is explained on the association's website and easily accessible for everyone (3).

An S3 guideline is based on systematic review, selection, and evaluation, of the literature, according to the criteria of evidence based medicine and on a structured consensus reached in a representative body.

If no robust evidence is found, substantial uncertainty remains, which has to be articulated and dealt with.

Transparency and decision aids are needed, which are based on an explanation of the evidence and recommendations that are clearly signposted as expert opinion.

The availability of evidence also has to be considered in providing information to patients and in medical practice. In order to satisfy the transparency criterion, the steps in the process are described in detail in the methods section on the guideline development.

DOI: 10.3238/arztebl.2013.0374b

REFERENCES

1. Brouwers MC, Kho ME, Browman GP, Burgers JS, Cluzeau F, Feder G, Fervers B, Graham ID, Grimshaw J, Hanna SE, et al.: AGREE II: advancing guideline development, reporting, and evaluation in health care. *Prev Med* 2010, 51: 421–4.
2. Deutsches Instrument zur methodischen Leitlinien-Bewertung (DELBI). Fassung 2005/2006 + Domäne 8 (2008). www.leitlinien.de/leitlinienmethodik/leitlinienbewertung/delbi.
3. Das AWMF-Regelwerk Leitlinien. Ständige Kommission „Leitlinien“ der Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF). 1st edition 2012. www.awmf.org/leitlinien/awmf-regelwerk/II-entwicklung/awmf-regelwerk-01-planung-und-organisation.html.
4. Rüttermann M, Maier-Hasselmann A, Nink-Grebe B, Burckhardt M: Local treatment of chronic wounds in patients with peripheral vascular disease, chronic venous insufficiency and diabetes. *Dtsch Arztebl Int* 2013; 110(3): 25–31.

Dr. med. Mike Rüttermann

Consultant Plastic Surgeon
University Medical Center Groningen
9700 RB Groningen/NL
mikeruettermann@yahoo.com

Conflict of interest statement

The author declares that no conflict of interest exists.