Published in final edited form as:

Mutat Res. 2009 April 30; 675(0): 77–80. doi:10.1016/j.mrgentox.2009.02.002.

Reticulocyte and micronucleated reticulocyte responses to gamma irradiation: effect of age

Stephen D. Dertinger^{1,*}, Jeffrey C. Bemis¹, Souk Phonethepswath¹, Ying Tsai², Irena Nowak², Ollivier Hyrien³, James Palis⁴, and Yuhchayau Chen²

¹Litron Laboratories, Rochester, NY

²University of Rochester Medical Center, Dept. Radiation Oncology

³University of Rochester Medical Center, Dept. Biostatistics and Computational Biology

⁴University of Rochester Medical Center, Dept. Pediatrics and Center for Pediatric Biomedical Research

Abstract

The effect of age on the formation of radiation-induced micronucleated reticulocytes (MN-RETs) and reticulocytes (RETs) was investigated by exposing female C57BL/6J mice to graded doses of gamma rays from a ¹³⁷Cs source. Age at time of irradiation was 6-, 16-, or 32-weeks, and doses ranged from 0.5 to 3 Gy. A flow cytometric technique based on anti-CD71 labeling was used to measure RET and MN-RET frequencies in blood specimens collected 43 hrs post-irradiation. Mean RET frequencies declined in a dose-dependent manner for each age group. There was only one significant difference among the ages, that is, %RET were not significantly reduced in the oldest animals at 0.5 Gy, whereas this dose did have a significant effect on the other age groups. MN-RET data were more complex. Age was observed to influence the baseline frequency of MN-RET, with the oldest mice exhibiting a significantly higher mean value. Each group's %MN-RET values increased up to 1 Gy, but past this dose the frequencies plateaued or decreased. Age was observed to influence micronucleus frequency, with older mice exhibiting higher mean MN-RET values, especially at the high doses where the response was saturated (2 to 3 Gy). We hypothesize that these dissimilar responses can largely be explained by an age-related down-regulation of apoptosis whereby younger animals eliminate damaged bone marrow erythroid precursors with a greater efficiency compared with aged mice.

Keywords

Micronuclei; Radiation; Chromosomal damage; Apoptosis; Automation; Age-dependence

© 2009 Elsevier B.V. All rights reserved.

*CORRESPONDING AUTHOR S.D.D., Litron Laboratories, 200 Canal View Blvd., Rochester, NY 14623, Tele: 585-442-0930; fax: 585-442-0934; sdertinger@litronlabs.com.

Conflict of Interest Statement

Disclosure Statement: Several of the authors are employed by Litron Laboratories, a company that owns patents regarding flow cytometry-based scoring of micronucleated erythrocytes as described herein. Litron sells kits related to these patented methods.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

The *in vivo* erythrocyte-based micronucleus assay is extensively used to evaluate the potential of chemicals to induce cytogenetic damage. Indeed, regulatory agencies throughout the world base their assessments of chemicals' genotoxic risk on data from this assay [1-3]. Although the endpoint has been applied to the study of physical agents on a more limited basis, it is nonetheless well appreciated to be a valuable tool for studying radiation's clastogenic activity [4-8].

This laboratory has recently contributed to the radiobiology literature by reporting dose-responses and kinetic data for peripheral blood reticulocytes (RETs) and micronucleated reticulocytes (MN-RETs) in mice exposed to gamma rays from a ^{137}Cs source [9]. That study demonstrated the sensitivity of the MN-RET endpoint, as marked increases were observed at the lowest dose evaluated (0.125 Gy). We also reported an interesting phenomenon whereby a dose-related increase in the percentage of MN-RET was followed by a downturn once exposures exceeded approximately 1 Gy. Cell cycle delay leading to a suboptimal blood harvesting time was excluded as a possible reason for this observation, since varying harvest time did not provide specimens with MN-RET frequencies that exceeded those observed at 43 hrs for the 1 Gy exposed mice.

The current study was designed to extend our previous work with gamma irradiation in two respects. First, we wished to evaluate the effect of age on radiation-induced MN-RET yield. Secondly, to explore the above-noted downturn phenomenon further, we included a range of doses that were expected to saturate and even diminish the MN-RET response. The resulting data should be of interest to radiation biologists as well as investigators who utilize the micronucleus assay to study the genotoxic potential of chemicals.

2. Materials and Methods

2.1 Reagents

Reagents required for blood collection, fixation, and staining included a heparin-based anticoagulant solution, buffer solution, methanol fixative, anti-CD71-FITC, anti-CD61-PE, RNase, and propidium iodide. These reagents are all components of the commercially available kit: Mouse MicroFlow^{PLUS} Kit (Litron Laboratories, Rochester, NY). The instrument calibration standard, fixed *Plasmodium berghei*-infected erythrocytes ("malaria biostandard"), was also provided in these kits.

2.2 Animals

Female C57BL/6J mice were purchased from the Jackson Laboratories (Bar Harbor, Maine). Mice were allowed to acclimate for at least one week prior to treatment. Rodents were handled in accordance with the standards established by the U.S. Animal Welfare Acts set forth in the National Institutes of Health (NIH) guidelines. All procedures were reviewed and approved by the University of Rochester's Committee on Animal Resources. Purina Mills Rodent Chow 5001 and water were available *ad libitum*.

2.3 Treatment

At the time of treatment mice were approximately 6-, 16-, or 32-weeks old. Doses were 0.5, 1, 2, and 3 Gy, and sham irradiation was used as the negative control. Five mice were randomly selected for each group, and total body irradiation was performed with a single dose of ^{137}Cs gamma rays. The dose rate was approximately 2 Gy per minute. After irradiation, the mice were maintained in groups of 5 per cage in a pathogen-free room.

2.4 Blood Collection and Fixation

Blood was collected 43 hrs post-exposure via an incision to the tail vein after mice were warmed briefly under a heat lamp. Approximately 50 μ L of free-flowing blood was collected into tubes containing 200 μ L heparin solution. Heparinized blood was maintained at room temperature until fixation occurred (within 4 hrs).

Heparinized blood samples were fixed with ultracold methanol according to Mouse MicroFlowPLUS Kit instructions. These fixed blood specimens were returned to the ultracold freezer for storage until flow cytometric analyses were performed as described below.

2.5 Staining and Flow Cytometric Analyses

Methanol-fixed blood samples were washed and labeled for flow cytometric analysis according to procedures described in the Mouse MicroFlowPLUS Kit manual (vP4.3M). Samples were analyzed by a flow cytometer equipped with a 488-nm laser (FACSCalibur, Becton Dickinson, San Jose, CA). Anti-CD71-FITC, anti-CD61-PE, and propidium iodide fluorescence signals were detected in the FL1, FL2, and FL3 channels, respectively (stock filter sets). The gating logic used to ensure that quantitative analyses of erythrocyte subpopulations were not contaminated by other cellular or noncellular events has been described previously [10]. Calibration of the flow cytometer was accomplished by staining malaria-infected erythrocytes in parallel with test samples on each day of analysis [11-12].

Data were acquired with CellQuest Pro software (v5.2, BD-Immunocytometry Systems, San Jose, CA). The stop mode was set so that 20,000 CD71-expressing RETs per blood sample were evaluated for micronuclei. The following measurements were calculated from the resulting flow cytometric data: percent reticulocytes (%RET, an index of erythropoiesis function), and percent micronucleated reticulocytes (%MN-RET, an index of recent chromosomal damage).

2.6 Statistics

Each animal's %RET and %MN-RET measurements were used to calculate group mean and standard error values. The nonparametric Wilcoxon rank sum and Kruskal and Wallis tests were used for one-way ANOVA. The effect of age across multiple dose groups was assessed using classical two-way ANOVA, including dose and age as factors, as well as their interaction. All tests were two-sided at the 5% level of significance. All statistical analyses were conducted using SAS version 9.1 (SAS Institute, Cary, NC, USA).

3. Results

3.1 Reticulocyte Responses

The effect of total body gamma irradiation on mean %RET is shown in Figure 1. Each group exhibited dose-related reductions of RET. At the highest dose level tested, mean values were reduced by approximately 90% relative to sham-exposed mice. Age was not observed to significantly affect the RET endpoint, except in the case of 0.5 Gy. At this dose, %RET values for the 32-week old mice were indistinguishable from those exhibited by the sham controls ($P=0.60$), whereas younger mice exhibited marked reductions ($P=0.016$ for both 6-week and 16-week old mice).

3.2 Micronucleated Reticulocyte Responses

Mean MN-RET frequencies for sham-exposed animals were 0.18, 0.24, and 0.53% for 6-, 16-, and 32-week old mice, respectively. The mean value for the oldest group of mice is significantly higher than the other two when compared by the Wilcoxon test ($P=0.001$).

The induction of MN-RET by gamma rays is shown in Figure 2. Irrespective of age, MN-RET values were observed to rise with increasing dose up to 1 Gy. Whereas the frequencies for 6- and 16-week old mice were equivalent across these moderate dose levels, significantly higher values were observed for the oldest group of animals ($P < 0.0001$).

Consistent with earlier findings, at doses greater than 1 Gy we observed MN-RET frequencies to plateau or actually decline with increasing dose. In this region of the dose-response curve, MN-RET frequencies were significantly different among each of the three age groups, with successively older mice yielding higher frequencies ($P < 0.0001$).

4. Discussion

The reproducibility of flow cytometry-based MN-RET measurements is well documented [13-16]. Furthermore, our previous report on gamma ray-exposed mice demonstrated that inter-experimental reproducibility is high [9]. Thus, with this backdrop, we are confident that the differences described herein are attributable to age-related effects as opposed to inter-experimental variation.

The relationship that we observed between baseline MN-RET frequency and age is consistent with other reports that have shown a direct correlation. For instance, it is well established in the cytokinesis block lymphocyte literature that micronucleus frequency is age-dependent. For example, Peace and Succop [17] noted that in 10 out of the 11 studies they surveyed, age was positively correlated with micronucleus frequency. As to the mechanism responsible for age-related increases in spontaneous micronucleus frequencies, there is evidence that whole chromosome loss is a major factor [18,19].

Regarding radiation-induced MN-RET, the findings presented here confirmed our earlier observations that the frequency of MN-RET becomes saturated at approximately 1 Gy in C57BL mice. Cell cycle delay is not a compelling explanation for this result, since delayed blood harvest times did not lead to higher MN-RET values [9]. Rather, we attribute the downturn phenomenon primarily to apoptosis. That is, we speculate that relatively less damaged erythroblasts (with or without micronuclei) are able to proceed through the process of terminal differentiation. Conversely, those that are more severely affected, and thus more likely to develop micronuclei, are disproportionally eliminated. Support for this hypothesis comes from work by Shimura *et al.*, [20] who reported that *p53* null allele mice are hypersensitive to X-rays for induction of peripheral blood MN-RETs.

Interestingly, we observed the greatest age-related differences as MN-RET responses became saturated. Thus, if apoptosis is indeed a major factor, one is led to conclude that age-related differences in MN-RET frequency are related to aged animals' reduced capacity to apoptose radiation-damaged erythroblasts. Support for this hypothesis comes from Suh *et al.* [20], who treated rats with a direct-acting clastogen, methyl methanesulfonate. Whereas aged rats exhibited a modest increase in apoptotic hepatocytes, young rats showed a robust response. Also, Polyak and colleagues [21] observed a reduced apoptotic response of mouse lymphocytes to 5 Gy radiation as age increased.

The present data set cannot rule out alternative explanations for dissimilar MN-RET yields. These include age-related changes to the kinetics by which MN-RET appear in circulation, and/or altered efficiencies of double-strand break repair. However, these mechanisms do not explain the downturn phenomenon as readily apoptosis does. Firstly, our previous report, wherein blood harvest times were varied, does not lend support to the kinetics explanation [9]. Secondly, the notion of age-related differences in efficiency of repair is not consistent with our observations that radiation-induced frequencies of micronucleated lymphocytes are

similar across the 6 to 32 week age range (unpublished observations in this same mouse strain, C57BL/6J).

In conclusion, data presented herein confirm our observations that the MN-RET endpoint becomes saturated in female C57BL mice at a total body dose of approximately 1 Gy. Age is observed to affect MN-RET frequencies, especially at doses ≤ 1 Gy. We hypothesize that these response profiles can largely be explained by an age-related down-regulation of apoptosis whereby younger animals eliminate damaged bone marrow erythroid precursors with a greater efficiency compared with aged mice. Direct measures of apoptosis in the bone marrow-derived cells, especially erythroblasts, are needed to test this hypothesis further.

Acknowledgments

This work was funded by a Center for Medical Countermeasures against Radiation Program (CMCR) grant from the National Institute of Health/National Institute of Allergy and Infectious Disease (NIAID) (Y.C., No. U19AI067733). The contents are solely the responsibility of the authors, and do not necessarily represent the official views of the NIAID.

References

1. OECD. OECD guideline 474. Guideline for the testing of chemicals. Mammalian erythrocyte micronucleus test. 1997
2. Hayashi M, MacGregor JT, Gatehouse DG, Adler ID, Blakey DH, Dertinger SD, Krishna G, Morita T, Russo A, Sutou S. In vivo rodent erythrocyte micronucleus assay. II. Some aspects of protocol design including repeat treatments, integration with toxicity testing, and automated scoring. *Environ Mol Mutagen*. 2000; 35:234–252. [PubMed: 10737958]
3. FDA (U.S. Food and Drug Administration), Office of Food Additive Safety, Redbook. Toxicological principles for safety assessment of food ingredients. 2000. updated November 2003, www.cfsan.fda.gov/~redbook/red-toca.html
4. Walker JA, Boreham DR, Unrau P, Duncan AMV. Chromosome content and ultrastructure of radiation-induced micronuclei. *Mutagenesis*. 1996; 11:419–424. [PubMed: 8921502]
5. Lenarczyk M, Goddu SM, Rao DV, Howell RW. Biologic dosimetry of bone marrow: Induction of micronuclei in reticulocytes after exposure to 32P and 90Y. *J Nuclear Med*. 2001; 42:162–169.
6. Chang PY, Torous D, Lutze-Mann L, Winegar R. Impact of p53 status on heavy-ion radiation-induced micronuclei in circulating erythrocytes. *Mutat Res*. 2000; 466:87–96. [PubMed: 10751730]
7. Abramsson-Zetterberg L, Grawe J, Zetterberg G. The micronucleus test in rat erythrocytes from bone marrow, spleen and peripheral blood: the response to low doses of ionizing radiation, cyclophosphamide and vincristine determined by flow cytometry. *Mutat Res*. 1999; 423:113–124. [PubMed: 10029688]
8. Kagawa N, Shimura M, Takai A, Endo S, Fujikawa K. Relative biological effectiveness of fission neutrons for induction of micronucleus formation in mouse reticulocytes in vivo. *Mutat Res*. 2004; 556:93–99. [PubMed: 15491636]
9. Dertinger SD, Tsai Y, Nowak I, Hyrien O, Sun H, Bemis JC, Torous DK, Keng P, Palis J, Chen Y. Reticulocyte and micronucleated reticulocyte responses to gamma irradiation: dose-response and time-course profiles measured by flow cytometry. *Mutat Res*. 2007; 634:119–125. [PubMed: 17686648]
10. Dertinger SD, Camphausen K, MacGregor JT, Bishop ME, Torous DK, Avlasevich S, et al. Three-color labeling method for flow cytometric measurement of cytogenetic damage in rodent and human blood. *Environ Molec Mutagen*. 2004; 44:427–435. [PubMed: 15517570]
11. Tometsko AM, Torous DK, Dertinger SD. Analysis of micronucleated cells by flow cytometry. 1. Achieving high resolution with a malaria model. *Mutat Res*. 1993; 292:129–135. [PubMed: 7692249]
12. Dertinger SD, Torous DK, Hall N, Tometsko CR, Gasiewicz T. Malaria-infected erythrocytes serve as biological standards to ensure reliable and consistent scoring of micronucleated erythrocytes by flow cytometry. *Mutat Res*. 2000; 464:195–200. [PubMed: 10648906]

13. Torous DK, Hall NE, Dertinger SD, Diehl MS, Illi-Love AH, Cederbrant K, Sandelin K, Bolcsfoldi G, Ferguson LR, Pearson A, Majeska JB, Tarca JP, Hewish DR, Doughty L, Fenech M, Weaver JL, Broud DD, Gatehouse DG, Hynes GM, Kwanyuen P, McLean J, McNamee JP, Parenteau M, Van Hoof V, Vanparys P, Lenarczyk M, Siennicka J, Litwinska B, Slowikowska MG, Harbach PR, Johnson CW, Zhao S, Aaron CS, Lynch AM, Marshall IC, Rodgers B, Tometsko CR. Flow cytometric enumeration of micronucleated reticulocytes: High transferability among 14 laboratories. *Environ Mol Mutagen*. 2001; 38:59–68. [PubMed: 11473389]
14. Torous DK, Hall NE, Illi-Love AH, Diehl MS, Cederbrant K, Sandelin K, Pontén I, Bolcsfoldi G, Ferguson LR, Pearson A, Majeska JB, Tarca JP, Hynes GM, Lynch AM, McNamee JP, Bellier PV, Parenteau M, Blakey D, Bayley J, van der Leede BM, Vanparys P, Harbach PR, Zhao S, Filipunas AL, Johnson CW, Tometsko CR, Dertinger SD. Interlaboratory validation of a CD71-based flow cytometric method (MicroFlow®) for the scoring of micronucleated reticulocytes in mouse peripheral blood. *Environ Mol Mutagen*. 2005; 45:44–55. [PubMed: 15605355]
15. Dertinger SD, Bishop ME, McNamee JP, Hayashi M, Zuzuki T, Asano N, Nakajima M, Saito J, Moore M, Torous DK, MacGregor JT. Flow cytometric analysis of micronuclei in peripheral blood reticulocytes: I. Intra- and inter-laboratory comparison with microscopic scoring. *Tox Sci*. 2006; 94:83–91.
16. Torous D, Asano N, Tometsko C, Sugunan S, Dertinger S, Morita T, Hayashi M. Performance of flow cytometric analysis for the micronucleus assay—a reconstruction model using serial dilutions of malaria infected cells with normal mouse peripheral blood. *Mutagenesis*. 2006; 21:11–13. [PubMed: 16188876]
17. Peace BE, Succop P. Spontaneous micronucleus frequency and age: what are normal values? *Mutat Res*. 1999; 425:225–230. [PubMed: 10216215]
18. Catalan J, Autio K, Wessman M, Lindholm C, Knuutila S, Sorsa M, Norppa H. Age-associated micronuclei containing centromeres and the X chromosome in lymphocytes of women. *Cytogenetics and Cell Genetics*. 1995; 68:11–16. [PubMed: 7956347]
19. Norppa H, Falck GC-M. What do human micronuclei contain? *Mutagenesis*. 2003; 18:221–233. [PubMed: 12714687]
20. Shimura M, Kagawa N, Fujikawa K. High sensitivity of p53(-/-) mice to spontaneous and X-ray induction of micronucleated reticulocytes. *J Radiat Res*. 2003; 44:449.
21. Suh Y, Lee K-A, Kim W-H, Han B-G, Vijg J, Park SC. Aging alters the apoptotic response to genotoxic stress. *Nat Med*. 1997; 8:3–4. [PubMed: 11786880]
22. Polyak K, Wu TT, Hamilton SR, Kinzler KW, Vogelstein B. Less death in the dying. *Cell Death Differ*. 1997; 4:242–246. [PubMed: 16465234]

Figure 1. Mean percentage peripheral blood reticulocyte frequencies (%RET) with standard error bars are graphed as a function of gamma radiation dose. Blood samples from each of the three age groups were collected 43 hrs post-total body irradiation.

Figure 2. Mean percentage peripheral blood micronucleated reticulocyte frequencies (%MN-RET) with standard error bars are graphed as a function of gamma radiation dose. Blood samples from each of the three age groups were collected 43 hrs post-total body irradiation.