Published in final edited form as:

Ann Biomed Eng. 2012 April ; 40(4): 849–859. doi:10.1007/s10439-011-0507-y.

E-selectin ligands as mechanosensitive receptors on neutrophils in health and disease

S. D. Chase¹, J. L. Magnani², and S. I. Simon^{1,*}

¹University of California Davis, CA

²GlycoMimetics, Inc., Gaithersburg, MD

Abstract

Application of mechanical force to bonds between selectins and their ligands is a requirement for these adhesion receptors to optimally perform functions that include leukocyte tethering and activation of stable adhesion. Although all three selectins are reported to signal from the outside-in subsequent to ligand binding, E-selectin is unique in its capacity to bind multiple sialyl Lewis^x presenting ligands and mediate slow rolling on the order of a micron per second. A diverse set of ligands are recognized by E-selectin in the mouse, including ESL-1, CD44, and PSGL-1 which are critical in transition from slow rolling to arrest and for efficient transendothelial migration. The molecular recognition process is different in humans as L-selectin is a major ligand, which along with glycolipids constitute more than half of the E-selectin receptors on human polymorphonuclear neutrophils (PMN). In addition, E-selectin is most efficient at raising the affinity and avidity of CD18 integrins that supports PMN deceleration and trafficking to sites of acute inflammation. The mechanism is only partially understood but known to involve a rise in cytosolic calcium and tyrosine phosphorylation that activates p38 MAP kinase and Syk kinase, both of which transduce signals from clustered E-selectin ligands. In this review we highlight the molecular recognition and mechanical requirements of this process to reveal how E-selectin confers selectivity and efficiency of signaling for extravasation at sites of inflammation and the mechanism of action of a new glycomimetic antagonist targeted to the lectin domain that has shown efficacy in blocking neutrophil activation and adhesion on inflamed endothelium.

Keywords

E-selectin; L-selectin; Neutrophils; PSGL-1; Cell Signaling

Introduction

PMN roll slowly on E-selectin in inflamed mouse microvasculature, which is crucial for optimal immune functions of cell arrest and transmigration to sites of injury. To support this function, E-selectin ligands (ESLs) on PMN transduce signals in response to tensional forces exerted on them during rolling under the shear forces of blood flow. What is unique about E-selectin, as compared to L-selectin and P-selectin, is the redundancy of ligands through which it tethers PMN and executes a slow rolling immunosurveillance at velocities on the order of one micron per second. Another remarkable feature is that PMN rolling on E-

*Corresponding author. Tel/Fax (530) 752-1430, (530) 754-5739.

Conflict of Interest

J.L.M. is the head of research at Glycomimetics, Inc. and has a financial interest in developing the selectin antagonist GMI-1070. S.I.S. and UC Davis are participating in a Phase I–II clinical study of GMI-1070 for the treatment of vaso-occlusion in sickle cell patients, but have no other financial interest in Glycomimetics, Inc.

selectin and subsequent ESL clustering transduce signals that elicit calcium flux, activate kinases, and rapidly upshift affinity of β_2 -integrins that supports PMN deceleration and arrest. These aspects of E-selectin function make it a fascinating receptor system to study vascular mechanobiology and an intriguing receptor to target for remediation of inflammatory diseases. Sickle cell disease has emerged as one in which E-selectin plays a central role in PMN mediated vasculopathy that characterizes the painful episodes that individuals suffer. In this review we will focus on the biophysical mechanisms by which ESLs are bound and mechanotransduce signaling of integrin activation and how this process is being targeted for therapeutic advantage.

E-selectin ligands recognized on PMN

All three selectins are transmembrane proteins of similar structure with three extracellular domains: an amino-terminal lectin domain, followed by an epidermal growth factor domain, and short consensus repeat units (1–5). The glycobiologist Ajit Varki proposed a litmus test for designating a glycosylated E-selectin ligand with the predominant stipulation that they are located at the right place at the right time to provide adequate affinity or avidity to mediate specific biological recognition under physiological conditions (6). Here, we revise it to focus on the PMN and add a functional criteria: 1) it is expressed on the PMN membrane at the proper time of myeloid differentiation; 2) its removal or blockade diminishes the capacity for PMN capture and rolling under shear stress; and 3) its binding selectivity and affinity are consistent with ligand clustering and signal transduction that elicits calcium release and the onset of integrin activation (7). Common strategies to assess these criteria are to remove specific ligands on PMN, or genetically modify or remove ESLs in mouse models and then measure the alteration in capacity of E-selectin to bind to the surface and support subsequent recruitment. Such studies have demonstrated that the C-type lectin domain of E-, P-, and L-selectin all interact in a calcium-dependent manner with the $\alpha_{2,3}$ sialylated and $\alpha_{1,3}$ or $\alpha_{1,4}$ fucosylated tetrasaccharide sialyl Lewis x (sLe^x), which decorate the terminal branches of N- and O-glycans of cell-surface glycoconjugates (8–16). E-selectin is the most permissive in recognition of a wide variety of fucosylated and sialylated glycolipids and glycoproteins that can function as high affinity ligands. In contrast, P- and L-selectin preferentially recognize sugars that are expressed with sulfated tyrosine residues on ligands (17–23). On PMN, E-selectin engages P-selectin glycoprotein ligand (PSGL)-1 (24–27), L-selectin (on human and not mouse) (28, 29), E-selectin ligand (ESL)-1 (on mouse and not human) (22, 30–32), CD44 (33, 34), and $\alpha_{M}\beta_2$ integrin (CD11b/CD18, Mac-1) (35–38). In addition, E-selectin binds CD43 on lymphocytes, but it remains controversial whether it is recognized on PMN (39–43). These major ESLs are briefly reviewed.

Human L-selectin is an 80–100 kDa glycoprotein that when glycosylated properly is bound by E-selectin (28, 29, 44, 45) and P-selectin (28). L-selectin is expressed on the surface of almost all circulating leukocytes, but only that on human PMN and not lymphocytes is specifically bound by E-selectin (28). Since the lectin binding domain from either mouse or human E-selectin can recognize human L-selectin, it appears to be the glycosylation pattern that confers this specificity (28, 29). L-selectin on human PMN is decorated with sLe^x that supports interaction with E-selectin (28), but most lymphocyte L-selectin is not recognized by sLe^x-reactive mAbs (28). E-selectin binding L-selectin requires sialic acid but is not dependent on N-glycans and this may explain the inability of most lymphocytes to tether and roll (29). Treating human PMN with anti-L-selectin mAbs partially inhibits the adhesion of the PMN to E-selectin in shear flow assays (28, 29, 45–49). Thus, L-selectin supports sLe^x dependent slow rolling on E-selectin substrates, which is restricted to human PMN (29, 50). L-selectin on both human and mouse PMN can bind to PSGL-1 on an adjacent neutrophil to support homotypic adhesion and this process enables a PMN rolling on an inflamed vessel wall to capture a PMN flowing in the free stream (51, 52). This prompted us

to examine the respective efficiencies for PMN homotypic adhesion versus capture via E-selectin expressed on a cell line (300.19 cells) (53). At comparable selectin site densities, PMN capture by 300.19 cells expressing E-selectin was four-fold more efficient than L-selectin. In contrast, L-selectin was observed to support more robust PMN capture over a greater range of shear rate and shear stress than does E-selectin. These studies revealed that this higher capture efficiency of L-selectin was not mediated by a longer bond lifetime during cell collision since doublet formation via E-selectin formed bonds that lasted twice as long. Considering the primary role of E-selectin in PMN rolling and L-selectin in secondary capture of PMN from the blood stream, the long lifetime of E-selectin bonds and the rapid bond formation and shear resistance of L-selectin appear to be well-adapted to these distinct functions in PMN recruitment.

PSGL-1 is a disulfide-linked mucin-like homodimeric glycoprotein consisting of two 120 kDa transmembrane subunits with amino-terminal binding regions that protrudes above most of the cell surface glycocalyx (23, 26). It is expressed on all leukocytes and is bound by all three selectins (21–24, 27, 54, 55). The amino acid sequences of the human and murine transmembrane and cytoplasmic domains are highly conserved, but the extracellular domains share little sequence similarity (56, 57). PSGL-1 has few N-glycans and many more O-glycans distributed into clusters along the extracellular domain (26). E-selectin binding to PSGL-1 requires sialylated and fucosylated core-2 O-glycans (23). Unlike P-selectin, which can only bind the amino-terminal region of PSGL-1, E-selectin binds to the amino-terminal region (24, 55) and one or more additional sites on PSGL-1 (48, 55). E- and P-selectin have essentially normal binding to PSGL-1 expressed by PMN from mice deficient in fucosyltransferase (Fuc-T) IV, but PSGL-1 expressed on Fuc-TVII-deficient PMN is not bound by E- or P-selectin (58). E-selectin-mediated rolling via PSGL-1 is mostly dependent on core-2 O-glycans since sLe^x is shown to mediate more than 75% of rolling (58). However, recent studies also suggest that PMN use both core 2 and extended core 1 O-glycans for capture and rolling, which extend the variety of O-glycan modified proteins that function as essential E-selectin ligands (43).

The CD44 protein family is a polymorphic group of transmembrane glycoproteins (80–200 kDa) present on most vertebrate cells (59). Some of this heterogeneity stems from cell-specific posttranslational modifications (59), which are required for endowing E-selectin binding ability, as not all of these CD44 expressing cell types bind E-selectin (34). The sialofucosylated glycoform of CD44 known as hematopoietic cell E-/L-selectin ligand (HCELL) originally described on human hematopoietic stem cells is the most potent E-selectin and L-selectin ligand expressed on human cells (60, 61). CD44 from peripheral blood PMN of both mice and humans is bound by E-selectin if properly glycosylated (34, 62). There is no binding to CD44 from PMN of mice deficient in α 1,3 fucosyltransferase IV and VII (Fuc-TIV/VII^{-/-}) (34). CD44 is decorated with both N- and O-linked glycans (59), but E-selectin binding is mediated only by the N-glycans (22, 34, 43). Unlike PSGL-1, L-selectin, and ESL-1, which are localized to the microvilli on the neutrophil surface, CD44 is expressed on the cell body (28, 63).

Function of E-selectin ligands

The membrane organization of ESLs, typically on the tips of microvilli, is critical for their dynamic recognition under shear flow. For instance, disruption of lipid rafts abrogates PMN rolling on E-selectin without altering membrane expression levels of PSGL-1, L-selectin, or sLe^x (64). An antibody against the epitope in the amino-terminal region where PSGL-1 is bound by P-selectin significantly inhibits attachment of human PMN to E-selectin expressing Chinese hamster ovary (CHO) cells under shear flow, but has no effect on adhesion under static conditions (48). The absence of PSGL-1 alone does not increase

leukocyte rolling velocity on E-selectin *in vitro* or under inflammatory conditions *in vivo* (27, 62), but when both PSGL-1 and ESL-1 are absent rolling velocity is increased further than the absence of ESL-1 alone (62). PSGL-1 is sufficient to observe slow rolling of murine and human PMN on E-selectin and ICAM-1 co-immobilized in a shear flow chamber (65, 66). Likewise, PSGL-1, supports slow rolling of PMN on L-cells expressing E-selectin and ICAM-1, which is significantly increased at a low shear stress of 0.1 dyne/cm², but not at 2 dynes/cm² (49). In contrast, PSGL-1 does not significantly contribute to the arrest of leukocytes that are already rolling in TNF- α stimulated venules *in vivo* (62). PSGL-1 is also partly responsible for the capacity of murine PMN to extravasate into thioglycollate-induced peritonitis (62, 65). These functions are related to PSGL-1's capacity to transduce signals, since engagement was shown to increase tyrosine phosphorylation of its cytoplasmic domain and activate MAP kinases in human PMN (67). The cytoplasmic domain of PSGL-1 is linked to the cytoskeleton by ERM (ezrin-moesin-radixin) family proteins (68), which mediate the association of PSGL-1 with spleen tyrosine kinase (Syk) in an ITAM-dependent manner (69). Engagement of PSGL-1 induces tyrosine phosphorylation of Syk in the human myeloid cell line U937 (69). Similarly, E-selectin-induced LFA-1-mediated slow rolling of both murine (65) and human PMN (66) is PSGL-1 and Syk-dependent. Following activation-induced shape change and polarization of human PMN, PSGL-1 becomes clustered at the uropod and co-localizes with moesin on stimulated human PMN (68). This process is associated with weakened adhesion to P-selectin that may facilitate integrin control of adhesion and transendothelial migration (19, 50, 56, 70). L-selectin is also localized in lipid rafts (64, 71) and clustered on the tips of microvilli or membrane ruffles (23, 28, 47, 72). The epitope of L-selectin that mediates PMN tethering to E-selectin under flow conditions is in the amino-terminal lectin domain and not on the short consensus repeats (47). Thus, the accessibility of ESLs is likely critical considering that capture via E-selectin on inflamed endothelium at high shear rates ($\sim 100 \text{ s}^{-1}$) must occur within the duration of collision, perhaps as brief as ~ 10 milliseconds.

Unlike PSGL-1 and L-selectin, which can be bound by both P- and E-selectin (21, 28), ESL-1 is not bound by P-selectin and is specific to E-selectin (29, 32, 62). Immobilized ESL-1, but not murine L-selectin, supports adhesion of murine E-selectin transfected CHO cells in a calcium dependent manner (32). ESL-1 interacts with E-selectin through presentation of sLe^x on some of its five N-glycans (22, 29) rather than via O-glycans like PSGL-1 and L-selectin (23, 29). In contrast to PSGL-1 and L-selectin, which are expressed only on the microvilli, most ESL-1 is found in the Golgi apparatus and a relatively smaller fraction is expressed on the cell surface where it is localized on the microvilli (19, 23, 28, 31, 47, 56, 64, 72). Fucosylation of ESL-1 is essential for E-selectin binding (32). CHO cells transfected with cDNA for ESL-1 alone or in combination with cDNA for human Fuc-TIII only become competent to bind E-selectin if Fuc-TIII is expressed in these cells (32). This selective generation of ligand activity correlates with the upregulation of the HECA-452-reactive carbohydrate epitope, which is functionally correlated with E-selectin ligand activity (33, 73–75). Although the absence of ESL-1 alone does not affect leukocyte rolling frequency, it is dramatically reduced when both ESL-1 and PSGL-1 are absent suggesting both are important in mediating the initial leukocyte tethering and rolling *in vivo* (62). In fact, the ability of soluble E-selectin to bind murine PMN deficient in both ESL-1 and PSGL-1 is completely abrogated suggesting cooperation for high-affinity binding (62). ESL-1 also cooperates with PSGL-1 and CD44 in controlling rolling velocities and is responsible for steady rolling; without ESL-1 murine leukocytes exhibit a “skipping” behavior (62). ESL-1 is important in the transition from rolling to firm arrest and the absence of ESL-1 shows a significant but moderate reduction in the ability of murine neutrophils to extravasate (62). PSGL-1 is required for E-selectin mediated slow rolling in an autoperfused flow-chamber system (65), however on E-selectin expressed on inflamed endothelium *in vivo*, CD44 controls rolling velocity to enable slow leukocyte rolling (34,

62). CD44 also plays a major role in the transition of rolling leukocytes to firm arrest *in vivo* (62) and cooperates with PSGL-1 in PMN extravasation into thioglycollate-induced peritonitis and staphylococcal enterotoxin A-injected skin pouch (34). Less is known regarding the cooperative process by which L-selectin, PSGL-1 and other ESLs support rolling and transduce signals in human PMN, especially *in vivo*.

Tethering and clustering of ESLs signal under shear flow

E-selectin engagement under shear flow induces the redistribution and co-localization of PSGL-1 and L-selectin into clusters on the trailing edge of human PMN (50). This has also been observed in mice treated with TNF- α where a large fraction of rolling leukocytes exhibit L-selectin polarization and co-localization with PSGL-1 in membrane clusters (62). This E-selectin-dependent redistribution of L-selectin on rolling leukocytes *in vivo* appears to involve CD44, which upon engagement can itself trigger p38 MAPK dependent clustering of L-selectin and PSGL-1 (62). CD44 crosslinking also induces PSGL-1 polarization, which is completely abolished by preincubation of PMN with an inhibitor of p38 MAPK activation (62). Unlike PSGL-1 and L-selectin, which cluster on the majority of slow-rolling leukocytes, CD44 remains evenly distributed on rolling cells and in some instances is more densely distributed in areas of contact with the endothelium (30, 62). Our studies suggest that the clustering of ESLs increases the valence of E-selectin recognition and provides a means for slow rolling (50, 76). Clustering of ESLs also provides for amplification of outside-in signaling that contributes to various post-rolling innate immune functions including oxidative burst, degranulation, actin polymerization, and an upshift in CD18 affinity. Dynamic fluorescence video microscopy and total internal reflection fluorescence (TIRF) analysis of rolling and migrating PMN demonstrate that rolling on E-selectin elicits a striking focal co-localization and membrane clustering of L-selectin, PSGL-1, and ADAM-17 that progress to the trailing edge of arresting PMN, which is not apparent on cells activated in suspension (Figure 1) (76). These receptors diffuse at distinct rates under a shear stress of 2 dyne/cm², thus highlighting a well coordinated dynamic process involving raft formation, calcium transients, and cytoskeletal linkage (76). Since it is known that PSGL-1 and moesin redistribute towards the cell uropod where activated ERM proteins also assemble, it is likely that the dynamics of the membrane determinants (e.g. lipid rafts) and signaling molecules (e.g. DAG, PLC, PIP_{2,3}, PI₃K, and MAPK) coordinate the precise redistribution of ESLs. Recently it was reported that L-selectin and PSGL-1 ligation by E-selectin activate PI₃K that in turn catalyzes the phosphorylation of Vav1 and downstream F-actin redistribution (77). These data suggest that PI₃K is required for the F-actin-based cytoskeleton changes that occur during neutrophil rolling on E-selectin.

A rise in intracellular calcium concentration plays a central role in signaling downstream of chemokine ligation and adhesion function and we and others have detected Ca²⁺ flux to be an indicator of the level of PMN activation during recruitment on inflamed endothelium under shear stress (78–85). A combination of shear stress and E-selectin engagement is a requirement for transition of LFA-1 to an extended conformation and subsequent maximal activation of β_2 -integrins in rolling PMN (49)(66). Moreover, it is the superposition of activation via ESLs and G-protein coupled receptors that elicit optimal levels of calcium flux on arrested PMN (86). We have imaged intracellular Ca²⁺ flux in real time where PMN were labeled with the dye Fura-2 and perfused at 2 dyne/cm² over L-cell monolayers expressing E-selectin. The kinetics of intracellular calcium release was recorded for PMN rolling to arrest over a dose range of IL-8, added as a bolus to the inlet of the microfluidic channel as depicted in Figure 2. In the absence of chemokine, or at concentrations of IL-8 below 0.01 nM, rolling PMN elicited a small but significant Ca²⁺ flux (i.e. ~50 nM). From this baseline, the level of cortical calcium rapidly increased with IL-8 concentration for PMN rolling to arrest under 2 dynes/cm² of shear stress. For example, stimulation with 0.05

nM IL-8 in suspension yielded ~150 nM of Ca^{2+} flux in arrested PMN under shear stress (Figure 2). To activate this same extent of Ca^{2+} flux on PMN stimulated in static suspension required a 100-fold higher concentration of IL-8 (i.e. 5 nM). Thus, ESL clustering under shear stress functions to amplify the sensitivity to chemokine activation and significantly increase the extent of Ca^{2+} flux in PMN arrested via high affinity β_2 -integrins (86).

Mechanical forces acting on E-selectin ligands function in signaling via cytosolic kinase activation

E-selectin bond strength exhibits a weak dependence on the rate of loading (e.g. PMN roll at a constant velocity of ~6 $\mu\text{m/s}$ over a wide range of fluid shear stress), but these bonds are more durable than those of L- or P-selectin (47). Shear forces of blood flow acting on membrane tethers with the substrate transmit force to the bonds with ESLs and are a critical determinant in catalyzing kinase phosphorylation and down-stream integrin activation. Thus, the mechanics and kinetics of E-selectin optimize it for recognition of multiple ligands and to function as a dynamic substrate for assembly of cytoskeletal proteins and signaling kinases. An important question is how the forces acting on ESL bonds influence assembly to adaptor molecules in a process summarily referred to as mechanotransduction. We have begun to dissect how force catalyzes kinase mediated activation using dynamic real time imaging of Ca^{2+} flux, upshift in CD18 affinity, and assembly and phosphorylation of kinases with the cyto-domains of ESLs in human PMN within a region of contact we denote the inflammatory synapse (see Figure 1) (87). The importance of this process in human disease was recently reported in patients with sepsis-induced acute kidney injury that exhibited elimination of selectin-mediated slow leukocyte rolling that was associated with reduced phosphorylation of spleen tyrosine kinase, Akt, phospholipase C- γ 2, and p38 MAPK that correlated with impaired PMN rolling and transmigration (88). The mechanism by which shear forces actively transport ESLs to sites of active MAPK and Src kinase activation have yet to be elucidated. It should be possible to image the redistribution of L-selectin, PSGL-1, CD44 and other ESLs under well defined conditions of shear stress in order to directly correlate with PMN activation as detected by Ca^{2+} flux, assembly of phospho-MAP and Src kinases, and activated CD18. The shortcoming of approaches that rely on imaging phospho-specific antibodies against Src (i.e. Tyr-419) and MAPK (p38MAPK or ERK2) in fixed and permeabilized cells is that the difficulty in identifying those signals elicited by integrins or other membrane receptors engaged. One promising strategy to detect mechanosensing is to disassemble the actin cytoskeleton under defined conditions of receptor ligation. Cuvelier et al. (89) adopted such an approach and found that ligation of leukocyte adhesion molecules under flow conditions leads to mechanotransduction in endothelial cells. Treatment of endothelium with latrunculin A prevented ERK2 phosphorylation following ligation of either vascular cell adhesion molecule-1 or E-selectin with antibody coated beads under flow conditions (89). Rapid phosphorylation of focal adhesion kinase and paxillin as observed for engagement of biomimetic beads resembles that for eosinophil binding to inflamed endothelium (89). This suggests that focal adhesions are involved in signaling to endothelium, but do not elucidate how leukocyte engagement mechanotransduce such intracellular signals. An intriguing question is whether membrane redistribution of ESLs upon stimulation requires their initial release from cortical actin associated proteins such as talin, vinculin, and α -actinin under the control of calpain, as compared with other receptors (e.g. CD18, CD45). ESL diffusion into discrete membrane rafts known to be enriched with members of the Ras and Src family also should be analyzed in the context of intracellular signaling under dynamic conditions of shear stress. For example, diffusion of L-selectin/PSGL-1 and CD44 in the membrane can be assessed for PMN incubated with the chelators methyl β -cyclodextrin and filipin to disrupt cholesterol rafts. To directly image selectin-ligands redistributing within rafts, PMN

can be labeled with DiI-C18, a fluorescent indocarbocyanine dye that preferentially partitions into lipid ordered domains, and ESL co-localization analyzed using 3-color immunofluorescence as depicted above in Figure 1.

E-selectin plays a dominant role in the cellular events of vaso-occlusive crisis in sickle cell disease

Sickle cell disease (SCD) is among the most common monogenetic hematological disorders in the United States (90). It is characterized by a single amino acid substitution in the β -chain of hemoglobin, which during hypoxic conditions induces its polymerization and causes the characteristic rigid sickle shape of the red blood cells. SCD sufferers undergo periodic vaso-occlusive events in which PMN adhere to the endothelium and bind and entrap circulating sickle red blood cells eventually leading to blockage of blood flow, ischemic damage to organs and intense pain. While a genetic disorder, the underlying cause of the vaso-occlusive crisis (VOC) is a heightened inflammatory response to stimuli ranging from general stress to bacterial and viral infections. Signs of chronic vascular inflammation are common in sickle cell disease. Numerous markers of inflammation such as TNF- α (91, 92), C-reactive protein (93), and interleukins 1 and 8 (94) are elevated in sickle cell individuals and result in activation of the endothelium. Clinical studies have also shown that elevation of circulating leukocytes in patients correlates with episodes of VOC, stroke, and acute chest syndrome (95). PMN activation is also pronounced in SCD and results in increased cell adhesion, while the activation of monocytes results in the formation of platelet-monocyte aggregates (96).

Mice expressing human sickle hemoglobin in their red blood cells mimic the human disease by undergoing VOC upon inflammatory stimuli. The cellular events that initiate VOC in these mice were directly observed using the technique of intravital microscopy (90). PMN initiate the process by rolling and achieving stable arrest on the vascular endothelium. Sickle erythrocytes are observed to then bind to the immobilized PMN eventually causing further entrapment and resulting in vaso-occlusion. To explore the possible role of adhesion molecules in VOC, a sickle cell mouse model was created by bone marrow transplantation and crossed with a genetic background lacking E- or P-selectins. Leukocyte recruitment to the vessel wall is defective in these knockout mice, which affords protection from VOC (90). These studies demonstrate the critical roles of E- and P-selectins, as all of the other adhesion molecules in these sickle cell mice are present yet they do not undergo VOC.

Hidalgo et al. (97) explored the relative roles of E-selectin and P-selectin during vaso-occlusive events in a mouse model of SCD. Two critical events that occur during the crisis are the foundation of the occlusions created by the binding of RBC to immobilized leukocytes and reduced blood flow. Blocking E-selectin with antibodies or genetic deletion in knockout mice fully inhibits the adhesion of RBCs to immobilized leukocytes over four hours of observation during a vaso-occlusive event. In contrast, blocking P-selectin exhibits transient inhibitory effects on the binding of RBCs to leukocytes, which return to control values over this time period. Likewise, blocking E-selectin, but not P-selectin, has a significant effect on increasing blood flow during VOC (97). The prevalence of E-selectin function over P-selectin is also seen in sickle cell patients. Kato and colleagues at the NIH (98) measured the level of adhesion molecules in the plasma of 160 sickle cell patients over a four year period and found a strong correlation of soluble E-selectin, but not P-selectin, with mortality over that time period. Similarly, in the sickle cell mouse model a genetic knockout of E-selectin, but not P-selectin, correlates with increased survival after induction of VOC (97). In mice, endothelial E-selectin, rather than P-selectin, engagement of ligand ESL-1 on the PMN surface induces secondary signaling events that generate activated Mac-1 integrin microdomains at the leading edge of adherent PMN in TNF- α stimulated

post-capillary venules. These activated Mac-1 clusters participate in heterotypic interactions with circulating RBCs (97). This suggests that E-selectin is responsible for signaling the adherent neutrophil to capture red blood cells and leukocytes, which is a critical step in the formation of an occlusion. A mechanism by which E-selectin affects the adhesion of either RBCs or platelets to leukocytes, as gleaned from the studies of Frenette, is depicted in Figure 3. Rolling and adhesion of leukocytes on E-selectin mechanotransduces signals including an upshift of CD11b/CD18 (Mac-1) into a high affinity conformation capable of binding RBCs or platelets. This model is congruent with the findings that elimination of either E-selectin or activated Mac-1 in knockout mice reduces the formation of RBC or platelet/leukocyte aggregates (97).

Inhibition of VOC by small molecule pan selectin antagonist GMI-1070

GMI-1070 is a small molecule pan selectin antagonist currently in Phase II clinical trials for VOC in sickle cell patients. It was designed to mimic the bioactive conformation of the functional carbohydrate ligands of the selectins and to incorporate the charged groups required for a pan selectin antagonist. Although GMI-1070 is bound by all three selectins, it demonstrates an order of magnitude greater potency in binding to E-selectin and blocking ligand binding (99). Chang et al. (99) tested the ability of GMI-1070 to reverse ongoing acute VOC in the sickle cell mouse model. The protocol was designed to be a clinically relevant model in which treatment with GMI-1070 was delayed as long as possible to simulate the arrival of a sickle cell patient already in VOC at the clinic. After initiation of VOC in the sickle cell model, mice were allowed to progress in VOC for 110 minutes. Mice were then treated with GMI-1070 and shortly thereafter control untreated mice began to die. GMI-1070 was observed to reverse measures of VOC as determined by dramatic increases in blood flow and the elimination of the capture of sickle RBCs by arrested leukocytes in the treated cohort of mice. This reversal of ongoing VOC in the treated mice also increased survival. Treatment with GMI-1070 saved half of the cohort and extended survival for the remaining half compared to untreated mice. These data suggest that although both E- and P-selectin have been associated with VOC, the effects of E-selectin clearly predominate, in the mouse where inhibition of E-selectin is sufficient for full inhibitory effects of VOC. Ongoing human studies on blood samples obtained from sickle cell disease subjects participating in Phase I and II clinical studies of the safety and efficacy of GMI-1070 reveal that it is well tolerated by patients following infusion at serum concentrations shown effective in the mouse VOC studies. Moreover, in-vitro assays of blood samples to assay PMN capture on E-selectin and integrin activation show a clear inhibitory benefit of GMI-1070 at micromolar concentrations.

Summary

E-selectin is the most promiscuous of selectins in recognition of a variety of sLe^x expressing glycoprotein and glycolipid ligands on leukocytes. Its role in immune surveillance may be to amplify the sensitivity to activation with chemokine at sites of vascular inflammation and locally activate PMN arrest and migratory function. Key issues to be elucidated regarding mechanotransduction of these functions are how the direction, magnitude, and duration of the hydrodynamic forces acting on each ESL influence their respective capacity to signal in the cytosol. Remaining to be discovered is the distinction between how distinct ESLs signal during rolling on E-selectin as compared to dynamic interaction via P-selectin and L-selectin, which are less efficient at mechanotransducing a calcium flux and integrin activation. In addition, how FAK/Src-dependent and ESL-mediated p38 MAPK and ERK activation is central to the organization and localization of active Raf-MEK1-MAPK signaling complexes and other guanine nucleotide exchange factors (i.e. Vav-1) within the PMN-endothelial inflammatory synapse. The role of E-selectin in various inflammatory

diseases is coming to light and in particular its function in VOC associated with SCD. The capacity of the glycomimetic antagonist GMI-1070 to block both PMN recruitment by effectively antagonize signaling of CD18 activation and RBC capture, highlights a complex process by which ESL recognition and reorganization are associated with mechanotransduction of PMN activation.

Acknowledgments

These studies were funded by a grant to S.I.S. from the NIH AI47294.

References

1. Bevilacqua MP, Stengelin S, Gimbrone MA Jr, Seed B. Endothelial leukocyte adhesion molecule 1: an inducible receptor for neutrophils related to complement regulatory proteins and lectins. *Science*. 1989 Mar 3.243:1160. [PubMed: 2466335]
2. Johnston GI, Cook RG, McEver RP. Cloning of GMP-140, a granule membrane protein of platelets and endothelium: sequence similarity to proteins involved in cell adhesion and inflammation. *Cell*. 1989 Mar 24.56:1033. [PubMed: 2466574]
3. Lasky LA, et al. Cloning of a lymphocyte homing receptor reveals a lectin domain. *Cell*. 1989 Mar 24.56:1045. [PubMed: 2647302]
4. Siegelman MH, van de Rijn M, Weissman IL. Mouse lymph node homing receptor cDNA clone encodes a glycoprotein revealing tandem interaction domains. *Science*. 1989 Mar 3.243:1165. [PubMed: 2646713]
5. Tedder TF, et al. Isolation and chromosomal localization of cDNAs encoding a novel human lymphocyte cell surface molecule, LAM-1. Homology with the mouse lymphocyte homing receptor and other human adhesion proteins. *The Journal of experimental medicine*. 1989 Jul 1.170:123. [PubMed: 2473156]
6. Varki A. Selectin ligands. *Proceedings of the National Academy of Sciences of the United States of America*. 1994 Aug 2.91:7390. [PubMed: 7519775]
7. Nimrichter L, et al. E-selectin receptors on human leukocytes. *Blood*. 2008 Nov 1.112:3744. [PubMed: 18579791]
8. Phillips ML, et al. ELAM-1 mediates cell adhesion by recognition of a carbohydrate ligand, sialyl-Lex. *Science*. 1990 Nov 23.250:1130. [PubMed: 1701274]
9. Walz G, Aruffo A, Kolanus W, Bevilacqua M, Seed B. Recognition by ELAM-1 of the sialyl-Lex determinant on myeloid and tumor cells. *Science*. 1990 Nov 23.250:1132. [PubMed: 1701275]
10. Lowe JB, et al. ELAM-1--dependent cell adhesion to vascular endothelium determined by a transfected human fucosyltransferase cDNA. *Cell*. 1990 Nov 2.63:475. [PubMed: 1699667]
11. Tiemeyer M, et al. Carbohydrate ligands for endothelial-leukocyte adhesion molecule 1. *Proceedings of the National Academy of Sciences of the United States of America*. 1991 Feb 15.88:1138. [PubMed: 1705026]
12. Goelz SE, et al. ELFT: a gene that directs the expression of an ELAM-1 ligand. *Cell*. 1990 Dec 21.63:1349. [PubMed: 1702034]
13. Polley MJ, et al. CD62 and endothelial cell-leukocyte adhesion molecule 1 (ELAM-1) recognize the same carbohydrate ligand, sialyl-Lewis x. *Proceedings of the National Academy of Sciences of the United States of America*. 1991 Jul 15.88:6224. [PubMed: 1712483]
14. Zhou Q, et al. The selectin GMP-140 binds to sialylated, fucosylated lactosaminoglycans on both myeloid and nonmyeloid cells. *The Journal of cell biology*. 1991 Oct.115:557. [PubMed: 1717488]
15. Hynes RO, Lander AD. Contact and adhesive specificities in the associations, migrations, and targeting of cells and axons. *Cell*. 1992 Jan 24.68:303. [PubMed: 1733501]
16. Foxall C, et al. The three members of the selectin receptor family recognize a common carbohydrate epitope, the sialyl Lewis(x) oligosaccharide. *The Journal of cell biology*. 1992 May. 117:895. [PubMed: 1374413]

17. Pouyani T, Seed B. PSGL-1 recognition of P-selectin is controlled by a tyrosine sulfation consensus at the PSGL-1 amino terminus. *Cell*. 1995 Oct 20.83:333. [PubMed: 7585950]
18. Sako D, et al. A sulfated peptide segment at the amino terminus of PSGL-1 is critical for P-selection binding. *Cell*. 1995 Oct 20.83:323. [PubMed: 7585949]
19. Lorant DE, et al. Activation of polymorphonuclear leukocytes reduces their adhesion to P-selectin and causes redistribution of ligands for P-selectin on their surfaces. *J Clin Invest*. 1995 Jul.96:171. [PubMed: 7542276]
20. Imai Y, Lasky LA, Rosen SD. Sulphation requirement for GlyCAM-1, an endothelial ligand for L-selectin. *Nature*. 1993 Feb 11.361:555. [PubMed: 7679207]
21. Asa D, et al. The P-selectin glycoprotein ligand functions as a common human leukocyte ligand for P- and E-selectins. *The Journal of biological chemistry*. 1995 May 12.270:11662. [PubMed: 7538120]
22. Lenter M, Levinovitz A, Isenmann S, Vestweber D. Monospecific and common glycoprotein ligands for E- and P-selectin on myeloid cells. *The Journal of cell biology*. 1994 Apr.125:471. [PubMed: 7512971]
23. Bruehl RE, Springer TA, Bainton DF. Quantitation of L-selectin distribution on human leukocyte microvilli by immunogold labeling and electron microscopy. *J Histochem Cytochem*. 1996 Aug. 44:835. [PubMed: 8756756]
24. Moore KL, et al. The P-selectin glycoprotein ligand from human neutrophils displays sialylated, fucosylated, O-linked poly-N-acetylglucosamine. *The Journal of biological chemistry*. 1994 Sep 16.269:23318. [PubMed: 7521878]
25. Moore KL, et al. Identification of a specific glycoprotein ligand for P-selectin (CD62) on myeloid cells. *The Journal of cell biology*. 1992 Jul.118:445. [PubMed: 1378449]
26. Norgard KE, et al. Characterization of a specific ligand for P-selectin on myeloid cells. A minor glycoprotein with sialylated O-linked oligosaccharides. *The Journal of biological chemistry*. 1993 Jun 15.268:12764. [PubMed: 7685350]
27. Sperandio M, et al. P-selectin glycoprotein ligand-1 mediates L-selectin-dependent leukocyte rolling in venules. *The Journal of experimental medicine*. 2003 May 19.197:1355. [PubMed: 12756271]
28. Picker LJ, et al. The neutrophil selectin LECAM-1 presents carbohydrate ligands to the vascular selectins ELAM-1 and GMP-140. *Cell*. 1991 Sep 6.66:921. [PubMed: 1716182]
29. Zollner O, et al. L-selectin from human, but not from mouse neutrophils binds directly to E-selectin. *The Journal of cell biology*. 1997 Feb 10.136:707. [PubMed: 9024699]
30. Levinovitz A, Muhlhoff J, Isenmann S, Vestweber D. Identification of a glycoprotein ligand for E-selectin on mouse myeloid cells. *The Journal of cell biology*. 1993 Apr.121:449. [PubMed: 7682218]
31. Steegmaier M, Borges E, Berger J, Schwarz H, Vestweber D. The E-selectin-ligand ESL-1 is located in the Golgi as well as on microvilli on the cell surface. *Journal of cell science*. 1997 Mar. 110(Pt 6):687. [PubMed: 9099943]
32. Steegmaier M, et al. The E-selectin-ligand ESL-1 is a variant of a receptor for fibroblast growth factor. *Nature*. 1995 Feb 16.373:615. [PubMed: 7531823]
33. Dimitroff CJ, Lee JY, Rafii S, Fuhlbrigge RC, Sackstein R. CD44 is a major E-selectin ligand on human hematopoietic progenitor cells. *The Journal of cell biology*. 2001 Jun 11.153:1277. [PubMed: 11402070]
34. Katayama Y, Hidalgo A, Chang J, Peired A, Frenette PS. CD44 is a physiological E-selectin ligand on neutrophils. *The Journal of experimental medicine*. 2005 Apr 18.201:1183. [PubMed: 15824084]
35. Crutchfield KL, et al. CD11b/CD18-coated microspheres attach to E-selectin under flow. *Journal of leukocyte biology*. 2000 Feb.67:196. [PubMed: 10670580]
36. Kotovuori P, et al. The vascular E-selectin binds to the leukocyte integrins CD11/CD18. *Glycobiology*. 1993 Apr.3:131. [PubMed: 7683936]
37. Santos AL, et al. Biology and pathogenesis of *Fonsecaea pedrosoi*, the major etiologic agent of chromoblastomycosis. *FEMS Microbiol Rev*. 2007 Sep.31:570. [PubMed: 17645522]

38. Zen K, Cui LB, Zhang CY, Liu Y. Critical role of mac-1 sialyl lewis x moieties in regulating neutrophil degranulation and transmigration. *J Mol Biol.* 2007 Nov 16;374:54. [PubMed: 17919658]
39. Fuhlbrigge RC, King SL, Sackstein R, Kupper TS. CD43 is a ligand for E-selectin on CLA+ human T cells. *Blood.* 2006 Feb 15;107:1421. [PubMed: 16269612]
40. Matsumoto M, et al. CD43 functions as a ligand for E-Selectin on activated T cells. *Journal of immunology.* 2005 Dec 15;175:8042.
41. Matsumoto M, Shigeta A, Miyasaka M, Hirata T. CD43 plays both antiadhesive and proadhesive roles in neutrophil rolling in a context-dependent manner. *Journal of immunology.* 2008 Sep 1;181:3628.
42. Carlow DA, Ziltener HJ. CD43 deficiency has no impact in competitive in vivo assays of neutrophil or activated T cell recruitment efficiency. *Journal of immunology.* 2006 Nov 1;177:6450.
43. Yago T, et al. Core 1-derived O-glycans are essential E-selectin ligands on neutrophils. *Proc Natl Acad Sci U S A.* 2010 May 18;107:9204. [PubMed: 20439727]
44. Jones WM, Watts GM, Robinson MK, Vestweber D, Jutila MA. Comparison of E-selectin-binding glycoprotein ligands on human lymphocytes, neutrophils, and bovine gamma delta T cells. *Journal of immunology.* 1997 Oct 1;159:3574.
45. Kishimoto TK, et al. Antibodies against human neutrophil LECAM-1 (LAM-1/Leu-8/DREG-56 antigen) and endothelial cell ELAM-1 inhibit a common CD18-independent adhesion pathway in vitro. *Blood.* 1991 Aug 1;78:805. [PubMed: 1713515]
46. Abbassi O, Kishimoto TK, McIntire LV, Anderson DC, Smith CW. E-selectin supports neutrophil rolling in vitro under conditions of flow. *J Clin Invest.* 1993 Dec;92:2719. [PubMed: 7504692]
47. Lawrence MB, Bainton DF, Springer TA. Neutrophil tethering to and rolling on E-selectin are separable by requirement for L-selectin. *Immunity.* 1994 May;1:137. [PubMed: 7534197]
48. Moore KL, et al. P-selectin glycoprotein ligand-1 mediates rolling of human neutrophils on P-selectin. *The Journal of cell biology.* 1995 Feb;128:661. [PubMed: 7532174]
49. Simon SI, Hu Y, Vestweber D, Smith CW. Neutrophil tethering on E-selectin activates beta 2 integrin binding to ICAM-1 through a mitogen-activated protein kinase signal transduction pathway. *J Immunol.* 2000 Apr 15;164:4348. [PubMed: 10754335]
50. Green CE, Pearson DN, Camphausen RT, Staunton DE, Simon SI. Shear-dependent capping of L-selectin and P-selectin glycoprotein ligand 1 by E-selectin signals activation of high-avidity beta2-integrin on neutrophils. *J Immunol.* 2004 Jun 15;172:7780. [PubMed: 15187162]
51. Finger EB, et al. Adhesion through L-selectin requires a threshold hydrodynamic shear. *Nature.* 1996 Jan 18;379:266. [PubMed: 8538793]
52. Taylor AD, Neelamegham S, Hellums JD, Smith CW, Simon SI. Molecular dynamics of the transition from L-selectin- to beta 2-integrin-dependent neutrophil adhesion under defined hydrodynamic shear. *Biophys J.* 1996 Dec;71:3488. [PubMed: 8968618]
53. McDonough DB, et al. Cooperativity between selectins and beta2-integrins define neutrophil capture and stable adhesion in shear flow. *Ann Biomed Eng.* 2004 Sep;32:1179. [PubMed: 15493506]
54. Sako D, et al. Expression cloning of a functional glycoprotein ligand for P-selectin. *Cell.* 1993 Dec 17;75:1179. [PubMed: 7505206]
55. Goetz DJ, et al. Isolated P-selectin glycoprotein ligand-1 dynamic adhesion to P- and E-selectin. *The Journal of cell biology.* 1997 Apr 21;137:509. [PubMed: 9128259]
56. Bruehl RE, et al. Leukocyte activation induces surface redistribution of P-selectin glycoprotein ligand-1. *Journal of leukocyte biology.* 1997 Apr;61:489. [PubMed: 9103236]
57. Yang J, Galipeau J, Kozak CA, Furie BC, Furie B. Mouse P-selectin glycoprotein ligand-1: molecular cloning, chromosomal localization, and expression of a functional P-selectin receptor. *Blood.* 1996 May 15;87:4176. [PubMed: 8639776]
58. Huang MC, et al. P-selectin glycoprotein ligand-1 and E-selectin ligand-1 are differentially modified by fucosyltransferases Fuc-TIV and Fuc-TVII in mouse neutrophils. *The Journal of biological chemistry.* 2000 Oct 6;275:31353. [PubMed: 10882744]

59. Ponta H, Sherman L, Herrlich PA. CD44: from adhesion molecules to signalling regulators. *Nat Rev Mol Cell Biol.* 2003 Jan.4:33. [PubMed: 12511867]
60. Sackstein R. The bone marrow is akin to skin: HCELL and the biology of hematopoietic stem cell homing. *The Journal of investigative dermatology.* 2004 May.122:1061. [PubMed: 15140204]
61. Jacobs PP, Sackstein R. CD44 and HCELL: Preventing hematogenous metastasis at step 1. *FEBS Lett.* 2011 Aug 6.
62. Hidalgo A, Peired AJ, Wild MK, Vestweber D, Frenette PS. Complete identification of E-selectin ligands on neutrophils reveals distinct functions of PSGL-1, ESL-1, and CD44. *Immunity.* 2007 Apr.26:477. [PubMed: 17442598]
63. von Andrian UH, Hasslen SR, Nelson RD, Erlandsen SL, Butcher EC. A central role for microvillous receptor presentation in leukocyte adhesion under flow. *Cell.* 1995 Sep 22.82:989. [PubMed: 7553859]
64. Abbal C, et al. Lipid raft adhesion receptors and Syk regulate selectin-dependent rolling under flow conditions. *Blood.* 2006 Nov 15.108:3352. [PubMed: 16849645]
65. Zarbock A, Lowell CA, Ley K. Spleen tyrosine kinase Syk is necessary for E-selectin-induced alpha(L)beta(2) integrin-mediated rolling on intercellular adhesion molecule-1. *Immunity.* 2007 Jun.26:773. [PubMed: 17543554]
66. Kuwano Y, Spelten O, Zhang H, Ley K, Zarbock A. Rolling on E- or P-selectin induces the extended but not high-affinity conformation of LFA-1 in neutrophils. *Blood.* 2010 Jul 29.116:617. [PubMed: 20445017]
67. Hidari KI, Weyrich AS, Zimmerman GA, McEver RP. Engagement of P-selectin glycoprotein ligand-1 enhances tyrosine phosphorylation and activates mitogen-activated protein kinases in human neutrophils. *The Journal of biological chemistry.* 1997 Nov 7.272:28750. [PubMed: 9353345]
68. Alonso-Lebrero JL, et al. Polarization and interaction of adhesion molecules P-selectin glycoprotein ligand 1 and intercellular adhesion molecule 3 with moesin and ezrin in myeloid cells. *Blood.* 2000 Apr 1.95:2413. [PubMed: 10733515]
69. Urzainqui A, et al. ITAM-based interaction of ERM proteins with Syk mediates signaling by the leukocyte adhesion receptor PSGL-1. *Immunity.* 2002 Oct.17:401. [PubMed: 12387735]
70. Dore M, Burns AR, Hughes BJ, Entman ML, Smith CW. Chemoattractant-induced changes in surface expression and redistribution of a functional ligand for P-selectin on neutrophils. *Blood.* 1996 Mar 1.87:2029. [PubMed: 8634454]
71. Phong MC, et al. Molecular mechanisms of L-selectin-induced co-localization in rafts and shedding [corrected]. *Biochem Biophys Res Commun.* 2003 Jan 10.300:563. [PubMed: 12504120]
72. Erlandsen SL, Hasslen SR, Nelson RD. Detection and spatial distribution of the beta 2 integrin (Mac-1) and L-selectin (LECAM-1) adherence receptors on human neutrophils by high-resolution field emission SEM. *J Histochem Cytochem.* 1993 Mar.41:327. [PubMed: 7679125]
73. Zollner O, Vestweber D. The E-selectin ligand-1 is selectively activated in Chinese hamster ovary cells by the alpha(1,3)-fucosyltransferases IV and VII. *The Journal of biological chemistry.* 1996 Dec 20.271:33002. [PubMed: 8955145]
74. Fuhlbrigge RC, Kieffer JD, Armerding D, Kupper TS. Cutaneous lymphocyte antigen is a specialized form of PSGL-1 expressed on skin-homing T cells. *Nature.* 1997 Oct 30.389:978. [PubMed: 9353122]
75. Berg EL, Robinson MK, Mansson O, Butcher EC, Magnani JL. A carbohydrate domain common to both sialyl Le(a) and sialyl Le(X) is recognized by the endothelial cell leukocyte adhesion molecule ELAM-1. *The Journal of biological chemistry.* 1991 Aug 15.266:14869. [PubMed: 1714447]
76. Schaff U, Mattila PE, Simon SI, Walcheck B. Neutrophil adhesion to E-selectin under shear promotes the redistribution and co-clustering of ADAM17 and its proteolytic substrate L-selectin. *J Leukoc Biol.* 2008 Jan.83:99. [PubMed: 17928459]
77. Luo J, et al. PI3K is involved in L-selectin- and PSGL-1-mediated neutrophil rolling on E-selectin via F-actin redistribution and assembly. *J Cell Biochem.* 2010 Jul 1.110:910. [PubMed: 20564190]
78. Davies MP, Hallam TJ, Merritt JE. A role for calcium and protein kinase C in agonist-stimulated adhesion of human neutrophils. *Biochem J.* 1990 Apr 1.267:13. [PubMed: 2109601]

79. Ginis I, Tauber AI. Activation mechanisms of adherent human neutrophils. *Blood*. 1990 Sep 15;76:1233. [PubMed: 2400811]
80. Smith RJ, et al. Receptor-coupled signal transduction in human polymorphonuclear neutrophils: effects of a novel inhibitor of phospholipase C-dependent processes on cell responsiveness. *J Pharmacol Exp Ther*. 1990 May;253:688. [PubMed: 2338654]
81. Truneh A, Albert F, Golstein P, Schmitt-Verhulst AM. Calcium ionophore plus phorbol ester can substitute for antigen in the induction of cytolytic T lymphocytes from specifically primed precursors. *Journal of immunology*. 1985 Oct;135:2262.
82. Truneh A, Albert F, Golstein P, Schmitt-Verhulst AM. Early steps of lymphocyte activation bypassed by synergy between calcium ionophores and phorbol ester. *Nature*. 1985 Jan 24–30;313:318. [PubMed: 3918270]
83. Lawson MA, Maxfield FR. Ca(2+)- and calcineurin-dependent recycling of an integrin to the front of migrating neutrophils. *Nature*. 1995 Sep 7;377:75. [PubMed: 7544874]
84. Nüsse O, et al. Store-operated Ca²⁺ influx and stimulation of exocytosis in HL-60 granulocytes. *The Journal of biological chemistry*. 1997 Nov 7;272:28360. [PubMed: 9353293]
85. Nüsse O, Serrander L, Lew DP, Krause KH. Ca²⁺-induced exocytosis in individual human neutrophils: high- and low-affinity granule populations and submaximal responses. *Embo J*. 1998 Aug 10;17:1279. [PubMed: 9482725]
86. Schaff UY, et al. Calcium flux in neutrophils synchronizes beta2 integrin adhesive and signaling events that guide inflammatory recruitment. *Ann Biomed Eng*. 2008 Apr;36:632. [PubMed: 18278555]
87. Simon SI, Green CE. Molecular mechanics and dynamics of leukocyte recruitment during inflammation. *Annu Rev Biomed Eng*. 2005; 7:151. [PubMed: 16004569]
88. Rossaint J, et al. Acute loss of renal function attenuates slow leukocyte rolling and transmigration by interfering with intracellular signaling. *Kidney Int*. 2011 Sep;80:493. [PubMed: 21562471]
89. Cuvelier SL, Paul S, Shariat N, Colarusso P, Patel KD. Eosinophil adhesion under flow conditions activates mechanosensitive signaling pathways in human endothelial cells. *The Journal of experimental medicine*. 2005 Sep 19;202:865. [PubMed: 16172263]
90. Turhan A, Weiss LA, Mohandas N, Collier BS, Frenette PS. Primary role for adherent leukocytes in sickle cell vascular occlusion: a new paradigm. *Proceedings of the National Academy of Sciences of the United States of America*. 2002 Mar 5;99:3047. [PubMed: 11880644]
91. Belcher JD, Marker PH, Weber JP, Heibel RP, Vercellotti GM. Activated monocytes in sickle cell disease: potential role in the activation of vascular endothelium and vaso-occlusion. *Blood*. 2000 Oct 1;96:2451. [PubMed: 11001897]
92. Francis RB Jr, Haywood LJ. Elevated immunoreactive tumor necrosis factor and interleukin-1 in sickle cell disease. *J Natl Med Assoc*. 1992 Jul;84:611. [PubMed: 1629925]
93. Akohoue SA, et al. Energy expenditure, inflammation, and oxidative stress in steady-state adolescents with sickle cell anemia. *Pediatr Res*. 2007 Feb;61:233. [PubMed: 17237728]
94. Duits AJ, Schnog JB, Lard LR, Saleh AW, Rojer RA. Elevated IL-8 levels during sickle cell crisis. *Eur J Haematol*. 1998 Nov;61:302. [PubMed: 9855244]
95. Vichinsky EP, et al. Acute chest syndrome in sickle cell disease: clinical presentation and course. *Cooperative Study of Sickle Cell Disease*. *Blood*. 1997 Mar 1;89:1787. [PubMed: 9057664]
96. Wun T, Cordoba M, Rangaswami A, Cheung AW, Paglieroni T. Activated monocytes and platelet-monocyte aggregates in patients with sickle cell disease. *Clinical and laboratory haematology*. 2002 Apr;24:81. [PubMed: 11985552]
97. Hidalgo A, et al. Heterotypic interactions enabled by polarized neutrophil microdomains mediate thromboinflammatory injury. *Nat Med*. 2009 Apr;15:384. [PubMed: 19305412]
98. Kato GJ, et al. Levels of soluble endothelium-derived adhesion molecules in patients with sickle cell disease are associated with pulmonary hypertension, organ dysfunction, and mortality. *British journal of haematology*. 2005 Sep;130:943. [PubMed: 16156864]
99. Chang J, et al. GMI-1070, a novel pan-selectin antagonist, reverses acute vascular occlusions in sickle cell mice. *Blood*. 2010 Sep 9;116:1779. [PubMed: 20508165]

Figure 1. Dynamic imaging of adhesive and signaling events that support neutrophil recruitment and assembly of the inflammatory synapse

Membrane tethers during rolling on a substrate of E-selectin is observed by phase contrast microscopy as cells flux calcium, detected as a red fluorescence using the intracellular reporter Fluoro-5. E-selectin ligands are clustered in *lipid rafts* as detected by DiI-C18, a fluorescent indocarbocyanine dye that preferentially partitions into lipid ordered domains (Red fluorescence) and antibody to L-selectin (green fluorescence). The *inflammatory synapse* is imaged with antibodies to L-selectin (green fluorescence) and PSGL-1 (red fluorescence) as previously reported⁵⁰.

Figure 2. PMN rolling on E-selectin and arrested in response to infusion of IL-8 at the indicated concentration. Neutrophils were loaded with Fluo-4 and perfused over monolayers expressing E-selectin at a shear stress of 2 dynes/cm², then exposed to a dose range of IL-8 following 2 min of shear interaction. (a) Individual neutrophils that have rolled to arrest rapidly increase their intracellular calcium in response to IL-8 (0.1 nM) resulting in an increase in Fluo-4 emission. (b) On average, neutrophils exhibited a rapid increase in Fluo-4 emission indicative of calcium flux in response to IL-8 concentrations of 0.1 nM or higher, but did not significantly increase calcium in unstimulated or at low IL-8 of 0.01 nM. Plot is representative of 4 independent experiments with measurements from at least 60 neutrophils

at each labeled concentration of IL-8. c) Calcium concentration was measured by ratiometric imaging of neutrophils that sedimented onto the monolayer (Static) or rolling on the monolayer under shear stress (2 dynes/cm²) following exposure to a dose range of IL-8 from 0.001 to 5 nM. The average calcium concentration in all neutrophils in a field of view was measured over time, and the peak value was recorded. Calcium concentration was measured by ratiometric imaging in neutrophils sedimented onto the monolayer (Static) or rolling on the monolayer under shear stress (2 dynes/cm²) following exposure to a dose range of IL-8 from 0.001 to 5 nM. The average calcium concentration in all neutrophils in a field of view was measured over time, and the peak value was recorded. (adapted from Schaff et al., 2008. *ABME*)

Figure 3. Schematic of ESL ligation and signaling of Mac-1 dependent VOC

Following white blood cell tethering by E-selectin a signal is transduced resulting in a conformational upshift in Mac-1 to high affinity that facilitates binding to specific ligands on platelets captured from the blood stream.